

CONSTITUENTS OF *SENECIO HOLLERMAYERII*

S. SEPULVEDA*, S. DELHVI, F. ZILLIKEN

Universität Bonn, Physiologisch-Chemisches Institut, Nußallee 11, 5300 Bonn, Germany.

B.K. CASSELS*

Universidad de Chile, Facultad de Ciencias, Departamento de Química, Casilla 653, Santiago, Chile.

Received September 14, 1992 - Accepted November 2, 1992.

Key words: *Senecio hollermayerii*; acetophenones; flavonoids; coumarins.

Plant. *Senecio hollermayerii* Cabr. (Compositae), aerial parts, collected in January 1979 in Puyehue National Park,¹ Chile. A voucher sample is on deposit in the herbarium of the Natural History Museum, Santiago, Chile.

Uses in traditional medicine. Very toxic plant.

Previously isolated classes of constituents. None.

New-isolated constituents. Piceol (0.0002%), scopoletin (0.004), scopoletin-7-O- β -D-glucoside (scopolin) (0.008), piceol-4- β -O-D-glucoside (picein) (0.005), chrysoeriol-7- β -O-D-glucoside (0.001) and cosmosin (0.0002).

Acknowledgements. Authors are grateful to Prof. E. Breitmaier and Dr G. Eckhardt for NMR and mass spectra.

REFERENCES 1. Muñoz M., "Flora del Parque Nacional de Puyehue", Editorial Consorcio de seguros Empaat, Chile 1980.

CONSTITUENTS OF *CENTAUREA CHILENSIS*

S. SEPULVEDA*, S. DELHVI, B. KOCH, F. ZILLIKEN

Universität Bonn, Physiologisch-Chemisches Institut, Nußallee 11, 5300 Bonn, Germany.

B.K. CASSELS*

Universidad de Chile, Facultad de Ciencias, Departamento de Química, Casilla 653, Santiago, Chile.

Received September 14, 1992 - Accepted November 2, 1992.

Key words: *Centaurea chilensis*; flavonoids.

Plant. *Centaurea chilensis* Hook. et Arn. (Compositae), leaves and stems, collected in Salamanca, 31° 45' S, 71° W, during summer 1982. A voucher sample is on deposit in the herbarium of the Natural History Museum, Santiago, Chile.

Uses in traditional medicine. General infections, treatment of kidney stones, rheumatic disorders.¹

Previously described classes of constituents. Sesquiterpenes,^{2,3} flavonoids⁴ and guaianolides.⁵

New-isolated constituents. 8-Methoxy-apigenin (0.035%), 8-methoxy-luteolin (0.013%), and taxifolin (0.005%).

REFERENCES 1. Muñoz M., Barrera M.E., Meza I., "El uso medicinal y alimenticio de plantas nativas y naturalizadas en Chile", Publicación Ocasional N. 33, Museo Nacional de Historia Natural, Santiago, Chile, 1981. 2. Negrete R.E., Backhouse N., Avendaño S., San Martín A., Pl.

- méd. et phytotér.* 18, 226 (1984). 3. Sepúlveda S., Breitmaier E., *Chemiker-Zeitung* 111, 187 (1987). 4. Negrete R.E., Latorre I., Backhouse N., Peña R., Delporte C., *Pl. méd. et phytotér.* 22, 1 (1988). 5. Negrete R.E., Backhouse N., San Martin A., Cassels B.K., Hartmann R., Breitmaier E., *Chemiker-Zeitung* 112, 144 (1988).

FLAVONOIDIC CONSTITUENTS OF *GARCINIA XANTHOCHYmus* LEAVES

NAZNEEN PARVEEN, M.P. SINGH, NIZAM U. KHAN*

Department of Chemistry, Aligarh Muslim University, Aligarh-202002, India.

M.K. LOGANI

Department of Shared Resources, Temple University, School of Medicine, Philadelphia, Pennsylvania, 19140, U.S.A.

Received September 15, 1992 - Accepted September 30, 1992.

Key words: *Garcinia xanthochymus*; flavonoids.

Plant. *Garcinia xanthochymus* Hook f. (Guttiferae), dried leaves, collected in December, 1988 from Botanical Garden, Calcutta, India, identified by Prof. Wazahat Husain, Department of Botany, Aligarh Muslim University, Aligarh; voucher specimen No. Husain 3098, Aligarh Muslim University, Herbarium, Aligarh.

Previously isolated classes of constituents. Xanthones,¹⁻⁶ biflavonoids,^{3,5} benzophenones,^{3,4,6} terpenoids.⁷

New-isolated constituents. Agathisflavone (from 3 g, 35 mg), 7-O-methylameno-flavone (50 mg) and vitexin (40 mg).

Acknowledgements. Authors are thankful to Prof. M.A. Beg, Department of Chemistry, for facilities and to Prof. W. Husain, Department of Botany, for identification of plant material, respectively. One of us (N. Parveen) is thankful to CSIR, New Delhi for financial assistance.

- REFERENCES**
1. Karanjgoakar A.V., Rama R.A.V., Venkataraman K., Yemul S.S., Palmer K.J., *Tetrahedron Lett.*, 4977 (1973).
 2. Blount J.F., Wilham T.H., *Tetrahedron Lett.*, 2921 (1976).
 3. Baslas R.K., Kumar P., *Curr. Sci.* 48, 814 (1979).
 4. Rama R.A.V., Venkatswamy G., Yemul S.S., *Ind. J. Chem.* 19B, 627 (1980).
 5. Baslas R.K., Kumar P., *Acta Cienc. Indica Chem.* 7, 31 (1981).
 6. Rama R.A.V., Venkatswamy G., Yemul S.S., *Chem. Ind.* 3, 92 (1979).
 7. Singh M.P., Parveen N., Khan N.U., Achari B., Dutta P.K., *Fitoterapia* 62, 286 (1991).

ALKALOIDS OF *XYLOPIA AETHIOPICA*

KARAMOKO MAMOUDOU KANDÉ, STEFAN PHILIPOV*, HEDVINA DUTSCHEWSKA
Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, 1113 Sofia, Bulgaria.

Received September 18, 1992 - Accepted November 16, 1992.

Key words: *Xylopia aethiopica*; aporphine alkaloids.

Plant. *Xylopia aethiopica* A. Rich. (Annonaceae), air-dried leaves (1 kg), collected in April 1988 from the village Snagbandou (East Guinea), identified by Prof.