

Heteroarylnitrones as Drugs for Neurodegenerative Diseases: Synthesis, Neuroprotective Properties, and Free Radical Scavenger Properties

Williams Porcal,[‡] Paola Hernández,[‡] Mercedes González,[‡] Ana Ferreira,[§] Claudio Olea-Azar,[#] Hugo Cerecetto,^{*,‡} and Ana Castro^{*,II,†}

Departamento de Química Orgánica, Facultad de Ciencias-Facultad de Química, and Catedra de Inmunología, Universidad de la República, Montevideo, Uruguay, Departamento de Química Inorgánica y Analítica, Facultad de Ciencias Químicas y Farmacéuticas, Universidad de Chile, Santiago, Chile, and Instituto de Química Médica (CSIC), Madrid, Spain

New 1,2,4-thiadiazolynitrones and furoxanylnitrones were developed and evaluated as neuroprotective agents on a human neuroblastoma (SH-SY5Y) cells model. They inhibited at low micromolar concentrations the oxidative damage and the death induced by exposure to hydrogen peroxide. These heteroarylnitrones showed excellent peroxyl free radical absorbance capacities, analyzed by oxygen radical absorbance capacity (ORAC) assay with fluorescein as the fluorescent probe, ranging from 1.5- to 16.5-fold the value of the reference nitrone, α -phenyl-*N*-*tert*-butylnitron (PBN). The electron spin resonance spectroscopy (ESR) demonstrated the ability of these derivatives to directly trap and stabilize oxygen, carbon, and sulfur-centered free radicals. These results demonstrated the potential use of these heteroarylnitrones as neuroprotective agents in preventing the death of cells exposed to enhanced oxidative stress and damage.

Introduction

Reactive oxygen species (ROS) and reactive nitrogen species (RNS) produced by oxidative metabolism are capable of damaging cellular components through molecular modifications to a polyunsaturated membrane's lipids, proteins, and nucleic acids.^{1,2} Much evidence suggests that biological oxidation in the human body generates highly pathogenic ROS and RNS such as hydroxyl free radical ($\cdot\text{OH}$), superoxide anion ($\text{O}_2^{\cdot-}$), peroxynitrite (ONOO^-), and lipid peroxide free radicals (ROO^\cdot), causing cellular injury.³⁻⁵ These pathological events have important roles in many degenerative disorders, for example, atherosclerosis, rheumatoid arthritis, and several neurodegenerative diseases such as ischemic conditions, stroke, Parkinson's disease, and Alzheimer's disease.⁶⁻⁸ The central nervous system (CNS) is especially sensitive to oxidative damage for reasons such as (a) high use of oxygen during the metabolic process and therefore greater production of ROS, (b) fewer antioxidants defenses than other organs, for example, liver and heart, (c) enrichment in the more easily peroxidable fatty acids, (d) areas with high levels of the Fe^{3+} /ascorbate pro-oxidant system that reduce H_2O_2 to form $\cdot\text{OH}$, a potent oxidant.⁹ Recently, it has been reported that β -amyloid solutions liberate hydrogen peroxide and, subsequently, $\cdot\text{OH}$ converted via Fenton's reaction.¹⁰ Consequently, the search of effective treatments that prevent oxidative stress associated with premature aging and




Figure 1. Antioxidant and neuroprotective mechanisms of action postulated for PBN.

neurodegenerative diseases is an important area of neurochemical research.

Recently, it has been proved that nitron-free radical trap, i.e., α -phenyl-*N*-*tert*-butylnitron (PBN,^a Figure 1), could be used in the treatment of neurodegenerative diseases as well as in the prolongation of life span.¹¹ Initially, the neuroprotective activity was attributed to the nitron group because of its ability to act as free radical scavenger (FRS, Figure 1). Subsequently, observations that PBN has a good neuroprotective activity at lower doses than those necessary to act as an effective FRS demonstrated the ability of PBN to inhibit signal transduction processes such as suppression of proinflammatory cytokines, genes, and mediators associated with enhanced neuroinflammatory processes (Figure 1).¹² The neuroprotective activity of the nitron pharmacophore depends in great part on the connectivity and the nature of substituents on the nitron group. In this sense, the chemical and pharmacological aspects of different heteroarylnitrones have been reviewed,¹³ describing

* To whom correspondence should be addressed. Phone: +34 918061130. Fax: +34 918034660. E-mail: acastro@neuropharma.es.

[‡] Departamento de Química Orgánica, Universidad de la República.

[§] Catedra de Inmunología, Universidad de la República.

[#] Universidad de Chile.

^{II} Instituto de Química Médica (CSIC).

[†] Present address: NeuroPharma, S. A.; Avda. de la Industria 52, 28760 Tres Cantos, Madrid, Spain.

^a Abbreviations: BBB, blood-brain barrier; DMPO, 5,5-dimethyl-1-pyrroline *N*-oxide; FL, fluorescein; FRS, free radical scavenger; HFS, hyperfine splitting; LDH, lactate dehydrogenase; NAC, *N*-acetyl cysteine; ORAC, oxygen radical absorbance capacity; PAMPA, parallel artificial membrane permeation assay; PBL, porcine brain lipid; PBN, α -phenyl-*N*-*tert*-butylnitron; RP, reduction percentage.





Figure 2. Heteroaryl nitrones with neuroprotective and spin trapping properties.

Scheme 1. Heteroaryl nitrones Designed as Potential Drugs for Neurodegenerative Diseases



some patented nitrone containing furans (I–III, Figure 2) with good activity against the neuronal cell damage induced by β -amyloid.¹⁴ On the other hand, imidazolyl nitrones (IV and V, Figure 2) has been developed and biologically evaluated, showing *in vivo* neuroprotective properties.¹⁵ Additionally several nitrones has been widely used as spin trap for the specific detection of transient radicals (e.g., $\cdot\text{OH}$, $\text{O}_2^{\cdot-}$, $\text{ROO}\cdot$) or relatively stable radicals (e.g., NO) that are undetectable under normal conditions,¹⁶ in the electron spin resonance (ESR) spectroscopy. With this aim two nitrones have been commonly used in the spin trapping technique: the linear nitrone PBN (Figure 1) and the cyclic nitrone DMPO (5,5-dimethyl-1-pyrroline *N*-oxide VI, Figure 2).


On the basis of that, herein we describe the design and synthesis of new heteroaryl nitrones combining, in their structures, fragments able to show neuroprotection properties, such as nitrone moiety, antioxidant fragments, and heterocyclic groups able to stabilize the generated free radical (Scheme 1). In order to assess the protective effects of heteroaryl nitrones, the effects of H₂O₂-injured in neuronal cells (SH-SY5Y) and *in vitro* unspecific cytotoxicity against J774-mouse macrophages were evaluated. Additionally, the antioxidant activities and the spin trapping properties have been investigated by evaluating the direct reaction with oxygen-, carbon-, and sulfur-centered free radicals by antioxidant assays and ESR spectroscopy.

Results and Discussion

Chemistry. Four different families of heteroaryl nitrones were selected to study the influence on the biological activity: 1,2,3-

thiadiazole, 1,2,4-thiadiazole, 1,2,5-oxadiazole *N*-oxide, and benzo[1,2-*c*]1,2,5-oxadiazole *N*-oxide derivatives.^{17,18} The nitrones syntheses were planned through the condensation between the corresponding heteroaromatic aldehyde and *N*-substituted hydroxylamine,¹⁹ our first goal being the preparation of the corresponding aldehyde derivatives. For the synthesis of 3- and 5-formyl-1,2,4-thiadiazole aldehydes, **6** and **9**, respectively, the synthetic routes shown in Scheme 2a were employed. First, the reaction of benzothioamide with *N,N*-dimethylalkanamide dimethyl acetals produced the corresponding *N'*-(thioaroyl)-*N,N*-dimethylamidines **1** and **2** in excellent yield.²⁰ These intermediates were converted by amination–cyclization into the 1,2,4-thiadiazole **3** and **4**. The 3-formyl-1,2,4-thiadiazole **6** was obtained through bromination–oxidation of thiadiazole **4**. Attempts to obtain aldehyde **6** using formylation reaction between 5-phenyl-1,2,4-thiadiazole **3** and *n*-BuLi in DMF were unsuccessful. In these conditions, the ring-opening product, **5**, was mainly generated. Oxidation of the methyl substituent in compound **4** using selenium dioxide was also explored in the preparation of aldehyde **6**; however, no reaction was observed. Second, nitrile sulfide, obtained *in situ* through thermolysis of the 1,3,4-oxathiazol-2-one **7** (Scheme 2a), was treated with ethyl cyanofornate, yielding ester **8** via a 1,3-dipolar cycloaddition process.²¹ Subsequently, the desired aldehyde **9** was prepared by reduction of **8** with DIBAL-H and subsequent oxidation with MnO₂. In order to know the 1,2,4-thiadiazole-substituents stereoelectronic/antioxidant effects (Scheme 1), different 3-aryl substitutions were studied (Scheme 2b). Consequently, aldehydes **18**, **20**, and **21** were prepared following the same synthetic

Scheme 2. Synthesis of Formyl-Heterocycle Derivatives^a




^a Reagents and conditions: (a) room temp; (b) HSA, Py, EtOH, room temp; (c) *n*-BuLi, THF, -78 °C, then DMF, 0 °C; (d) SeO₂, dioxane or ClCH₂CH₂Cl, reflux; (e) NBS, PDBO, CCl₄, reflux, then DMSO, 105 °C; (f) Cl(C=O)SCL, toluene, reflux; (g) decaline, 140–160 °C; (h) EtO₂C-CN; (i) (1) DIBAL-H, THF, 0 °C; (2) MnO₂, CHCl₃, reflux; (j) (1) H₂NNHCO₂Met, *p*-TsOH, toluene, reflux; (2) SOCl₂, 60 °C; (k) NaBH₄, EtOH, room temp; (l) MnO₂, CHCl₃, reflux; (m) NBS, PDBO, CCl₄, reflux, then DMSO, 120 °C; (n) NaNO₂, AcOH, 0 °C; (o) MnO₂, CHCl₃, room temp; (p) (1) NaN₃, DMSO, 60–70 °C; (2) DMSO, 90–100 °C.

route for aldehyde **9** preparation. Attempts to obtain the desired 4-NO₂ analogue **19** were unsuccessful because of the low stability of ester **15** in the reduction conditions (i) (Scheme 2b). The 1,2,3-thiadiazole carbonylic reagents were prepared from ethyl acetoacetate following a condensation with methyl carbazate, forming the corresponding acylhydrazone, and subsequent cyclization via Hurd–Mori reaction in the presence of thionyl chloride (Scheme 2c).²² Then thiadiazole **22** was transformed into the key aldehydes **24** and **25** by reduction/oxidation for the first case and by benzylic-like bromination/oxidation for the second. Finally, the synthetic procedures for furoxan and benzofuroxan systems included, for the first heterocycle, addition to alkenes followed by intramolecular cyclization and thermointramolecular cyclization for the second one (Scheme 2d). Consequently, the reaction of cinnamyl

alcohol or crotonaldehyde with NaNO₂ in AcOH produced the furoxan derivatives **26** and **28**, respectively.²³ Alcohol **26** was converted into the aldehyde **27** by oxidation with MnO₂.²³ The formylbenzofuroxan **29** was obtained from 4-chloro-3-nitrobenzaldehyde in a one-pot process via nucleophilic substitution with the azide anion and subsequent cyclization by pyrolysis of the corresponding *o*-nitrophenylazide intermediate.²⁴ The desired nitrones, **30–41**, were obtained in good yield for reaction between the corresponding aldehydes **6**, **9**, **18**, **20**, **21**, **24**, **25**, **27–29** and *N*-substituted hydroxylamine hydrochloride in the presence of NaHCO₃ (Scheme 3). Taking into account that the nitron stability, as well as the stability of the corresponding spin-adduct (nitroxyl free radical), is highly influenced by the *N*-alkyl substituent,¹³ we developed derivatives **31–33** by

Scheme 3. Synthesis and Structure of New Heteroarylnitrones^a



^a Asterisk (*) indicates at room temperature.

varying R¹ (Scheme 3) in order to evaluate the effect of this modulation in the bioactivity studies.

All the structures of the desired nitrones, **30–41**, were established by NMR (¹H, ¹³C, HMQC, and HMBC experiments), IR, and MS. The purity was established by TLC and microanalysis. The stereochemistry around the olefinic carbon–nitrogen bond of the nitron function was established using NOE-diff experiments, and they are *Z*-isomers in all the cases.

Biology. Neuroprotective Activity. The neuroprotective activity of the new developed heteroarylnitrones, **30–41**, was studied using a human neuronal-like cellular system (SH-SY5Y cells) exposed to H₂O₂ as oxidative damaging agent.²⁵ First, SH-SY5Y cytotoxicity of the developed heteroarylnitrones and PBN was evaluated in vitro in the range 0.05–10 μM for 24 h. The cell viability was determined by measuring the activity of lactate dehydrogenase (LDH) released into the culture medium and expressed as the difference between nitrones-treated and untreated cells. All the nitrones displayed no cytotoxicity for neuronal cells, 100% of survival, at assayed neuroprotective doses (10 μM). Second, each one of the heteroarylnitrones, at 10 μM doses, was administered to the cells 1 h before the incubation with H₂O₂. The cell viability was determined by

measuring the activity of LDH released and expressed as the difference between nitrones-treated and untreated cells. The basal percentage of viable neurons after treatment with 100 μM H₂O₂ was 45%, and *N*-acetylcysteine (NAC) was used as positive test at 5 mM (Figure 3). The values summarized in Table 1 show that the 1,2,4-thiadiazolylnitrones **30–36** and the furoxanylnitron **40** protect similarly to or slightly higher against the cellular damage than the parent nitron PBN, with values between 33% and 46%. Interestingly, the treatment of SH-SY5Y cells with the 1,2,4-thiadiazolylnitron **36** at 1 μM significantly protects against H₂O₂ induced neurotoxicity, with a value of 20% (Figure 3). Furthermore, the benzofuroxanylnitron **41** showed lower capacity to protect neuroblastoma human cells from death induced by oxidative stress than PBN, while the 1,2,3-thiadiazolylnitrones (**37** and **38**) and furoxanylnitron **39** did not show neuroprotective effects. These results revealed the consequence of the connectivity and the nature of the nitron-pharmacophore substituents in the neuroprotective activity. The 1,2,4-thiadiazolylnitrones **31**, **32**, and **33**, substituted in the nitrogen atom of the nitron by a *tert*-butyl, cyclohexyl, and benzyl group, respectively, exhibited differential activity, indicating some *N*-alkyl moieties dependence.




Figure 3. Cellular viability in different experimental conditions: H₂O₂ (100 μM), nitrone **36** (at 10 and 1 μM), and NAC (5 mM) used as positive test. Viability was determined through LDH activity after 24 h of incubation (mean ± SD values from three independent experiments).

Table 1. Effect of HeteroarylNitrones on Cell viability in SH-SY5Y Cells Preincubated with H₂O₂ (100 μM) and HeteroarylNitrones Cytotoxicities (J774 Mouse Macrophages)

ref	difference in cell viability ^{a-c}	% neuroprotection ^{c,d}	J774 IC ₅₀ (μM) ^e
PBN	20.6 ± 0.2	37	>400
30	20.0 ± 1.9	36	>400
31	20.7 ± 9.4	38	>400
32	23.6 ± 10.7	43	100
33	18.1 ± 3.0	33	240
34	19.1 ± 2.1	34	200
35	25.0 ± 5.2	46	>400
36	25.2 ± 5.6	46	>400
37	0		>400
38	0		>400
39	0		100
40	23.0 ± 2.4	42	>400
41	13.4 ± 6.0	24	150

^a The cell viability was measured with LDH activity being 100% the basal value. The basal value was obtained from the cells that did not receive any treatment. Difference of cell viability was determined by the difference between LDH activity in nitrones–H₂O₂-treated cells and LDH activity in H₂O₂-treated cells and normalized to 100%. ^b Data are expressed as the mean ± SD ($p < 0.001$). Experiments were done in triplicate. ^c Doses of heteroaryl nitrones are 10 μM. ^d Basal percentage of viable neurons after treatment with 100 μM H₂O₂ is 45%. The percentages of neuroprotection are determined from the basal values. ^e IC₅₀: concentration that produces 50% inhibitory effect. The results are the mean values of two different experiments with SD values less than 10% in all cases.

Mammal Cytotoxicity. Unspecific mammalian cytotoxicity of the developed heteroaryl nitrones was evaluated in vitro in the range 50–400 μM, using J774 mouse macrophages as the cellular model (Table 1).²⁶ In the study PBN was included as a neuroprotective reference compound. The developed heteroaryl nitrones displayed no cytotoxicity for mammalian cells at its neuroprotective doses (10 μM). Furthermore, the most active heteroaryl nitrones, **35**, **36** and **40**, show low mammalian cytotoxicity in the assayed in vitro model, these compounds being as cytotoxic as the parent compound PBN. Moreover, the IC₅₀ values summarized in Table 1 show that the 1,2,4-thiadiazolyl, **32**–**34**, furoxanyl, **39**, and benzofuroxanyl nitrones, **41**, are a little more toxic against this cellular system than the parent compound PBN.

Blood–Brain Barrier Permeation Studies. For the 1,2,4-thiadiazolyl nitrones **35** and **36**, with the best neuroprotective




Figure 4. Prediction of the brain penetration using PAMPA-BBB assay by 1,2,4-thiadiazolyl nitrones **35** and **36** (CNS +: compounds able to cross the BBB).

Table 2. Free-Radical Scavenger/Antioxidant Activity of HeteroarylNitrones Derivatives Using ORAC_{FL} Method and DPPH Captured

ref	ORAC (PBN equiv) ^a	RP of DPPH (%) ^b
PBN	1.0	0 (20°)
30	2.0 ± 0.1	41 ± 3
31	5.5 ± 0.1	0
32	4.6 ± 0.2	11 ± 3
33	3.5 ± 0.3	0
34	1.5 ± 0.2	0
35	7.0 ± 0.4	10 ± 4
36	6.0 ± 0.2	40 ± 3
37	2.5 ± 0.1	0
38	3.0 ± 0.1	0
39	6.0 ± 0.2	0
40	2.5 ± 0.1	8 ± 1
41	16.5 ± 0.4	17 ± 5

^a Data are expressed as μmol of PBN equiv/μmol of tested heteroaryl nitrones and are the mean ($n = 3$) ± SD. ^b All heteroaryl nitrones were assayed at 0.1 mM. ^c Evaluated at 10 mM.

profile and the lowest mammalian cytotoxicity (Table 1), the capacity to cross the blood–brain barrier (BBB) by means of a parallel artificial membrane permeation assay for BBB (PAMPA-BBB) using a lipid extract of porcine brain was determined. The assay validation was made by comparing experimental permeabilities of 20 commercial drugs to reported values.²⁷ From this experiment and taking into account the limit established by Di et al. for BBB permeation, compounds with permeability (P_e) above $4.4 \times 10^{-6} \text{ cm s}^{-1}$ could cross the BBB (Figure 4).²⁸

Free-Radical Scavenger Properties Antioxidant Activity. The antioxidant activity of the newly developed heteroaryl nitrones was determined using two in vitro assays, the oxygen radical absorbance capacity (ORAC) and the measurement of reduction percentage (RP) of the stable diphenylpicrylhydrazyl free radical (DPPH). ORAC assay²⁹ measures peroxy free radical scavenger compounds capability and using fluorescein (FL) as the fluorescent probe. The peroxidation of FL yields nonfluorescent products at 520 nm, so the remaining fluorescence of FL in the presence of the free radical scavenger compound is a measure of the antioxidant activity of the test compound. In this assay the results were expressed in reference to PBN, as PBN equiv (μmoles of PBN equivalents per μmoles of tested compound) (Table 2). On the other hand, the RP of DPPH is based on the reaction between the deep-violet stable free radical (characterized by an absorption band in ethanol solution centered at 520 nm) and the studied free radical scavenger measuring the absorbance of nitrone-treated and untreated DPPH solutions (Table 2).³⁰ The peroxy free radical scavenger property of our nitrones, according to ORAC assay, was better than that of PBN, being 1.5–16.5 higher than PBN value. The benzofuroxanyl nitrone **41** showed the highest ORAC_{FL} value (16.5 PBN equiv), whereas the 1,2,4-thiadiazolyl nitrone **34** showed the lowest ORAC_{FL} value (1.5 PBN equiv). Similar results of antioxidant




Figure 5. ESR spectra obtained in the trapping of $\cdot\text{OH}$, $\text{CH}_3\cdot\text{CHOH}$, and $\text{SO}_3^{\cdot-}$ free radicals with nitrones **31**, **38**, and **40**. Superscript “a” in the heading entries indicates that $\cdot\text{OH}$, $\text{CH}_3\cdot\text{CHOH}$, and $\text{SO}_3^{\cdot-}$ free radicals are generated by Fenton’s reaction.

activity were obtained through DPPH assay. The heteroaryl nitrones were tested at 0.1 mM, being more potent antioxidants than PBN, which at 10 mM possess 20% of DPPH reduction. In this sense, the 1,2,4-thiadiazolyl nitrones **30** and **36** showed the highest antioxidant values, 41 and 40 DPPH RP, respectively.

Electron Spin Resonance Measurements. Initially the $\cdot\text{OH}$ -spin trapping ability of the developed heteroaryl nitrones was investigated using ESR spectroscopy. The $\cdot\text{OH}$ free radical was generated in situ from a Fenton reaction system. All the assayed heteroaryl nitrones, except the thiadiazolyl nitrones **32** and **33** and furoxanyl nitron **39**, gave strong ESR signals (Figure 5), and the hyperfine splitting (HFS) constants for the different spin adducts are reported in Table 3. These data indicate that in general our heteroaryl nitrones generate stable spin adducts with $\cdot\text{OH}$, like previously reported imidazolyl nitrones.³¹ The absence of signal in the cases of nitrones **32** and **33** indicated that the cyclohexyl and benzyl groups affect the stability of the spin adduct generated under the studied conditions.

In order to study the capability of the developed nitrones as spin trapping against other free radical species, we selected one nitron from each proposed family. Consequently, the 1,2,4-thiadiazolyl nitron **31**, the 1,2,3-thiadiazolyl nitron **38**, and the furoxanyl nitron **40** were studied as both ethanol and sulfite free radical spin trappings. The ESR signals obtained for these heteroaryl nitrones are given in Figure 5. The ESR signals for the free radical adducts with nitrones **31**, **38**, and **40** indicated

Table 3. HFS Constants for the Heteroaryl Nitron- $\cdot\text{OH}$ Spin Adduct

nitron- $\cdot\text{OH}$ ^{a,b} spin adduct, ref	a_N (G) ^c	$a_{H\beta}$ (G) ^c
30	14.8	2.4
31	15.7	0.9
32	ns ^d	ns ^d
33	ns ^d	ns ^d
34	15.3	0.6
35	15.7	0.7
36	14.0	2.6
37	14.5	2.6
38	15.5	4.0
39	ns ^d	ns ^d
40	15.8	2.1
41	14.1	2.2

^a $\cdot\text{OH}$ was generated from a standard Fenton system: $\text{Fe}(\text{NH}_4)_2(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$ (0.5 mM) was added to a solution of nitron (50 mM) and H_2O_2 (0.5 mM) in phosphate buffer (0.1 M). ^b The ESR spectrum of the spin adduct was recorded 30 s after addition of FeSO_4 . ^c G: gauss. ^d ns: no signals in the assayed conditions.

that these nitrones are also very efficient to trap and stabilize both carbon- and sulfur-centered free radicals, $\text{CH}_3\cdot\text{CHOH}$ and $\text{SO}_3^{\cdot-}$, respectively. These results suggest that our nitrones have great capacity to scavenge free radicals and potential use to prevent or reduce the damage caused by radicals in several degenerative disorders. However, because nitron **38** is very efficient in spin trapping (i.e., $\cdot\text{OH}$, $\text{CH}_3\cdot\text{CHOH}$, $\text{SO}_3^{\cdot-}$, Figure 5), it does not have neuroprotective effects in the neuroblastoma

human SH-SY5Y system (Table 1), showing that the neuroprotective activity might not exclusively reside in its capacity to scavenge free radicals. These results are similar to those recently reported for other heteroaryl nitrones.³²

Conclusion

New series of thiadiazolyl and furoxanyl nitrones have been synthesized and biologically evaluated as potential neuroprotective agents for the treatment of neurodegenerative disorders. The designed heteroaryl nitrones were efficiently obtained through the condensation between heteroaromatic aldehydes and N-monosubstituted hydroxylamines with good to excellent yields and chromatographically isolated as pure *Z*-isomers. They show excellent free radical scavenger capacity and good neuroprotective effects without cellular toxicity. The ESR spectroscopy demonstrated the ability of these heteroaryl nitrones to directly scavenge different free radicals (i.e., $\cdot\text{OH}$, $\text{CH}_3\cdot\text{CHOH}$, $\text{SO}_3\cdot^-$). Taking into account all the results, the designed heteroaryl nitrones have demonstrated a therapeutic potential as neuroprotective agents in preventing the death of cells exposed to enhanced oxidative stress and damage. Further structural optimization, QSAR, and in vivo neuroprotective property studies are currently underway.

Experimental Section

Compounds **3**, **4**,²⁰ **8**,²¹ **14–17**,^{33–36} and **26–29**²³ were prepared according to literature procedures. Melting points were determined with an electrothermal melting point apparatus (Electrothermal 9100) and are uncorrected. Proton and carbon NMR spectra were recorded on a Bruker DPX-400 spectrometer. The chemical shifts values are expressed in ppm relative to tetramethylsilane as internal standard. Mass spectra were determined in a MSD 5973 Hewlett-Packard or LC/MSD series 100 Hewlett-Packard spectrometer using electronic impact (EI) or electrospray ionization (ESI), respectively. Microanalyses were performed in a Fisons EA 1108 CHNS-O equipment and were within $\pm 0.4\%$ of the calculated compositions. Column chromatography was carried out using Merck silica gel (60–230 mesh). Most chemicals and solvents were analytical grade and used without further purification. All the reactions were carried out in a nitrogen atmosphere. The typical workup included washing with brine and drying the organic layer with sodium sulfate before concentration.

3-Formyl-5-phenyl-1,2,4-thiadiazole (6). A mixture of **4**²⁰ (0.4 g, 2.3 mmol), *N*-bromosuccinimide (0.45 g, 2.5 mmol), and dibenzoyl peroxide (0.08 g, 0.34 mmol) in dry CCl_4 (8.0 mL) was heated at reflux for 16 h. Then the mixture was allowed to cool to room temperature, the succinimide was collected and washed with CCl_4 (10.0 mL), and the organic layer was evaporated in vacuo. 3-Bromomethyl-5-phenyl-1,2,4-thiadiazole was obtained as an oil ($\sim 40\%$ by ^1H NMR signals) and was used in the next reaction without further purification. ^1H NMR (CDCl_3 , 400 MHz) δ : 4.69 (s, 2H), 7.45–7.55 (m, 3H), 7.90–8.00 (m, 2H). The crude of the bromination process in DMSO (6.0 mL) was stirred and warmed at 105 °C during 1.5 h. Subsequently, the mixture was poured into saturated sodium chloride solution (30.0 mL) and extracted with EtOAc (3 \times 20.0 mL). After the workup of the combined organic layers, the residue was purified by column chromatography (SiO_2 , petroleum ether/EtOAc (9:1)), yielding derivative **6** as a yellow oil (130 mg, 30% two steps). ^1H NMR (CDCl_3) δ : 7.45–7.65 (m, 3H), 7.90–8.10 (m, 2H), 10.30 (s, 1H). EI-MS, m/z (abundance, %): 190 (M^{++} , 27), 135 (100), 103 (15).

General Procedure for the Synthesis of 3-Aryl-5-formyl-1,2,4-thiadiazole Derivatives (9 and 18–21). A solution of the corresponding ester^{20,33–36} (1 equiv) in THF (4 mL/mmol) as solvent was cooled at 0 °C. Then a solution of DIBAL-H (1 M in hexane, 2–3 equiv) was added slowly and the final mixture was stirred during 3 h. The reaction mixture is diluted with EtOAc and washed with HCl (1 M). After the workup, the residue, corresponding to

the 3-aryl-5-hydroxymethyl-1,2,4-thiadiazole (pure by TLC), was used in the next reaction without further purification. Subsequently, a mixture of the corresponding alcohol (1 equiv) and MnO_2 (10 equiv) in CHCl_3 (12 mL/mmol) as solvent was heated at reflux until absence of the alcohol (checked by TLC). The resulting dispersion was then filtered through a short pad of Celite, the organic phase concentrated in vacuo and the residue purified by column chromatography (SiO_2 , petroleum ether/EtOAc (8:2)) to afford spectroscopically pure compounds.

5-Hydroxymethyl-3-phenyl-1,2,4-thiadiazole. ^1H NMR (CDCl_3) δ : 2.83 (t, 1H), 5.17 (d, 2H, $J = 5.5$ Hz), 7.45–7.51 (m, 3H), 8.24–8.30 (m, 2H). EI-MS, m/z (abundance, %): 192 (M^{++} , 38), 135 (100), 103 (24), 77 (20).

5-Formyl-3-phenyl-1,2,4-thiadiazole (9). Yellow solid, 0.13 g (25%, two steps). ^1H NMR (CDCl_3) δ : 7.52 (m, 3H), 8.33 (m, 2H), 10.19 (s, 1H). ^{13}C NMR (CDCl_3) δ : 184.3 (C5), 182.5 (CHO), 175.1 (C3), 131.9 (Cq), 131.0 (Co), 128.9 (Cm), 128.3 (Cp). EI-MS, m/z (abundance, %): 190 (M^{++} , 84), 135 (100), 103 (30), 77 (24).

3-(4-Chlorophenyl)-5-hydroxymethyl-1,2,4-thiadiazole. ^1H NMR (CDCl_3) δ : 2.70 (s, 1H), 5.05 (s, 2H), 7.38 (d, 2H, $J = 8.5$ Hz), 8.15 (d, 2H, $J = 8.5$ Hz). EI-MS, m/z (abundance, %): 226 (M^{++} , 51), 169 (100), 137 (35), 102 (15).

3-(4-Chlorophenyl)-5-formyl-1,2,4-thiadiazole (18). Yellow solid, 0.28 g (44%, two steps). ^1H NMR ($\text{DMSO}-d_6$) δ : 7.67 (d, 2H, $J = 8.5$ Hz), 8.28 (d, 2H, $J = 8.7$ Hz), 10.16 (s, 1H). ^{13}C NMR ($\text{DMSO}-d_6$) δ : 180.2 (CHO), 179.1 (C5), 173.52 (C3), 137.6 (Cq), 132.7 (Cq), 130.6 (Co), 130.0 (Cm). EI-MS, m/z (abundance, %): 224 (M^{++} , 75), 169 (100), 137 (39), 102 (18).

5-Hydroxymethyl-3-(4-methoxyphenyl)-1,2,4-thiadiazole. ^1H NMR (CDCl_3) δ : 2.75 (s, 1H), 3.87 (s, 3H), 5.15 (s, 2H), 7.01 (d, 2H, $J = 8.6$ Hz), 8.42 (d, 2H, $J = 8.6$ Hz). EI-MS, m/z (abundance, %): 222 (M^{++} , 49), 167 (100), 135 (57).

5-Formyl-3-(4-methoxyphenyl)-1,2,4-thiadiazole (20). Yellow solid, 0.13 g (31%, two steps). ^1H NMR (CDCl_3) δ : 3.85 (s, 3H), 7.04 (d, 2H, $J = 8.6$ Hz), 8.31 (d, 2H, $J = 8.7$ Hz), 10.19 (s, 1H). ^{13}C NMR (CDCl_3) δ : 181.2 (CHO), 178.3 (C5), 172.3 (C3), 160.9 (Cp), 130.1 (Co), 114.5 (Cm), 54.9 (OCH_3). EI-MS, m/z (abundance, %): 220 (M^{++} , 99), 165 (100), 150 (45), 133 (75).

3-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-5-hydroxymethyl-1,2,4-thiadiazole. ^1H NMR (CDCl_3) δ : 1.52 (s, 18H), 2.93 (s, 1H), 5.16 (s, 2H), 5.53 (s, 1H), 8.14 (s, 2H). ESI-MS, m/z : 321 ($\text{M}^{++} + \text{H}$).

3-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-5-formyl-1,2,4-thiadiazole (21). Brown solid, 0.14 g (27%, two steps). ^1H NMR (CDCl_3) δ : 1.54 (s, 18H), 5.61 (s, 1H), 8.21 (s, 2H), 10.23 (s, 1H). ^{13}C NMR (CDCl_3) δ : 183.9 (C5), 182.8 (CHO), 175.9 (C3), 156.6 (Cp), 136.4 (Cm), 125.7 (Co), 125.3 (Cq), 34.4 ($\text{C}(\text{CH}_3)_3$), 30.2 ($\text{C}(\text{CH}_3)_3$). ESI-MS, m/z : 319 ($\text{M}^{++} + \text{H}$).

5-Ethoxycarbonyl-4-methyl-1,2,3-thiadiazole (22). A mixture of ethyl acetoacetate (4.9 mL, 38.5 mmol), methyl carbazate (3.8 g, 42.4 mmol), and *p*-TsOH (catalytic amounts) in toluene (120.0 mL) was heated at reflux for 2 h. Then the solvent was concentrated in vacuo and the crude of the reaction was used in the next reaction without further purification. The acylhydrazone obtained was cooled at 0 °C, and SOCl_2 (20.0 mL, 270 mmol) was added dropwise with stirred during 10 min. The final mixture was stirred during 2 h at 60 °C, the excess of SOCl_2 was evaporated in vacuo, and the residue was neutralized with aqueous saturated NaHCO_3 and extracted with EtOAc (3 \times 10 mL). After the workup the organic solvent was evaporated in vacuo and the crude purified by column chromatography (SiO_2 , petroleum ether/EtOAc (8:2)), yellow oil, 1.63 g (25%). ^1H NMR (CDCl_3) δ : 1.55 (t, 3H), 2.95 (s, 3H), 4.42 (q, 2H). ^{13}C NMR (CDCl_3) δ : 162.5 (C4), 160.1 (C=O), 139.8 (C5), 62.8 (CH_2CH_3), 14.5 ($\text{CH}_2\text{CH}_3 + \text{CH}_3$). ESI-MS, m/z : 173.0 ($\text{M}^{++} + \text{H}$).

5-Hydroxymethyl-4-methyl-1,2,3-thiadiazole (23). A solution of **22** (0.8 g, 4.7 mmol) in EtOH (8.0 mL) was cooled at 0 °C. Then NaBH_4 (0.3 g, 7.9 mmol) was added slowly and the final mixture was stirred during 2 h at room temperature. The organic solvent was evaporated in vacuo, and the residue was treated with H_2O (50.0 mL) and extracted with EtOAc (3 \times 10.0 mL). After the

workup the organic solvent was evaporated in vacuo. The residue was purified by column chromatography (SiO₂, petroleum ether/EtOAc (7:3)), yielding derivative **23** as a yellow oil (0.42 g, 70%). ¹H NMR (CDCl₃) δ: 2.55 (s, 3H), 4.55 (bs, 1H), 4.94 (s, 2H). ¹³C NMR (CDCl₃) δ: 154.4 (C5), 152.9 (C4), 56.3 (CH₂OH), 12.1 (CH₃). ESI-MS, *m/z*: 131 (M⁺⁺ + H).

5-Formyl-4-methyl-1,2,3-thiadiazole (24). A mixture of **23** (0.49 g, 3.77 mmol) and MnO₂ (3.28 g, 37.7 mmol) in CHCl₃ (45 mL) as solvent is heated at reflux until absence of the alcohol **23** (checked by TLC). The resulting dispersion was then filtered through a short pad of Celite, the organic phase concentrated in vacuo, and the residue purified by column chromatography (SiO₂, petroleum ether/EtOAc (8:2)), yielding derivative **24** as a yellow oil (105 mg, 26%). ¹H NMR (CDCl₃) δ: 3.05 (s, 3H), 10.26 (s, 1H). ¹³C NMR (CDCl₃) δ: 184.3 (CHO), 156.2 (C5), 154.2 (C4), 14.4 (CH₃). ESI-MS, *m/z*: 129 (M⁺⁺ + H).

5-Ethoxycarbonyl-4-formyl-1,2,3-thiadiazole (25). A mixture of **22** (1.0 g, 5.8 mmol), *N*-bromosuccinimide (1.14 g, 6.4 mmol), and dibenzoyl peroxide (0.06 g, 0.23 mmol) in dry CCl₄ (40.0 mL) was heated at reflux for 16 h. Then the mixture was allowed to cool to room temperature, the succinimide was collected and washed with CCl₄, and the organic layer was evaporated in vacuo. DMSO (10.0 mL) was added, and the mixture was warmed at 120 °C during 3 h. Then EtOAc (50 mL) was added and the organic layer was washed with saturated sodium chloride solution (40.0 mL). After the workup the organic solvent was evaporated in vacuo and the residue purified by column chromatography (SiO₂, petroleum ether/EtOAc (8:2)), yielding a yellow oil that crystallized at 4 °C (0.4 g, 40%). ¹H NMR (CDCl₃) δ: 1.48 (t, 3H), 4.56 (q, 2H), 10.67 (s, 1H). ¹³C NMR (CDCl₃) δ: 186.1 (CHO), 157.2 (C4), 161.8 (C=O), 141.5 (C5), 64.2 (CH₂CH₃), 14.1 (CH₂CH₃). ESI-MS, *m/z*: 187 (M⁺⁺ + H).

General Procedure for the Synthesis of α-Heteroaryl-*N*-alkyl-nitron Derivatives (30–41). A mixture of the corresponding aldehyde (1 equiv), *N*-alkylhydroxylamine hydrochloride (1.5 equiv), and sodium bicarbonate (1.5 equiv) in absolute ethanol (6 mL/mmol) as solvent was heated at 60 °C until the carbonyl compound was not present (checked by TLC). The solvent was removed in vacuo and the reaction mixture diluted with H₂O and extracted with EtOAc. After the workup of the combined organic layers, the residue was purified by column chromatography (SiO₂, mixtures of petroleum ether/EtOAc).

α(Z)-(5-Phenyl-1,2,4-thiadiazol-3-yl)-*N*-tert-butylnitron (30). Yellow solid (87%); mp 111.0–113.0 °C. ¹H NMR (CDCl₃) δ: 1.63 (s, 9H), 7.48–7.56 (m, 3H), 7.97 (d, 2H), 8.18 (s, 1H). ¹³C NMR (CDCl₃) δ: 187.4 (C5), 166.5 (C3), 132.1 (Cp), 129.3 (Cm), 127.5 (Co), 123.2 (CH=N), 73.0 (C(CH₃)₃), 28.3 (C(CH₃)₃). ESI-MS, *m/z*: 262 (M⁺⁺ + H). IR (KBr) *v*: 1707, 1561, 1469, 1435, 1248, 1152, 773, 695. Anal. Calcd for (C₁₃H₁₅N₃O₃): C, 59.74; H, 5.79; N, 16.08; S, 12.27. Found: C, 59.46; H, 5.86; N, 15.97; S, 12.27.

α(Z)-(3-Phenyl-1,2,4-thiadiazol-5-yl)-*N*-tert-butylnitron (31). Yellow solid (82%); mp 107.0–108.0 °C. ¹H NMR (CDCl₃) δ: 1.66 (s, 9H), 7.44–7.50 (m, 3H), 8.29–8.32 (m, 2H), 8.63 (s, 1H). ¹³C NMR (CDCl₃) δ: 175.8 (C5), 172.0 (C3), 133.2 (Cq), 130 (Co), 128.9 (Cm), 128.2 (Cp), 126.1 (CH=N), 71.9 (C(CH₃)₃), 28.3 (C(CH₃)₃). EI-MS, *m/z* (abundance, %): 261 (M⁺⁺, 16), 205 (42), 135 (19), 103 (19), 57 (100). IR (KBr) *v*: 1544, 1470, 1414, 1344, 1234, 1112, 913, 716, 696, 558. Anal. Calcd for (C₁₃H₁₅N₃O₃): C, 59.74; H, 5.79; N, 16.08; S, 12.27. Found: C, 59.35; H, 5.74; N, 15.89; S, 12.08.

α(Z)-(3-Phenyl-1,2,4-thiadiazol-5-yl)-*N*-cyclohexylnitron (32). Yellow solid (86%); mp 194.0–196.0 °C. ¹H NMR (CDCl₃) δ: 1.21–1.50 (m, 5H), 1.99–2.20 (m, 5H), 4.16 (m, 1H), 7.52–7.47 (m, 3H), 8.34–8.31 (m, 2H), 8.51 (s, 1H). ¹³C NMR (CDCl₃) δ: 175.4 (C5), 172.2 (C3), 133.4 (Cq), 130.6 (Co), 129.1 (Cm), 128.5 (Cp), 128.0 (CH=N), 74.9 (CCH₂), 31.7 (CCH₂), 25.3 (CH₂CH₂). EI-MS, *m/z* (abundance, %): 287 (M⁺⁺, 79), 269 (32), 243 (25), 205 (86), 189 (31), 135 (60), 103 (42), 55 (100). IR (KBr) *v*: 1551, 1465, 1432, 1352, 926. Anal. Calcd for (C₁₃H₁₇N₃O₃): C, 62.69; H, 5.96; N, 14.62; S, 11.16. Found: C, 62.39; H, 5.78; N, 14.55; S, 10.98.

α(Z)-(3-Phenyl-1,2,4-thiadiazol-5-yl)-*N*-benzylnitron (33). White solid (65%); mp 156.0–158.0 °C. ¹H NMR (CDCl₃) δ: 5.21 (s, 2H), 7.46–7.45 (m, 3H), 7.48 (s, 5H), 8.30–8.27 (m, 2H), 8.35 (s, 1H). ¹³C NMR (CDCl₃) δ: 174.9 (C5), 171.9 (C3), 132.8 (Cq), 130.9 (Cq), 130.3 (Co), 129.9 (Cm), 128.8 (Co), 129.5 (Cm), 129.4 (Cp), 128.7 (Cp), 128.0 (CH=N), 69.5 (CH₂). EI-MS, *m/z* (abundance, %): 295 (M⁺⁺, 32), 278 (4), 135 (5), 103 (7), 91 (100). IR (KBr) *v*: 1534, 1465, 1423, 1323, 1234, 1130, 919, 726. Anal. Calcd for (C₁₆H₁₃N₃O₃): C, 65.06; H, 4.44; N, 14.23; S, 10.86. Found: C, 64.88; H, 4.30; N, 14.12; S, 10.61.

α(Z)-[3-(4-Chlorophenyl)-1,2,4-thiadiazol-5-yl]-*N*-tert-butylnitron (34). Yellow solid (68%); mp 129.0–131.0 °C. ¹H NMR (CDCl₃) δ: 1.68 (s, 9H), 7.46 (d, 2H, *J* = 8.5 Hz), 8.27 (d, 2H, *J* = 8.5 Hz), 8.63 (s, 1H). ¹³C NMR (CDCl₃) δ: 175.7 (C5), 170.7 (C3), 136.2 (Cp), 131.5 (Cq), 129.3 (Co), 128.9 (Cm), 125.8 (CH=N), 71.7 (C(CH₃)₃), 28.1 (C(CH₃)₃). EI-MS, *m/z* (abundance, %): 295 (M⁺⁺, 26), 239 (35), 169 (8), 137 (14), 57 (100). IR (KBr) *v*: 1548, 1471, 1420, 1404, 1346, 1236, 1118, 913, 838, 741, 558. Anal. Calcd for (C₁₃H₁₄ClN₃O₃): C, 52.79; H, 4.77; N, 14.21; S, 10.84. Found: C, 52.78; H, 5.07; N, 14.02; S, 11.01.

α(Z)-[3-(4-Methoxyphenyl)-1,2,4-thiadiazol-5-yl]-*N*-tert-butylnitron (35). Orange solid (88%); mp 130.0–132.0 °C. ¹H NMR (CDCl₃) δ: 1.67 (s, 9H), 3.87 (s, 3H), 7.01 (d, 2H, *J* = 8.7 Hz), 8.28 (m, 2H, *J* = 8.8 Hz), 8.63 (s, 1H). ¹³C NMR (CDCl₃) δ: 175.4 (C5), 171.6 (C3), 161.2 (Cp), 129.5 (Co), 125.9 (CH=N), 114.0 (Cm), 71.6 (C(CH₃)₃), 55.3 (OCH₃), 28.0 (C(CH₃)₃). EI-MS, *m/z* (abundance, %): 291 (M⁺⁺, 91), 235 (100), 165 (26), 133 (38), 57 (89). IR (KBr) *v*: 1609, 1547, 1408, 1314, 1246, 1180, 1026, 838, 750, 659, 556. Anal. Calcd for (C₁₄H₁₇N₃O₂S): C, 57.71; H, 5.88; N, 14.42; S, 11.00. Found: C, 56.56; H, 5.68; N, 14.23; S, 10.80.

α(Z)-[3-(3,5-Di-*tert*-butyl-4-hydroxyphenyl)-1,2,4-thiadiazol-5-yl]-*N*-tert-butylnitron (36). Yellow solid (92%); mp 292.0–294.0 °C. ¹H NMR (CDCl₃) δ: 1.51 (s, 18H), 1.68 (s, 9H), 5.48 (s, 1H), 8.19 (s, 2H), 8.64 (s, 1H). ¹³C NMR (CDCl₃) δ: 175.2 (C5), 172.7 (C3), 155.9 (Cp), 136.1 (Cq), 126.0 (Co), 125.2 (CH=N), 124.6 (Cm), 71.5 (C(CH₃)₃), 34.4 (C(CH₃)₃), 30.3 (C(CH₃)₃), 28.4 (C(CH₃)₃). EI-MS, *m/z* (abundance, %): 389 (M⁺⁺, 87), 374 (46), 333 (36), 318 (100), 57 (40). IR (KBr) *v*: 3593, 2961, 1544, 1398, 1254, 1236, 1121, 913, 747, 696, 555. Anal. Calcd for (C₂₁H₃₁N₃O₂S): C, 64.75; H, 8.02; N, 10.79; S, 8.23. Found: C, 64.50; H, 8.30; N, 10.63; S, 8.18.

α(Z)-(4-Methyl-1,2,3-thiadiazol-5-yl)-*N*-tert-butylnitron (37). Beige solid (77%); mp 154.0–156.0 °C. ¹H NMR (CDCl₃) δ: 1.67 (s, 9H), 2.85 (s, 3H), 8.20 (s, 1H). ¹³C NMR (CDCl₃) δ: 157.7 (C5), 138.4 (C4), 122.7 (CH=N), 71.2 (C(CH₃)₃), 28.5 (C(CH₃)₃), 13.9 (CH₃). EI-MS, *m/z* (abundance, %): 199 (M⁺⁺, 2), 171 (2), 123 (30), 108 (42), 57 (100). IR (KBr) *v*: 3136, 2997, 2972, 1547, 1493, 1439, 1361, 1323, 1201, 1140, 1035, 887, 827, 723, 572. Anal. Calcd for (C₈H₁₃N₃O₂S): C, 48.22; H, 6.58; N, 21.09; S, 16.09. Found: C, 48.09; H, 6.30; N, 20.82; S, 15.88.

α(Z)-(5-Ethoxycarbonyl-1,2,3-thiadiazol-4-yl)-*N*-tert-butylnitron (38). Yellow solid (48%); mp 93.0–95.0 °C. ¹H NMR (CDCl₃) δ: 1.44 (t, 3H), 1.65 (s, 9H), 4.48 (q, 2H), 8.52 (s, 1H). ¹³C NMR (CDCl₃) δ: 159.9 (C=O), 153.9 (C4), 143.2 (C5), 121.2 (CH=N), 72.8 (C(CH₃)₃), 63.3 (CH₂CH₃), 28.5 (C(CH₃)₃), 14.5 (CH₂CH₃). EI-MS, *m/z* (abundance, %): 257 (M⁺⁺, 1), 201 (9), 173 (9), 127 (6), 57 (100). IR (KBr) *v*: 3147, 299, 1732, 1562, 1535, 1394, 1313, 1209, 1153, 1087, 910, 860, 767, 559. Anal. Calcd for (C₁₀H₁₅N₃O₃S): C, 46.68; H, 5.88; N, 16.33; S, 12.46. Found: C, 46.33; H, 5.50; N, 16.20; S, 12.19.

α(Z)-(4-Phenylfuroxan-3-yl)-*N*-tert-butylnitron (39). Yellow solid (57%); mp 78.0–80.0 °C. ¹H NMR (CDCl₃) δ: 1.61 (s, 9H), 7.42–7.56 (m, 5H), 7.58 (s, 1H). ¹³C NMR (CDCl₃) δ: 155.8 (C4), 133.9 (Cm), 130.5 (Co), 126.7 (Cp), 114.9 (CH=N), 110.3 (C3), 73.2 (C(CH₃)₃), 28.3 (C(CH₃)₃). EI-MS, *m/z* (abundance, %): 261 (M⁺⁺, 2), 205 (18), 145 (10), 105 (18), 57 (100). IR (KBr) *v*: 3138, 2979, 2935, 1600, 1560, 1456, 1382, 1247, 1199, 1139, 1028, 983, 858, 812, 709, 556. Anal. Calcd for (C₁₃H₁₅N₃O₃): C, 59.76; H, 5.79; N, 16.08. Found: C, 59.45; H, 5.61; N, 15.87.

α(Z)-(3-Methylfuroxan-4-yl)-*N*-tert-butylnitron (40). Yellow solid (81%); mp 119.0–121.0 °C. ¹H NMR (CDCl₃) δ: 1.61 (s,

9H), 2.33 (s, 3H), 7.96(s, 1H). ¹³C NMR (CDCl₃) δ: 151.1 (C4), 117.9 (CH=N), 112.6 (C3), 72.9 (C(CH₃)₃), 28.6 (C(CH₃)₃), 10.4 (CH₃). EI-MS, *m/z* (abundance, %): 199 (M⁺, 3), 143 (9), 113 (6), 83 (4), 57 (100). IR (KBr) *v*: 2984, 1616, 1566, 1457, 1364, 1251, 1178, 1140, 834, 763, 549. Anal. Calcd for (C₈H₁₃N₃O₃): C, 48.23; H, 6.58; N, 21.09. Found: C, 48.02; H, 6.30; N, 20.92.

α(Z)-(Benzofuroxan-5(6)-yl)-N-tert-butylnitron (41). Yellow solid (61%); mp 112.0–114.0 °C. ¹H NMR (CDCl₃) δ: 1.60 (s, 9H), 7.32 (bs, 1H), 7.50 (bs, 1H), 7.60 (s, 1H), 9.33 (bs, 1H). ¹³C NMR (CDCl₃) δ: 127.4 (CH=N), 72.5 (C(CH₃)₃), 28.3 (C(CH₃)₃). EI-MS, *m/z* (abundance, %): 235 (M⁺, 9), 179 (17), 163 (5), 152 (6), 57 (100). IR (KBr) *v*: 2984, 1610, 1570, 1487, 1361, 1236, 1153, 1014, 868, 828, 585. Anal. Calcd for (C₁₁H₁₃N₃O₃): C, 56.16; H, 5.57; N, 17.86. Found: C, 55.99; H, 5.34; N, 17.70.

Measurement of Cell Viability. Culture of SH-SY5Y Cells. SH-SY5Y cells, at passages between 3 and 16 after defreezing, were maintained in a minimum essential medium (MEM) containing 15 nonessential amino acids (NEAAs) and supplemented with 10% fetal bovine serum, 1 mM glutamine, 50 units/mL penicillin, and 50 μg/mL streptomycin (reagents from GIBCO, Madrid, Spain). Cultures were seeded into flasks containing supplemented medium and maintained at 37 °C in 5% CO₂/humidified air.

Cell Viability Assays. The effects of nitrones on SH-SY5Y cell viability were determined by measuring the activity of LDH released into the culture medium (Cytotox 96 LDH assay, Promega, U.K.). The assay was performed in accordance with the manufacturer's protocol using SH-SY5Y cells seeded at 10⁴ cells/well in 96-well plates. Cells were exposed to 10, 1, and 0.05 μM of each one nitron for 24 h.

The effects of H₂O₂ on SH-SY5Y cell viability were determined by measuring the activity of LDH released into the culture medium (Cytotox 96 LDH assay, Promega, U.K.). The assay was performed in accordance with the manufacturer's protocol using SH-SY5Y cells seeded at 10⁴ cells/well in 96-well plates. Cells were exposed to 10 μM of each one heteroarylnitron for 1 h and subsequently exposed to 100 μM H₂O₂ for 24 h. An amount of 5 mM NAC was used as positive test.

Unspecific Mammalian Cytotoxicity. J774 murine macrophage-like cells (ATCC) were maintained by passage in Dulbecco's modified Eagle's medium (DMEM) containing 4 mM L-glutamine and supplemented with 10% heat-inactivated fetal calf serum. J774 cells were seeded (1 × 10⁵ cells/well) in 96-well microplates with 100 μL of RPMI medium supplemented with fetal serum. Cells were allowed to attach for 48 h in a humidified 5% CO₂/95% air atmosphere at 37 °C and then exposed to nitrones (50–500 μM) for 48 h. Afterward, cell viability was assessed by measuring the mitochondrial-dependent reduction of 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT, Sigma) to formazan. For that purpose, MTT was added to cells to a final concentration of 0.2 mg/mL and cells were incubated at 37 °C for 3 h. After removal of the media, formazan crystals were dissolved in DMSO, and the absorbance at 560 nm was read using a microplate spectrophotometer. Results are expressed as IC₅₀ (compound concentration that reduce 50% control absorbance at 560nm). Every IC₅₀ is the average of two different experiments.

Blood–Brain Barrier Permeation Assay. Brain penetration of new nitrones was evaluated using a parallel artificial membrane permeation assay (PAMPA) in a similar manner as described previously.²⁷ Commercial drugs were purchased from Sigma, Aldrich, Acros, and Fluka. Millex filter units (PVDF membrane, diameter 25 mm, pore size 0.45 μm) were acquired from Millipore. The porcine brain lipid (PBL) was obtained from Avanti Polar Lipids. The donor microplate was a 96-well filter plate (PVDF membrane, pore size 0.45 μm), and the acceptor microplate was an indented 96-well plate, both from Millipore. The acceptor 96-well microplate was filled with 170 μL of PBS/ethanol (9:1), and the filter surface of the donor microplate was impregnated with 4 μL of PBL in dodecane (20 mg/mL). Compounds were dissolved in PBS/ethanol (9:1) at 1 mg/mL, filtered through a Millex filter, and then added to the donor wells (170 μL). The donor filter plate was carefully put on the acceptor plate to form a sandwich, which

was left undisturbed for 120 min at 25 °C. After incubation, the donor plate was carefully removed and the concentration of compounds in the acceptor wells was determined by UV spectroscopy. Every sample was analyzed in four wells and at least in three independent runs, and the results are given as the mean ± standard deviation.

Determination of Antioxidant Activity. ORAC_{FL} Assay. A luminescence spectrometer LS 50B (PerkinElmer, Boston, MA), a heating circulator bath DC1-B3 (Haake Haake Fisons, Karlsruhe, Germany), and quartz cuvettes were used. For the ORAC_{FL} assay, the 490-P excitation and 515-P emission filters were used, and the fluorescence measurement was carried out at 60 °C. The method of Ou et al.²⁹ was modified as follows: the reaction was carried out in 75 mM phosphate buffer (pH 7.4), and the final reaction volume was 3000 μL. Studied compounds (10–1000 μM final concentrations) and fluorescein (215 μL, 70 nM final concentration) solutions were placed in the quartz cuvette. The mixture was preincubated for 30 s at 60 °C. 2,2'-Azobis(2-amidinopropane) dihydrochloride (AAPH) solution (240 μL; 12 mM, final concentration) was added rapidly using a single channel pipet. The quartz cuvette was immediately placed in the luminescence spectrometer and the fluorescence recorded every minute for 12 min. As a blank FL + AAPH in phosphate buffer instead of the studied compounds solutions were employed, and eight calibration solutions using Trolox (1–8 μM, final concentration) as antioxidant positive control were also carried out in each assay. All the reaction mixtures were prepared in duplicate, and at least three independent assays were performed for each sample. Raw data were exported to an OriginPro (OriginLab Corporation, Northampton, MA) sheet for further calculations. Blank and antioxidant curves (fluorescence versus time) were first normalized by dividing original data by fluorescence at *t* = 0 s. From the normalized curves, the area under the fluorescence decay curve (AUC) was calculated as

$$\text{AUC} = 1 + \sum_{i=1}^{i=12} \frac{f_i}{f_0}$$

where *f*₀ is the initial fluorescence reading at 0 min and *f*_{*i*} is the fluorescence reading at time *i*. The net AUC corresponding to each sample was calculated by subtracting the AUC corresponding to the blank. Regression equations between net AUC and antioxidant concentration were calculated for all the samples. ORAC_{FL} values were expressed as PBN equivalents by using the standard curve calculated for each assay. Final results were expressed in μmole of PBN equivalent per μmole of samples.

DPPH Assay. The reaction was carried out in 96-well microplates, and the final reaction mixture was 100 μL per well. Antioxidant (0.1 mM, final concentration) and DPPH (0.1 mM, final concentration) were dissolved in EtOH and preincubated for 5 min at 37 °C, and the absorbance at 520 nm was read using a microplate spectrophotometer. The antioxidant activity was determined as the RP of DPPH, calculated as follows: RP = 100 [(A₀ - A_c)/A₀], where A₀ is the untreated DPPH absorbance and A_c is the value for added sample concentration *c*.

ESR Spectroscopy. The spin trapping ability of the developed heteroarylnitrones was studied by ESR spectroscopy. ESR spectra were recorded in the X band (9.85 GHz) using a Bruker ECS 106 spectrometer with a rectangular cavity and 50 kHz field modulation. All the spectra were registered in the same scale after 15 scans. The hyperfine splitting constants were estimated to be accurate within 0.05 G.

Generation of Free Radicals. Hydroxyl radical (•OH) was generated from a standard Fenton system: Fe(NH₄)₂(SO₄)₂•6H₂O (50 μL, 0.5 mM) was added to a solution of nitron (50 mM, final concentration) and H₂O₂ (50 μL, 0.5 mM) in phosphate buffer (final reaction mixture was 250 μL). Cysteinyl radical (SO₃^{•-}) was generated from a standard Fenton system: Fe(NH₄)₂(SO₄)₂•6H₂O (50 μL, 0.5 mM) was added to a solution of nitron (50 mM, final concentration), Na₂SO₃ (50 μL, 250 mM), and H₂O₂ (50 μL, 0.5 mM) in phosphate buffer (final reaction mixture was 250 μL).

Ethanol radical (CH₃•CHOH) was generated from a standard Fenton system: Fe(NH₄)₂(SO₄)₂•6H₂O (50 μL, 0.5 mM) was added to a solution of nitron (50 mM, final concentration), EtOH (50 μL), and H₂O₂ (50 μL, 0.5 mM) in phosphate buffer (final reaction mixture was 250 μL).

Acknowledgment. The authors thank Collaborative Project UdelaR (Uruguay), CSIC (Spain) (Grant 2006UY0009), and Neuropharma S.A (Madrid, Spain). W.P. thanks Collaborative Project (Grant 2006UY0009) and CSIC (UdelaR) for fellowships. W.P. and P.H. are PEDECIBA students.

Supporting Information Available: Elemental analysis results for heteroaryl nitron derivatives. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Betteridge, D. J. What Is Oxidative Stress? *Metabolism* **2000**, *49* (Suppl. 1), 3–8.
- (2) Radi, R. Nitric Oxide, Oxidants, and Protein Tyrosine Nitration. *Proc. Nat. Acad. Sci. U.S.A.* **2004**, *101*, 4003–4008.
- (3) Kowald, A.; Kirkwood, T. B. Accumulation of Defective Mitochondria through Delayed Degradation of Damaged Organelles and Its Possible Role in the Ageing of Post-Mitotic and Dividing Cells. *J. Theor. Biol.* **2000**, *202*, 145–160.
- (4) Lu, C. Y.; Lee, H. C.; Fahh, H. J.; Wei, Y. H. Oxidative Damage Elicited by Imbalance of Free Radical Scavenging Enzymes Is Associated with Large-Scale mtDNA Deletions in Aging Human Skin. *Mutat. Res.* **1999**, *423*, 11–21.
- (5) Hensley, K.; Robinson, K. A.; Gabbita, P.; Salsman, S.; Floyd, R. A. Reactive Oxygen Species, Cell Signaling, and Cell Injury. *Free Radical Biol. Med.* **2000**, *28*, 1456–1462.
- (6) Floyd, R. A.; Hensley, K. Oxidative Stress in Brain Aging Implications for Therapeutics of Neurodegenerative Diseases. *Neurobiol. Aging* **2002**, *23*, 795–807.
- (7) Yokoyama, M. Oxidant Stress and Atherosclerosis. *Curr. Opin. Pharmacol.* **2004**, *4*, 110–115.
- (8) Aslan, M.; Ozben, T. Reactive Oxygen and Nitrogen Species in Alzheimer's Disease. *Curr. Alzheimer Res.* **2004**, *1*, 111–19.
- (9) Floyd, R. A. Antioxidant, Oxidative Stress, and Degenerative Neurological Disorders. *Proc. Soc. Exp. Biol. Med.* **1999**, *222*, 236–245.
- (10) Tabner, B. J.; Turnbull, S.; El-Agnaf, O. M. A.; Allsop, D. Formation of Hydrogen Peroxide and Hydroxyl Radicals from β and α-Synuclein as a Possible Mechanism of Cell Death in Alzheimer's Disease and Parkinson's Disease. *Free Radical Biol. Med.* **2002**, *32*, 1076–1083.
- (11) Floyd, R. A.; Hensley, K.; Forster, M.; Kelleher-Andersson, J. A.; Wood, P. L. Nitrones, Their Value as Therapeutics and Probes To Understand Aging. *Mech. Ageing Dev.* **2002**, *123*, 1021–1031.
- (12) Floyd, R. A.; Hensley, K.; Jaffery, F.; Maitt, L.; Robinson, K.; Pye, Q.; Stewart, C. Increased Oxidative Stress Brought on by Pro-Inflammatory Cytokines in Neurodegenerative Processes and the Protective Role of Nitron-Based Free Radical Traps. *Life Sci.* **1999**, *65*, 1893–1899.
- (13) Goldstein, S.; Lestage, P. Chemical and Pharmacological Aspects of Heteroaryl-Nitrones. *Curr. Med. Chem.* **2000**, *7*, 1255–1267.
- (14) Kelleher, J.; Maples, K. R.; Waterbury, L. D.; Wilcox, A. L.; Xu, Hong; Zhang, Y. K. WO 98/03496, 2002 (Centaur Pharmaceuticals, Inc., 484 Oakmead Parkway, Sunnyvale, CA 94086).
- (15) Dhainaut, A.; Tizot, A.; Raimbaud, E.; Lockhart, B.; Lestage, P.; Goldstein, S. Synthesis, Structure, and Neuroprotective Properties of Novel Imidazolyl Nitrones. *J. Med. Chem.* **2000**, *43*, 2165–2175.
- (16) Villamena, F. A.; Zweier, J. L. Detection of Reactive Oxygen and Nitrogen Species by EPR Spin Trapping. *Antioxid. Redox Signaling* **2004**, *6*, 619–629.
- (17) Castro, A.; Gil, C.; Encinas, A.; Castaño, T.; Porcal, W. Advances on the Synthesis and Recent Therapeutic Applications of 1,2,4-Thiadiazole Heterocycles. *Bioorg. Med. Chem.* **2006**, *14*, 1644–1652.
- (18) Porcal, W.; Cerecetto, H. Pharmacological Properties of Furoxans and Benzofuroxans: Recent Developments. *Mini-Rev. Med. Chem.* **2005**, *5*, 57–71.
- (19) Patai, A.; Rappoport, Z. *Nitrones, Nitronates and Nitroxides*; John Wiley & Sons Ltd.: New York, 1989.
- (20) Lin, Y.-i.; Lang, S. A.; Petty, S. R. New Synthesis of 1,2,4-Thiadiazole. *J. Org. Chem.* **1980**, *45*, 3750–3753.
- (21) Howe, R. K.; Franz, J. E. Nitrile Sulphides. Synthesis of 1,2,4-Thiadiazole. *J. Org. Chem.* **1974**, *39*, 962–964.
- (22) Bakulev, V. A.; Dehaen, W. Synthesis of 1,2,3-Thiadiazoles. *The Chemistry of 1,2,3-Thiadiazoles*; John Wiley & Sons, Inc.: Hoboken, NJ, 2004.
- (23) Cerecetto, H.; Di Maio, R.; González, M.; Risso, M.; Saenz, P.; Seoane, G.; Denicola, A.; Peluffo, G.; Quijano, C.; Olea-Azar, C. 1,2,5-Oxadiazole N-Oxide Derivatives and Related Compounds as Potential Antitrypanosomal Drugs: Structure–Activity Relationships. *J. Med. Chem.* **1999**, *42*, 1941–1950.
- (24) Porcal, W.; Boiani, M.; Gerpe, A.; Merlino, A.; González, M.; Cerecetto, H. Arylethylbenzofuroxan Derivatives as Drugs for Chagas Disease: Multigram-Batch Synthesis using Wittig–Boden Process. *Org. Process Res. Dev.* **2008**, *12*, 156–162.
- (25) Marco-Contelles, J.; León, R.; de los Ríos, C.; Guglietta, A.; Terencio, J.; López, M. G.; García, A. G.; Villarroya, M. Novel Multipotent Tacrine–Dihydropyridine Hybrids with Improved Acetylcholinesterase Inhibitory and Neuroprotective Activities as Potential Drugs for the Treatment of Alzheimer's Disease. *J. Med. Chem.* **2006**, *49*, 7607–7610.
- (26) López, G. V.; Blanco, F.; Hernández, P.; Ferreira, A.; Piro, O. E.; Bathyány, C.; González, M.; Rubbo, H.; Cerecetto, H. Second Generation of α-Tocopherol Analogs–Nitric Oxide Donors: Synthesis, Physicochemical and Biological Characterization. *Bioorg. Med. Chem.* **2007**, *15*, 6262–6272.
- (27) Rodríguez-Franco, M. I.; Fernández-Bachiller, M. I.; Pérez, C.; Hernández-Ledesma, B.; Bartolomé, B. Novel Tacrine–Melatonin Hybrids as Dual-Acting Drugs for Alzheimer Disease, with Improved Acetylcholinesterase Inhibitory and Antioxidant Properties. *J. Med. Chem.* **2006**, *49*, 459–462.
- (28) Di, L.; Kerns, E. H.; Fan, K.; McConnell, O. J.; Carter, G. T. High Throughput Artificial Membrane Permeability Assay for Blood–Brain Barrier. *Eur. J. Med. Chem.* **2003**, *38*, 223–232.
- (29) Ou, B.; Hampsch-Woodill, M.; Prior, R. L. Development and Validation of an Improved Oxygen Radical Absorbance Capacity Assay Using Fluorescein as the Fluorescent Probe. *J. Agric. Food Chem.* **2001**, *49*, 4619–4626.
- (30) Molyneux, P. The Use of the Stable Free Radical Diphenylpicrylhydrazyl (DPPH) for Estimating Antioxidant Activity. *J. Sci. Technol.* **2004**, *26*, 211–219.
- (31) Reybier, K.; Boyer, J.; Farines, V.; Camus, F.; Souchard, J.; Monje, M.-C.; Bernardes-genisson, V.; Goldstein, S.; Nepveu, F. Radical Trapping Properties of Imidazolyl Nitrones. *Free Radical Res.* **2006**, *40*, 11–20.
- (32) Kim, S.; Vilela, G. V. M.; Bouajila, J.; Dias, A. G.; Cyrino, F. Z.; Bouskela, E.; Costa, P. R. R.; Nepveu, F. α-Phenyl-*N-tert*-butyl nitron (PBN) Derivatives: Synthesis and Protective Action against Microvascular Damages Induced by Ischemia/Reperfusion. *Bioorg. Med. Chem.* **2007**, *15*, 3572–3578.
- (33) Howe, R. K.; Gruner, T. A.; Franz, J. E. Nitrile Sulphides. Synthesis of 5-Aryl-1,2,4-thiadiazole-3-carboxylates. *J. Org. Chem.* **1977**, *42*, 1813–1814.
- (34) Castro, A.; Encinas, A.; Gil, C.; Bräse, S.; Porcal, W.; Pérez, C.; Moreno, F. J.; Martínez, M. Non ATP-Competitive Glycogen Synthase Kinase β (GSK-3β) Inhibitors: Study of Structural Requirements for Thiadiazolidinone Derivatives. *Bioorg. Med. Chem.* **2008**, *16*, 495–510.
- (35) Paton, R. M.; Robertson, F. M.; Ross, J. F. Nitrile Sulphides. Part 3. Thermal Fragmentation of 1,3,4-Oxathiazoles: Formation of Nitriles Sulphides in a Retro-1,3-Dipolar Cycloaddition Reaction. *J. Chem. Soc., Perkin Trans. 1* **1985**, *151*, 7–1521.
- (36) Unangst, P. C.; Shrum, G. P.; Connor, D. T.; Dyer, R. D.; Schriert, D. J. Novel 1,2,4-Oxadiazoles and 1,2,4-Thiadiazoles as Dual 5-Lipoxygenase and Cyclooxygenase Inhibitors. *J. Med. Chem.* **1992**, *35*, 3691–3698.

JM8006432