

Nonacarbonyl[μ_3 -(1,2-diphenylethyl)-phenylphosphinito]triosmium(2 Os—Os)

MARIO V. CAPPARELLI,^a YSAURA DE SANCTIS,^a
ALEJANDRO J. ARCE^a AND EVGENIA SPODINE^b

^a*Centro de Química, Instituto Venezolano de Investigaciones Científicas (IVIC), Apartado 21827, Caracas 1020-A, Venezuela,* and ^b*Facultad de Ciencias Químicas y Farmacéuticas, Universidad de Chile, Casilla 233, Santiago 1, Chile. E-mail: mcappare@dino.conicit.ve*

(Received 10 April 1996; accepted 20 June 1996)


Abstract

The reaction of 1,2,3-triphenylphosphirene with $[\text{H}_2\text{Os}_3(\text{CO})_{10}]$ leads to the title compound, $[\text{Os}_3(\text{CO})_9\{(\text{C}_6\text{H}_5)_3\text{C}_2\text{PO}\}]$, which displays an open $\text{Os}_3(\text{CO})_9$ cluster [Os—Os 2.703 (1) and 2.867 (2) Å, and Os—Os—Os 124.24 (4)°]. This moiety is μ_3, η^3 -bonded to a phosphirene ligand with a cleaved P—C bond, PhC—C(Ph)—P . An O atom was incorporated into the molecule resulting in an unusual five-membered metallacycle,

Os—C—P—O, with distances Os—C 2.10 (1), C—C 1.46 (2), C—P 1.76 (1), P—O 1.628 (9) and O—Os 2.136 (9) Å. This ring makes a dihedral angle of 57.3 (2)° with the plane of the metal atoms.

Comment

Three-membered phosphirene rings, (RC=CR)PR, are potentially very useful precursors to a range of organometallic ligands such as acetylenes, phosphine-indenes and various other linked systems (Marinetti, Fischer & Mathey, 1985; Svava, Marinetti & Mathey, 1986; Carmichael, Hitchcock, Nixon, Mathey & Pidcock, 1986). We recently studied the reaction of 1,2,3-triphenylphosphirene, (PhC=CPh)PPh, with [Ru₃(CO)₁₂] (Arce *et al.*, 1995), which was found to produce the complex [Ru₃(CO)₈(μ₃-C₄Ph₆P₂)] containing two opened and coupled phosphirene rings. We report here the results of a similar reaction with [H₂Os₃(CO)₁₀], which afforded a different and totally unexpected compound, (I).



The structure analysis of the reaction product showed that the crystals contain discrete molecules of [Os₃(CO)₉(Ph₃C₂PO)] (Fig. 1). The title compound displays an open Os₃(CO)₉ cluster μ₃,η³-bonded to a phosphirene ligand in which one of the P—C bonds was cleaved, as in the Ru complex reaction mentioned above. In this case, however, an O atom (O1) was incorporated into the molecule resulting in an unusual five-membered Os1—C2—C1—P—O1 ring. The source of this atom could be either a carbonyl group, molecular oxygen or moisture. The first possibility is very unlikely due to the stability of the C≡O bond, so that it is probable that the O1 atom comes from a contaminant, presumably moisture. In spite of repeated efforts (see *Experimental*), we were unable to establish and/or eliminate the source of the O atom.

We have no doubt that O1 is a single O atom (not OH). The evidence is both crystallographic (*U*_{eq} value




Fig. 1. The molecular structure of the title compound showing the atomic numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

for O1 is quite sensible as opposed to refinement as C or N) and chemical spectroscopic (the electron count for the cluster is precise); the CPh—CPh—PPh—O ligand acts as an eight-electron donor group for a 50-electron cluster, there are only two Os—Os bonds and no evidence of Os—H in the ¹H NMR spectrum.

The Os2—Os3 bond length is within the range of distances observed in other open Os₃ clusters, e.g. 2.819 (2)—2.930 (2) Å in [HOs₃(CO)₉(PC₄H₄Me)-(C₆H₄)] (Deeming, Powell, Arce, De Sanctis & Manzur, 1991) and 2.813 (1)—2.948 (1) Å in [Os₃(CO)₉(PhPC₄H₄)] (Arce, De Sanctis, Manzur, Deeming & Powell, 1991). On the other hand, the O1 atom bridging the Os1 and Os2 atoms produces a significant shortening of the Os1—Os2 distance. The Os3—P bond length is within the range of distances (ca 2.32—2.41 Å) observed for Os—P bonds in tertiary phosphines (Orpen *et al.*, 1989) and similar to an equivalent Ru—P distance [2.337 (1) Å] found in [Ru₃(CO)₈(μ₃-C₄Ph₆P₂)] (Arce *et al.*, 1995). The Os2...P non-bonded distance is 2.581 (4) Å. The Os—C bonds to the vinyl moiety which bridges Os1 and Os2 (i.e. Os1—C2, Os2—C1 and Os2—C2) have lengths comparable to the equivalent Ru—C bonds of [Ru₃(CO)₈(μ₃-C₄Ph₆P₂)], i.e. 2.093 (4), 2.316 (4) and 2.217 (4) Å, respectively. In the present structure, however, both π bonds are equal, while in [Ru₃(CO)₈(μ₃-C₄Ph₆P₂)], they are significantly asymmetric, probably due to steric distortion. As expected, the Os1—O1 σ bond is significantly shorter than the Os2—O1 π bond.

The P atom lies 0.995 (4) Å from the plane of the metal atoms and displays a distorted tetrahedral environment, with angles in the range 96.9 (6)—126.8 (5)°.

The P—C and P—O distances are within normal ranges (Allen *et al.*, 1987).

The most interesting feature of this compound is the five-membered Os—C—C—P—O metallacycle. To the best of our knowledge, no compound containing a similar ring with any transition metal has been structurally characterized so far. The ring is nearly planar, but is better described as an envelope, with the O1 atom at a distance of 0.324 (9) Å (towards Os2) from the plane of the other four atoms (mean deviation from plane is 0.015 Å). The C51 and C61 atoms are approximately coplanar with that plane [distances to plane are 0.04 (1) and 0.24 (1) Å, respectively], while it makes a dihedral angle of 57.3 (2)° with the Os₃ plane.

Experimental

1,2,3-Triphenylphosphirene, (PhC=CPh)PPh, was prepared according to the method of Lochsmitz, Mathey & Schmidpeter (1986). To a solution of [H₂Os₃(CO)₁₀] (0.10 g, 0.12 mmol) in dry THF (40 ml) was added an equimolar amount of triphenylphosphirene (0.34 g, 0.12 mmol) and the resulting solution heated under reflux under nitrogen for 24 h. Removal of the solvent *in vacuo* and TLC separation (SiO₂, *n*-hexane) yielded the title compound (0.011 g; yield 8%) as a yellow solid. Crystals suitable for X-ray analysis were obtained by slow evaporation of a cyclohexane solution. To establish the source of a possible contaminant, and eventually to eliminate it, we repeated the reaction three times, but in all cases we obtained the same results. To avoid molecular oxygen, the nitrogen gas was passed through a trap containing Cr²⁺/Zn—Hg and to ensure the dryness of the solvents they were distilled over Na wire (using benzophenone as an indicator).

Crystal data

[Os₃(C₂₀H₁₅OP)(CO)₉]

M_r = 1125.01

Monoclinic

*P*2₁/*n*

a = 11.049 (6) Å

b = 17.638 (3) Å

c = 15.894 (2) Å

β = 91.12 (2)°

V = 3097 (1) Å³

Z = 4

D_x = 2.413 Mg m⁻³

D_m not measured

Data collection

Rigaku AFC-7S diffractometer

ω -2 θ scans

Absorption correction: refined from ΔF (DIFABS; Walker & Stuart, 1983)

T_{min} = 0.488, *T_{max}* = 1.000

5624 measured reflections

5316 independent reflections

Mo *K* α radiation

λ = 0.71069 Å

Cell parameters from 25 reflections

θ = 23.5–28.9°

μ = 12.377 mm⁻¹

T = 295 (1) K

Prism

0.33 × 0.15 × 0.13 mm

Yellow

4133 reflections with

I > 3 σ (*I*)

R_{int} = 0.0236

θ_{max} = 25°

h = 0 → 13

k = 0 → 20

l = -18 → 18

3 standard reflections

every 150 reflections intensity decay: 18.2%

Refinement

Refinement on *F*²

R = 0.0454

wR = 0.0552

S = 3.200

4133 reflections

403 parameters

Only H-atom *U*'s refined

Weighting scheme based on counting statistics

(Δ/σ)_{max} = 0.0007

$\Delta\rho_{max}$ = 1.67 e Å⁻³

$\Delta\rho_{min}$ = -1.61 e Å⁻³

Extinction correction: none

Scattering factors from International Tables for X-ray Crystallography (Vol. IV)

Table 1. Selected geometric parameters (Å, °)

Os1—Os2	2.703 (1)	Os3—P	2.367 (4)
Os1—O1	2.136 (9)	Os3—C31	1.94 (2)
Os1—C2	2.10 (1)	Os3—C32	1.96 (2)
Os1—C11	1.96 (2)	Os3—C33	2.00 (2)
Os1—C12	1.82 (2)	Os3—C34	1.95 (2)
Os1—C13	1.93 (2)	P—O1	1.628 (9)
Os2—Os3	2.867 (2)	P—C1	1.76 (1)
Os2—O1	2.188 (9)	P—C41	1.81 (1)
Os2—C1	2.31 (1)	C1—C2	1.46 (2)
Os2—C2	2.32 (1)	C1—C51	1.50 (2)
Os2—C21	1.87 (2)	C2—C61	1.47 (2)
Os2—C22	1.87 (2)		
Os2—Os1—O1	52.2 (2)	O1—P—C41	105.6 (6)
Os2—Os1—C2	56.0 (4)	C1—P—C41	115.4 (6)
O1—Os1—C2	78.5 (5)	Os1—O1—Os2	77.4 (3)
Os1—Os2—Os3	124.24 (4)	Os1—O1—P	123.7 (5)
Os1—Os2—O1	50.5 (2)	Os2—O1—P	83.7 (4)
Os1—Os2—C1	75.7 (3)	Os2—C1—P	77.3 (4)
Os1—Os2—C2	48.8 (3)	Os2—C1—C2	71.7 (7)
Os3—Os2—O1	74.1 (2)	Os2—C1—C51	126.0 (9)
Os3—Os2—C1	79.3 (3)	P—C1—C2	117.8 (9)
Os3—Os2—C2	115.1 (3)	P—C1—C51	121 (1)
O1—Os2—C1	68.6 (4)	C2—C1—C51	120 (1)
O1—Os2—C2	73.0 (4)	Os1—C2—Os2	75.2 (4)
C1—Os2—C2	36.8 (5)	Os1—C2—C1	119.4 (9)
Os2—Os3—P	58.14 (9)	Os1—C2—C61	119.3 (9)
Os3—P—O1	99.9 (4)	Os2—C2—C1	71.5 (7)
Os3—P—C1	106.7 (5)	Os2—C2—C61	130.8 (9)
Os3—P—C41	126.8 (5)	C1—C2—C61	120 (1)
O1—P—C1	96.9 (6)		

An absorption correction based on ψ -scan data gave unsatisfactory results. Better results were obtained with the DIFABS (Walker & Stuart, 1983) procedure. H atoms were placed in calculated positions (C—H 0.96 Å) and assigned fixed coordinates and refined with individual isotropic displacement parameters. The structure contains two symmetry-related solvent accessible voids (0.0, 0, and $\frac{1}{2}, \frac{1}{2}, \frac{1}{2}$) of ≈ 60 Å³ each, however, no significant peaks were found there. The 12 largest peaks in the final difference map were near osmium and were interpreted as residual absorption effects.

Data collection: MSCIAFC Diffractometer Control Software (Molecular Structure Corporation, 1993a). Cell refinement: MSCIAFC Diffractometer Control Software. Data reduction: TEXSAN (Molecular Structure Corporation, 1993b). Program(s) used to solve structure: SHELXS86 (Sheldrick, 1990). Program(s) used to refine structure: TEXSAN. Software used to prepare material for publication: TEXSAN.

The authors acknowledge the Venezuelan National Research Council (CONICIT) for providing funds for the purchase of the Rigaku AFC-7S diffractometer and auxiliary equipment (Project PI-092).

Lists of atomic coordinates, displacement parameters, structure factors and complete geometry have been deposited with the IUCr (Reference: SK1030). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–S19.
- Arce, A. J., De Sanctis, Y., Machado, R., Capparelli, M. V., Manzur, J. & Deeming, A. J. (1995). *Organometallics*, **14**, 3592–3595.
- Arce, A. J., De Sanctis, Y., Manzur, J., Deeming, A. J. & Powell, N. I. (1991). *J. Organomet. Chem.* **408**, C18–C21.
- Carmichael, D., Hitchcock, P. B., Nixon, J. F., Mathey, F. & Pidcock, A. (1986). *J. Chem. Soc. Chem. Commun.* pp. 762–763.
- Deeming, A. J., Powell, N. I., Arce, A. J., De Sanctis, Y. & Manzur, J. (1991). *J. Chem. Soc. Dalton Trans.* pp. 3381–3386.
- Lochsmidt, S., Mathey, F. & Schmidpeter, A. (1986). *Tetrahedron Lett.* **27**, 2635–2638.
- Marinetti, A., Fischer, J. & Mathey, F. (1985). *J. Am. Chem. Soc.* **107**, 5001–5002.
- Molecular Structure Corporation (1993a). *MSCIAFX Diffractometer Control Software*. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
- Molecular Structure Corporation (1993b). *TEXSAN, Single Crystal Structure Analysis Software*, Version 1.6. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
- Orpen, A. G., Brammer, L., Allen, F. H., Kennard, O., Watson, D. G. & Taylor, R. (1989). *J. Chem. Soc. Dalton Trans.* pp. S1–S83.
- Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.
- Svara, J., Marinetti, A. & Mathey, F. (1986). *Organometallics*, **5**, 1161–1167.
- Walker, N. & Stuart, D. (1983). *Acta Cryst.* **A39**, 158–166.

Acta Cryst. (1997). **C53**, 305–309

Hexakis(dimethylformamide)bis(hexaphenylcyclohexasiloxanehexaolato)-hexacopper(II) Dimethylformamide Solvate

SERGEY V. LINDEMAN, OLGA I. SHCHEGOLIKHINA,
YULIA A. MOLODTSOVA AND ALEXANDR A. ZHIDANOV

*A. N. Nesmeyanov Institute of Organoelement Compounds,
Russian Academy of Sciences, 28 Vavilov St., Moscow
117813, Russia. E-mail: serg@xray.ineos.ac.ru*

(Received 28 May 1996; accepted 2 September 1996)

Abstract

The sandwich-like title complex, hexakis(dimethylformamide)- $1\kappa O, 2\kappa O, 3\kappa O, 4\kappa O, 5\kappa O, 6\kappa O$ -bis(2,4,6,8,10,12-hexaphenylsiloxane-2,4,6,8,10,12-hexaolato(6-))-

$1:2\kappa^2 O^1, 2:3\kappa^2 O^2, 3:4\kappa^2 O^3, 4:5\kappa^2 O^4, 5:6\kappa^2 O^5, 1:6\kappa^2 O^6$]-hexacopper(II) tetrakis(dimethylformamide) solvate, $[Cu_6(C_6H_7NO)_6\{(C_6H_5)_6O_{12}Si_6\}_2] \cdot 4C_3H_7NO$, is comprised of two regular crown-shaped macrocyclic hexadentate organosiloxanolate ligands chelating a flat Cu_6 hexagon, as in the ethanol-solvated analogue investigated previously. The title complex has a more distorted shape than the trigonal ethanol-solvated analogue, being slightly side-oblated, but still contains a large empty inner channel accessible by small molecules (the diameter of the free cross-section being about 2.5 Å). Each Cu^{II} ion has a square-pyramidal coordination comprised of four basal siloxanolate O atoms and an apical dimethylformamide (DMFA) molecule (coordinated through its carbonyl group). The average bond lengths are $Cu-O(Si)$ 1.964 (11) and $Cu-O(DMFA)$ 2.215 (10) Å. The structure contains four additional DMFA molecules per complex unit, linked by weak $C-H \cdots O$ hydrogen bonds. Unexpectedly, the $C=O$ bond length is longer (1.248 (10) and 1.255 (9) Å) in the uncoordinated DMFA molecules than in the coordinated molecules [1.214–1.227 (7) Å].

Comment

Recently, a new class of polynuclear metal complexes has been obtained (involving alkaline metals, Mn, Co, Ni and other bivalent transition metals, Cu and trivalent lanthanides) with polydentate regular macrocyclic organosiloxanolate $[-RSi(O^-)O^-]_n$ ligands ($n = 6, 8$ or 12), which are known as polymetalloorganosiloxanates (PMOS) and have been structurally characterized by Struchkov & Lindeman (1995). The size and structure of the organocyclosiloxanolate ligands formed seem to be dependent mainly on the coordination characteristics (size, charge and coordination number) of the metal. The Cu^{II} complexes are of special interest since at least two quite different structural types have been obtained using different reaction conditions, i.e. sandwich-like complexes, such as $[Cu_6\{(C_6H_5)_6O_{12}Si_6\}(\text{EtOH})_6]$ (1); Igonin *et al.*, 1992) or $K_2[K_2Cu_4\{(C_6H_5)_6O_{12}Si_6\}] \cdot 4n\text{-BuOH}$ (2); Igonin *et al.*, 1993), and globular-like complexes, such as $[Na_4Cu_4\{(C_6H_5)_6O_{12}Si_6\}] \cdot 8n\text{-BuOH}$ (3), and $K_4[Cu_4(O_2Si_2V_12)] \cdot 6n\text{-BuOH}$ (4); Igonin *et al.*, 1991).

Complex (1) is rather unique as a result of its large empty inner channel and square-pyramidal coordination of the Cu atoms (by four siloxanolate groups and an axial solvent hydroxy group); other complexes have more compact structures and contain square-planar-coordinated Cu ions without the additional coordination by solvent molecules. To obtain additional evidence on the stability and reproducibility of such an open structure [in related PMOS compounds based on bivalent d -group metals or trivalent lanthanides, the central cavity is occupied by an additional small anion (Cl^- , OH^- or O^{2-}) (Struchkov & Lindeman, 1995)], a close ana-