

ORIGINAL ARTICLE

Stimulatory response of neutrophils from periodontitis patients with periodontal pathogens

CG Restaño¹, A Chaparro², MA Valenzuela¹, AM Kettlun¹, R Vernal², A Silva³, J Puente¹, MP Jaque¹, R León², J Gamonal²

¹Departamento de Bioquímica y Biología Molecular, Facultad de Ciencias Químicas y Farmacéuticas, Universidad de Chile, Santiago, Chile; ²Departamento de Odontología Conservadora, Área de Periodoncia, Facultad de Odontología, Universidad de Chile, Santiago, Chile; ³Departamento de Inmunología, Centro de Investigaciones Biológicas (CIB), Madrid, Spain

OBJECTIVE: Neutrophils play a crucial role in the defense of invading bacteria by releasing biologically active molecules. The response of peripheral blood neutrophils was studied in periodontitis-affected patients and in healthy controls towards stimulation to *Porphyromonas gingivalis* (*Pg*) and *Actinobacillus actinomycetemcomitans* (*Aa*) extracts.

MATERIALS AND METHODS: Peripheral venous blood was drawn from 23 adult patients with moderate to advanced chronic periodontitis (probing depth ≥5 mm, attachment loss ≥3 mm), and 30 healthy volunteers. Neutrophil response followed by metalloproteinase-9 (MMP-9) and interleukin-8 (IL-8) secretion was assayed by zymography and enzyme-linked immunosorbent assay, respectively, on both whole blood and purified neutrophils. In addition to periodontal pathogen extracts, known stimulating agents were tested, such as *Escherichia coli*-lipopolysaccharide (LPS), phytohemagglutinin, and zymosan A.

RESULTS: Neutrophil response, expressed as a secretion ratio under stimulated and non-stimulated conditions, measured in whole blood, showed no differences between periodontitis and healthy controls. Instead, in purified neutrophils from patients, MMP-9 exhibited a significantly higher secretion ratio with LPS and Pg (1.5- to 2-fold), whereas IL-8 showed a larger increase in secretion ratio (3- to 7-fold) in the presence of Pg, Aa, LPS, and zymosan A.

CONCLUSION: Peripheral neutrophils of periodontitis-affected patients are more reactive as suggested by their significantly higher response toward periodontal pathogen extracts and other stimulating agents.

Oral Diseases (2007) 13, 474–481

Keywords: periodontitis; neutrophils; matrix metalloproteinases; MMP-9; gelatinase B; interleukin-8

Introduction

Periodontitis is a chronic inflammatory disease characterized by connective tissue and alveolar bone destruction, eventually leading to tooth loss. The two major pathogens in periodontal disease are the Gram-negative bacteria *Porphyromonas gingivalis* (*Pg*) and *Actinobacillus actinomycetemcomitans* (*Aa*) (Socransky *et al.*, 1998; Eley and Cox, 2003). These microorganisms induce the release of cytokines from fibroblasts and inflammatory cells (Yoshimura *et al.*, 1997; Johansson *et al.*, 2005). Neutrophils, macrophages, B and T lymphocyte cells accumulate in periodontal lesions and tissue destruction takes place (Assuma *et al.*, 1998; Gamonal *et al.*, 2001). The microorganisms seem to produce destruction directly through their endotoxins (LPS) and indirectly through activation of the host cells to produce a variety of biologically active substances, i.e. cytokines, arachidonic acid metabolites, and proteolytic enzymes (Trevani *et al.*, 2003). Pg stimulates the expression of interleukin-8 (IL-8) in peripheral neutrophils (Yoshimura *et al.*, 1997; Sugita *et al.*, 1998). IL-8 is a potent cytokine that leads to the induction of chemotaxis, migration, and activation of neutrophils (Kuhns *et al.*, 1998; Godaly *et al.*, 2001).

When exposed to bacteria-derived products or inflammatory mediators, neutrophils respond by chemotaxis, phagocytosis, and microbial killing through oxygen radical- and non-oxygen-dependent mechanisms (Kobayashi *et al.*, 2003; Moraes and Downey, 2003). During phagocytosis, neutrophils produce reactive oxygen species (ROS), together with the release of cytotoxic granule components (Kobayashi *et al.*, 2003). Gelatinase and specific granules are reservoirs of metalloproteinases (MMPs), while azurophil granules contain a neutral

Correspondence: J Gamonal, Facultad de Odontología, Universidad de Chile, Olivos 943, Independencia, Santiago, Chile. Tel: +562 978 1833, Fax: +562 978 1815, E-mail: jgamonal@uchile.cl
Received 6 February 2006; revised 29 April 2006, 14 June 2006; accepted 1 August 2006

protease called elastase (Faurschou and Borregaard, 2003; Uitto *et al*, 2003).

In response to stimulation, the gelatinase granules are more easily exocytosed followed by specific granules. The former contain MMP-9, or gelatinase B as well as leukolysin or MMP-25; these MMPs occur in inactive form, undergoing proteolytic activation following exocytosis. The latter, namely, specific granules, contain MMP-8 and some smaller amounts of gelatinase as well. On the other hand, azurophil granules undergo limited exocytosis in response to stimulation (Faurschou and Borregaard, 2003; Uitto *et al*, 2003).

Microbial infection is detected by the family of innate immune signaling receptors known as the Toll-like receptors (TLRs) present in neutrophils, that recognize conserved microbial structures called pathogen-associated molecular patterns (Akira, 2003; Kopp and Medzhitov, 2003). CR3 and Fc gamma Rs are receptors involved in the phagocytic process leading to bacterial destruction by the production of reactive oxygen intermediates and degranulation (Forsberg *et al*, 2001; Kobayashi *et al*, 2003).

Activation of neutrophils and the cleavage of cell matrix molecules by MMP-8, MMP-9 and elastase, generate peptide fragments that are chemotactic for monocytes, promoting the recruitment of inflammatory cells to the site of injury and thus releasing proinflammatory mediators that amplify the local inflammatory response (Faurschou and Borregaard, 2003; Figueredo *et al*, 2005). The role of neutrophils in the pathogenesis of periodontal disease is unclear. The protective functions of neutrophils have been studied in relation to defense mechanisms against periodontal pathogens (Kobayashi *et al*, 2003). However, this increased neutrophil activity could be responsible for the tissue destruction observed in periodontitis (Kantarci and Van Dyke, 2002; Fredriksson *et al*, 2003; Kantarci *et al*, 2003; Zekonis and Zekonis, 2004). Deguchi *et al* (1990) showed that LPS-stimulated human neutrophils get adhered to damaged periodontal tissue.

The sites of chronic periodontitis are rich sources of virulence agents, such as LPS, proteases from *Pg* and a leukotoxin produced by *Aa* that induces neutrophil degranulation (Geerts *et al*, 2002; Deas *et al*, 2003; Johansson *et al*, 2005). Reports on healthy subject neutrophils described an increased activity of MMP-9 and MMP-8, followed by stimulation with *Pg* or *Aa* (Ding *et al*, 1997; Claesson *et al*, 2002). In another report, the stimulation of neutrophils with LPS from *Escherichia coli* and *Aa* showed the release of significantly greater amounts of IL-8 (Yoshimura *et al*, 1997).

The effects of periodontal pathogen extracts and some known stimulating agents were studied on peripheral whole blood and on purified neutrophils of both control and periodontitis-affected patients. The effects were followed by (a) MMP-9 release from gelatinase granules required during neutrophil extravasations (Faurschou and Borregaard, 2003) and (b) IL-8 release, one of the most abundant cytokines produced by neutrophils, reported in high amounts in the plasma of rapidly progressive periodontitis patients, which suggests a

systemic effect of this periodontal disease (Gainet *et al*, 1998).

Materials and methods

The study group comprised 23 adult patients with moderate to advanced chronic periodontitis selected from the Center of Diagnostic and Treatment of Northern Metropolitan Health Services. The criteria for entry were a minimum of 14 natural teeth, excluding third molars, and including at least 10 posterior teeth. Patients with chronic periodontitis had moderate to advanced periodontitis (at least five to six teeth had sites with probing depth (PD) ≥ 5 mm and with attachment loss ≥ 3 mm and extensive bone loss in radiography), and had received no periodontal treatment at the time of examination. Subjects did not suffer from systemic illness and had received no antibiotics or non-steroid anti-inflammatory therapy in the 6-month period prior to the study. The control group comprised 30 subjects with no clinical signs of periodontal [absence of clinical attachment loss (CAL) or increased probing pocket depths] and systemic diseases. Prior to the study, all the subjects received supragingival prophylaxis to remove gross calculus and allow PD. The protocol was clearly explained to all patients and controls, and Institutional Review Board-approved informed consents were obtained. The protocol stated that, within 2 weeks of the detection of disease activity, all the patients would be provided with periodontal treatment. Periodontal therapy consisted of scaling, root planing, and instruction in oral hygiene.

Clinical measurements

Clinical parameters were evaluated by a single calibrated investigator in all teeth (excluding third molars) and included PD, CAL and dichotomous measurements of supragingival plaque accumulation or plaque index (PI) and bleeding on probing to the base of the crevice (BOP). Six sites were examined for each tooth: mesiobuccal, buccal, distobuccal, distolingual, lingual, and mesiolingual positions. The Florida Disk Probe was used for relative attachment level recordings and the Florida Probing Depth for PD recording (Florida Probe Corporation, Gainesville, FL, USA).

Whole blood/purified neutrophil stimulation assay

Whole blood from healthy human volunteers and periodontitis-affected patients was collected on the day of the assay, and was drawn into heparinized syringes (10 U ml^{-1} heparin sulfate; Liquemine; Roche Pharma, Reinach, Switzerland). Neutrophils were purified from whole blood samples as described previously (Rose *et al*, 1992), using a Hystopaque 1077 gradient (Sigma Chemical Co, St Louis, MO, USA), and were diluted to 10^6 cells ml^{-1} in RPMI 1640 with 10% bovine fetal serum (Gibco, Basel, Switzerland), 2 mm glutamine and $5\text{ }\mu\text{g ml}^{-1}$ streptomycin. Before assay, 1 ml of whole blood was diluted 1:4 with RPMI 1640 containing 10% bovine fetal serum or 1 ml purified neutrophils was incubated at 37°C for 1 h. Assays were started by the

addition of up to 10% of the assay volume of the stimulating agents previously diluted in phosphate-buffered saline (PBS; 0.137 M NaCl, 12 mM Na₂HPO₄, 2 mM KH₂PO₄ adjusted to pH 7.4); assays were then maintained at 37°C for 1 h. Different concentrations of stimulatory agents were initially tested in order to choose appropriate conditions, which include both neutrophil viability and stimulatory effect. Chosen concentrations were in the range of those used by other authors (Pugin *et al*, 1999; Hayashi *et al*, 2003). *Escherichia coli* LPS (O111:B4) (tests ranging from 0.05 to 0.8 µg ml⁻¹) was used at a final concentration of 0.8 µg ml⁻¹, phytohemagglutinin (PHA; 1.0–5.0 µg ml⁻¹) at 2.0 µg ml⁻¹ and zymosan A (5.0–100 µg ml⁻¹) at 100 µg ml⁻¹ (all were from Sigma Chemical Co). Pg and Aa extracts (prepared in our laboratory) were used at 0.36 and 0.07 µg ml⁻¹, respectively. Tested dilutions of bacterial extracts were in the range of 0.09–0.360 µg ml⁻¹ for Pg and 0.02–0.07 µg ml⁻¹ for Aa. Concentrations used were the maximum allowed for the viability of the neutrophils. Addition of PBS alone was used as a non-stimulated condition. After incubation cell suspensions were spun at 2000 g for 10 min, and the supernatant of blood/neutrophil samples was stored at -20°C until use.

Preparation of bacterial extracts

Actinobacillus actinomycetemcomitans (Y4) was grown in enriched tryptic soy agar and tryptone soy broth containing Na₂CO₃, L-cysteine, hemin and vitamin K. *Porphyromonas gingivalis* (A7A1-28) was cultured in enriched Todd Hewitt broth and butylated hydroxyanisole containing the same supplements. Anaerobic bacteria were cultured under strictly anaerobic conditions according to Loomer *et al* (1995). After culturing, the bacteria were killed by 10 min boiling, then washed thrice with PBS, and stored at -20°C. Boiling does not destroy the LPS component. Amounts of bacterial extracts were expressed according to their protein content measured by the bicinchoninic acid method (Pierce Biotechnology, Rockford, IL, USA) following the manufacturer's instructions.

Gelatin zymography

Gelatin zymography was performed according to Pozo *et al* (2005). Whole blood or purified peripheral neutrophils were resuspended in non-reducing sodium dodecyl sulfate (SDS) sample buffer (0.4 M Tris-HCl, pH 6.8, 5% SDS, 20% glycerol, 0.05% bromophenol blue); boiling was not permitted to allow further enzyme reactivation, and loaded onto a 10% SDS-polyacrylamide gel, including 1 mg ml⁻¹ of gelatin (Merck, Darmstadt, Germany). In order to work in a linear zone of gelatinase activity, adequate conditions for sample dilution and volume were tested. The best conditions were: sample dilution 1:20 in upper SDS-PAGE buffer (Tris-HCl, pH 6.8), and a volume of 5 µl of the diluted samples previously treated with the non-reducing Laemmli buffer (Pozo *et al*, 2005). After the electrophoretic run, SDS was removed from the gel before incubation for 20 h at 37°C with the renaturating buffer (50 mM Tris-HCl, 5 mM CaCl₂, 0.02% NaN₃, pH

7.5). After Coomassie brilliant blue R-250 staining, and destaining (7% acetic acid) (Ejeil *et al*, 2003), proteolysis areas were estimated by densitometric analysis using the 'Uni-Scan-it Gel-Digitizing System' software (Silk Scientific Corporation, Orem, UT, USA). The arbitrary density units (in pixels) of the stimulating conditions were referred to as the basal or non-stimulated condition. Through zymography, both MMP and proMMP (non-enzymatically active form) can be detected because the presence of SDS in the gel and its later removal through Triton X-100 exchange activates zymogen without changing the molecular weight (Pugin *et al*, 1999).

Determination of IL-8 secretion upon stimulation

Interleukin-8 was measured in 50–100 µl portions of an incubation medium with an enzyme-linked immunosorbent assay kit (BioSource Europe S. A., Nivelles, Belgium) following the manufacturer's instructions.

Statistical analysis

When necessary (IL-8 results), data from patient and control groups were normalized using the logarithmic function. The differences between values of both groups were assessed by a two-tailed Student's *t*-test, using the statistic software Systat 11.0. Statistical significance was set at a value of *P* < 0.05.

Results

Clinical characteristics of all the 23 patients under study are shown in Table 1. We used whole blood from 13 patients and purified neutrophils from the remaining 10. The mean values ± s.d. of PI, PD, BOP, and CAL are significantly higher than those of healthy individuals who exhibited PD < 3 mm, no attachment loss, and no clinical inflammation.

A representative gelatin zymogram of purified neutrophil secretion from a periodontitis-affected patient stimulated with LPS, PHA, Pg, Aa, and zymosan A, together with the non-stimulated condition (lane 1), and the presence of a positive standard of proMMP-9 is shown in Figure 1a, where gelatinolytic zones are visualized as white bands on a dark background. In Figure 1b we compare the effect of the stimulating agents on both a patient (lanes 1–5) and a healthy

Table 1 Clinical characteristics of periodontitis and healthy groups (mean ± s.d.)

	Periodontitis group (n = 23)	Healthy group (n = 30)
Age (years)	36.69 ± 9.49	38.42 ± 8.42
Women (n)	17	21
Mean probing depth (mm)	3.35 ± 0.55*	1.65 ± 0.48*
Mean attachment level (mm)	3.30 ± 0.57*	0.26 ± 0.26*
% sites with plaque	61.34**	25.60**
% sites with bleeding on probing	51.87*	1.42*

*P < 0.0001; **P < 0.001.




Figure 1 Zymogram of various forms of metalloproteinase-9 (MMP-9) released from active neutrophils. **(a)** Neutrophil sample from a periodontitis-affected patient. Lane 1: neutrophils without stimulation; lane 2: neutrophils stimulated with lipopolysaccharide (LPS; $0.8 \mu\text{g ml}^{-1}$); lane 3: neutrophils stimulated with phytohemagglutinin ($2 \mu\text{g ml}^{-1}$); lanes 4 and 5: neutrophils in the presence of *Porphyromonas gingivalis* (*Pg*; $0.36 \mu\text{g ml}^{-1}$) or *Actinobacillus actinomycetemcomitans* (*Aa*; $0.07 \mu\text{g ml}^{-1}$), respectively; lane 6: neutrophils stimulated with zymosan A ($100 \mu\text{g ml}^{-1}$); and lane 7: a positive proMMP-9 control. **(b)** Neutrophil sample from a patient (lanes 1–5). Lane 1: without stimulation; lane 2: stimulated with LPS ($0.8 \mu\text{g ml}^{-1}$); lane 3: stimulated with *Pg* ($0.36 \mu\text{g ml}^{-1}$); lane 4: stimulated with *Aa* ($0.07 \mu\text{g ml}^{-1}$); lane 5: stimulated with zymosan A ($100 \mu\text{g ml}^{-1}$). Neutrophil sample from a healthy subject (lanes 6–9) – lane 6: without stimulation; lane 7: stimulated with LPS ($0.8 \mu\text{g ml}^{-1}$); lane 8: stimulated with *Pg* ($0.36 \mu\text{g ml}^{-1}$); lane 9: stimulated with zymosan A ($100 \mu\text{g ml}^{-1}$). MMP-9 dimer (175 kDa), MMP-9-lipocalin (130 kDa), and proMMP-9 (92 kDa)

control sample (lanes 6–9). The 92 kDa band corresponds to proMMP-9, a 130-kDa band to the reported neutrophil-MMP-9-lipocalin complex (Westerlund *et al*, 1996), and a gelatinolytic band over 175 kDa, described as the dimeric form of an MMP-9 (Olson *et al*, 2000). No active form for MMP-9 was detected because of the absence of gelatinolytic activity at 84 kDa. Whole blood samples after incubation showed a similar pattern.

Measurements of the proMMP-9 band gave more reproducible results than when also including the other two MMP-9-related bands (lipocalin complex and dimer). The stimulating agents did not produce a significant effect on whole blood in both healthy and patient groups. Results with purified neutrophils from patients with either moderate or advanced periodontitis were not statistically different, so all the results were aggregated. To compare the effect of these agents on the secretory response, data were expressed as the secretion ratio of the gelatinolytic area under stimulated and non-stimulated conditions (S-proMMP-9/N-proMMP-9). Figure 2 shows that purified neutrophils from patients display significantly higher levels of proMMP-9 than those of controls ($P < 0.05$), between 1.5- and 2-fold, using LPS or Pg treatments. The dispersion of the results with zymosan A hinders significant statistical differences (Figure 2).

Similar to MMP-9 results, IL-8 release from the healthy group samples (whole blood and purified neutrophil samples) was not increased by the stimulating agents. IL-8 secretion promoted by these agents in whole blood samples did not show differences between both groups (data not shown). Instead, in the purified neutrophils of periodontitis-affected patients, LPS, Pg, *Aa*, and zymosan A triggered a significant response to




Figure 2 Comparison of the effect of stimulating agents on metalloproteinase-9 (MMP-9) secretion from purified neutrophils of periodontitis-patients and controls. Results are expressed as the secretion ratio under stimulated and non-stimulated conditions (S-MMP9/N-MMP9). Stimulant concentrations are similar to those mentioned in Figure 1. Data correspond to mean values \pm standard error of the mean (s.e.m.). * $P < 0.05$ between patients and healthy controls

IL-8 release when compared with the non-stimulated condition (S-IL-8/N-IL-8) (Figure 3a).

Unlike the healthy group, the level of secreted IL-8 (expressed as pg ml^{-1}) for the patient group was significantly lower either in the presence or absence of stimulating agents (Figure 3b). These results differ from those obtained for proMMP-9 secretion from gelatinase granules, where no differences were observed in proMMP-9 levels for both groups in the non-stimulated condition (incubation with PBS).

Discussion

The enhancement of proMMP-9 (1.5- to 2-fold) and of IL-8 (3- to 7-fold) secretion under the stimulated




Figure 3 Comparison of the effect of stimulating agents on interleukin-8 (IL-8) secretion from purified neutrophils of periodontitis patients and controls. (a) Ratio of IL-8 secretion in stimulated/non-stimulated conditions (S-IL8/N-IL8). (b) Levels of IL-8 secreted from neutrophils. Data are expressed as mean values \pm s.e.m. Stimulant concentrations are similar to those mentioned in Figure 1. * $P < 0.05$ between patients and healthy controls

condition by periodontal pathogen extracts and other proinflammatory mediators suggests that neutrophils are more reactive in periodontitis-affected patients than in controls. Purified neutrophil samples proved to be more adequate than whole blood samples for neutrophil stimulation as they helped in detecting differences between the patients and controls. This hyper-reactivity may be secondary to the inflammatory reaction occurring in the gingival tissue during periodontal disease; however, such hyper-reactivity could also be the result of constitutive differences in neutrophils from controls and periodontitis patients. This latter situation could be of relevance in the pathogenesis of the periodontal disease (Fredriksson et al, 2003; Gustafsson et al, 2006); hence it seems important to determine whether the hyper-reactivity followed by proMMP-9 and IL-8 release is reverted after non-surgical periodontal treatment. Gainet et al (1998) reported normalization of IL-8 concentration in plasma from progressive periodontitis patients after treatment, and we observed a reduction of

MMP-activity to control values in gingival crevicular fluid after a 3-month treatment (Pozo et al, 2005).

The significantly reduced levels of IL-8 secretion in periodontitis-affected patients could be attributed to the known endotoxin tolerance phenomenon (West and Heagy, 2002). Marie et al (1998) observed a significantly decreased release of IL-8 in neutrophils from patients with infectious and non-infectious systemic inflammatory response syndrome; the observed decrease in periodontitis patients did not, however, reflect tolerance because neutrophils were not desensitized to a second challenge. The downregulation of the synthesis and release of excessive proinflammatory cytokines after infection could be an adaptive mechanism to avoid deleterious effects for the host. Plasma samples from periodontitis-affected patients showed significantly reduced systemic levels of IL-1 β compared with healthy controls (Johansson et al, 2005).

Different forms of MMP-9 (proMMP-9, MMP-9-lipocalin complex, and a dimeric form) are released after 60 min incubation either from blood or purified neutrophils. This gelatinase activity can be attributed to neutrophils and monocytes/macrophages. A secretion within 60 min cannot be the result of *de novo* protein synthesis by monocytes/macrophages, requiring at least 7 h of incubation, hence this secretion should come from neutrophil granules (Pugin et al, 1999). In addition, the presence of the 130 kD proMMP-9-lipocalin complex in non-reducing SDS-PAGE zymography indicates a PMN origin for this gelatinase (Olson et al, 2000; Faurschou and Borregaard, 2003). A report on interaction between neutrophils and whole cells of *P. gingivalis* and other periodontopathogens showed that azurophil, specific, and gelatinase granules are triggered during this interaction (Ding et al, 1997).

Elevation of systemic inflammatory mediators including C-reactive protein, IL-6, neutrophils, and bacterial endotoxins occurs in patients with periodontitis (Loos et al, 2000; Geerts et al, 2002; Deas et al, 2003; Johansson et al, 2005). This suggests that the infected periodontum serves as a source of circulating endotoxin. Low levels of endotoxin or bacterial products derived from periodontitis may therefore, contribute to immune activation and could be associated with general systemic conditions such as miscarriage or cardiovascular disease (Farell et al, 2006; Spahr et al, 2006; Taylor et al, 2006). In this disease the persistent accumulation of bacteria in the periodontal pocket maintains the recruitment of inflammatory cells and a prolonged and sustained release of inflammatory mediators (Eley and Cox, 2003; Sela et al, 2003; Pozo et al, 2005). These products can amplify the local inflammatory response, further promoting leukocyte and platelet recruitment and thus perpetuating the inflammation. Additionally, they can lead to elevated amounts of these mediators into the bloodstream.

As the factors released by activated neutrophils are potentially toxic to host tissues, neutrophils are subsequently removed by the process of apoptosis (Moraes and Downey, 2003; Scheel-Toellner et al, 2004). This mechanism prevents the secretion of pro-inflammatory

mediators into the tissues avoiding the inflammatory response. Thus, modulation of the activation status of the neutrophils is essential in determining the balance between the immune defense and host injury (Kobayashi *et al*, 2003; Moraes and Downey, 2003). MMPs and other enzymes released from activated neutrophils modify constituents of the extracellular matrix (ECM) through which cells migrate to reach target organs (Vaday and Lider, 2000). Growing evidence suggests that mediators such as MMPs and cytokines secreted from immune cells invading into tissues through ECM induce distinctive cellular responses, being probably part of a regulatory mechanism of perpetuation or arrest of inflammation (Vaday and Lider, 2000).

Neutrophils recognize bacterial structure mainly via the TLRs (Akira, 2003). TLRs involved in the recognition of pathogen-associated molecular patterns, activate intracellular signaling and transcription of protein essential for the induction of the inflammatory response and elimination of invading pathogens (Sabroe *et al*, 2003; Netea *et al*, 2004; Parker *et al*, 2005). It is quite probable that similar to *E. coli* LPS, the stimulating effects of *Pg*, *Aa*, and zymosan depend on their interaction with TLR-2 and/or TLR-4 (Armant and Fenton, 2002; Akira, 2003; Kopp and Medzhitov, 2003; Hashimoto *et al*, 2004). The response to zymosan A, a *Saccharomyces cerevisiae* cell wall particle, may occur through both the TLR-2 receptor (Hayashi *et al*, 2003), and complement C3bi (CR3) receptor for the opsonized particle of zymosan. TLRs detect structurally different components in pathogens. In *Pg* both receptors are involved: lipid A, a component of LPS is the main inmunomodulator acting through TLR-4, while the lipoprotein activates cells through a TLR-2-dependent pathway (Akira, 2003; Hashimoto *et al*, 2004). In chronic periodontitis, the gingival tissue is infiltrated by large numbers of TLR-2 and TLR-4 cells (Muthukuru *et al*, 2005). However, real-time PCR analysis showed a significant downregulation of TLR-2 and TLR-4 mRNA suggesting that the sustained exposure to bacterial structure modulates local immune response (Muthukuru *et al*, 2005). However, we have to bear in mind that in the boiling process used during the bacterial extracts preparation in this study, some important pathogenic agents, such as leukotoxin produced by *A. actinomycetemcomitans* are lost. This toxin is recognized by the lymphocyte function-associated antigen 1 receptor of leukocytes and induces the lysis of neutrophils and monocytes (Baehni *et al*, 1979; Yamaguchi *et al*, 2004).

Attempts to find the differential expression of either selected adhesion molecules or secretion of IL-4 and IL-6 in peripheral blood leukocytes in patients with various forms of periodontal diseases have been unsuccessful (Pietruska *et al*, 2004, 2005). Previous reports have studied only the stimulation of neutrophils with non-opsonized *E. coli*. However, *E. coli* LPS and opsonized *Staphylococcus aureus* demonstrated a significant increase in reactive oxygen metabolites release in periodontitis-affected patients with respect to healthy controls (Fredriksson *et al*, 2003; Zekonis and Zekonis, 2004;

Sadzevieciene *et al*, 2005; Gustafsson *et al*, 2006). In addition, elastase release after stimulation with *S. aureus* has been reported to be significantly higher in patients in relation to the controls (Figueroedo *et al*, 1999). Hence, the relevance of the current study lies in the demonstration of the hyper-reactivity of peripheral neutrophils in patients with advanced or moderate periodontitis toward periodontal pathogens such as *P. gingivalis* and *A. actinomycetemcomitans* based on MMP-9 and IL-8 release.

Acknowledgements

This study was supported by the project grant University of Chile-CSIC and DENTAID® Laboratory. We are gratefully indebted to Dr Christopher I. Pogson and Mr Claudio Telha for their critical reading of this manuscript.

References

- Akira S (2003). Mammalian Toll-like receptors. *Curr Opin Immunol* **15**: 5–11.
- Armant MA, Fenton MJ (2002). Toll-like receptors: a family of pattern-recognition receptors in mammals. *Genome Biol* **3**: 3011.1–3011.6.
- Assuma R, Oates T, Cochran D, Amar S, Graves DT (1998). IL-1 and TNF antagonists inhibit the inflammatory response and bone loss in experimental periodontitis. *J Immunol* **160**: 403–409.
- Baehni P, Tsai C, McArthur WP, Hammond BF, Taichman NS (1979). Interaction of inflammatory cells and oral microorganisms. VIII. Detection of leukotoxic activity of a plaque-derived gram-negative microorganism. *Infect Immun* **24**: 233–243.
- Claesson R, Jojansson A, Belibasakis G, Hänström L, Kalfas S (2002). Release and activation of matrix metalloproteinase 8 from human neutrophils triggered by the leukotoxin of *Actinobacillus actinomycetemcomitans*. *J Periodontal Res* **37**: 353–359.
- Deas DE, Mackey SA, McDonell HT (2003). Systemic disease and periodontitis: manifestations of neutrophil dysfunction. *Periodontol 2000* **32**: 82–104.
- Deguchi S, Hori T, Creamer H, Gabler W (1990). Neutrophil-mediated damage to human periodontal ligament-derived fibroblasts: role of lipopolysaccharide. *J Periodontal Res* **25**: 293–299.
- Ding Y, Haapasalo M, Kerosuo E, Lounatmaa K, Kotiranta A, Sorsa T (1997). Release and activation of human neutrophil matrix metallo- and serine proteinases during phagocytosis of *Fusobacterium nucleatum*, *Porphyromonas gingivalis* and *Treponema denticola*. *J Clin Periodontol* **24**: 237–248.
- Ejeil AL, Igondjo-Tchen S, Ghomrasseni S, Pellat B, Godeau G, Gogly B (2003). Expression of matrix metalloproteinases (MMPs) and tissue inhibitors of metalloproteinases (TIMPs) in healthy and diseased human gingival. *J Periodontol* **74**: 188–195.
- Eley B, Cox SW (2003). Proteolytic and hydrolytic enzymes from putative periodontal pathogens: characterization, molecular genetics, effects on host defenses and tissues and detection in gingival crevice fluid. *Periodontol 2000* **31**: 105–124.
- Farell S, Ide M, Wilson RF (2006). The relationship between maternal periodontitis, adverse pregnancy outcome and miscarriage in never smokers. *J Clin Periodontol* **33**: 115–120.

- Faurschou M, Borregaard N (2003). Neutrophil granules and secretory vesicles in inflammation. *Microbes Infect* **5**: 1317–1327.
- Figueredo CM, Gustafsson A, Asman B, Bergström K (1999). Increased release of elastase from *in vitro* activated peripheral neutrophils in patients with adult periodontitis. *J Clin Periodontol* **26**: 206–211.
- Figueredo CM, Fischer RG, Gustafsson A (2005). Aberrant neutrophil reactions in periodontitis. *J Periodontol* **76**: 951–955.
- Forsberg M, Lofgren R, Zheng L, Stendahl O (2001). Tumour necrosis factor-alpha potentiates CR3-induced respiratory burst by activating p38 MAP kinase in human neutrophils. *Immunology* **104**: 466–472.
- Fredriksson MI, Gustafsson AK, Bergström KG, Asman BE (2003). Constitutionally hyperreactive neutrophils in periodontitis. *J Periodontol* **74**: 219–224.
- Gainet J, Chollet-Martin S, Brion M, Hakim J, Gougerot-Pocidalo MA, Elbim C (1998). Interleukin-8 production by polymorphonuclear neutrophils in patients with rapidly progressive periodontitis: an amplifying loop of polymorphonuclear neutrophil activation. *Lab Invest* **78**: 755–762.
- Gamonal J, Acevedo A, Bascones A, Jorge O, Silva A (2001). Characterization of cellular infiltrate, detection of chemokine receptor CCR5 and interleukin-8 and RANTES chemokines in adult periodontitis. *J Periodontal Res* **36**: 194–203.
- Geerts SO, Nys M, De MP et al. (2002). Systemic release of endotoxins induced by gentle mastication: association with periodontitis severity. *J Periodontol* **73**: 73–78.
- Godaly G, Bergsten G, Hang L et al. (2001). Neutrophil recruitment, chemokine receptors, and resistance to mucosal infection. *J Leukoc Biol* **69**: 899–906.
- Gustafsson A, Ito H, Asman B, Bergström K (2006). Hyper-reactive mononuclear cells and neutrophils in chronic periodontitis. *J Clin Periodontol* **33**: 126–129.
- Hashimoto M, Asai Y, Ogawa T (2004). Separation and structural analysis of lipoprotein in a lipopolysaccharide preparation from *Porphyromonas gingivalis*. *Int Immunol* **16**: 1431–1437.
- Hayashi F, Means TK, Luster AD (2003). Toll-like receptors stimulate human neutrophil function. *Blood* **102**: 2660–2669.
- Johansson A, Buhlin K, Koski R, Gustafsson A (2005). The immunoreactivity of systemic antibodies to *Actinobacillus actinomycetemcomitans* and *Porphyromonas gingivalis* in adult periodontitis. *Eur J Oral Sci* **113**: 197–202.
- Kantarci A, Van Dyke TE (2002). Neutrophil-mediated host response to *Porphyromonas gingivalis*. *J Int Acad Periodontol* **4**: 119–125.
- Kantarci A, Oyaizu K, Van Dyke TE (2003). Neutrophil-mediated tissue injury in periodontal disease pathogenesis. *J Periodontol* **74**: 66–75.
- Kobayashi SD, Voyich JM, De Leo FR (2003). Regulation of the neutrophil-mediated inflammatory response to infection. *Microbes Infect* **5**: 1337–1344.
- Kopp E, Medzhitov R (2003). Recognition of microbial infection by Toll-like receptors. *Curr Opin Immunol* **15**: 396–401.
- Kuhns DB, Young HA, Gallin EK, Gallin JI (1998). Ca²⁺-dependent production and release of IL-8 in human neutrophils. *J Immunol* **161**: 4332–4339.
- Loomer PM, Ellen RP, Tenenbaum HC (1995). Characterization of inhibitory effects of suspected periodontopathogens on osteogenesis *in vitro*. *Infect Immun* **63**: 3287–3296.
- Loos BG, Craandijk J, Hoek FJ, Wertheim-van Dillen PM, van der Velven U (2000). Elevation of systemic markers related to cardiovascular diseases in the peripheral blood of periodontitis patients. *J Periodontol* **71**: 1528–1534.
- Marie C, Muret J, Fitting C, Losser MR, Payen D, Cavaillon JM (1998). Reduced *ex vivo* interleukin-8 production by neutrophils in septic and nonseptic systemic inflammatory response syndrome. *Blood* **91**: 3439–3446.
- Moraes T, Downey GP (2003). Neutrophil cell signaling in infection: role of phosphatidylinositide 3-kinase. *Microbes Infect* **5**: 1293–1298.
- Muthukuru M, Jotwani R, Cutler CW (2005). Oral mucosal endotoxin tolerance induction in chronic periodontitis. *Infect Immun* **73**: 687–694.
- Netea MG, van der Graaf C, van der Meer JW, Kullberg BJ (2004). Toll-like receptors and the host defense against microbial pathogens: bringing specificity to the innate-immune system. *J Leukoc Biol* **75**: 749–755.
- Olson MW, Bernardo M, Pietila M et al. (2000). Characterization of the monomeric and dimeric forms of latent and active matrix metalloproteinase-9. *J Biol Chem* **275**: 2661–2668.
- Parker LC, Whyte MK, Dower SK, Sabroe I (2005). The expression and roles of Toll-like receptors in the biology of the human neutrophil. *J Leukoc Biol* **77**: 886–892.
- Pietruska M, Zak J, Wysocka J et al. (2004). Evaluation of selected peripheral blood leukocyte functions in patients with various forms of periodontal disease. *Arch Immunol Ther Exp* **52**: 208–212.
- Pietruska M, Zak J, Pietruski J, Wysocka J (2005). Expressions of selected adhesion molecules on peripheral blood leukocytes in patients with aggressive periodontitis. *Arch Immunol Ther Exp* **53**: 266–271.
- Pozo P, Valenzuela MA, Melej C et al. (2005). Longitudinal analysis of metalloproteinases, tissue inhibitors of metalloproteinases and clinical parameters in gingival crevicular fluid from periodontitis-affected patients. *J Periodontal Res* **40**: 199–207.
- Pugin J, Widmer MC, Kossodo S, Liang CM, Preas HL, Suffredini AF (1999). Human neutrophils secrete gelatinase B *in vitro* and *in vivo* in response to endotoxin and proinflammatory mediators. *Am J Respir Cell Mol Biol* **20**: 458–464.
- Rose N, De Macario EC, Fohey J, Friedman H, Penn G (1992). *Manual of clinical laboratory immunology*, 4th edn. Washington, DC: American Society for Microbiology.
- Sabroe I, Prince LR, Jones EC et al. (2003). Selective roles for Toll-like receptor (TLR)2 and TLR4 in the regulation of neutrophil activation and life span. *J Immunol* **170**: 5268–5275.
- Sadzheviciene R, Zekonis J, Zekonis G (2005). Generation of superoxide anion by peripheral blood leukocytes in periodontitis patients with type 1 diabetes mellitus. *Medicina (Kaunas)* **41**: 477–481.
- Scheel-Toellner D, Wang KQ, Webb PR et al. (2004). Early events in spontaneous neutrophil apoptosis. *Biochem Soc Trans* **32**: 461–464.
- Sela MN, Kohavi D, Krausz E, Steinberg D, Rosen G (2003). Enzymatic degradation of collagen-guided tissue regeneration membranes by periodontal bacteria. *Clin Oral Implants Res* **14**: 263–268.
- Socransky SS, Haffajee AD, Cugini MA, Smith C, Kent RL Jr (1998). Microbial complexes in subgingival plaque. *J Clin Periodontol* **25**: 134–144.

- Spahr A, Klein E, Khuseyinova N et al. (2006). Periodontal infections and coronary heart disease: role of periodontal bacteria and importance of total pathogen burden in the coronary event and periodontal disease (CORODON) study. *Arch Intern Med* **166**: 554–559.
- Sugita N, Kimura A, Matsuki Y, Yamamoto T, Yoshie H, Hara K (1998). Activation of transcription agents and IL-8 expression in neutrophils stimulated with lipopolysaccharide from *Porphyromonas gingivalis*. *Inflammation* **22**: 253–266.
- Taylor BA, Tofler GH, Carey HM et al. (2006). Full-mouth tooth extraction lowers systemic inflammatory and thrombotic markers of cardiovascular risk. *J Dent Res* **85**: 74–78.
- Trevani AS, Chorny A, Salamone G et al. (2003). Bacterial DNA activates human neutrophils by a CpG-independent pathway. *Eur J Immunol* **33**: 3164–3174.
- Uitto VK, Overall CM, Culloch C (2003). Proteolytic host cell enzymes in gingival crevice fluid. *Periodontology* **31**: 77–104.
- Vaday GG, Lider O (2000). Extracellular matrix moieties, cytokines, and enzymes: dynamic effects on immune cell behavior and inflammation. *J Leukoc Biol* **67**: 149–159.
- West MA, Heagy W (2002). Endotoxin tolerance: a review. *Crit Care Med* **30**: S64–S73.
- Westerlund U, Ingman T, Lukinmaa PL et al. (1996). Human neutrophil gelatinase and associated lipocalin in adult and localized juvenile periodontitis. *J Dental Res* **75**: 1553–1563.
- Yamaguchi N, Kubo C, Masuhiro Y, Lally ET, Koga T, Hanazawa S (2004). Tumor necrosis factor alpha enhances *Actinobacillus actinomycetemcomitans* leukotoxin-induced HL-60 cell apoptosis by stimulating lymphocyte function-associated antigen 1 expression. *Infect Immun* **72**: 269–276.
- Yoshimura A, Hara Y, Kaneko T, Kato I (1997). Secretion of IL-1 β , TNF- α , IL-8 and IL-1ra by human polymorphonuclear leukocytes in response to lipopolysaccharides from periodontopathic bacteria. *J Periodontal Res* **32**: 279–286.
- Zekonis G, Zekonis J (2004). Effect of bacterial stimulants on release of reactive oxygen metabolites from peripheral blood neutrophils in periodontitis. *Medicina (Kaunas)* **40**: 260–264.