

Xestospongine B, a competitive inhibitor of IP₃-mediated Ca²⁺ signalling in cultured rat myotubes, isolated myonuclei, and neuroblastoma (NG108-15) cells

Enrique Jaimovich^a, César Mattei^b, José Luis Liberona^a, Cesar Cardenas^a, Manuel Estrada^a, Julien Barbier^b, Cécile Debitus^c, Dominique Laurent^d, Jordi Molgó^{b,*}

^a Instituto de Ciencias Biomédicas, Facultad de Medicina, Universidad de Chile, casilla 70005, Santiago 7, Chile

^b Laboratoire de Neurobiologie Cellulaire et Moléculaire, U.P.R. 9040, Institut Fédératif de Neurobiologie Alfred Fessard, C.N.R.S., 1 avenue de la Terrasse, 91198 Gif-sur-Yvette Cedex, France

^c I.R.D.-C.R.S.N.-Pierre Fabre, 3 Rue Ariane, 31527 Ramonville Saint Agne, France

^d Laboratoire des Substances Naturelles, I.R.D., 98848-Noumea, New Caledonia

Abstract Xestospongine B, a macrocyclic bis-1-oxaquinolizidine alkaloid extracted from the marine sponge *Xestospongia exigua*, was highly purified and tested for its ability to block inositol 1,4,5-trisphosphate (IP₃)-induced Ca²⁺ release. In a concentration-dependent manner xestospongine B displaced [³H]IP₃ from both rat cerebellar membranes and rat skeletal myotube homogenates with an EC₅₀ of 44.6 ± 1.1 μM and 27.4 ± 1.1 μM, respectively. Xestospongine B, depending on the dose, suppressed bradykinin-induced Ca²⁺ signals in neuroblastoma (NG108-15) cells, and also selectively blocked the slow intracellular Ca²⁺ signal induced by membrane depolarization with high external K⁺ (47 mM) in rat skeletal myotubes. This slow Ca²⁺ signal is unrelated to muscle contraction, and involves IP₃ receptors. In highly purified isolated nuclei from rat skeletal myotubes, Xestospongine B reduced, or suppressed IP₃-induced Ca²⁺ oscillations with an EC₅₀ = 18.9 ± 1.35 μM. In rat myotubes exposed to a Ca²⁺-free medium, Xestospongine B neither depleted sarcoplasmic reticulum Ca²⁺ stores, nor modified thapsigargin action and did not affect capacitative Ca²⁺ entry after thapsigargin-induced depletion of Ca²⁺ stores. Ca²⁺-ATPase activity measured in skeletal myotube homogenates remained unaffected by Xestospongine B. It is concluded that xestospongine B is an effective cell-permeant, competitive inhibitor of IP₃ receptors in cultured rat myotubes, isolated myonuclei, and neuroblastoma (NG108-15) cells.

Keywords: Xestospongine B; Inositol 1,4,5-trisphosphate; Intracellular Ca²⁺ signals; Bradykinin; Neuroblastoma (NG108-15) cell; Rat skeletal myotubes; Cerebellar membranes; Isolated myonuclei; Calcium ATPase

1. Introduction

Xestospongines are a group of macrocyclic bis-1-oxaquinolizidines alkaloids isolated from marine sponges [1–6]. These

compounds have been described to be potent membrane-permeable inhibitors of inositol 1,4,5-trisphosphate (IP₃)-mediated Ca²⁺ release from endoplasmic reticulum stores [7,8]. Although various compounds of the xestospongine's family were reported to inhibit IP₃-dependent processes in distinct cell types [7,9], their use has been mainly limited to xestospongine C (XeC) [10–13].

The precise site of action for xestospongines at the cellular level remains elusive. In fact, XeC was reported to block IP₃-mediated Ca²⁺ signalling without interacting with the IP₃-binding site, since the alkaloid did not affect the ability of [³H]IP₃ to bind to cerebellar microsomes [7]. Furthermore, several reports provided evidence indicating that XeC lacks specificity, and acts by inhibiting the calcium pump in smooth muscle [14], cultured dorsal root ganglia neurons [15] and Schwann cells of the neuromuscular junction *in situ* [16]. In neuronal cells, IP₃ is known to exert an important second messenger role in the release of Ca²⁺ from intracellular stores in response to membrane-receptor activation (for a review see [17]). Furthermore, recent studies suggest a role for IP₃ receptors in complex spatial and temporal intracellular Ca²⁺ signals in cultured skeletal muscle cells, and mature mouse neuromuscular junctions [18–23].

The purpose of the present study was to test the specificity, the site of action and the ability of Xestospongine B (XeB) (Fig. 1), to block IP₃-mediated Ca²⁺ signalling. For this purpose, binding studies were carried out first, in both rat cerebellum membranes, and myotube homogenates. Second, we tested the ability of XeB to block bradykinin-induced increase of intracellular Ca²⁺ levels in differentiated rodent neuroblastoma (NG108-15) cells. Bradykinin has been shown in NG108-15 cells to increase the degradation of phosphatidylinositol-4,5-bisphosphate leading to the production IP₃ and diacylglycerol [24]. Third, we tested the effect of XeB in rat myotubes, where the involvement of IP₃ receptors in the generation of a slow intracellular Ca²⁺ transient involved in the regulation of gene expression has been described [18,19,23,25]. Finally, we investigated the ability of XeB to affect Ca²⁺ signals in highly purified isolated myonuclei.

*Corresponding author. Fax: +33 169 82 41 41.

E-mail address: jordi.molgo@nbcn.cnrs-gif.fr (J. Molgó).




Fig. 1. Chemical structure of XeB and XeC. Note that XeB differs from XeC by the stereochemistry of the two *cis*-oxaquinolizidine rings (*cis* and *trans* in XeC). XeB also bears a hydroxyl group on one of the oxaquinolizidine rings, and a methyl on the other one.

2. Materials and methods

2.1. Cell cultures

Mouse neuroblastoma × rat glioma NG108-15 hybrid cells were grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with 5% fetal calf serum, 100 μ M hypoxanthine, 0.4 μ M aminopterin, 16 μ M thymidine, 2 mM glutamine, 3 μ M glycine, 100 IU/ml penicillin and 100 μ g/ml streptomycin. The cultures were maintained at 37 °C in a humidified atmosphere containing 95% air–5% CO₂. Cells were seeded on sterile glass coverslips placed in plastic dishes (35 mm) at a density of 30×10^3 cells per dish. Cells were differentiated by reducing to 1% fetal calf serum and adding 0.5 μ M dibutyryl-cyclic-adenosine-monophosphate (Sigma–Aldrich Chimie, Saint Quentin Fallavier, France). This medium maintained cell differentiation for 1–5 days.

Neonatal rat skeletal muscle cells in primary culture were prepared as follows: myoblasts were obtained from posterior hind limbs of 12–24 h-old new-born rats. After dissection, the tissue was mechanically dispersed and then treated with 0.2% (w/v) collagenase type IV (Sigma) for 15 min at 37 °C under mild agitation. The suspension was filtered through nytex membranes or lens tissue paper and spun down at low speed. After a 10–15 min pre-plating on a 150 mm dish to partially eliminate the faster sedimenting fibroblasts, cells were plated on round coverslips at a density of ca. 350×10^3 per dish (35 mm). Culture medium was DMEM/F12, 10% bovine serum, 2.5% fetal calf serum, 100 mg/l penicillin, 50 mg/l streptomycin and 2.5 mg/l amphoterycin B. To eliminate remaining fibroblasts, 10 μ M cytosine arabinoside (Sigma) was added at the 3rd day of culture for 36 h. Then, the medium was replaced by a cytosine arabinoside-free medium containing 1.8% fetal calf serum concentration. Myotubes after the 5th day of culture had an estimated purity of more than 90%.

2.2. Myonuclei isolation

Highly purified myonuclei were obtained by the combined use of a hypotonic shock and mechanical disruption in a Dounce homogenizer, as previously reported [26]. Briefly, myotubes (10×10^6) were washed in phosphate-buffered saline (PBS), and incubated in 0.1% (v/v) trypsin (Sigma) for 20 min at 25–30 °C and later scrapped off using a rubber policeman. Cells were spun down at $4000 \times g$ (Heraeus Biofuge 15R) to eliminate the trypsin. The resulting cell pellet was suspended in hypotonic buffer containing 10 mM Tris–HCl, pH 7.8, 10 mM β -mercaptoethanol, 0.5 mM phenylmethylsulfonyl fluoride (Calbiochem), 1 μ g ml⁻¹ aprotinin, leupeptin and pepstatin (Calbiochem). After 2–3 min in ice the swollen cells were broken in a Dounce homogenizer. The nuclear pellet was obtained by centrifugation at $500 \times g$ for 6 min at 4 °C, and washed in 10 mM Tris–HCl, pH 7.2, 2 mM MgCl₂ plus protease inhibitors. Finally, the nuclear pellet was resuspended in 10 mM HEPES–Tris, pH 7.6, 110 mM KCl, 1 mM MgCl₂ and the protease inhibitors. Nuclear integrity was routinely checked by electron and confocal microscopy, both in unstained and acridine orange (2.5% v/v; Sigma) stained samples, as will be published elsewhere.

2.3. [³H]IP₃ binding assays

Equilibrium competition assays were performed using increasing concentrations of XeB (0.1–100 μ M) both in the absence and in the presence of a constant low concentration of tritiated IP₃ (³H]IP₃) in the reaction medium, as described previously [19]. Cerebellar membrane vesicles were incubated 30 min at 4 °C in 50 mM Tris–HCl (pH 8.4) containing 1 mM EDTA, 2 mM β -mercaptoethanol, and 1.6 nM of [³H]IP₃ (22 Ci/mmol, American Radiolabelled Chemicals Inc., USA). Myotubes homogenates were incubated at 4 °C for 30–40 min in a medium containing 50 mM Tris–HCl (pH 8.4), 1 mM EDTA, and 2 mM β -mercaptoethanol. 50 nM (from a 1 μ M stock) of [³H]IP₃ (*D*-myo-[2-³H]inositol 1,4,5-trisphosphate, specific activity 21.0 Ci/mmol; NEN-Dupont; 800–1000 cpm/pmol). The suspensions were centrifuged (Heraeus Biofuge 15R) at $10000 \times g$ for 10 min, the supernatant was aspirated, and the pellets were washed with PBS, and dissolved in NaOH (1 M). The [³H]IP₃ radioactivity remaining bound to the membranes was measured by liquid scintillation. Non-specific binding was determined in the presence of 1–2 μ M IP₃ (Sigma).

Competition binding experiments were analyzed by the Prism software (GraphPad Software Inc., USA) using a non-linear Hill equation for EC₅₀ determination (the concentration of competitor that competes for half the specific binding). The concentration of the competing ligand that will bind to half the binding sites at equilibrium, in the absence of radioligand, or other competitors (*K*_i values) were calculated by the Cheng and Prusoff equation (*K*_i = EC₅₀ / (1 + ([ligand*]/*K*_d)), where [ligand*] is the concentration of the hot ligand) [27].

2.4. Calcium signal measurements

NG108-15 cells were incubated for 45 min with 4 μ M fluo-3/AM (Molecular Probes Europe, The Netherlands) in standard solution (containing in mM: 154 NaCl, 5 KCl, 2 CaCl₂, 1 MgCl₂, 5 HEPES, and 11 glucose, pH 7.4) at 25 °C. After fluo-3/AM was loaded into cells, the dye was washout from the medium and the coverslips containing the cells were transferred to a recording chamber (1.5 ml volume). Ca²⁺ signals were imaged with an upright microscope equipped with an epifluorescence unit and an extended ISIS cooled CCD, video camera (Photonic Science, UK). For video imaging, illumination was provided by a quartz-halogen bulb (12 V/100 W) and fluorescence recorded through a plan 40 \times (0.7 numerical aperture) long-working-distance, water-immersion lens, using an interference set of filters (excitation: 485 nm; emission: 535 nm) and a dichroic mirror (505 nm), as previously reported [28]. Digitizing and analysis of fluorescence images was performed using a frame grabber (DT-3155, Data Translation, USA) and imaging workbench 2.1 software (Axon Instruments, Union City, CA, USA).

For measuring intracellular Ca²⁺ signals in rat myotubes, the cells were pre-incubated in a resting solution of the following composition (in mM): 145 NaCl, 5 KCl, 2.6 CaCl₂, 1 MgCl₂, 10 Na HEPES and 5.6 glucose (pH 7.4), containing 5.4 μ M fluo-3/AM for 30 min at 25 °C. Cells attached to coverslips were placed in a perfusion chamber (1 ml capacity) and mounted on an inverted fluorescence microscope (Olympus, Japan). Images were acquired with a cooled CCD camera (Spectra-Source MCD 600, USA). A filter wheel (Lambda-10-2, Sutter Instrument Co, USA) was used as a shutter in order to avoid unnecessary light exposure to the cells. The fluorescent images were collected every 0.1–2.0 s and analyzed frame by frame. Cells were depolarized by a fast perfusion system using a high K⁺ (47 mM) isotonic solution. Images from each experiment were processed identically by outlining the cell's fluorescence and determining their mean fluorescence before (*F*₀) and during various treatments (*F*). The relative fluorescence ($\Delta F/F_0$) was calculated as $(F - F_0)/F_0$.

Purified intact nuclei were incubated with fluo-3/AM (5.4 μ M) for 30 min in the presence or absence of known concentration of XeB at room temperature. Dye-loaded nuclei were washed, and centrifuged ($1000 \times g$ for 1 min) to remove the excess of dye, and were re-suspended in an internal buffer containing: 110 mM KCl, 10 mM HEPES–Tris (pH 7.0), 3 mM MgCl₂, 0.1 mM EGTA, 2.5 mM ATP, 25 μ M phosphocreatine (Sigma) and 5 UI ml⁻¹ of creatine phosphokinase (Sigma). Thereafter, nuclei were stimulated with 10 μ M IP₃ (Calbiochem). The photometric measurements were made in a Fluoromax-2 (ISA Instruments S.A. Inc. Jobin Yvon Spex). Excitation and emission wavelengths were 506 and 526 nm, respectively.

2.5. ATPase determination

For determining ATPase activity, 6–7 days-old rat-myotube cultures were used. Cells were rinsed in PBS solution and incubated for 15–20 min at room temperature with a solution containing 145 mM NaCl, 5 mM KCl, 1 mM MgCl₂, 5.6 mM glucose, 10 mM HEPES–Na (pH 7.4) and 1–100 μM XeB. Then, the reaction was stopped by rapid aspiration of the medium and the cells were homogenized using a Dounce homogenizer and re-suspended in a solution containing 50 mM Tris–HCl, pH 7.4, 1 mM EDTA, 1 mM β-mercaptoethanol and 0.3 M sucrose. Ca²⁺-ATPase activity was calculated as the difference between total ATPase and Mg²⁺-ATPase. The reaction medium to determine Mg²⁺-ATPase activity contained 50 mM L-histidine (pH 7.4), 3 mM MgCl₂, 100 mM KCl, 1 mM EGTA, 4 mM ATP and 0.05–0.1 mg/ml of protein in a final volume of 1.0 ml. To measure the total ATPase activity, EGTA was excluded from the medium and 0.1 mM CaCl₂ was added. The reaction was started by addition of ATP and the incubation was carried out at 25 °C for 15 min. The reaction was stopped by addition of 0.25 ml of 10% SDS, and then, 0.5 ml of 2.5% ammonium molybdate and 0.1 ml of ELON (*p*-methyl amino phenol and 3% sodium sulphite) was added and mixed thoroughly. The amount of inorganic Pi formed was calculated from a colorimetric calibration curve [29].

2.6. Xestospongins B

Xestospongia exigua specimens were collected in Prony Bay (New Caledonia). The isolation procedure of xestospongins B was the same as previously described [2]. Samples were controlled with electrospray ionization mass spectrometry (Bruker Esquire-LC ion trap; positive mode M⁺H 477.4 and M⁺Na 499.6; negative mode M–H 475.5) and TLC (Merck silica gel 60 F₂₅₄; hexane–Et₂O–MeOH–N–H₄OH 20/70/10/0.5).

3. Results

3.1. Binding competition between XeB and radiolabelled IP₃

The precise site of action for xestospongins has been a matter of controversy, probably in part because no reports of a direct interaction between the alkaloids and IP₃ receptors are available. As shown in Fig. 2, we measured displacement of [³H]IP₃ from its receptor by XeB, both in rat cerebellar membranes (Fig. 2A) and in rat myotube homogenates (Fig. 2B). In both cases, XeB displaced [³H]IP₃ from its receptor in a concentration-dependent manner. The curves were fitted to a first order kinetics and the concentration of XeB that competes for half the specific binding was calculated. Thus, the XeB EC₅₀ was 44.6 ± 1.1 μM (*n* = 3 independent preparations) for cerebellar membranes, and 27.4 ± 1.09 μM (*n* = 3) for myotube homogenates, with *K*_i values of 31 and 16 μM, respectively. Interestingly, xestospongins C, reported as unable to displace binding of IP₃ from brain preparations at low concentrations [7], also tended to displace IP₃ binding from myotube homogenates in a range of concentrations similar to those of XeB (Fig. 2B).

3.2. XeB blocks bradykinin-induced Ca²⁺ signalling in NG108-15 cells

Differentiated NG108-15 cells loaded with fluo-3/AM and bathed with standard physiological solution responded to the external application of bradykinin (1 μM) with a transient increase in the cytosolic fluorescence, as shown in a typical experiment (Fig. 3A–C). The Ca²⁺ signal induced by bradykinin (Fig. 3G) was characterized by a rapid rise in fluorescence that reached a maximum followed by a decay phase that attained the basal fluorescence level of the cells. When NG108-15 cells were pre-treated with 27 μM XeB for 20 min, bradykinin was unable to induce the characteristic transient Ca²⁺




Fig. 2. Dose–response relationship for the effect of XeB on the [³H]IP₃ binding to cerebellar rat membranes (A) and rat myotube homogenates (B). The data in A and B were corrected for non-specific binding, measured in the presence of 1 μM IP₃. The 100% value represents the [³H]IP₃ binding in the absence of XeB or XeC. For comparison, data obtained with 3 concentrations of XeC on rat myotube homogenates are shown. Means ± S.D. of three different determinations are shown in A and B.

signal observed in control experiments as shown in Fig. 3D–F, H, which is representative of nine different experiments with 36 independent coverslips. The blockade of bradykinin response by XeB was not accompanied by changes in the basal level of fluorescence when continuously monitored during a 20 min period (*n* = 6 independent coverslips, data not shown). However, the blockade of bradykinin responses by XeB depended on the concentration used. Two different XeB concentrations (5 and 13.5 μM), applied for 20 min, reduced the number of cells in a given coverslip that responded to bradykinin by 12 ± 3.2 and 66 ± 12%, respectively (*n* = 6 independent coverslips).

3.3. XeB blocks selectively a delayed Ca²⁺ signalling in rat myotubes

Cultured skeletal myotubes respond to membrane depolarization induced by a high K⁺ (47 mM) medium with two kinetically distinct Ca²⁺ signals (Fig. 4A): (i) a fast Ca²⁺ transient (lasting 1–2 s) related to excitation-contraction coupling, and (ii) a slow Ca²⁺ signal, starting seconds after the fast one, that is unrelated to contraction, and lasts several seconds [18,19].

XeB (10 and 40 μM), had little, or no effect on the fast Ca²⁺ transient evoked by a high K⁺ medium (*n* = 4) (Fig. 4C–E). The fact that the fast Ca²⁺ transient remained unaltered suggests that neither ryanodine receptors, nor Ca²⁺ storage pools were affected by the drug. Interestingly, XeB depending on the dose, either reduced and largely delayed the onset of the slow Ca²⁺ signal (Fig. 4D), or completely blocked the slow Ca²⁺ signal evoked by the high K⁺ medium (Fig. 4B and E). These results indicate that the slow Ca²⁺-signal, which is unrelated to




Fig. 3. Ca^{2+} signals induced by the external application of $1\ \mu\text{M}$ bradykinin in NG108-15 cells and its suppression by pre-treatment with $27\ \mu\text{M}$ XeB. Images show the basal fluorescence level of the cells before (A, D), during bradykinin exposure (B, E), and 5 min after bradykinin exposure (C, F). Time course of the relative fluorescence change of a cell exposed to bradykinin (G), and pre-treated with XeB before the addition of bradykinin to the standard medium (H). Distinct NG108-15 cells were used in A–C, D–F, G and H. Calibration in F applies to all images.

excitation-contraction coupling, can be effectively blocked by XeB.

3.4. XeB neither affects the Ca^{2+} pump nor capacitative Ca^{2+} entry in rat myotubes

It has been reported that XeC not only inhibited IP_3 -induced Ca^{2+} release, but also blocked the Ca^{2+} pump of the endoplasmic reticulum in various cell systems [14–16]. Therefore, in the present study we further investigated the specificity of XeB on Ca^{2+} release induced by thapsigargin, a well known blocker of the sarcoplasmic/endoplasmic reticulum Ca^{2+} ATPase pumps [30,31]. In control rat myotubes, exposed to a Ca^{2+} -free medium, the addition of thapsigargin ($0.5\ \mu\text{M}$) elicited a slow and transient calcium signal, that reflects the release and depletion of Ca^{2+} from the sarcoplasmic reticulum (Fig. 5A). This prolonged but transient Ca^{2+} release induced by thapsigargin was not altered when myotubes were previously incubated, for 20 min, with either 10 or $40\ \mu\text{M}$ XeB, as shown in representative recordings (Fig. 5B and C). Similar results to those shown in Fig. 5 have been observed in three additional experiments. These results indicate that in rat myotubes XeB neither depleted the sarcoplasmic reticulum Ca^{2+} stores nor prevented thapsigargin action.

In recent years, much research has been devoted to the mechanism linking depletion of intracellular Ca^{2+} stores to the activation of plasma membrane channels (for a review




Fig. 4. XeB inhibits the slow Ca^{2+} signal evoked by membrane depolarization with high K^+ ($47\ \text{mM}$) medium in rat myotubes. A, series of fluorescence images in myotubes previously loaded with fluo-3/AM, before (b) and after depolarization with high K^+ at the times indicated (s). The basal fluorescence (b) is shown on the top of the panels and the following images were taken immediately after the addition of the high K^+ medium, which remained in the bath throughout the recording. Images were acquired every 2 s and are presented in pseudocolors. Note the fast- and slow- Ca^{2+} signal transients. B, effect of pre-treatment with $40\ \mu\text{M}$ XeB for 20 min, in another myotube on the fast- and slow- Ca^{2+} signals triggered by high K^+ . Notice that XeB suppressed the slow- Ca^{2+} transient. C, D, and E, time course of the relative fluorescence changes, expressed with respect to the basal fluorescence, obtained in three different myotubes after addition of the high K^+ medium at time = 0. Myotubes in D and E, were pre-treated with 10 and $40\ \mu\text{M}$ XeB, respectively. Note both the reduction in amplitude and delayed onset of the first component of the slow signal (D) and complete blockade (E) of the slow- Ca^{2+} signal, together with the lack of significant effect of XeB on the fast Ca^{2+} signal.

see [33]). Therefore, it was of interest also to determine whether XeB affected capacitative Ca^{2+} entry through store operated channels after depletion of Ca^{2+} stores by thapsigargin. As shown in Fig. 5A, depletion of sarcoplasmic reticulum Ca^{2+} stores by $0.5\ \mu\text{M}$ thapsigargin, in Ca^{2+} -free medium, was accompanied by a fast increase in fluorescence, upon addition of $2\ \text{mM}$ Ca^{2+} to the external medium ($n = 3$). Furthermore, as shown in Fig. 5A–C, XeB (10 and $40\ \mu\text{M}$) did not appear to modify Ca^{2+} influx (as far as it can be judged by the fluorescence measurements), which reflect Ca^{2+} entry through store operated Ca^{2+} channels.

In order to further explore a possible interaction of XeB with the calcium pump, we measured the calcium-dependent ATPase activity on muscle cell homogenates. Ca^{2+} stimulated activity, measured as the difference between total activity, in the presence of calcium and magnesium, minus the activity in the absence of calcium, was $0.3\ \mu\text{mol Pi mg protein}^{-1}\ \text{min}^{-1}$, and was not significantly modified when cell homogenates were incubated with 1, 50 and $100\ \mu\text{M}$ XeB (Table 1).




Fig. 5. Lack of effect of XeB on Ca^{2+} transients evoked by thapsigargin ($0.5 \mu\text{M}$) in rat myotubes. A, relative fluorescence variation in a region of interest before and after addition of thapsigargin to the Ca^{2+} -free medium supplemented with EGTA (0.5 mM). When indicated by an arrow, the external medium was replaced by one containing 2 mM Ca^{2+} . B, same experiment as in A, using a myotube previously treated for 20 min with $10 \mu\text{M}$ XeB. C, same experiment as in A, using a myotube previously treated (20 min) with $40 \mu\text{M}$ XeB.

3.5. XeB blocks Ca^{2+} oscillations induced by IP_3 in isolated myonuclei

We have investigated the effect of XeB on Ca^{2+} mobilization induced by IP_3 , in a population of isolated rat myotube nuclei. The advantage of this model is that isolated myotube nuclei contain the nuclear envelope and some perinuclear endoplasmic reticulum, but are devoid of other cellular organelles, thus simplifying the interpretation of results. Previous work has shown that IP_3 can release Ca^{2+} from the nuclear envelope suggesting that IP_3 receptors may be localized predominantly in the inner nuclear membrane [33,34]. The addition of $10 \mu\text{M}$ IP_3 to the external medium of isolated myonuclei, previously loaded with fluo-3/AM, evoked oscillatory Ca^{2+} signals that could be measured during a given time period. These Ca^{2+} transients which reflect the activation of IP_3 receptors in the




Fig. 6. XeB blocks IP_3 -induced Ca^{2+} oscillations in highly purified isolated myonuclei. A, isolated myonuclei images obtained with contrast-phase (left panel) and fluorescence microscopy (right panel) after staining with acridine orange. Calibration in A = $10 \mu\text{m}$. B, dose-response relationship for the effect of XeB on calcium oscillations induced by IP_3 ($10 \mu\text{M}$) in a population of isolated myonuclei. The 100% value represents the frequency of IP_3 -induced calcium oscillations in the absence of XeB. The curve was fitted to a first order kinetics, and the calculated EC_{50} was $18.9 \pm 1.35 \mu\text{M}$ ($n = 3$ different determinations), with $R^2 = 0.9761$ for a goodness-of-fit.

population of nuclei were quantified under control conditions, and after incubation with various concentrations of XeB. As shown in Fig. 6, the alkaloid, depending on the concentration, either reduced the frequency, or completely abolished the calcium signal oscillations induced by $10 \mu\text{M}$ IP_3 . The curve was fitted to a single binding site and this fitting was better than that to a two-site model, indicating that the response is fairly specific.

4. Discussion

To the best of our knowledge, this is the first study to demonstrate that highly purified XeB blocks IP_3 -mediated Ca^{2+} signalling in differentiated rodent NG108-15 cells, intact rat

Table 1
Effect of xestospongine B on total, Mg^{2+} - and Ca^{2+} -ATPase activity determined in rat myotube homogenates

Condition	Total ATPase	Mg^{2+} -ATPase	Ca^{2+} -ATPase
Control	0.383 ± 0.100	0.062 ± 0.026	0.321 ± 0.096
$1 \mu\text{M}$ Xestospongine B	0.349 ± 0.084	0.058 ± 0.026	0.291 ± 0.097
$50 \mu\text{M}$ Xestospongine B	0.489 ± 0.047 (*)	0.153 ± 0.041 (**)	0.336 ± 0.071
$100 \mu\text{M}$ Xestospongine B	0.449 ± 0.054	0.134 ± 0.039 (***)	0.315 ± 0.109

ATPase activities were determined as described in Section 2 and are expressed as $\mu\text{mol Pi mg protein}^{-1} \text{ min}^{-1}$. Values are presented as the means \pm S.D., of $n = 6$ different determinations from different myotube cultures. (*), (**), (***) Denote values significantly different from controls ($P < 0.05$, 0.01 and 0.03 , respectively).

skeletal myotubes, and isolated myonuclei. Previous studies have considered xestospongins as allosteric antagonists for IP₃ receptor, since XeC (1–10 μM) did not affect the ability of [³H]IP₃ (1 nM) to bind to cerebellar microsomes, despite blocking IP₃-induced Ca²⁺ release [7]. In our hands, XeC had little or no effect on IP₃ binding in myotube homogenates at sub-micro molar concentrations (0.5–1 μM) at which pharmacological effects have been reported in other systems; nevertheless, some binding displacement was evident at higher concentrations. Our results on the inhibition of Ca²⁺ signals in intact cultured cells, and the binding competition experiments, in both rat cerebellar membranes and skeletal myotube homogenates (Fig. 2), strongly suggest that XeB effects can be explained by competitive inhibition of IP₃ receptors. The fact that the apparent K_i values for displacement of [³H]IP₃ by XeB are in the same concentration range than the EC₅₀ for inhibition of Ca²⁺ signals is indicative that the alkaloid has a low affinity for IP₃ receptors. Since results obtained in intact cultured cells lack precision in terms of the effective concentration of the blocker at the active intracellular binding site, experiments were performed on isolated rat skeletal myonuclei in which XeB concentration could be more effectively monitored. Isolated myonuclei have been shown to produce Ca²⁺ transients upon addition of IP₃ (Cardenas, Liberona, Molgó, Colasante, Mignery and Jaimovich, to be published). Under this condition, XeB inhibition of Ca²⁺ oscillations occurred with similar EC₅₀ as the ones obtained for IP₃ binding-displacement experiments. These results further confirm the low affinity XeB has for IP₃ receptors.

In NG108-15 cells XeB completely blocked bradykinin-induced intracellular Ca²⁺ signals (Fig. 3) that are known to be mediated by IP₃ [24,35]. A similar blockade of bradykinin-induced Ca²⁺ signal in pheochromocytoma (PC12) cells was previously shown with equivalent concentrations of XeC [7]. However, in those cells, XeC was reported not only to inhibit Ca²⁺ release from IP₃-sensitive Ca²⁺ stores, but also from ryanodine-sensitive Ca²⁺ stores [7]. In the present study we were able to test in rat skeletal myotubes depolarized with a high K⁺ medium the effect of XeB in the two Ca²⁺ stores, finding no reduction on the fast Ca²⁺ signal relating Ca²⁺ release from ryanodine-sensitive stores (Fig. 4). Only the slow Ca²⁺ signal occurring during K⁺-induced membrane depolarization was affected by XeB. This slow Ca²⁺ signal has been shown to be related to IP₃ transients and IP₃ receptors [18–20], and recent work suggests its implication in regulating gene expression [21,22,25]. Thus, XeB appears rather selective for IP₃-sensitive Ca²⁺ stores in skeletal myotubes. Cultured skeletal muscle cells are endowed with all three types of IP₃-receptor [21], but their relative role in Ca²⁺ signalling remain at present unknown, in part due to the lack of selective antagonists for IP₃ receptor subtypes.

Despite its popularity XeC has been reported to have some limitations in regard to its specificity. Thus, it has been suggested that XeC not only inhibited IP₃-induced Ca²⁺ release in permeabilized A7r5 smooth muscle cells, but also was an equally potent blocker of the endoplasmic-reticulum calcium pump [14]. However, subsequent studies were unable to observe any inhibitory effect of XeC on caffeine-induced Ca²⁺ release, suggesting that the calcium pump was unaffected by the alkaloid [36]. In the present study a lack of effect of XeB on the calcium pump of skeletal myotubes was actually shown in two independent ways. Thus, prolonged incubation of cells with

40 μM XeB did not alter Ca²⁺ transients induced by thapsigargin in Ca²⁺-free medium (Fig. 5). These transients reflect the Ca²⁺ content of intracellular stores, which are completely depleted by thapsigargin [31,32]. The fact that the content of the stores remained unchanged clearly indicates that XeB does not inhibit the calcium pump. Furthermore, direct measurements of the Ca-ATPase activity in muscle cell homogenates (Table 1) indicated that it was not inhibited by XeB. The sole non-specific effect so far detected with XeB appeared to be a slight enhancement of basal Mg-dependent ATPase activity.

We conclude that XeB is an effective inhibitor of IP₃-dependent Ca²⁺ signals in both skeletal muscle and neuronal cells, and isolated myonuclei and constitutes a new membrane-permeable pharmacological tool for studying IP₃-dependent signal transduction in living cells and subcellular organelles.

Acknowledgment: This work was made possible by an ECOS Sud-CONY CIT exchange program (C03S02), and was supported in part by FONDAP (grant no. 10500006 to E.J.), the Association Française contre les Myopathies (AFM), and the Direction des Systèmes de Forces et de la Prospective (grant no. 02 60 65 093 to J. M.). J.B. and C.M. were supported by D.G.A.-C.N.R.S fellowships. C.C. was supported by a fellowship from the AFM. We are grateful to Miss Isabelle Pouny (I.R.D.-C.R.S.N. Pierre Fabre) for running the electrospray mass spectrometry analysis of xestospongins B, and Dr. Béatrice Rouzère-Dubois for providing the neuroblastoma (NG108-15) cells used in the present study.

References

- [1] Nakagawa, M., Endo, M., Tanaka, N. and Gen-Pei, L. (1984) Structures of xestospongins A, B, C, D, novel vasodilative compounds from marine sponge *Xestospongia exigua*. *Tetrahedron Lett.* 25, 3227–3230.
- [2] Quirion, J.C., Sevenet, T., Husson, H.P., Weniger, B. and Debitus, C. (1992) Two new alkaloids from *Xestospongia* Sp., a new caledonian sponge. *J. Nat. Prod.* 55, 1505–1508.
- [3] Venkateswarlu, Y., Reddy, M.V.R. and Rao, J.V. (1994) Bis-1-oxaquinolizidines from the sponge *Haliclona exigua*. *J. Nat. Prod.* 57, 1283–1285.
- [4] Pettit, G.R., Orr, B., Herald, D.L., Doubek, D.L., Tackett, L., Schmidt, J.M., Boyd, M.R., Pettit, R.K. and Hooper, J.N.A. (1996) Antineoplastic agents. 357. Isolation and X-ray crystal structure of racemic xestospongins D from the Singapore marine sponge *Niphates* sp. *Bioorg. Med. Chem. Lett.* 6, 1313–1318.
- [5] Reddy, M.V.R. and Faulkner, D.J. (1997) 3,3'-Dimethylxestospongins C, a new bis-1-oxaquinolizidine alkaloid from the Palauan sponge *Xestospongia* Sp. *Nat. Prod. Lett.* 11, 53–59.
- [6] Orabi, K.Y., El Sayed, K.A., Hamann, M.T., Dunbar, D.C., Al-Said, M.S., Higa, T. and Kelly, M. (2002) Araguspongins K and L, new bioactive bis-1-oxaquinolizidine N-oxide alkaloids from Red Sea specimens of *Xestospongia exigua*. *J. Nat. Prod.* 65, 1782–1785.
- [7] Gafni, J., Munsch, J.A., Lam, T.H., Catlin, M.C., Costa, L.G., Molinski, T.F. and Pessah, I.N. (1997) Xestospongins: Potent membrane permeable blockers of the inositol 1,4,5-trisphosphate receptor. *Neuron* 19, 723–733.
- [8] Wilcox, R.A., Primrose, W.U., Nahorski, S.R. and Challiss, R.A.J. (1998) New developments in the molecular pharmacology of the myo-inositol 1,4,5-trisphosphate receptor. *Trends Pharmacol. Sci.* 19, 467–475.
- [9] Venkateswarlu Rao, J., Desai, D., Vig, P.J. and Venkateswarlu, Y. (1998) Marine biomolecules inhibit rat brain nitric oxide synthase activity. *Toxicology* 129, 103–112.
- [10] Miyamoto, S., Izumi, M., Hori, M., Kobayashi, M., Ozaki, H. and Karaki, H. (2000) Xestospongins C, a selective and membrane-permeable inhibitor of IP₃ receptor, attenuates the positive inotropic effect of alpha-adrenergic stimulation in guinea-pig papillary muscle. *Br. J. Pharmacol.* 130, 650–654.

- [11] Oka, T., Sato, K., Hori, M., Ozaki, H. and Karaki, H. (2002) Xestospongin C, a novel blocker of IP₃ receptor, attenuates the increase in cytosolic calcium level and degranulation that is induced by antigen in RBL-2H3 mast cells. *Br. J. Pharmacol.* 135, 1959–1966.
- [12] Dajas-Bailador, F.A., Mogg, A.J. and Wonnacott, S. (2002) Intracellular Ca²⁺ signals evoked by stimulation of nicotinic acetylcholine receptors in SH-SY5Y cells: Contribution of voltage-operated Ca²⁺ channels and Ca²⁺ stores. *J. Neurochem.* 81, 606–614.
- [13] Suen, K.C., Lin, K.F., Elyaman, W., So, K.F., Chang, R.C. and Hugon, J. (2003) Reduction of calcium release from the endoplasmic reticulum could only provide partial neuroprotection against beta-amyloid peptide toxicity. *J. Neurochem.* 87, 1413–1426.
- [14] De Smet, P., Parys, J.B., Callewaert, G., Weidema, A.F., Hill, E., De Smedt, H., Erneux, C., Sorrentino, V. and Missiaen, L. (1999) Xestospongin C is an equally potent inhibitor of the inositol 1,4,5-trisphosphate receptor and the endoplasmic-reticulum Ca²⁺ pumps. *Cell Calcium* 26, 9–13.
- [15] Solovyova, N., Fernyhough, P., Glazner, G. and Verkhratsky, A. (2002) Xestospongin C empties the ER calcium store but does not inhibit InsP₃-induced Ca²⁺ release in cultured dorsal root ganglia neurons. *Cell Calcium* 32, 49–52.
- [16] Castonguay, A. and Robitaille, R. (2002) Xestospongin C is a potent inhibitor of SERCA at a vertebrate Synapse. *Cell Calcium* 32, 39–47.
- [17] Bootman, M.D., Lipp, P. and Berridge, M.J. (2001) The organisation and functions of local Ca²⁺ signals. *J. Cell Sci.* 114, 2213–2222.
- [18] Jaimovich, E., Estrada, M., Mattei, C., Debitus, C., Laurent, D. and Molgó, J. (2000) Xestospongins reveal the involvement of IP₃ receptors in slow calcium signals of skeletal myotubes. *Biophys. J.* 78, 432A, (Abstract).
- [19] Jaimovich, E., Reyes, R., Liberona, J.L. and Powell, J.A. (2000) IP₃ receptors, IP₃ transients, and nucleus-associated Ca²⁺ signals in cultured skeletal muscle. *Am. J. Physiol.* 278, C998–C1010.
- [20] Estrada, M., Cárdenas, C., Liberona, J.L., Carrasco, M.A., Mignery, G.A., Allen, P.D. and Jaimovich, E. (2001) Calcium transients in 1B5 myotubes lacking ryanodine receptors are related to inositol trisphosphate receptors. *J. Biol. Chem.* 276, 22868–22874.
- [21] Powell, J.A., Carrasco, M.A., Adams, D.S., Drouet, B., Rios, J., Müller, M., Estrada, M. and Jaimovich, E. (2001) IP₃ receptor function and localization in myotubes: An unexplored Ca²⁺ signaling pathway in skeletal muscle. *J. Cell Sci.* 114, 3673–3683.
- [22] Araya, R., Liberona, J.L., Cardenas, J.C., Riveros, N., Estrada, M., Powell, J.A., Carrasco, M.A. and Jaimovich, E. (2003) Dihydropyridine receptors as voltage sensors for a depolarization-evoked, IP₃R-mediated, slow calcium signal in skeletal muscle cells. *J. Gen. Physiol.* 121, 3–16.
- [23] Powell, J.A., Molgó, J., Adams, D.S., Colasante, C., Williams, A., Bohlen, M. and Jaimovich, E. (2003) IP₃ receptors and associated Ca²⁺ signals localize to satellite cells and to components of the neuromuscular junction in skeletal muscle. *J. Neurosci.* 23, 8185–8192.
- [24] Yano, K., Higashida, H., Inoue, R. and Nozawa, Y. (1984) Bradykinin-induced rapid breakdown of phosphatidylinositol 4,5-bisphosphate in neuroblastoma × glioma hybrid NG108-15 cells. *J. Biol. Chem.* 259, 10201–10207.
- [25] Carrasco, M.A., Riveros, N., Rios, J., Müller, M., Torres, F., Pineda, J., Lantadilla, S. and Jaimovich, E. (2003) Depolarization-induced slow calcium transients activate early genes in skeletal muscle cells. *Am. J. Physiol. Cell. Physiol.* 284, C1438–C1447.
- [26] Martelli, A.M., Gilmour, R.S., Bertagnolo, V., Neri, L.M., Manzoli, L. and Cocco, L. (1992) Nuclear localization and signalling activity of phosphoinositidase C beta in Swiss 3T3 cells. *Nature* 358, 242–245.
- [27] Cheng, Y.C. and Prusoff, W.H. (1973) Relationship between the inhibition constant (K_i) and the concentration of inhibitor which causes 50 per cent inhibition (IC₅₀) of an enzymatic reaction. *Biochem. Pharmacol.* 22, 3099–3108.
- [28] Meunier, F.A., Mattei, C., Chameau, P., Lawrence, G., Colasante, C., Kreger, A.S., Dolly, J.O. and Molgó, J. (2000) Trachinylisin mediates SNARE-dependent release of catecholamines from chromaffin cells via external and stored Ca²⁺. *J. Cell Sci.* 113, 1119–1125.
- [29] Fiske, C.H. and Subbarow, Y. (1925) The colorimetric determination of phosphorus. *J. Biol. Chem.* 66, 375–400.
- [30] Lytton, J., Westlin, M. and Hanley, M.R. (1991) Thapsigargin inhibits the sarcoplasmic or endoplasmic reticulum Ca-ATPase family of calcium pumps. *J. Biol. Chem.* 266, 17067–17071.
- [31] Treiman, M., Caspersen, C. and Christense, S.B. (1998) A tool coming of age: Thapsigargin as an inhibitor of sarco-endoplasmic reticulum Ca²⁺-ATPases. *Trends Pharmacol. Sci.* 19, 131–135.
- [32] Putney Jr., J.W., Broad, L.M., Braun, F.J., Lievreumont, J.P. and Bird, G.S. (2001) Mechanisms of capacitative calcium entry. *J. Cell Sci.* 114, 2223–2229.
- [33] Gerasimenko, O.V., Gerasimenko, J.V., Tepikin, A.V. and Petersen, O.H. (1995) ATP dependent accumulation and inositol trisphosphate- or cyclic ADP-ribose-mediated release of Ca²⁺ from the nuclear envelope. *Cell* 80, 439–444.
- [34] Gerasimenko, J., Maruyama, Y., Tepikin, A., Petersen, O.H. and Gerasimenko, O. (2003) Calcium signalling in and around the nuclear envelope. *Biochem. Soc. Trans.* 31, 76–78.
- [35] Higashida, H. and Ogura, A. (1991) Inositol trisphosphate/calcium-dependent acetylcholine release evoked by bradykinin in NG108-15 rodent hybrid cells. *Ann. NY Acad. Sci.* 635, 153–166.
- [36] Ozaki, H., Hori, M., Kim, Y.S., Kwon, S.C., Ahn, D.S., Nakazawa, H., Kobayashi, M. and Karaki, H. (2002) Inhibitory mechanism of xestospongin-C on contraction and ion channels in the intestinal smooth muscle. *Br. J. Pharmacol.* 137, 1207–1212.