

Significance of Cerebral Small-Vessel Disease in Acute Intracerebral Hemorrhage

Por: [Sato, S](#) (Sato, Shoichiro)^[1,2]; [Delcourt, C](#) (Delcourt, Candice)^[1,3,4]; [Heeley, E](#) (Heeley, Emma)^[1,3]; [Arima, H](#) (Arima, Hisatomi)^[1,3,5]; [Zhang, SH](#) (Zhang, Shihong)^[6]; [Salman, RA](#) (Salman, Rustam Al-Shahi)^[7]; [Stapf, C](#) (Stapf, Christian)^[8,9]; [Woo, D](#) (Woo, Daniel)^[10]; [Flaherty, ML](#) (Flaherty, Matthew L.)^[10]; [Vagal, A](#) (Vagal, Achala)^[11] ...[Más](#)

Autoría conjunta: [INTERACT2 Investigators](#)

STROKE

Volumen: 47

Número: 3

Páginas: 701-707

DOI: 10.1161/STROKEAHA.115.012147

Fecha de publicación: MAR 2016

[Ver información de revista](#)

Resumen

Background and Purpose-

The significance of structural changes associated with cerebral small-vessel disease (SVD), including white matter lesions (WML), lacunes, and brain atrophy, to outcome from acute intracerebral hemorrhage is uncertain. We determined associations of computed tomographic radiological manifestations of cerebral SVD and outcomes, and in terms of any differential effect of early intensive blood pressure-lowering treatment, in the large-scale Intensive Blood Pressure Reduction in Acute Cerebral Hemorrhage Trial (INTERACT2).

Methods-

We graded WML (van Swieten scale), the presence of lacunes, and brain atrophy (2 linear measurements and visual rating) for 2069 of 2839 patients with available baseline brain computed tomography (< 6 hours of intracerebral hemorrhage onset) by 3 independent neurologists blind to clinical data.

Results-

WML grade and 2 linear measurements of brain atrophy were associated with death or major disability at 90 days: multivariable-adjusted odds ratios for WML (grade 3 and 4 versus 0), frontal ratio, and third ventricle Sylvian fissure distance (most versus least severe atrophy quartile) were 1.42 (95% confidence interval, 1.02-1.98), 1.47 (1.08-1.99), and 1.64 (1.21-2.22), respectively (all P for trend < 0.05). There was no association between lacunes and outcomes. There were no

significant differences in the effects of intensive blood pressure-lowering across subgroups of cerebral SVD.

Conclusions-

Preexisting cerebral SVD manifestations of WML and brain atrophy predict poor outcome in acute intracerebral hemorrhage. There is no apparent hazard of early intensive lowering of blood pressure according to the INTERACT2 protocol, in patients with underlying cerebral SVD.

Clinical Trial Registration-

URL:

[GRAPHICS]

. Unique identifier: NCT00716079.

Palabras clave

Palabras clave de autor: cerebral small vessel disease; stroke; hypertension; cerebral hemorrhage; blood pressure

KeyWords Plus: WHITE-MATTER LESIONS; VASOMOTOR REACTIVITY; COMPUTED-TOMOGRAPHY; STROKE TRIAL; LEUKOARAIOSIS; BRAIN; CT; MRI; ASSOCIATION; IMPAIRMENT

Información del autor

Dirección para petición de copias: Anderson, CS (autor para petición de copias)

+ George Inst Global Hlth, POB M201, Missenden Rd, Camperdown, NSW 2050, Australia.

Direcciones:

- + [1] George Inst Global Hlth, Neurol & Mental Hlth Div, Sydney, NSW, Australia
- + [2] Natl Cerebral & Cardiovasc Ctr, Dept Cerebrovasc Med, Suita, Osaka, Japan
- + [3] Univ Sydney, Sydney Med Sch, Sydney, NSW 2006, Australia
- + [4] Royal Prince Alfred Hosp, Dept Neurol, Camperdown, NSW 2050, Australia
- + [5] Shiga Univ Med Sci, Ctr Epidemiol Res Asia, Otsu, Shiga, Japan
- + [6] Sichuan Univ, West China Hosp, Dept Neurol, Chengdu 610064, Peoples R China
- + [7] Univ Edinburgh, Ctr Clin Brain Sci, Edinburgh, Midlothian, Scotland
- [8] Hop Lariboisiere, Dept Hosp Univ DHU NeuroVasc, F-75475 Paris, France

- + [9] Univ Montreal, Dept Neurosci, Ctr Hosp Univ Montreal CRCHUM, Ctr Rech, Montreal, PQ, Canada
- + [10] Univ Cincinnati, Acad Hlth Ctr, Dept Neurol, Cincinnati, OH 45221 USA
- + [11] Univ Cincinnati, Acad Hlth Ctr, Dept Radiol, Cincinnati, OH 45221 USA
- + [12] Univ Newcastle, Hunter Med Res Inst, John Hunter Hosp, Dept Neurol, Newcastle, NSW 2300, Australia
- + [13] Shanghai Jiao Tong Univ, Shanghai Inst Hypertens, Ruijin Hosp, Shanghai 200030, Peoples R China
- + [14] Univ Leicester, Dept Cardiovasc Sci, Leicester, Leics, England
- + [15] Univ Leicester, NIHR Biomed Res Unit Cardiovasc Dis, Leicester, Leics, England
- + [16] Univ Desarrollo, Serv Neurol, Dept Med, Clin Alemana Santiago, Fac Med Clin Alemana, Santiago, Chile
- + [17] Univ Chile, Fac Med, Santiago 7, Chile
- + [18] Westmead Hosp, Sch Clin, Dept Med, Westmead, NSW 2145, Australia

Direcciones de correo electrónico: canderson@georgeinstitute.org.au

Financiación

Entidad financiadora	Número de concesión
National Health and Medical Research Council (NHMRC) of Australia	571281 512402 1004170

[Ver texto de financiación](#)

Editorial

LIPPINCOTT WILLIAMS & WILKINS, TWO COMMERCE SQ, 2001 MARKET ST, PHILADELPHIA, PA 19103 USA

Categorías / Clasificación

Áreas de investigación:Neurosciences & Neurology; Cardiovascular System & Cardiology

Categorías de Web of Science:Clinical Neurology; Peripheral Vascular Disease

Información del documento

Tipo de documento:Article

Idioma:English

Número de acceso: [WOS:000371496200010](#)

ID de PubMed: 26846860

ISSN: 0039-2499

eISSN: 1524-4628

Información de la revista

- **Impact Factor:** [Journal Citation Reports®](#)

Otra información

Número IDS: DF6VT

Referencias citadas en la Colección principal de Web of Science: **30**

Veces citado en la Colección principal de Web of Science: **0**