

## TABLE OF CONTENTS

INTRODUCTORY REMARKS.....	15
CHAPTER I:	
FACTORS CONTROLLING ALPINE GLACIATIONS IN THE SIERRA BAGUALES MOUNTAIN RANGE OF SOUTHERN PATAGONIA (50° S), INFERRED FROM THE MORPHOMETRIC ANALYSIS OF GLACIAL CIRQUES.....	28
Abstract.....	28
Resumen.....	29
Introduction.....	30
Study area	
Geologic and climatical setting.....	31
Geomorphology.....	32
Methodology.....	33
Results	
Cirque classification according to their activity.....	41
Cirque shape.....	42
Elevation, slope and aspect.....	44
L/H W/H and L/W ratios.....	45
Morphological cirque types.....	46
Composite map.....	48
Discussion.....	49
Cirque distribution, elevation and dimensions.....	51
Paleoglacial significance.....	51
Conclusions.....	52
Acknowledgements.....	53
Bibliography.....	54

CHAPTER II:	
RELICT GLACIAL LANDSCAPE IN THE SIERRA BAGUALES MOUNTAIN RANGE (50°-51° S): EVIDENCE OF GLACIATION DYNAMICS AND STYLES IN THE EASTERN FOOTHILLS OF THE SOUTHERN PATAGONIAN ANDES.....	61
Abstract.....	61
Introduction.....	62
Study area.....	63
Methods	
Geomorphological mapping.....	66
Radiocarbon dates.....	67
ELA estimation and reconstruction of surface profile of former glaciers.....	67
SBMR temperature variation based on former ELA position.....	68
Results	
Glacial geomorphology.....	69
Radiocarbon dates.....	75
ELA behavior and temperature variation.....	76
Surface profiles of former glaciers.....	76
Discussion.....	77
Conclusions.....	79
Acknowledgements.....	80
Bibliography.....	80
FINAL REMARKS.....	88
APPEND 1.....	94
APPEND 2.....	98

## INDEX OF FIGURES

Figure 1.1. Sketch of southern Patagonian glaciations during LGM of the major paleo-glacier lobes from 46° to 54° S. Modified from Rabassa *et al.* (2011), and references therein. Last 5 Kyr B.P. shows neoglacial chronologies proposed from the fluctuation of Patagonian glaciers in the Holocene by Mercer and by Aniya, modified from Glasser *et al.* (2004)..... 17

Figure 1.2. SRTM Digital Elevation Model of Southern Patagonia showing major lakes and structural regions (FTB: Magallanic Fold-and-Thrust Belt, FB: Magellanes Foreland Basin; DM: Deseado Masiff). Contour lines show equal precipitation (DGA, 1987) and the orographic effect of the Andean Main Range and Sierra Baguales Mountain Range. PA: Punta Arenas, PNat: Puerto Natales, White dotted box shows study area..... 20

Figure 2.1a. Right box shows the location of the study area. Elevations were obtained from the SRTM model. The brown dotted line shows the major moraine systems east of Lago Argentino, Brazo Rico and Brazo Sur (Mercer, 1976; Strelin and Malagnino, 2000; and references therein), and south and east of Lago Toro and Lago Sarmiento respectively (Sagredo *et al.*, 2011; Garcia *et al.*, 2013). Cirque activity corresponds to 1). current glacier activity, 2). perennial snow activity, 3). no glacier or snow activity, 4). not categorized. Figure 2.1b. Mapped cirque floor elevations, which show the increase in elevation towards the east. A – A' shows a topographic profile from the SPIF to the easternmost SBMR..... 34

Figure 2.2. Google Earth © image showing the study area from the SW. Photographs a) and b) (see location on Google image) show a hanging valley and glacial cirque, and photo c) shows a glacial valley. Current snow and glacier processes are particularly active along the high elevations on the eastern side of the SBMR (i.e., see Figure. 2.1)..... 35

Figure 2.3. Lateral moraine (dark lines) below the cirques shown in figure 2a (F3 in Fig. 2.1b).....	35
Figure 2.4. Striated erratic (F4 in Fig. 2.1b).....	36
Figure 2.5. Definition of key morphometric attributes. Left box shows the length (L) and azimuth (for aspect) in a grey line and width (W) in a white dashed line. i - ii correspond to a topographic profile on ASTERGDEM. The right box shows the characteristics of range (H) and the topographic profile.....	37
Figure 2.6. Cirque showing evidence of glacier activity, at the easternmost section of the SBMR (F6 in Fig. 2.1b).....	42
Figure 2.7. Cirque showing perennial snow activity (F7 in Fig. 2.1 b).....	42
Figure 2.8. Frequency distribution of L, W and H. Cirque mean surface (km <sup>2</sup> ) decreases with an increasing in depth (H). Otherwise, when the average slope becomes gentle (i.e. decrease of H), cirques are most likely to grow longitudinally and laterally. As shown in figure 2.1, large cirques are concentrated along the eastern margin of the SPIF and gradually get smaller towards the interior of the continent around the SBMR.....	43
Figure 2.9. Rose diagram showing frequency distribution of aspect. Cirques with a SW aspect tend to be located in the intermediate zone and altitudinal range E (1,472 – 1,741 m a.s.l), while cirques with S aspects are preferably located at the SPIF. Otherwise cirques with a S to SE aspect show a homogeneous distribution within the study area and an altitudinal range between C to D (1,126 –1,292 and 1,293 – 1,471 m a.s.l.). Moreover, cirques with W to SW aspects tend to show larger surface areas.....	45

Figure 2.10. Tree cluster of cirques (Euclidean distance measurements and amalgamation conducted by Ward`s method). The analysis shows two major morphological types where the differences are based on the cirque activity (i.e. glacial, perennial snow, or absence of both), surface, shape and spatial distribution. Morphological type 2 tends to show largest cirques compared with the morphological type 1..... 47

Figure 2.11. Composite map of study area. Cirques were categorized using 10 parameters extracted from the morphometric analysis (Table 1.2). Black dotted lines show outlines of LGM lobes. Cirque areas closest to the SPIF, such as in the far SW, were partly covered or adjacent to the large ice sheet during the LGM (Garcia *et al.*, 2012; Strelin *et al.*, 2011). This contributed to their higher areal development in relation to those located in the eastern SBMR, which were isolated from the ice sheet outlet lobes. Note that high values show the spatial distribution of currently active snow/ice processes, which particularly for the eastern SBMR, are conditioned by cirque floor elevations (Fig. 2.1) and the rainshadow effect. The lowest values spatially match with evidence of areas covered or adjacent to former outlet glaciers of the SPIF, and cirques with no evidence of current glacier or snow activity..... 50

Figure 3.1. The right box shows the general location of the study area, and the blackstippled box shows the SBMR. Major lakes appear in dark blue, SPIF is shown in light blue. The black stippled line shows an outline of the mapped moraine systems around Lago Argentino (Strelin and Malagnino, 2009, and references therein, Kaplan *et al.*, 2005; 2008; 2011; Strelin *et al.*, 2014), and Lagos Toro and Sarmiento (Marden, 1997, Sagredo *et al.*, 2011; Garcia *et al.*, 2012). CG: Cerro Guido; RC: Rio Las Chinas; TdP: Torres del Paine, which correspond to meteorological stations of the Direccion General de Aguas (DGA). Elevations were obtained from an SRTM model..... 65

Figure 3.2. SBMR glacial geomorphology. White boxes show the location of

samples for radiocarbon dating. RB-RB` shows the location of the topographic profile (Figure 3.8). AA`, BB`, CC` show the former surface topography of alpine glaciers (Figure 3.10)..... 71

Figure 3.3. U-shaped valley in the area where the former Alpine glaciers coalesced to form a small Icefield (F3 in Fig. 3.2)..... 72

Figure 3.4. Remnant of lateral moraine and boulders deposited and aligned with the moraine (F4 in Fig. 3.2)..... 72

Figure 3.5. Lateral moraine and boulders below the U-shaped valley shown in Figure 3 (F5 in Fig. 3.2)..... 73

Figure 3.6. Figure 3.6. Hummocky terrain and remnants of lateral moraine. These deposits, located in the middle section of the Baguales River basin, constitute evidence of a former Icefield (F6 in Fig. 3.2)..... 73

Figure 3.7. Eroded kame terraces along the Baguales River (see also Figure 3.8) (F7 in Fig. 3.2)..... 74

Figure 3.8. RB-RB' topographic profile..... 74

Figure 3.9. Spatial distribution of rock mass strength in SBMR, the box on the left shows that at greater rock resistance cirque areas tend to decrease..... 75

Figure 3.10. Former surface topography profiles of alpine glaciers. Topography and ice thickness vary depending on changes in the bedrock slope..... 77

## INDEX OF TABLES

Table 1.1. Data source for cirque mapping .....	36
Table 1.2. Summary of cirque morphometric characteristics.....	38
Table 1.3. Correlation between analyzed morphometric characteristics. S, P, L, and W parameters have a high positive correlation followed by the altitude range (H). These variables are dependent and determine mainly the shape and cirque development. On the other hand, the $E_{med}$ , $E_{min}$ and $S_{mean}$ parameters have a weak negative correlation with the shape parameters. Particularly M has a strong negative correlation with the planimetric development and cirque incision. See Table 1.2 for definition of attributes....	44
Table 1.4. Distribution of cirque attributes with aspect. See Table 2 for definition of attributes.....	46
Table 1.5. Summary statistics of morphometric characteristics for cirque types obtained from cluster analysis. See Table 2 for definition of attributes.....	48
Table 2.1. Meteorological stations and data obtained from the DGA of the Chilean Ministerio de Obras Publicas website ( <a href="http://snia.dga.cl/BNAConsultas/reportes">http://snia.dga.cl/BNAConsultas/reportes</a> ).....	69