

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	Objectives . . . . .	2
1.2	Adsorption . . . . .	2
1.2.1	Physisorption . . . . .	4
1.2.2	Chemisorption . . . . .	4
1.3	Electroadsorptive effect . . . . .	6
1.4	Other adsorbed species . . . . .	9
1.5	Atomic Force Microscopy . . . . .	9
1.6	Low Energy Ion Spectroscopy . . . . .	10
1.7	X-ray photoelectron spectroscopy (XPS) . . . . .	11
1.7.1	Time resolved measurements . . . . .	12
1.8	Mass Spectrometer . . . . .	12
<b>2</b>	<b>Experimental setup</b>	<b>13</b>
2.1	General description . . . . .	13
2.1.1	Insertion arm . . . . .	14
2.1.2	Sample holder . . . . .	14
2.1.3	Gas Source . . . . .	15
2.1.4	Mass Spectrometer . . . . .	15
2.1.5	Gas atmosphere . . . . .	15
2.2	Sample . . . . .	16
2.2.1	Surface potential measurement . . . . .	17
2.2.2	Circuit . . . . .	17
2.3	Other experiments . . . . .	18
2.3.1	Titanium oxide . . . . .	18
2.3.2	Thiols . . . . .	18
2.3.3	Topographical characterization . . . . .	19
<b>3</b>	<b>Results</b>	<b>20</b>
3.1	Preliminary measurements . . . . .	20
3.1.1	XPS spectra at $\phi = 0$ . . . . .	20
3.1.2	Grounding of the surface . . . . .	22
3.1.3	Verification of surface contaminants . . . . .	26
3.1.4	Direct measurement of the surface potential . . . . .	27
3.2	Exposition to the NO <sub>2</sub> /O <sub>2</sub> gas mixture. . . . .	32
3.2.1	Sample never polarized . . . . .	33

3.2.2	Sample polarized only during gas exposition . . . . .	36
3.2.3	Sample permanently polarized . . . . .	39
3.3	Additional experiments . . . . .	41
3.3.1	Thiols on the SnO <sub>2</sub> surface . . . . .	41
3.3.2	Titanium dioxide film on the SnO <sub>2</sub> surface . . . . .	43
3.3.3	Topographical characterization . . . . .	45
<b>4</b>	<b>Discussion</b>	<b>46</b>
4.1	Grounding of the surface of the sample . . . . .	46
4.2	Argon erosion . . . . .	47
4.3	QMA . . . . .	48
4.4	NO <sub>x</sub> and H <sub>2</sub> O interaction . . . . .	48
4.5	Adsorption and desorption kinetics . . . . .	49
4.6	Desorption and temperature . . . . .	50
4.7	Thiols on SnO <sub>2</sub> . . . . .	50
4.8	TiO <sub>2</sub> film on SnO <sub>2</sub> . . . . .	51
4.9	Future work . . . . .	51
	<b>Conclusion</b>	<b>52</b>
	<b>Bibliography</b>	<b>53</b>

# List of Tables

1.1	Values for the various constants used on equation (1.6) . . . . .	7
2.1	Relative percentages of gases in chamber . . . . .	16
2.2	Relative percentages of gases in chamber during flooding . . . . .	16
3.1	Peak positions, FWHM and chemical contribution at 0 V . . . . .	22
3.2	Peak shift as function of the polarization voltage $\phi$ . . . . .	25
3.3	Binding energy (eV) of the N 1s peaks. . . . .	33
3.4	Fitted values for the exponential function parameters. The function used was $f(t) = y_0 + A \exp(-R_0 t)$ . . . . .	35
3.5	N 1s, NO <sub>2</sub> and NO <sub>3</sub> relative % present on the sample at t=2.5 min as function of V and the current through the sample . . . . .	38
3.6	Values of the fitting parameters. The function used was $f(t) = y_0 + Ae^{-R_0 t}$ . . . . .	40
3.7	Binding energy for each Gauss-Lorenz adjustment curve, its relative percentage compared to the whole S signal and its interpretation as found in the literature . . . . .	43
3.8	Ti and Sn peak positions . . . . .	44



# List of Figures

1.1	Band bending for different semiconductors. <b>a)</b> and <b>b)</b> show the energy diagram for a n-type semiconductor before and after adsorption equilibrium, while <b>c)</b> and <b>d)</b> present the energy diagram for a p-type semiconductor before and after adsorption equilibrium . . . . .	6
1.2	Sketch of the electroadsorptive effect for <b>a</b> induced electric field $E$ pointing towards the sample bottom and <b>b</b> induced electric field $E$ pointing towards the sample surface. . . . .	8
1.3	Diagram of a thiol on a substrate. The head group (red), tail group (blue) and terminal group (green) are shown. . . . .	9
1.4	Diagram of AFM working principle . . . . .	10
1.5	Diagram of the geometry used for LEIS. . . . .	11
1.6	Diagram of XPS working principle . . . . .	11
2.1	Experimental setup schematics . . . . .	13
2.2	Side view of the insertion arm. . . . .	14
2.3	Pictures of the sample holder. The red line shows its scale. . . . .	15
2.4	Sample, right after contact evaporation . . . . .	17
2.5	Sample with the point contacts used for the surface potential measurement . . . . .	17
2.6	Schematics of the circuit used. . . . .	18
3.1	Survey spectrum when the sample is not polarized. . . . .	20
3.2	High resolution spectra of the non polarized sample. . . . .	21
3.3	Expected behavior for a MOS device with small contacts on top . . . . .	22
3.4	Schematics of the different circuits tested . . . . .	23
3.5	XPS spectra of <b>a</b> C 1s , <b>b</b> O 1s and <b>c</b> Cu 2p <sub>3/2</sub> at different polarization $\phi$ values. . . . .	24
3.6	Sn 3d spectra at -15, 0 and +15 V. The pink dashed line corresponds to the Gauss-Lorenz curve used for adjustment. . . . .	26
3.7	Sn 3d spectra at -12, 0 and +12V. The pink dashed line corresponds to the Gauss-Lorenz curve used for adjustment. . . . .	26
3.8	LEIS spectra of a sample with evaporated contacts. The associated element to each peak is labelled on top. . . . .	26
3.9	Contour line plot for $\phi = -5$ (top) and $\phi = +5$ (bottom). . . . .	28
3.10	Contour line plot for $\phi = -9$ (top) and $\phi = +9$ (bottom). . . . .	29
3.11	Contour line plot for $\phi = -12$ (top) and $\phi = +12$ (bottom). The voltage is measured in mV . . . . .	30
3.12	Contour line plot for $\phi = -15$ (top) and $\phi = +15$ (bottom). . . . .	31

3.13	N 1s XPS spectra recorded at <b>a</b> $t=2,5$ min, <b>b</b> $t=20$ min and <b>c</b> $t=42,5$ min after the measurement started. . . . .	34
3.14	Mean curve obtained for $\text{NO}_2 + \text{NO}_3$ concentration onto the $\text{SnO}_2$ surface as function of time. The exponential fit is shown as a red line . . . . .	35
3.15	N 1s XPS spectra recorded after exposition to the gas mixture while polarized at <b>a)</b> $\phi = +15$ V, <b>b)</b> $\phi = 0$ V and <b>c)</b> $\phi = -15$ V. . . . .	36
3.16	N percentage as a function of initial polarization $\phi$ . . . . .	37
3.17	$\text{NO}_2$ percentage as a function of initial polarization $\phi$ . . . . .	37
3.18	$\text{NO}_3$ percentage as a function of initial polarization $\phi$ . . . . .	37
3.19	$\text{NO}_2 + \text{NO}_3$ percentage as a function of initial polarization $\phi$ . . . . .	37
3.20	N 1s XPS spectra recorded after exposition to the gas mixture while maintaining $\phi$ at <b>a)</b> $+8$ V, <b>(b)</b> $-8$ V, <b>(c)</b> $+15$ V, <b>(d)</b> $-15$ V. . . . .	39
3.21	$\text{NO}_2 + \text{NO}_3$ relative area as function of time at $\phi = \{\pm 8, +10, \pm 12 \text{ and } \pm 15\}$ . The colored line shows the respective exponential fit. . . . .	40
3.22	XPS high-resolution spectra of the S signal on the sample right after preparation, after a $+15$ V potential was applied to the sample backside and after a $-15$ V potential was applied to the sample backside. . . . .	42
3.23	LEIS spectra of a sample with a 1 nm Ti film on top. . . . .	43
3.24	XPS spectra of the $\text{TiO}_2$ coated sample at $\phi = +15$ (red), 0 (black) and $-15$ V (blue) . . . . .	44
3.25	AFM image of the $\text{SnO}_2$ sample surface before <b>(a)</b> ) and after <b>(b)</b> ) gas exposure while its back was polarized at different $\phi$ . . . . .	45
3.26	Height histogram of a sample before <b>(a)</b> ) and after <b>(b)</b> ) gas exposure while its back was polarized at different $\phi$ . . . . .	45
4.1	<b>a)</b> Highest deviation from zero surface potential as function of $\phi$ . <b>b)</b> Point where the highest deviation was found for all $\phi$ (area marked in blue) . . . .	46
4.2	Sketch of the proposed grid contact for grounding the surface of the sample .	47
4.3	Left: This work proposed $\text{SnO}_2$ surface after adsorption of $\text{NO}_x$ species. Right: Top view of the two possible configurations for water adsorption on a saturated $\text{SnO}_2$ (110) surface [1]. . . . .	49