

Frequency, determinants, and effects of early seizures after thrombolysis for acute ischemic stroke The ENCHANTED trial

Por: [Xu, Y](#) (Xu, Ying)^[1,2]; [Hackett, ML](#) (Hackett, Maree L.)^[1,2]; [Chalmers, J](#) (Chalmers, John)^[1,2]; [Lindley, RI](#) (Lindley, Richard I.)^[1,2]; [Wang, X](#) (Wang, Xia)^[1]; [Li, Q](#) (Li, Qiang)^[1]; [Robinson, T](#) (Robinson, Thompson)^[3,4]; [Arima, H](#) (Arima, Hisatomi)^[1,5]; [Lavados, PM](#) (Lavados, Pablo M.)^[6,7]; [Anderson, CS](#) (Anderson, Craig S.)^[1,2,8,9]

Autoría conjunta: [ENCHANTED Study Grp](#)

[Ver ResearcherID y ORCID](#)

NEUROLOGY-CLINICAL PRACTICE

Volumen: 7

Número: 4

Páginas: 324-332

DOI: 10.1212/CPJ.0000000000000384

Fecha de publicación: AUG 2017

Tipo de documento: Article

Resumen

Background: Seizures after ischemic stroke have not been well-studied. We aim to determine the frequency, determinants, and significance of early seizures after thrombolysis for acute ischemic stroke. Methods: Data are from the Enhanced Control of Hypertension and Thrombolysis Stroke Study (ENCHANTED), an international, multicenter, randomized controlled trial where patients with acute ischemic stroke were randomized to low-dose (0.6 mg/kg) or standard-dose (0.9 mg/kg) IV alteplase. The protocol prespecified prospective data collection on in-hospital seizures over 7 days postrandomization. Logistic regression models were used to determine variables associated with seizures and their significance on poor outcomes of death or disability (modified Rankin scale scores 3-6), symptomatic intracerebral hemorrhage (sICH), and European Quality of Life 5-Dimensions questionnaire [EQ-5D] over 90 days. Results: Data were available for 3,139 acute ischemic stroke participants, of whom 42 (1.3%) had seizures at a median 22.7 hours after the onset of symptoms. Baseline variables associated with seizures were male sex (odds ratio [OR] 2.19, 95% confidence interval [CI] 1.07-4.50), severe neurologic impairment (NIH Stroke Scale score \geq 10; OR 2.16, 95% CI 1.06-4.40), and fever (OR 4.55, 95% CI 2.37-8.71). Seizures independently predicted poor recovery: death or major disability (OR 2.88, 95% CI 1.28-6.47), unfavorable ordinal shift of mRS scores (OR 1.94, 95% CI 1.10-3.39), and lower than median EQ-5D health utility index score (OR 3.50, 95% CI 1.37-8.91). There was no association of seizures with sICH in adjusted analysis. Conclusions: In thrombolysis-treated patients with acute ischemic stroke, seizures are uncommon, occur early, and predict poor recovery.

Palabras clave

KeyWords Plus:[POSTSTROKE SEIZURES](#); [PREDICTIVE FACTORS](#); [RISK-FACTORS](#); [MULTICENTER](#); [ALTEPLASE](#); [EPILEPSY](#); [IMPACT](#)

Información del autor

Dirección para petición de copias: Anderson, CS (autor para petición de copias)

+ Univ New South Wales, George Inst Global Hlth, Sydney, NSW, Australia.

Dirección para petición de copias: Anderson, CS (autor para petición de copias)

+ Univ Sydney, Sydney Med Sch, Sydney, NSW, Australia.

Dirección para petición de copias: Anderson, CS (autor para petición de copias)

+ Royal Prince Alfred Hosp, Dept Neurol, Sydney, NSW, Australia.

Dirección para petición de copias: Anderson, CS (autor para petición de copias)

+ Peking Univ, Hlth Sci Ctr, George Inst China, Beijing, Peoples R China.

Direcciones:

+ [1] Univ New South Wales, George Inst Global Hlth, Sydney, NSW, Australia

+ [2] Univ Sydney, Sydney Med Sch, Sydney, NSW, Australia

+ [3] Univ Leicester, Dept Cardiovasc Sci, Leicester, Leics, England

+ [4] Univ Leicester, NIHR Biomed Res Unit Cardiovasc Dis, Leicester, Leics, England

+ [5] Fukuoka Univ, Fac Med, Dept Prevent Med & Publ Hlth, Fukuoka, Japan

+ [6] Univ Desarrollo, Clin Alemana, Fac Med, Clin Alemana Santiago, Concepcion, Chile

+ [7] Univ Chile, Fac Med, Dept Ciencias Neurol, Santiago, Chile

+ [8] Royal Prince Alfred Hosp, Dept Neurol, Sydney, NSW, Australia

+ [9] Peking Univ, Hlth Sci Ctr, George Inst China, Beijing, Peoples R China

Direcciones de correo electrónico:canderson@georgeinstitute.org.cn

Financiación

Entidad financiadora	Número de concesión
National Health and Medical Research Council (NHMRC) of Australia	1020462
Stroke Association of the United Kingdom	TSA 2012/01
National Council for Scientific and Technological Development of Brazil (CNPq)	467322/2014-7
Ministry for Health, Welfare, and Family Affairs of South Korea	

[Ver texto de financiación](#)

Editorial

LIPPINCOTT WILLIAMS & WILKINS, TWO COMMERCE SQ, 2001 MARKET ST,
PHILADELPHIA, PA 19103 USA

Categorías / Clasificación

Áreas de investigación:Neurosciences & Neurology

Categorías de Web of Science:Clinical Neurology

Información del documento

Idioma:English

Número de acceso: WOS:000423488100012

ID de PubMed: 29185557

ISSN: 2163-0402

eISSN: 2163-0933