

# Changes in cyclic AMP dependent protein kinase and active stiffness in the rat volume overload model of heart hypertrophy

Lavandero, Sergio

Cartagena, Gastón

Guarda, Eduardo

Corbalán, Ramón

Godoy, Iván

Sapag-Hagar, Mario

Jalil, Jorge E.

**Objective:** The aim was to clarify the role of cyclic AMP dependent protein kinase (PKA) and changes in mechanical heart function during development of cardiac hypertrophy induced by volume overload. **Methods:** Protein and DNA contents, PKA activity, and peak systolic stress-strain relationships in hearts from animals submitted to aortocaval shunt were assessed as a function of time. Sham operated (control) rats were used as controls. **Results:** Heart weight to body weight ratio and cardiac protein content per heart increased from d 7 ( $p < 0.005$  and  $p < 0.01$ , respectively) reaching their highest values by d 56; the same occurred with cardiac DNA content. PKA activity·g<sup>-1</sup> tissue in soluble extracts of hearts from rats with aortocaval shunt increased by 2.7-fold on d 2 ( $p < 0.005$ ), reached a ninefold peak increase by d 7 ( $p < 0.0001$ ) and declined to fourfold by d 56 with respect to control values. The end peak systolic stress-strain relation slopes were: control, 368(SEM 14) g·cm<sup>-2</sup> (n=16); ao