

An Efficient Synthesis of Bis(calix[4]arenes), Bis(crown ether)-Substituted Calix[4]arenes, Aza-Crown Calix[4]arenes, and Thiaza-Crown Calix[4]arenes

Francisco Santoyo-González,^{*[a]} Antonio Torres-Pinedo,^[a] and Claudio Saitz Barria^[b]

Keywords: Bis(calixarenes) / Calixarenes / Crown compounds / Isothiocyanates / Macrocycles

The easy transformation of 5,11,17,23-tetra-*tert*-butyl-25,27-bis(aminoethoxy)-26,28-dihydroxycalix[4]arene (**2**) into 5,11,17,23-tetra-*tert*-butyl-26,28-dihydroxy-25,27-bis(2-isothiocyanoethoxy)calix[4]arene (**3**) and 5,11,17,23-tetra-*tert*-butyl-25,27-bis(chloroacetamidoethoxy)-26,28-dihydroxycalix[4]arene (**4**) has been exploited for the development of an efficient and expeditious synthesis of a variety of calix[4]arene derivatives [bis(calix[4]arenes), bis(crown ether) calix[4]arenes, aza-crown calix[4]arenes, and thiaza-crown

calix[4]arenes]. The functionality of compounds **2** and **3** allowed the formation of intramolecular bridges, leading to capped calix[4]arenes (compounds **5**, **10**, and **14**) as well as to the construction of double calix[4]arene units by means of spacers containing thiourea or amide-sulfur groups (compounds **6** and **15**, respectively). In addition, the bis(isothiocyanate) derivative **3** gave access to a high-yield preparation of heteroditopic bis(crown ether) calix[4]arenes (**7–9**) from commercial amino crown ethers.

Introduction

The development of supramolecular chemistry has led to a growing interest in the design and synthesis of macrocyclic molecules incorporating intramolecular cavities.^[1] Calixarenes^[2–17] have been used as building blocks for the synthesis of large host molecules with different supramolecular functions. Like crown ethers and cyclodextrins, calixarenes are oligomeric macrocyclic molecules made up of phenol units and incorporating a cavity. They offer the advantage of their easy synthesis on a large scale in a simple one-pot procedure from inexpensive starting material^[14–17] and, in addition, are readily amenable to chemical modification on both smaller (lower) and larger (upper) rims, by attachment of a wide range of potential ligating groups.

The versatility of calix[4]arenes as host molecules suggested that they might serve as potential building blocks for designing more elaborate structures, consisting of double calix[4]arenes.^[18,19] Depending on the connecting site, there are three possible modes of assembly for these double calix[4]arenes: *head-to-head*, *tail-to-tail*, or *head-to-tail*. Or, put another way, by both larger (upper) rims, by both smaller (lower) rims, or by the larger (upper) rim of one and the smaller (lower) rim of another, respectively. Shinkai et al.^[20–22] have reported the synthesis of a series of bis(calix[4]arenes) possessing two metal binding sites, each containing four ester or four ether groups. Double and triple calix[4]arenes connected through the oxygen atoms have

been described,^[23] produced by reactions between calix[4]arenes 1,3-difunctionalized at the smaller (lower) rim and difunctional reagents such as diacid dichlorides or diamines.^[23,24] Double calixarenes linked through one bridge at the smaller (lower) rim and bearing urea groups on the larger (upper) rims have also been reported.^[25,26] Recently, Rebek et al. have described^[27] the synthesis and the encapsulation behaviour of bis(calix[4]arenes) linked by one bridge at the larger (upper) rims and bearing urea groups in the same rims. Bis(calix[*n*]arenes) singly and doubly bridged at the larger (upper) rims by 2-butenyl or 2-methylenepropyl moieties have been prepared by tandem Claisen rearrangement of bis(calix[*n*]arenes) singly and doubly bridged by ether linkages at the smaller (lower) rims with the same spacers. The complexation behaviour of these bis(calixarenes) toward fullerenes C60 and C70 has been studied.^[28] A series of bis(calix[4]arene) derivatives linked through their phenolic oxygens with the aid of a single aliphatic or aromatic chain (*tail-to-tail*) have been obtained by alkylation of 28-hydroxy-25,26,27-triisopropoxycalix[4]arenes with α,ω -dibromoalkanes in the presence of NaH,^[29] and by condensation of *p-tert*-butylcalix[4]arene with methyl 2,6-bis(bromomethyl)benzoate,^[30] 2,6-bis(bromomethyl)-4-methylanisole,^[31] or with 5,5'-bis(bromomethyl)-2,2'-bipyridine *N,N'*-dioxide.^[32] Bis- and oligocalix[4]arenes have been obtained by intermolecular metathesis of dialkenylcalix[4]arene derivatives.^[33]

The term “calixcrowns” refers to the family of macrocyclic molecules in which a monocyclic calixarene structure is linked through its phenolic oxygens to a cyclic polyether moiety, hence combining calixarene and polyether elements in their structures^[8,10,11,19]. Since first reported by Ungaro,^[34] these compounds have emerged as a new, exciting class of ionophores, thanks to their tendency – strongly dependent on their conformation – to bind alkali and alkaline earth metal cations selectively.^[35–43] In the past dec-

^[a] Instituto de Biotecnología, Departamento de Química Orgánica, Facultad de Ciencias, Campus Fuentenueva s/n, Universidad de Granada, Granada, 18071-Spain
Fax: (internat.) +34-95/824-3187
E-mail: fsantoyo@ugr.es

^[b] Visiting Professor at the University of Granada, Departamento de Química Orgánica y Físico Química, Facultad de Ciencias Químicas y Farmacéuticas, Universidad de Chile, Casilla 233, Santiago, Chile


ade, the design of new ditopic receptors for the simultaneous complexation of anionic and cationic guest species has become an active new area of coordination chemistry, of interest for the selective extraction and/or transportation across lipophobic membranes, of metal salts. In this area, Beer et al.^[44] reported the synthesis and coordination chemistry of new heteroditopic calix[4]arene compounds containing benzo-15-crown-5 units linked through amides at the smaller (lower) rim. As well as this, it should be also mentioned that hydrogen-bonding donor thiourea functions have been used for the design of anion receptors.^[45–62]

In this paper, we report the synthesis from the easily accessible 25,27-bis(aminoethoxy)calix[4]arene (**2**) of a series of *tail-to-tail* bis(calix[4]arenes), bis(crown ether) calix[4]arenes, aza-crown calix[4]arenes and thiaza-crown calix[4]arenes.

Results and Discussion


In a project directed towards synthetic applications of the isothiocyanate function,^[63,64] it was proposed that isothiocyanate derivatives of calixarenes could be suitable precursors for the easy construction of calixarene-based heterotopic receptors. As the starting material, 25,27-bis(aminoethoxy)calix[4]arene (**2**) was chosen. This is readily obtained in two steps from *p*-*tert*-butylcalix[4]arene (**1**), by reaction with bromoacetonitrile and reduction of the cyano function with LiAlH_4 .^[65] We performed the transformation of **2** into the corresponding diisothiocyanate derivative **3** in excellent yields, using thiophosgene as reported by García-Fernández et al.^[66] (see Scheme 1). Previous observations, from our group and others, have demonstrated^[64] that isocyanates and isothiocyanates are readily transformed into the corresponding symmetric *N,N'*-disubstituted ureas and thioureas by a simple treatment with pyridine–water. Application of those conditions to compound **3** resulted in the isolation of the capped calix[4]arene **5**, in 52% yield (see Scheme 2). The formation of this compound is explained by the intermolecular reaction of the isothiocyanate functions, competing with the intramolecular process.

Assembly of double calixarene units linked by thiourea bridges could be effected by direct reaction of the diamino derivative **2** with the diisothiocyanate derivative **3**. The reaction was performed in pyridine under high dilution conditions. In this way, the bis(calix[4]arene) **6** was the only reaction product formed, isolated in good yields (60%) (see Scheme 2). The reactivity of isothiocyanates with amines also found application in the construction of heteroditopic structures base on compound **3**. The synthesis of bis(crown ether)-substituted calix[4]arenes was accomplished by treatment of that compound with commercial amino crown-ether derivatives (4'-aminobenzo-15-crown-5, 4'-aminobenzo-18-crown-6, and 2-(aminomethyl)-18-crown-6). We carried out these reactions in ethyl acetate in the presence of triethylamine, allowing easy coupling (94–98% yield) of the two different subunits, followed by the isolation of compounds **7–9** (see Scheme 2).




Scheme 1. Reagents and conditions: (i) see ref.^[67]; (ii) Cl_2CS , BaCO_3 , CH_2Cl_2 , room temp.; (iii) $(\text{ClCH}_2\text{CO})_2\text{O}$, AcOEt, room temp.

Because of the faculty of those groups to suffer nucleophilic displacement reactions, the transformation of the amino functions of compound **2** into the corresponding chloroacetamido groups was then investigated, as a potential gateway to the preparation of different calix[4]arene derivatives. Treatment of **2** with chloroacetic anhydride, using ethyl acetate as solvent, led to the dichloroacetamido derivative **4** in high yield (84%) (see Scheme 1). The two arms of that compound were linked by treatment with Na_2S in refluxing methanol at high dilution, yielding the capped calixarene derivative **10** in moderate yield (47%) (see Scheme 3). The next goal was the construction of an analogous compound, but with a disulfide bridge. Bis(crown ether)s containing disulfide bridges have attracted attention as redox-switchable systems, for their utility in ion extraction and ion transport across liquid membranes.^[67,68] Cyclic disulfides are typically prepared by oxidative dimerisation of α,ω -dithiols,^[69–75] or α,ω -bisthiocyanates.^[76–79] For this reason, we first carried out the synthesis of the bisthiocyanate derivative **11** (85%), by treatment of **4** with KSCN in refluxing acetone (see Scheme 3). Our attempts to obtain the disulfide bridge using the conditions previously described^[79] were unsuccessful, however, in spite of the good yields reported by Field et al. for this reaction. Therefore, we considered a new strategy, entailing the synthesis of the bisdithiol derivative **13** via the corresponding bis(acetylthio) derivative **12**. That compound was easily obtained by reaction of **4** with KSAc in refluxing acetone (73%) and was then *S*-deacetylated, using NaOMe in MeOH (95%). Complete spontaneous transformation of compound **13** into the desired disulfide **14** was then observed during the purification process of **13** by standard column chromatography. Compound **14** was isolated in 60% overall yield from **12** (see Scheme 3).



Scheme 2. Reagents and conditions: (i) Py, H₂O, Et₃N, 60 °C; (ii) **2**, Py, room temp.; (iii) amino crown-ether, AcOEt, room temp.



Scheme 3. Reagents and conditions: (i) Na₂S · 9 H₂O, MeOH, reflux; (ii) KSCN or KSAc, acetone, reflux; (iii) NaOMe/MeOH, room temp.

The in situ generation of the dithiol **13** from the bis(acylthio) derivative **12** in the presence of the dichloroacetamide derivative **4** gave rise to bis(calix[4]arene) **15**. The reaction was performed using NaOMe/MeOH at room temp., yielding **15** in 52% yield (see Scheme 3).

In all cases the structures of the new calix[4]arene derivatives **3–15** were established by elemental analysis and/or high resolution FAB spectroscopy and from spectroscopic data (IR and NMR) (see Experimental Section).

Conclusions

In summary, the construction of a variety of “tailor-made” calix[4]arene derivatives (biscalix[4]arenes, bis(crown ether) calix[4]arenes, aza-crown calix[4]arenes, and thiaza-crown calix[4]arenes) has been achieved easily from 25,27-bis(aminoethoxy)calix[4]arene **2**. The key step in these synthesis was the direct transformation of compound **2** into the corresponding bis(isothiocyanate) and bis(chloroacet-

amido) derivatives **3** and **4**. The functionality of these compounds permitted the formation of intramolecular bridges, leading to capped calix[4]arenes (compounds **5**, **10**, and **14**) as well as to the assembly of double calix[4]arene units by means of spacers containing thiourea or amide-sulfur groups (compounds **6** and **15**, respectively). In addition, the bis(isothiocyanate) derivative **3** gave access to a high-yielding preparation of heteroditopic bis(crown ether) calix[4]arenes (**7–9**) from commercial amino crown ethers.

Experimental Section

General Experimental Details: TLC was performed on Merck silica gel 60F₂₄₅ aluminium sheets, with detection by Mostain reagent [ceric sulfate (1% w/v and ammonium molybdate (2.5% w/v) in 10% (v/v) aqueous sulfuric acid] and by UV light when applicable. Flash column chromatography on Merck or Scharlau silica gel (230–400 mesh, ASTM). – All evaporations were carried out under reduced pressure at 40 °C. – Melting points were determined with a Gallenkamp apparatus and are uncorrected. – NMR spectra were recorded at room temperature on a Bruker AM-300 spectrometer. ¹H chemical shifts are given in ppm and referenced to internal CHCl₃ (δ = 7.26) for CDCl₃ solutions. ¹³C chemical shifts are given in ppm and referenced to CDCl₃ (δ = 77.0). *J* values are given in Hz. – FAB mass spectra were obtained on a Fisons VG Autospec-Q spectrometer, using *m*-nitrobenzyl alcohol or thioglycerol as matrix. – Anhydrous solvents were prepared according to standard procedures and were freshly distilled prior to use. For reasons of clarity and to reduce space, the names calix[4]arene (pentacyclo-[19.3.1.1^{3,7}.1^{9,13}.1^{15,19}]octacos-1(25),3,5,7(28),9,11,13(27),15,17,19(26),21,23-dodecaene), benzo-15-crown-5 (2,3,5,6,8,9,11,12-octahydro-1,4,7,10,13-benzopentaoxacyclopentadecine), and benzo-18-crown-6 (2,3,5,6,8,9,11,12,14,15-decahydro-1,4,7,10,13,16-benzohexaoxacyclooctadecine) have been used instead of the original IUPAC names.

5,11,17,23-Tetra-*tert*-butyl-26,28-dihydroxy-25,27-bis(2-isothiocyanatoethoxy)calix[4]arene (3): To a solution of **2** (0.5 g, 0.68 mmol) in CH₂Cl₂ (10 mL) was added thiophosgene (0.15 mL, 2.04 mmol), BaCO₃ (0.8 g, 4.1 mmol), and H₂O (3 mL). The reaction was stirred for 24 h at room temp. After filtration, CH₂Cl₂ (10 mL) was added and the system was extracted with H₂O (10 mL). The organic phase was dried and evaporated, yielding **3** (0.535 g, 96%) as a solid: M.p. 267–270 °C (dec). – IR (KBr): $\tilde{\nu}$ = 3393, 2193, 2097, 1485, 1346, 1302, 1202, 1126, 1045, 874 cm⁻¹. – ¹H NMR (300 MHz, CDCl₃): δ = 7.07 (s, 4 H; ArH), 6.92 (s, 2 H; OH), 6.82 (s, 4 H; ArH), 4.25 (d, *J* = 13.0 Hz, 4 H; ArCH₂Ar), 4.20–4.14 (m, 8 H; 2 OCH₂CH₂NCS), 3.36 (d, *J* = 13.0 Hz, 4 H; ArCH₂Ar), 1.26, 0.97 (2 s, 36 H; 4 Me₃C). – ¹³C NMR (75 MHz, CDCl₃): δ = 150.6, 149.3, 147.5, 141.8, 132.3, 127.7, 125.9, 125.3 (Ar, NCS), 73.4 (OCH₂), 45.1 (CH₂NCS), 33.9 (CMe₃), 31.8, 31.1 (ArCH₂Ar, Me₃C). – HR-MS (FAB+) C₅₀H₆₂N₂O₄NaS₂: calcd for [M + Na]⁺ 841.4048; found: 841.4051. – C₅₀H₆₂N₂O₄S₂ (819.2): calcd. C 73.31, H 7.63, N 3.42, S 7.83; found C 73.30, H 8.10, N 3.53, S 8.10.

5,11,17,23-Tetra-*tert*-butyl-25,27-bis{(chloromethyl)carbonylamino}ethoxy-26,28-dihydroxycalix[4]arene (4): To a solution of **2** (0.7 g, 0.95 mmol) in AcOEt (50 mL) was added chloroacetic anhydride (0.36 g, 2.09 mmol). The reaction was kept at room temp. and monitored by TLC (AcOEt) until complete disappearance of **2** (1 h). The reaction mixture was washed with aqueous

saturated NaHCO₃ solution (2 × 15 mL). The organic phase was dried and evaporated, giving a crude product that was purified by short column chromatography (CH₂Cl₂/hexane 2:1), yielding **4** (0.71 g, 84%) as a solid: M.p. 246–247 °C. – IR (KBr): $\tilde{\nu}$ = 3422, 1665, 1541, 1485, 1362, 1202 cm⁻¹. – ¹H NMR (300 MHz, CDCl₃): δ = 8.37 (br s, 2 H; 2 NH), 8.11 (s, 2 H, 2 OH), 7.05, 6.95 (2 s, 8 H; Ar), 4.20 (d, *J* = 13.0 Hz, 4 H; ArCH₂Ar), 4.12 (t, *J* = 4.7 Hz, 4 H; 2 OCH₂), 4.11 (s, 4 H; 2 CH₂Cl), 3.97 (m, 4 H; 2 CH₂NH), 3.39 (d, *J* = 13.0 Hz, 4 H; ArCH₂Ar) 1.26, 1.07 (2 s, 36 H; 4 Me₃C). – ¹³C NMR (75 MHz, CDCl₃): δ = 166.9 (CO), 149.8, 148.6, 148.1, 142.6, 132.9, 127.6, 126.1, 125.6 (Ar), 75.1 (OCH₂), 42.7 (NHCH₂), 40.0 (CH₂Cl), 34.2, 33.9 (Me₃C), 32.1 (Ar-CH₂Ar), 31.7, 31.1 (Me₃C). – HR-MS (FAB+) C₅₂H₆₈Cl₂N₂O₆Na: calcd. for [M + Na]⁺ 909.4352; found 909.4357. – C₅₂H₆₈Cl₂N₂O₆ (888.0): calcd. C 70.33, H 7.72, N 3.15; found C 70.15, H 7.99, N 2.87.

1,7-(5,11,17,23-Tetra-*tert*-butyl-26,28-dihydroxy-25,27-dioxy-calix[4]arene)-3,5-diaza-4-heptanethione (5): To a solution of **3** (0.164 g, 0.2 mmol) in Py/H₂O (18:2 mL) was added triethylamine (0.1 mL). The reaction mixture was heated at 60 °C for 24 h. Evaporation and coevaporation with toluene gave a crude product that was triturated with ether/hexane (1:1), giving **5** (0.08 g, 52.5%) as a solid after filtration: M.p. 140–142 °C. – IR (KBr): $\tilde{\nu}$ = 3388, 3239, 3180, 1554, 1485, 1295, 1252, 1199, 1034 cm⁻¹. – ¹H NMR (300 MHz, CDCl₃): δ = 8.90 (s, 2 H; 2 OH), 7.30 (br s, 2 H; 2 NH), 7.07, 7.01 (2 s, 8 H; Ar), 4.40 (br s, 4 H; 2 OCH₂), 4.30 (d, *J* = 12.9 Hz, 4 H; ArCH₂Ar), 4.15 (br s, 4 H; 2 CH₂NH), 3.41 (d, *J* = 12.9 Hz, 4 H; ArCH₂Ar), 1.21, 1.19 (2 s, 36 H; 4 Me₃C). – ¹³C NMR (75 MHz, CDCl₃): δ = 188.5 (CS), 150.0, 149.5, 148.0, 142.5, 133.5, 127.8, 126.6, 125.5 (Ar), 73.5 (CH₂O), 45.4 (CH₂NH), 34.3, 34.0 (Me₃C), 33.2 (ArCH₂Ar) 31.6, 31.3 (Me₃C). – HR-MS (FAB+): C₄₉H₆₄N₂O₄NaS: calcd. for [M + Na]⁺ 799.4484; found: 799.4487. – C₄₉H₆₄N₂O₄S (777.1): calcd. C 75.13, H 8.30, N 3.6; found C 75.3, H 8.41, N 3.44.

25,25',27,27'-Bis[*N,N'*-bis(oxyethyl)thioureido]-bis(5,11,17,23-tetra-*tert*-butyl-26,28-dihydroxycalix[4]arene) (6): To a solution of **3** (0.082, 1 mmol) in Py (25 mL) was added **2** (0.073, 0.1 mmol). The reaction mixture was kept at room temp. for 30 h and monitored by TLC (chloroform/ether 3:1). Evaporation of the solvents gave a crude product that was triturated with MeOH to give **6** (0.093 g, 60%) as a solid: M.p. 165–167 °C. – IR (KBr): $\tilde{\nu}$ = 3385, 1637, 1545, 1475, 1298, 1201, 1037 cm⁻¹. – ¹H NMR (400 MHz, CDCl₃): δ = 7.65 (br s, 8 H, 4 OH; 4 NH), 7.04, 6.81 (2 s, 16 H, ArH), 4.40–4.00 (m, 24 H; 4 OCH₂CH₂NH, ArCH₂Ar), 3.26 (d, *J* = 13.0 Hz, 8 H; ArCH₂Ar), 1.29, 1.05 (2 s, 72 H, 8 Me₃C). – ¹³C NMR (CDCl₃, 100 MHz): δ = 149.5, 148.9, 147.8, 142.8, 132.5, 127.9, 125.9, 125.3 (Ar), 77.3 (CH₂O), 45.3 (CH₂NH), 34.1, 34.0 (Me₃C), 31.7, 31.1 (ArCH₂Ar, Me₃C). – HR-MS (FAB+) C₉₈H₁₂₈N₄O₈S₂Na: calcd. for [M + Na]⁺ 1554.928 found 1554.933. – C₉₈H₁₂₈N₄O₈S₂ (1554.23): calcd. C 75.73, H 8.31, N 3.61, S 4.12; found C 75.30, H 8.58, N 3.66, S 4.08.

General Procedure for the Synthesis of Bis(crown ether)-Substituted Calix[4]arenes 7–9: To a solution of **3** (0.146 g, 0.178 mmol) in AcOEt (5 mL) was added the corresponding amino crown ether (0.39 mmol) and triethylamine (0.05 mL). The reaction was monitored by TLC (CH₂Cl₂/MeOH 10:1) until complete disappearance of the starting materials (24 h for **7** and **8**, and 3 h for **9**). AcOEt (50 mL) was then added and the system was washed with 5% HCl (25 mL) and H₂O (25 mL). The organic phase was dried and evaporated, giving a crude product.

5,11,17,23-Tetra-*tert*-butyl-25,27-bis[*N'*-(4'-benzo-15-crown-5)-thioureido]ethoxy-26,28-dihydroxycalix[4]arene (7): Column chro-

matography of the crude product (CH₂Cl₂/hexane 7:1) gave **7** (0.231 g, 94%) as a solid: M.p. 134–137 °C. – IR (KBr): $\tilde{\nu}$ = 3383, 1560, 1540, 1508, 1240, 1132 cm⁻¹. – ¹H NMR (300 MHz, CDCl₃): δ = 7.89 (s, 2 H; 2 OH), 7.80 (br s, 2 H; 2 × NH), 7.73 (s, 2 H; 2 × NH), 6.93 (s, 4 H; ArH), 6.90 (s, 4 H; ArH), 6.75 (d, J = 2.2 Hz, 2 H; HNC₆H₃), 6.64 (dd, J = 8.4, 2.2 Hz, 2 H; HNC₆H₃), 6.27 (d, J = 8.4 Hz, 2 H; HNC₆H₃), 4.14–3.71 [m, 44 H; 2 × OCH₂CH₂N, 2 × –O(CH₂CH₂O)₄–, ArCH₂Ar], 3.28 (d, J = 13.1 Hz, 4 H; ArCH₂Ar), 1.21, 1.08 (2 s, 36 H; 4 × Me₃C). – ¹³C NMR (75 MHz, CDCl₃): δ = 182.3 (CS), 149.6, 149.4, 148.5, 148.2, 147.9, 142.3, 133.1, 128.3, 127.8, 125.9, 125.5, 119.3, 113.4, 112.4 (Ar, C₆H₃), 75.2, 71.3, 71.2, 70.7, 69.6, 69.5, 69.0, 68.5 [OCH₂CH₂N, –O(CH₂CH₂O)₄–], 46.3 (CH₂N), 34.2, 33.8 (Me₃C), 32.1, 31.5 (ArCH₂Ar), 31.7 and 31.1 (Me₃C). – HR-MS (FAB+) C₇₈H₁₀₄N₄O₁₀NaS₂: calcd. for [M + Na]⁺ 1407.689; found 1407.686. – C₇₈H₁₀₄N₄O₁₄S₂ (1385.81): calcd. C 67.6, H 7.56, N 4.04; found C 67.12, H 7.81, N 3.91.

5,11,17,23-Tetra-tert-butyl-25,27-bis{[N'-(4'-benzo-18-crown-6)-thioureido]ethoxy}-26,28-dihydroxycalix[4]arene (8): Column chromatography of the crude product (CH₂Cl₂:hexane 7:1) gave **8** (0.257 g, 94%) as a solid (98%): M.p. 142–144 °C. – IR (KBr): $\tilde{\nu}$ = 3300, 1558, 1541, 1516, 1508, 1238, 1126 cm⁻¹. – ¹H NMR (300 MHz, CDCl₃): δ = 8.07 (s, 2 H; 2 OH), 7.90 (bt, J = 5.5 Hz, 2 H; 2 HNCH₂), 7.71 (s, 2 H; 2 NHC₆H₃), 6.93 (s, 4 H; ArH), 6.92 (s, 4 H; ArH), 6.77 (d, J = 2.2 Hz, 2 H; 2 C₆H₃), 6.61 (dd, J = 8.5, 2.2 Hz, 2 H; 2 C₆H₃), 6.20 (d, J = 8.5 Hz, 2 H; 2 C₆H₃), 4.17 (m, 4 H; 2 HNCH₂), 4.08 (m, 4 H; 2 OCH₂CH₂N), 4.00–3.97, 3.87–3.83, 3.79–3.71, 3.66–3.56 [4 m, 44 H; 2 –O(CH₂CH₂O)₅–, ArCH₂Ar], 3.30 (d, J = 13.0 Hz, 4 H; ArCH₂Ar), 1.21, 1.10 (2 s, 36 H; 4 Me₃C). – ¹³C NMR (75 MHz, CDCl₃): δ = 182.4 (CS), 149.6, 148.6, 148.0, 142.3, 133.3, 127.9, 126.0, 125.6, 119.3, 112.9, 112.2 (Ar, C₆H₃), 75.4, 71.0, 70.9, 70.8, 70.7, 69.6, 69.0, 68.4 [OCH₂CH₂N, –O(CH₂CH₂O)₅–], 46.6 (CH₂N), 34.2, 33.3 (Me₃C), 32.1 (ArCH₂Ar), 31.7, 31.2 (Me₃C). – HR-MS (FAB+) C₈₂H₁₁₂N₄O₁₆NaS₂: calcd. for [M + Na]⁺ 1495.7406; found 1495.7412. – C₈₂H₁₁₂N₄O₁₆S₂ (1473.92): calcd. C 66.82, H 7.66, N 3.80, S 4.35; found C 66.63, H 7.89, N 3.71, S 3.75.

5,11,17,23-Tetra-tert-butyl-26,28-dihydroxy-25,27-bis{[N'-(2-methyl-18-crown-6)thioureido]ethoxy}calix[4]arene (9): Treatment of the crude product with ether:hexane yielded **9** (0.242 g, 97%) as an amorphous solid: M.p. 70–72 °C. – ¹H NMR (300 MHz, CDCl₃): δ 8.15 (br s, 2 H, 2 NH) 7.05 (br s, 2 H, 2 NH), 7.03 (s, 6 H, 2 OH, ArH), 6.89 (s, 4 H, ArH), 4.23 (d, J = 13.0 Hz, 4 H; ArCH₂Ar), 4.18 (m, 4 H; 2 CH₂O), 3.77–3.44 [m, 54 H; 2 CH₂NHC(S)NHCH₂, 2 × (CH₂CH₂O)₅CH₂CHO], 3.36 (d, 4 H, J = 12.8 Hz, 2 H; ArCH₂Ar), and 1.24, 1.04 (2 s, 36 H; 4 Me₃C). – ¹³C NMR (300 MHz, CDCl₃): δ 183.0 (CS), 149.5, 149.0, 147.7, 142.7, 132.8, 128.0, 125.9, 125.5 (Ar), 77.0, 75.5, 71.7, 70.7, 70.4, 70.2, 68.9 [(CH₂CH₂O)₅CH₂CHO, OCH₂CH₂N], 45.4, 44.5 (OCH₂CH₂N, CH₂N), 34.1, 33.9 (Me₃C), 32.0, 29.7 (ArCH₂Ar), 31.7, 31.0 (Me₃C). – HR-MS (FAB+) C₇₆H₁₁₆N₄O₁₆NaS₂: calcd. for [M + Na]⁺ 1427.8; found 1427.9.

1,11-(5,11,17,23-Tetra-tert-butyl-26,28-dihydroxy-25,27-dioxycalix[4]arene)-6-thia-3,9-diaza-4,8-undecanedione (10): A solution of **4** (0.1 g, 0.113 mmol) and Na₂S · 9 H₂O (0.02 g, 0.250 mmol) in MeOH (60 mL) was refluxed for 24 h. Evaporation of the solvent gave a crude product that was dissolved in AcOEt (50 mL) and washed with H₂O (2 × 10 mL). The organic phase was dried and evaporated, giving a crude product that was purified by column chromatography (AcOEt) to yield **10** (0.045 g, 47%) as a solid: M.p. 297–230 °C (dec) (from ether/hexane). – IR (KBr): $\tilde{\nu}$ = 3424, 3298, 1672, 1650, 1560, 1485, 1461, 1362, 1202, 1103 cm⁻¹. – ¹H

NMR (300 MHz): δ = 8.25 (br s, 2 H; 2 NH), 7.81 (s, 2 H; 2 OH), 7.08, 6.94 (2 s, 8 H; Ar), 4.23 (d, J = 13.0 Hz, 4 H; ArCH₂Ar), 4.15 (t, J = 4.4 Hz, 4 H; 2 OCH₂), 3.97 (m, 4 H; 2 CH₂N), 3.45 [s, 4 H; (CH₂)₂S], 3.40 (d, J = 13.0 Hz, 4 H; ArCH₂Ar), 1.28, 1.06 (2 s, 36 H; 4 Me₃C). – ¹³C NMR (CDCl₃, 75 MHz): δ = 168.6 (CO), 149.9, 148.8, 147.9, 142.6, 132.7, 127.8, 126.0, 125.5, (Ar), 75.1 (OCH₂), 40.2, 37.5 (CH₂NHCOCH₂), 34.2, 34.0 (Me₃C), 32.1, 32.0 (ArCH₂Ar), 31.7, 31.1 (Me₃C). – HR-MS C₅₂H₆₈N₂O₆NaS: calcd. for [M + Na]⁺ 871.4696; found. 871.4674. – C₅₂H₆₈N₂O₆S (849.18): calcd. C 73.55, H 8.07, N 3.30; found C 73.14, H 8.30, N 3.22.

5,11,17,23-Tetra-tert-butyl-26,28-dihydroxy-25,27-bis{(thiocyanomethyl)carbonylamino]ethoxy}calix[4]arene (11): A solution of **4** (0.420 g, 0.474 mmol) and KSCN (0.230 g) in dry acetone (25 mL) was refluxed for 18 h. After cooling, the solvent was evaporated, yielding a crude product that was dissolved in AcOEt (75 mL) and washed with H₂O (10 mL). The organic phase was dried and evaporated, giving a crude product that was purified by column chromatography (AcOEt/hexane 1:1) to give **11** (0.378 g, 85.5%) as a solid: M.p. 240–242 °C. – ¹H NMR (300 MHz, CDCl₃): δ = 8.91 (br s, 2 H; 2 NH), 8.68 (s, 2 H; 2 OH), 7.07, 7.02 (2 s, 8 H; Ar), 4.23 (d, J = 13.1 Hz, 4 H; ArCH₂Ar), 4.18–4.10, 4.01–3.93 (2 m, 8 H; 2 NHCH₂CH₂O), 3.78 (s, 4 H; 2 CH₂S), 3.44 (d, J = 13.1 Hz, 4 H; ArCH₂Ar), 1.24, 1.12 (2 s, 36 H; 4 Me₃C). – ¹³C NMR (75 MHz, CDCl₃): δ = 165.6 (CO), 148.6, 143.8, 133.2, 127.9, 126.4, 126.1 (Ar), 112.0 (SCN), 75.7 (OCH₂), 40.3, 37.3 (CH₂NHCOCH₂S), 34.3, 34.0 (Me₃C), 32.3 (ArCH₂Ar), 31.6, 31.1 (Me₃C). – HR-MS (FAB+) C₅₄H₆₈N₄O₆NaS₂: calcd. for [M + Na]⁺ 955.4478; found 955.4476.

5,11,17,23-Tetra-tert-butyl-25,27-bis{(acetylthiomethyl)carbonylamino]ethoxy}-26,28-dihydroxycalix[4]arene (12): A solution of **4** (0.1 g, 0.113 mmol) and KSAc (0.150 g) in dry acetone (10 mL) was refluxed for 1 h. After cooling, the solvent was evaporated, yielding a crude product that was dissolved in AcOEt (50 mL) and washed with H₂O (10 mL). The organic phase was dried and evaporated, giving a crude product that was purified by column chromatography (AcOEt/hexane 3:1) to produce **12** (0.08 g, 73%) as a solid: M.p. 220–222 °C (from ether/hexane). – IR (KBr): $\tilde{\nu}$ = 3410, 3335, 1707, 1651, 1532, 1485, 1362 cm⁻¹. – ¹H NMR (300 MHz, Cl₃CD): δ = 8.69 (s, 2 H; 2 OH), 8.51 (t, J = 5.4 Hz, 2 H; 2 NH), 7.07, 7.02 (2 s, 8 H; ArH), 4.27 (d, J = 13.0 Hz, 4 H; ArCH₂Ar), 4.14 (t, J = 4.5 Hz, 4 H; 2 OCH₂), 3.93 (m, 4 H; 2 CH₂N), 3.72 (s, 4 H; 2 CH₂S), 3.40 (d, J = 13.0 Hz, 4 H; ArCH₂Ar), 2.09 (s, 6 H; AcS), 1.23, 1.13 (2 s, 36 H; Me₃C). – ¹³C NMR (75 MHz, Cl₃CD): δ = 194.5 [C(O)S], 168.3 [C(O)N], 149.5, 148.7, 148.3, 143.1, 133.2, 127.8, 126.2, 125.8 (Ar), 75.8 (OCH₂), 39.9 (CH₂N), 34.3, 33.9 (Me₃C), 33.2, 32.3 (CH₂S, ArCH₂Ar), 31.6, 31.2 (Me₃C), 29.8 (MeC(O)S). – HR-MS (FAB+) C₅₆H₇₄N₂O₈NaS₂: calcd. for [M + Na]⁺ 989.4784; found 989.4773. – C₅₆H₇₄N₂O₈S₂ (967.33): calcd. C 69.53, H 7.71, N 2.89, S 6.63; found C 69.15, H 7.35, N 2.98, S 6.54.

5,11,17,23-Tetra-tert-butyl-26,28-dihydroxy-25,27-bis{(thiomethyl)carbonylamino]ethoxy}calix[4]arene (13): To a solution of **12** (0.110 g, 0.114 mmol) in MeOH (25 mL) was added a 1 N NaOMe solution in MeOH until pH 9–10 was reached. The reaction mixture was kept at room temp. for 1 h. Addition of AcOH (0.25 mL) was followed by evaporation of the solvent, yielding a crude product that was dissolved in CH₂Cl₂ (50 mL) and washed with H₂O (5 mL). The organic phase was dried and evaporated, giving **13** (0.095 g, 95%) as a solid foam: ¹H NMR (300 MHz, Cl₃CD): δ = 8.43 (t, J = 5.5 Hz, 2 H; 2 NH), 8.29 (s, 2 H; 2 OH), 7.01, 6.66 (2 s, 8 H; ArH), 4.17 (d, J = 13.0 Hz, 4 H; ArCH₂Ar), 4.07 (t, J =

4.6 Hz, 4 H; 2 OCH₂), 3.91 (m, 4 H; 2 CH₂N), 3.62 (d, *J* = 13.0 Hz, 4 H; ArCH₂Ar), 3.24 (d, *J* = 8.4 Hz, 4 H; 2 CH₂SH), 2.09 (t, *J* = 8.4 Hz, 2 H; 2 SH), 1.21, 1.04 (2 s, 36 H; 4 Me₃C). – ¹³C NMR (75 MHz, Cl₃CD): δ = 170.2 (CO), 149.3, 148.5, 1498.1, 149.9, 132.8, 127.7, 126.0, 125.8 (Ar), 75.4 (CH₂O), 39.8 (CH₂N, CH₂S), 34.1, 33.8 (Me₃C), 32.1 (ArCH₂Ar), 31.6, 31.1 (Me₃C).

1,12-(5,11,17,23-Tetra-tert-butyl-26,28-dihydroxy-25,27-dioxy-calix[4]arene)-3,10-diaza-6,7-dithia-4,9-dodecanedione (14): Preparation of **13** as described above, followed by column chromatography (AcOEt), yielded **14** (60% from **12**) as a solid: M.p. 260–262 °C. – IR (KBr): $\tilde{\nu}$ = 3426, 1655, 1551, 1485, 1204 cm⁻¹. – ¹H NMR (300 MHz, Cl₃CD): δ = 7.82 (t, *J* = 5.4 Hz, 2 H; 2 NH), 7.50 (s, 2 H; 2 OH), 7.07, 6.86 (2 s, 8 H; ArH), 4.22 (d, *J* = 13.0 Hz, 4 H; ArCH₂Ar), 4.13 (t, *J* = 3.7 Hz, 4 H; 2 OCH₂), 4.03 (m, 4 H; 2 CH₂N), 3.50 (s, 4 H; 2 CH₂S), 3.38 (d, *J* = 13.0 Hz, 4 H; ArCH₂Ar), 1.29, 0.99 (2 s, 36 H; 4 Me₃C). – ¹³C NMR (75 MHz, Cl₃CD): δ = 169.1 (CON), 149.9, 149.2, 147.6, 132.6, 132.5, 127.7, 125.9, 125.5 (Ar), 75.7 (OCH₂), 42.5, 40.3 (CH₂N, CH₂S), 33.9 (Me₃C), 31.8 (ArCH₂Ar), 31.7, 31.0 (Me₃C). – HR-MS (FAB+) C₅₂H₆₈N₂O₆Na₂: calcd. for [M + Na]⁺ 903.4416; found 903.4412. – C₅₂H₆₈N₂O₆S₂ (881.24): calcd. C 70.87, H 7.78, N 3.18; found C 70.98, H 7.41, N 3.41.

Bis(5,11,17,23-tetra-tert-butyl-26,28-dihydroxycalix[4]arene)-25,25',27,27'-bis[(oxyethylamino)carbonylmethyl]sulfide (15): To a solution of **4** (0.050 g, 0.056 mmol) and **12** (0.055 g, 0.057 mmol) in dry MeOH (75 mL) was added 1 N NaOMe solution in MeOH (2 mL). The reaction mixture was kept at room temp. for 24 h. AcOH (1 mL) was added and then the solvent was evaporated, giving a crude product that was dissolved in CH₂Cl₂ (50 mL) and washed with H₂O (5 mL). The organic phase was dried and evaporated. The crude product obtained was purified by column chromatography (AcOEt/hexane 5:1), yielding **15** (0.050 g, 52%) as a solid: M.p. 285–288 °C. – IR (KBr): $\tilde{\nu}$ = 3474, 3350, 3277, 1736, 1650, 1529, 1485, 1465, 1286, 1203 cm⁻¹. – ¹H NMR (300 MHz, CDCl₃): δ = 8.75 (t, *J* = 5.4 Hz, 4 H; 4 NH), 7.17, 6.85 (2s, 16 H; ArH), 7.14 (s, 4 H; 4 OH), 4.29 (d, *J* = 13.0 Hz, 8 H; ArCH₂Ar), 4.10 (m, 8 H; 4 CH₂O), 3.98 (m, 8 H; 4 CH₂N), 3.64 (s, 8 H; 4 CH₂S), 3.41 (d, *J* = 13.0 Hz, 8 H; ArCH₂Ar), 1.41, 1.01 (2s, 72 H; 8 Me₃C). – ¹³C NMR 170.0, 150.4, 148.9, 147.3, 141.8, 132.3, 127.9, 125.5, 125.2, 75.0, 39.3, 37.4, 34.0, 33.9, 31.8, 31.2, 31.0. – HR-MS (FAB+) C₁₀₄H₁₃₆N₄O₁₂Na₂: calcd. for [M + Na]⁺ 1719.9493. – Found 1719.95.

Acknowledgments

We thank the Dirección General de Investigación Científica y Técnica for financial support (PB95-1207 and PB98-1320).

- [1] *Comprehensive Supramolecular Chemistry* (Eds.: J. M. Lehn, J. L. Atwood, J. E. D. Davies, D. D. McNicol, F. Vogtle), Pergamon, Elsevier Science, Oxford, **1996**.
 [2] C. D. Gutsche, *Calixarenes* (Ed: J. F. Stoddart), Royal Society of Chemistry, London, **1989**.
 [3] *Calixarenes. A Versatile Class of Macrocyclic Compounds* (Eds.: J. Vicens, V. Böhmer), Kluwer Academic Publishers, Dordrecht, The Netherlands, **1991**.
 [4] S. Shinkai, *Biorg. Chem. Front.* **1990**, *1*, 161.
 [5] S. Shinkai, *Tetrahedron* **1993**, *49*, 8933.
 [6] H. Ohtsuka, S. Shinkai, *Supramolecular Sci.* **1996**, *3*, 189.
 [7] J. L. Atwood, G. W. Orr, K. D. Robinson, F. Hamada, *Supramol. Chem.* **1993**, *2*, 309.
 [8] V. Böhmer, *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 713.
 [9] A. Ikeda, S. Shinkai, *Chem. Rev.* **1997**, *97*, 1713.
 [10] A. Casnati, *Gazz. Chim. Ital.* **1997**, *127*, 637.

- [11] A. F. Danil de Namor, R. M. Cleverley, M. L. Zapata-Ormaechea, *Chem. Rev.* **1998**, *98*, 2495.
 [12] A. Pochini, R. Ungaro, in: *Comprehensive Supramolecular Chemistry* (Eds.: F. Vogtle), Pergamon Press, Oxford, **1996**; pp. 103–142.
 [13] C. D. Gutsche, *Aldrichimica Acta* **1995**, *28*, 3.
 [14] C. D. Gutsche, M. Iqbal, *Org. Synth., Coll.* **1993**, *Vol VIII*, 75.
 [15] C. D. Gutsche, B. Dhawan, M. Leonis, D. Stewart, *Org. Synth., Coll.* **1993**, *Vol VIII*, 77.
 [16] S. H. Munch, C. D. Gutsche, *Org. Synth., Coll.* **1993**, *Vol VIII*, 80.
 [17] A. Arduini, A. Casnati, in: *Calixarenes in Macrocyclic Synthesis: A Practical Approach* (Eds.: D. Parker), Oxford University Press, Oxford, **1996**; pp. 145–173.
 [18] I. Higler, P. Timmerman, W. Verboom, D. N. Reinhoudt, *Eur. J. Org. Chem.* **1998**, 2689.
 [19] A. Ikeda, S. Shinkai, *Chem. Rev.* **1997**, *97*, 1713.
 [20] F. Ohseto, S. Shinkai, *Chem. Lett.* **1993**, 2045.
 [21] F. Ohseto, T. Sakaki, K. Araki, S. Shinkai, *Tetrahedron Lett.* **1993**, *32*, 2149.
 [22] F. Ohseto, S. Shinkai, *J. Chem. Soc., Perkin Trans. 2* **1993**, 2045.
 [23] D. Kraft, J. D. van Loon, M. Owens, W. Verboom, W. Vogt, M. A. McKerverey, V. Böhmer, D. N. Reinhoudt, *Tetrahedron Lett.* **1990**, *31*, 4941.
 [24] M. A. McKerverey, M. Owens, H.-R. Schulten, W. Vogt, V. Böhmer, *Angew. Chem. Int. Ed. Engl.* **1990**, *29*, 280.
 [25] C. A. Schalley, R. K. Castellano, M. S. Brody, D. M. Rudkevich, G. Siuzdak, J. Rebek, *J. Am. Chem. Soc.* **1999**, *121*, 4568.
 [26] R. K. Castellano, J. Rebek, *J. Am. Chem. Soc.* **1998**, *120*, 3657.
 [27] M. S. Brody, C. A. Schalley, D. M. Schalley, D. M. Rudkevich, J. Rebek, *Angew. Chem. Int. Ed.* **1999**, *38*, 1640.
 [28] J. Wang, C. D. Gutsche, *J. Am. Chem. Soc.* **1998**, *120*, 12226.
 [29] P. Lhotak, S. Shinkai, *Tetrahedron* **1995**, *51*, 7681.
 [30] P. D. Schmitt, P. D. Beer, M. G. B. Drew, P. D. Sheen, *Tetrahedron Lett.* **1998**, *39*, 6383.
 [31] Z. L. Zhong, Y.-Y. Chen, X.-R. Lu, *Tetrahedron Lett.* **1995**, *36*, 6735.
 [32] L. Prodi, S. Pivari, F. Bolleta, M. Hissler, R. Ziessel, *Eur. J. Inorg. Chem.* **1998**, 1958.
 [33] M. Pitarch, V. Mckee, M. Nieuwenhuyzen, M. A. McKerverey, *J. Org. Chem.* **1998**, *63*, 946.
 [34] C. Alfieri, E. Dradi, A. Pochini, R. Ungaro, C. D. Andreotti, *J. Chem. Soc., Chem. Commun.* **1983**, 1075.
 [35] P. J. Dijkstra, J. A. Brunink, K.-E. Bugge, D. N. Reinhoudt, S. Harkema, R. Ungaro, F. Ugozzoli, E. Ghidini, *J. Am. Chem. Soc.* **1989**, *111*, 7567.
 [36] E. Ghidini, F. Ugozzoli, R. Ungaro, S. Harkema, A. A. El-Fadd, D. N. Reinhoudt, *J. Am. Chem. Soc.* **1990**, *112*, 6979.
 [37] R. Ungaro, A. Casnati, F. Ugozzoli, A. Pochini, J.-F. Dozol, C. Hill, H. Rouquette, *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 1506.
 [38] A. Casnati, A. Pochini, R. Ungaro, F. Ugozzoli, F. Arnaud, S. Fanni, M.-J. Schwing, R. J. M. Egberink, F. de Jong, D. N. Reinhoudt, *J. Am. Chem. Soc.* **1995**, *117*, 2767.
 [39] Z. Asfari, S. Wenger, J. Vicens, *J. Inclusion. Phenom.* **1994**, *19*, 137.
 [40] R. A. Sachleben, A. Urvoas, J. C. Bryan, T. J. Haverlock, B. P. Hay, B. A. Moyer, *Chem. Commun.* **1999**, 1751.
 [41] Z. Asfari, C. Bressot, J. Vicens, C. Hill, J.-F. Dozol, H. Rouquette, S. Lamare, B. Tournois, *Anal. Chem.* **1995**, *67*, 3133.
 [42] M. Grunder, J.-F. Dozol, Z. Asfari, J. Vicens, *J. Radionucl. Chem.* **1999**, *241*, 59.
 [43] S. Pappalardo, A. Petringa, M. F. Parisi, G. Ferguson, *Tetrahedron Lett.* **1996**, *37*, 3907.
 [44] P. D. Beer, M. G. Drew, R. J. Knubley, M. I. Ogden, *J. Chem. Soc., Dalton Trans.* **1995**, 3117.
 [45] J. Scheerder, J. F. Engbersen, D. N. Reinhoudt, *Recl. Trav. Chim. Pays-Bas* **1996**, *115*, 307.
 [46] F. P. Schmidtchen, M. Berger, *Chem. Rev.* **1997**, *97*, 1609.
 [47] L. A. J. Chrisstoffels, F. de Jong, D. N. Reinhoudt, S. Sivelli, L. Gazzola, A. Casnati, R. Ungaro, *J. Am. Chem. Soc.* **1999**, *121*, 10142.
 [48] J. C. Adriam, Jr., C. S. Wilcox, *J. Am. Chem. Soc.* **1992**, *114*, 1398.
 [49] T. R. Kelly, M. H. Kim, *J. Am. Chem. Soc.* **1994**, *116*, 7072.

- [50] E. Fan, S. A. V. Arman, S. Kincaid, A. D. Hamilton, *J. Am. Chem. Soc.* **1993**, *115*, 369.
- [51] J. Scheerder, M. Fochi, J. F. Engbersen, D. N. Reinhoudt, *J. Org. Chem.* **1994**, *59*, 7815.
- [52] C. Raposo, M. Almaraz, M. Martin, V. Weinrich, L. Mussons, V. Alcaraz, C. Caballero, J. R. Mosan, *Chem. Lett.* **1995**, 759.
- [53] C. S. Wilcox, E.-I. Kim, S. Romano, L. H. Kuo, A. L. Burt, D. P. Curran, *Tetrahedron* **1995**, *51*, 621.
- [54] P. Bühlmann, S. Nishizawa, K. P. Xiao, Y. Umezawa, *Tetrahedron* **1997**, *53*, 1647.
- [55] R. C. Jagessar, D. H. Burns, *Chem. Commun.* **1997**, 1997.
- [56] M. S. Goodman, V. Jubian, B. Linton, A. D. Hamilton, *J. Am. Chem. Soc.* **1995**, *117*, 11610.
- [57] S. Nishikawa, P. Bühlmann, M. Iwao, Y. Umezawa, *Tetrahedron Lett.* **1995**, *36*, 6483.
- [58] J. L. Jiménez-Blanco, J. L. Benito, C. Ortiz-Mellet, J. M. García-Fernández, *Org. Lett.* **1999**, *1*, 1217.
- [59] J. Scheerder, J. F. J. Engbersen, W. Verboom, J. P. M. van Duynhoven, D. N. Reinhoudt, *J. Org. Chem.* **1996**, *61*, 3476.
- [60] G. J. Pernia, J. D. Kilburn, J. W. Essex, R. J. Mortisshire-Smith, M. Rowley, *J. Am. Chem. Soc.* **1996**, *118*, 10220.
- [61] Y. Tobe, A. Sasaki, M. Mizuno, K. Hirose, K. Naemura, *J. Org. Chem.* **1998**, *63*, 7481.
- [62] S. Sasaki, M. Mizuno, K. Naemura, T. Tobe, *J. Org. Chem.* **2000**, *65*, 275.
- [63] C. Saitz-Barria, A. Torres-Pinedo, F. Santoyo-González, *Synlett* **1999**, 1891.
- [64] J. L. Blanco, C. S. Barria, J. M. Benito, C. O. Mellet, J. Fuentes, F. Santoyo-González, J. M. Fernández, *Synthesis* **1999**, 1907.
- [65] W. C. Zhang, Z. T. Huang, *Synthesis* **1997**, 1073.
- [66] J. M. Fernández, C. O. Mellet, J. Fuentes, *J. Org. Chem.* **1993**, *58*, 5192.
- [67] T. Minami, S. Shinkai, O. Manabe, *Tetrahedron Lett.* **1982**, *23*, 5167.
- [68] S. Shinkai, T. Minami, Y. Araragi, O. Manabe, *J. Chem. Soc., Perkin Trans. 2* **1985**, 503.
- [69] D. N. Harpp, J. G. Gleason, *J. Org. Chem.* **1970**, *35*, 3259.
- [70] X. Wu, R. D. Rieke, L. Zhu, *Synth. Commun.* **1996**, *26*, 191.
- [71] W. W. Cleland, *Biochemistry* **1964**, *3*, 480.
- [72] E. Wenschuh, M. Heydeinreich, R. Runtge, S. Fischer, *Sulfur Lett.* **1988**, *8*, 251.
- [73] L. Field, Y. H. Khim, *J. Org. Chem.* **1972**, *37*, 2710.
- [74] R. H. Cragg, A. F. Weston, *Tetrahedron Lett.* **1973**, *14*, 655.
- [75] G. Crank, M. I. H. Makin, *Aust. J. Chem.* **1984**, *37*, 2331.
- [76] R. G. Guy, *The Chemistry of Cyanates and Their Thio Derivatives. Part II* (Ed: S. Patai), Wiley, New York, **1977**, pp 867.
- [77] X. Jia, Y. Zhang, X. Zhou, *Tetrahedron Lett.* **1994**, *35*, 8833.
- [78] K. R. Prabhu, A. R. Ramesha, S. Chandrasekaran, *J. Org. Chem.* **1995**, *60*, 7142.
- [79] C. J. Burns, L. D. Field, J. Morgan, D. D. Ridley, V. Vignevich, *Tetrahedron Lett.* **1999**, *40*, 6489.