

Singular Limits of a Two-Dimensional Boundary Value Problem Arising in Corrosion Modelling

JUAN DÁVILA, MANUEL DEL PINO, MONICA MUSSO
& JUNCHENG WEI

Abstract

We consider the boundary value problem

$$\Delta u = 0 \text{ in } \Omega, \quad \frac{\partial u}{\partial \nu} = 2\lambda \sinh u \text{ on } \partial\Omega$$

where Ω is a smooth and bounded domain in \mathbb{R}^2 and $\lambda > 0$. We prove that for any integer $k \geq 1$ there exist at least two solutions u_λ with the property that the boundary flux satisfies up to subsequences $\lambda \rightarrow 0$,

$$2\lambda \sinh(u_\lambda) \rightharpoonup 2\pi \sum_{j=1}^{2k} (-1)^{j-1} \delta_{\xi_j},$$

where the ξ_j are points of $\partial\Omega$ ordered clockwise in j .

1. Introduction

Let Ω be a bounded domain in \mathbb{R}^2 with smooth boundary $\partial\Omega$. A very common boundary condition arising in corrosion modelling in a planar sample represented by Ω is associated with the names of Butler and Volmer. In its simplest form it asserts an exponential relationship between boundary voltages and boundary normal currents which takes the form

$$\frac{\partial u}{\partial \nu} = \lambda(e^{2\beta u} - e^{-2(1-\beta)u}) + g \text{ on } \partial\Omega,$$

where the constant $0 < \beta < 1$ depends on the constituents of the electrochemical system but not on their concentrations. Here λ is a constant highly dependent on their concentrations and g is an externally imposed current. Assuming the presence

of no sources or sinks in Ω , the balanced situation $\beta = \frac{1}{2}$, and $g = 0$, the boundary value problem satisfied by the voltage potential becomes in an ideal situation

$$\Delta u = 0 \text{ in } \Omega, \quad \frac{\partial u}{\partial \nu} = 2\lambda \sinh u \text{ on } \partial\Omega. \tag{1.1}$$

We refer the reader to [13] and [5] for the derivation of this and related corrosion models and references to the applied literature.

We assume throughout this paper that $\lambda > 0$. We are interested in solutions to this problem when λ assumes very small values. A surprising example of an explicit solution when $\Omega = D$, the unit disk in \mathbb{R}^2 , was given by BRYAN and VOGELIUS in [3]. Consider $2k$ points on ∂D , ξ_1, \dots, ξ_{2k} corresponding to vertices of a regular polygon, ordered clockwise, and set

$$u_\lambda(x) = \sum_{j=1}^{2k} (-1)^{j-1} \log \frac{1}{|x - \alpha_k \xi_j|^2} \tag{1.2}$$

where $\alpha_k = [(k + 2\lambda)/(k - 2\lambda)]^{\frac{1}{2k}}$. As $\lambda \rightarrow 0$, we observe a singular concentration behavior in the nonlinear boundary condition. Indeed we have that

$$2\lambda \sinh u_\lambda \rightarrow 2\pi \sum_{j=1}^{2k} (-1)^{j-1} \delta_{\xi_j}$$

in the sense of measures in ∂D , where δ_{ξ_j} denote Dirac masses at points ξ_j . In [10] the possible behavior of solutions u_λ with boundary condition $2\lambda \sinh(u_\lambda)$ of uniformly bounded mass is established: the limit of the boundary flux along subsequences is a sum of Dirac masses located at a finite set of points with weights greater than or equal to 4π , potentially accompanied by a regular part which has one sign. Solutions to the problem with this property were found by KAVIAN and VOGELIUS in [8] via Ljusternik–Schnirelmann theory. However, their asymptotic behavior is only partly understood by virtue of the above result. It remains an open question if solutions of the form (1.2) exist in general two-dimensional domains.

The purpose of this paper is to show that in any domain Ω , there are at least two distinct families of solutions which exhibit exactly the qualitative behavior of the explicit solution (1.2), namely with limiting boundary flux given by an array of $2k$ Dirac masses with weight 2π and alternate signs. The location of the $2k$ concentration points can be accurately characterized as special critical points of a functional φ_k defined explicitly in terms of $G(x, y)$ the Green’s function for the Neumann problem

$$\begin{cases} \Delta_x G(x, y) &= 0 & \text{in } \Omega \\ \frac{\partial G}{\partial \nu_x}(x, y) &= 2\pi \delta_y(x) - \frac{2\pi}{|\partial\Omega|} & \text{on } \partial\Omega \\ \int_{\partial\Omega} G(x, y) dx &= 0. \end{cases} \tag{1.3}$$

We denote by $H(x, y)$ its regular part:

$$H(x, y) = G(x, y) - \log \frac{1}{|x - y|^2}. \tag{1.4}$$

For $m \geq 1$ and points ξ_1, \dots, ξ_m on $\partial\Omega$ ordered clockwise we define

$$\varphi_m(\xi_1, \dots, \xi_m) = \sum_{l=1}^m H(\xi_l, \xi_l) + \sum_{j \neq l} (-1)^{l+j} G(\xi_j, \xi_l). \quad (1.5)$$

Our main result states as follows.

Theorem 1. *Let $k \geq 1$ be a positive integer. There is a number $\lambda_k > 0$ such that for any $0 < \lambda < \lambda_k$ there are two solutions $u_{\lambda 1} \neq -u_{\lambda 2}$ with*

$$\lambda \int_{\partial\Omega} |\sinh u_{\lambda l}| \rightarrow 8k\pi \quad \text{as } \lambda \rightarrow 0.$$

More precisely, given any sequence $\lambda = \lambda_n \rightarrow 0$, there is a subsequence, two arrays of $2k$ points of $\partial\Omega$ ($\xi_{l1}, \xi_{l2}, \dots, \xi_{l2k}$) ordered clockwise and distinct modulo-cyclic permutations, positive constants $\mu_j = \mu_{lj}$, for $j = 1, \dots, 2k$, and two solutions $u_{\lambda l}$ of (1.1), $l = 1, 2$ such that, omitting the subindex l ,

$$u_\lambda(x) = \sum_{j=1}^{2k} (-1)^{j-1} \log \frac{2\mu_j}{|x - (\xi_j + \lambda\mu_j v_j)|^2} + O(1),$$

where v_j denotes unit outer normal to $\partial\Omega$ at ξ_j and

$$2\lambda \sinh u_\lambda \rightarrow 2\pi \sum_{j=1}^{2k} (-1)^{j-1} \delta_{\xi_j}.$$

Moreover, the $2k$ -tuples $(\xi_{l1}, \dots, \xi_{l2k})$ are critical points of φ_{2k} , and the constants μ_j are explicitly given by

$$\log 2\mu_j = H(\xi_j, \xi_j) + \sum_{l \neq j} (-1)^{l+j} G(\xi_j, \xi_l).$$

It is easily checked that the solutions (1.2) correspond exactly to this description in the case $\Omega = D$. MEDVILLE and VOGELIUS [10] have established that if the limit boundary flux has no regular part, then the concentration points $(\xi_{l1}, \dots, \xi_{l2k})$ necessarily constitute a critical point of φ_{2k} , and that the weights of the delta functions are equal to 2π . They provide numerical evidence that solutions with a boundary flux having a non-trivial regular part exist, but this remains an open question. Let us mention that in [11] MEDVILLE and VOGELIUS considered the nonlinear boundary condition

$$\frac{\partial u}{\partial \nu} = Du + 2\lambda \sinh u \quad \text{on } \partial\Omega,$$

where $D > 0$. They analyzed the difference in blow-up as λ approaches 0 from the right (pointwise blow-up) and from the left (blow-up ‘‘almost everywhere’’).

It is interesting to mention the analogy existing between this result and the problem $-\Delta u = \lambda e^u$ under Dirichlet boundary conditions, whose solutions with $\lambda \int_\Omega e^u$ uniformly bounded have become well understood after the works [12, 2, 9]. It follows from those results that concentration occurs in the form $\lambda e^u \rightarrow 8\pi \sum \delta_{\xi_j}$. In [1, 6, 7] solutions with these properties have been built. In [4] the problem

$$\Delta u - u = 0, \quad \frac{\partial u}{\partial \nu} = \lambda e^u \tag{1.6}$$

was analyzed and given any $k \geq 1$, a solution peaking in such a way that $\lambda e^u \rightarrow 2\pi \sum_{j=1}^k \delta_{\xi_j}$ was built up, using as basic cells (after suitable blowing-up) explicit solutions of

$$\begin{cases} \Delta v = 0 & \text{in } \mathbb{R}_+^2 \\ \frac{\partial v}{\partial \nu} = e^v & \text{on } \partial \mathbb{R}_+^2, \end{cases} \tag{1.7}$$

where \mathbb{R}_+^2 denotes the upper half plane $\{(x_1, x_2) : x_2 > 0\}$ and ν the unit exterior normal to $\partial \mathbb{R}_+^2$, given by

$$w_{t, \mu}(x_1, x_2) = \log \frac{2\mu}{(x_1 - t)^2 + (x_2 + \mu)^2} \tag{1.8}$$

where $t \in \mathbb{R}$ and $\mu > 0$ are parameters. The solutions whose existence are asserted in Theorem 1 are also constructed using the above solutions scaled up and projected to make them of one right order for the boundary condition. Solutions are found as a small additive perturbation of these initial approximations. A linearization procedure leads to a finite dimensional reduction, where the reduced problem corresponds to that of adjusting variationally the location of the concentration points. An important element in the reduction procedure is the non-degeneracy of these solutions up to variations of the parameters t and μ in (1.8). Problem (1.1) has a basic difficulty in comparison to (1.6) which is linked to the fact that the limiting equation formally satisfied by the sought solution u_λ is

$$\Delta u = 0, \quad \frac{\partial u}{\partial \nu} = 2\pi \sum_{j=1}^{2k} (-1)^{j-1} \delta_{\xi_j},$$

whose solution is not unique but invariant under the addition of constants. This is not such an innocent matter since this hidden limiting invariance is not present in the equation itself, and unlike the other “obvious” elements of the limiting kernel (see (2.1), (2.2) below), it is not localized near the points of concentration. It must be mentioned that the simple use of an additive constant as an extra parameter in the solution does not suffice, basically because the constant itself is not a good approximation of an element of the kernel before reaching the limit. We are able to overcome this difficulty by identifying an extra element of the approximate kernel (see (3.5) below), which introduces another parameter to be adjusted in the problem. We will devote the rest of this paper to the proof of Theorem 1.

2. Preliminaries

Let us define

$$z_0 = 1 - 2\mu \frac{x_2 + \mu}{x_1^2 + (x_2 + \mu)^2}, \tag{2.1}$$

and

$$z_1 = -2 \frac{x_1}{x_1^2 + (x_2 + \mu)^2}, \quad (2.2)$$

which correspond to derivatives of the basic solutions $w_{t,\mu}$ with respect to its parameters of translation and dilation respectively. These objects obviously lie in the kernel of the linearization of problem (1.7) at the solution $w_{0,\mu}$, namely they solve the problem

$$\begin{cases} \Delta \phi = 0 & \text{in } \mathbb{R}_+^2 \\ \frac{\partial \phi}{\partial \nu} - \frac{2\mu}{x_1^2 + \mu^2} \phi = 0 & \text{on } \partial \mathbb{R}_+^2. \end{cases} \quad (2.3)$$

Reciprocally, we have the following.

Lemma 1. *Any bounded solution of (2.3) is a linear combination of z_0 and z_1 .*

Proof. This result was established in [4]. For the sake of completeness, the proof is presented here. Let ϕ be a solution to (2.3) and set

$$w(y) = \phi \left(\frac{y}{|y|^2} - (0, \mu) \right).$$

The function w is just the Kelvin transform of ϕ about the point $(0, -\mu)$. The domain of w is the disk $D = B \left(\left(0, \frac{1}{2\mu}\right), \frac{1}{2\mu} \right)$ and w is a bounded function that satisfies $\Delta w = 0$ in D ,

$$\frac{\partial w}{\partial \nu'} = 2\mu w \quad \text{on } \partial D \setminus \{0\}, \quad (2.4)$$

where ν' is the exterior unit normal to D . To see this observe that the map $y \mapsto K(y) = \frac{y}{|y|^2} - (0, \mu)$ is anti-conformal (preserves angles and reverses orientation) and maps the normal vector to D to a normal vector to $\partial \mathbb{R}_+^2$. More precisely, if ν' is the exterior unit normal vector to D then

$$\frac{\partial w}{\partial \nu'} = \frac{1}{|y|^2} \frac{\partial \phi}{\partial \nu}.$$

Thus on ∂D

$$\frac{\partial w}{\partial \nu'} = \frac{1}{|y|^2} e^{w_{0,\mu}(K(y))} w$$

and a calculation shows that

$$\frac{1}{|y|^2} e^{w_{0,\mu}(K(y))} = \frac{1}{|y|^2} \frac{2\mu}{\frac{y_1^2}{|y|^4} + \mu^2} = 2\mu.$$

Since w is bounded, by elliptic regularity (2.4) holds in all ∂D .

By translating in the y_2 direction we can assume that D is the disk centered at the origin with radius $\frac{1}{2\mu}$. We think of w as the real part of an analytic function w and write

$$w(y) = \sum_{k=0}^{\infty} a_k r^k e^{ik\theta}$$

with $y = r e^{i\theta}$. Condition (2.4) is equivalent to

$$\operatorname{Re} \left(\sum_{k=0}^{\infty} a_k (k-1) e^{ik\theta} \right) = 0 \quad \forall \theta,$$

and hence $a_0 = 0$, $a_k = 0$ for all $k > 1$. Looking at the real part of w , and recalling that we shifted in the y_2 direction we see that it is a linear combination of $y_1 = \frac{x_1}{x_1^2 + (x_2 + \mu)^2}$ and $y_2 - \frac{1}{2\mu} = \frac{x_2 + \mu}{x_1^2 + (x_2 + \mu)^2} - \frac{1}{2\mu}$. \square

In what remains of this paper we fix $k \geq 1$ and denote $m = 2k$. We will provide a first approximation for the solutions of problem (1.1) predicted in Theorem 1. For $j = 1, \dots, m$, let ξ_j be clockwise ordered points on the boundary of Ω and μ_j positive numbers. Define, for $x \in \Omega$,

$$u_j^\lambda(x) = \log \frac{2\mu_j}{|x - \xi_j - \lambda\mu_j v_j|^2} \tag{2.5}$$

and $H_j^\lambda(x)$ to be the unique solution of

$$\Delta H_j^\lambda = 0 \text{ in } \Omega, \tag{2.6}$$

$$\frac{\partial H_j^\lambda}{\partial \nu} = -\frac{\partial u_j^\lambda}{\partial \nu} + \lambda e^{u_j^\lambda} - \lambda \frac{1}{|\partial\Omega|} \int_{\partial\Omega} e^{u_j^\lambda} \text{ on } \partial\Omega \tag{2.7}$$

with the property that

$$\int_{\partial\Omega} H_j^\lambda(x) dx = - \int_{\partial\Omega} u_j^\lambda(x) dx. \tag{2.8}$$

We look for a solution to (1.1) of the form

$$u(x) = U(x) + \Phi(x) \tag{2.9}$$

where

$$U(x) = \sum_{j=1}^m (-1)^{j-1} (u_j^\lambda(x) + H_j^\lambda(x)) \tag{2.10}$$

while Φ is a lower-order term with respect to U .

The function H_j^λ above resembles the shape of the regular part of the Green's function. Indeed, the following estimate for H_j^λ holds true.

Lemma 2. For any $0 < \alpha < 1$,

$$H_j^\lambda(x) = H(x, \xi_j) - \log 2\mu_j + O(\lambda^\alpha) \quad (2.11)$$

uniformly in $\bar{\Omega}$.

Proof. The normal derivative of H_j^λ on the boundary of Ω can be computed explicitly, namely

$$\begin{aligned} \frac{\partial H_j^\lambda}{\partial \nu}(x) &= 2\lambda\mu_j \frac{1 - \nu(\xi_j) \cdot \nu(x)}{|x - \xi_j - \lambda\mu_j \nu(\xi_j)|^2} + 2 \frac{(x - \xi_j) \cdot \nu(x)}{|x - \xi_j - \lambda\mu_j \nu(\xi_j)|^2} \\ &\quad - \lambda \frac{1}{|\partial\Omega|} \int_{\partial\Omega} e^{u_j^\lambda(x)}. \end{aligned}$$

Thus,

$$\lim_{\lambda \rightarrow 0} \frac{\partial H_j^\lambda}{\partial \nu}(x) = 2 \frac{(x - \xi_j) \cdot \nu(x)}{|x - \xi_j|^2} - \frac{2\pi}{|\partial\Omega|} \quad \forall x \neq \xi_j \quad (2.12)$$

since

$$\begin{aligned} \lambda \int_{\partial\Omega} e^{u_j^\lambda(x)} &= \lambda \int_{\partial\Omega} \frac{2\mu_j}{|x - \xi_j - \lambda\mu_j \nu(\xi_j)|^2} = 2 \int_{\frac{\partial\Omega - \xi_j}{\lambda\mu_j}} \frac{1}{|y - \nu(0)|^2} \\ &= 2 \left(\int_{-\infty}^{\infty} \frac{dt}{1+t^2} - O \left(\int_{\lambda^{-1}\mu_j^{-1}}^{\infty} \frac{dt}{1+t^2} \right) \right) \\ &= 2\pi + O \left(\arctan(\lambda\mu_j)^{-1} - \frac{\pi}{2} \right) \\ &= 2\pi + O(\arctan(\lambda\mu_j)) \\ &= 2\pi + O(\lambda\mu_j). \end{aligned} \quad (2.13)$$

Define $z_\lambda(x) = H_j^\lambda(x) + \log 2\mu_j - H(x, \xi_j)$. Since the regular part of the Green's function $H(x, \xi_j)$ is harmonic in Ω and satisfies the Neumann boundary condition

$$\frac{\partial H}{\partial \nu_x}(x, y) = 2 \frac{(x - y) \cdot \nu(x)}{|x - y|^2} - \frac{2\pi}{|\partial\Omega|} \quad x \in \partial\Omega,$$

the difference z_λ solves the problem

$$\begin{cases} -\Delta z_\lambda = 0 & \text{in } \Omega \\ \frac{\partial z_\lambda}{\partial \nu} = \frac{\partial H_j^\lambda}{\partial \nu} - 2 \frac{(x - \xi_j) \cdot \nu(x)}{|x - \xi_j|^2} + \frac{2\pi}{|\partial\Omega|} & \text{on } \partial\Omega. \end{cases}$$

Since z_λ is harmonic in Ω , for any $1 \leq p \leq \infty$, $z_\lambda \in W^{1,p}(\Omega)$ and, by the Poincaré inequality, we get

$$\left\| z_\lambda - \frac{1}{|\partial\Omega|} \int_{\partial\Omega} z_\lambda \right\|_{L^p(\Omega)} \leq \|Dz_\lambda\|_{L^p(\Omega)}.$$

Hence, by L^p theory, we have for any $0 < s < \frac{1}{p}$

$$\left\| z_\lambda - \frac{1}{|\partial\Omega|} \int_{\partial\Omega} z_\lambda \right\|_{W^{1+s,p}(\Omega)} \leq C \left\| \frac{\partial z_\lambda}{\partial \nu} \right\|_{L^p(\partial\Omega)} \leq C \lambda^{\frac{1}{p}},$$

where the last inequality can be obtained by arguing as in Lemma 3.1 in [4] and using (2.12). This implies the existence of a constant l such that, for any $\alpha \in (0, 1)$,

$$z_\lambda(x) = l + O(\lambda^\alpha)$$

uniformly in $\bar{\Omega}$, where $l = \lim_{\lambda \rightarrow 0} \frac{1}{|\partial\Omega|} \int_{\partial\Omega} z_\lambda dx$.

In order to get the result, we are left to show that $l = 0$. We have

$$l = \lim_{\lambda \rightarrow 0} \left[\frac{1}{|\partial\Omega|} \int_{\partial\Omega} H_j^\lambda(x) dx + \log 2\mu_j - \frac{1}{|\partial\Omega|} \int_{\Omega} H(x, \xi_j) dx \right]. \quad (2.14)$$

We directly compute from (2.8)

$$\begin{aligned} \frac{1}{|\partial\Omega|} \int_{\partial\Omega} H_j^\lambda(x) dx &= -\frac{1}{|\partial\Omega|} \int_{\partial\Omega} \log \frac{2\mu_j}{|x - \xi_j - \lambda\mu_j\nu(\xi_j)|^2} dx \\ &= -\log 2\mu_j - \frac{1}{|\partial\Omega|} \int_{\partial\Omega} \log \frac{1}{|x - \xi_j|^2} dx \\ &\quad + \frac{1}{|\partial\Omega|} \int_{\partial\Omega} \log \left(1 + 2\lambda\mu_j\nu(\xi_j) \cdot \frac{(x - \xi_j)}{|x - \xi_j|^2} + \frac{\lambda^2\mu_j^2}{|x - \xi_j|^2} \right) dx \\ &= -\log 2\mu_j + \frac{1}{|\partial\Omega|} \int_{\partial\Omega} H(x, \xi_j) dx + O(\lambda), \end{aligned}$$

where the last equality is a consequence of the definition of the regular part of the Green's function. Hence (2.14) yields that $l = 0$. \square

By the following scaling,

$$x = \lambda y, \quad y \in \Omega_\lambda \equiv \frac{\Omega}{\lambda}, \quad v(y) = u(\lambda y)$$

solving problem (1.1) is equivalent to solving

$$\Delta v = 0 \text{ in } \Omega_\lambda, \quad \frac{\partial v}{\partial \nu} = 2\lambda^2 \sinh v \text{ on } \partial\Omega_\lambda. \quad (2.15)$$

In the expanded domain Ω_λ , the main term (2.10) of the ansatz (2.9) now looks like

$$V(y) = \sum_{j=1}^m (-1)^{j-1} \left[\log \frac{2\mu_j}{|y - \xi'_j - \mu_j\nu'_j|^2} - 2\log \lambda + H_j^\lambda(\lambda y) \right], \quad (2.16)$$

where $\xi'_j = \lambda^{-1}\xi_j$ and $\nu'_j = \nu(\xi'_j)$.

We call

$$v_j(y) = u_j^\lambda(\lambda y) + 2 \log \lambda = \log \frac{2\mu_j}{|y - \xi'_j - \mu_j v'_j|^2}$$

and

$$V_j(y) = v_j(y) - 2 \log \lambda + H_j^\lambda(y).$$

A function v of the form

$$v(y) = V(y) + \phi(y), \quad y \in \Omega_\lambda$$

is a solution for (2.15) if and only if ϕ solves

$$\begin{cases} \Delta \phi = 0 & \text{in } \Omega_\lambda, \\ \frac{\partial \phi}{\partial \nu} - W \phi = R + N(\phi) & \text{on } \partial \Omega_\lambda, \end{cases} \quad (2.17)$$

where

$$W = 2\lambda^2 \cosh V, \quad (2.18)$$

$$R = - \left[\frac{\partial V}{\partial \nu} - 2\lambda^2 \sinh V \right] \quad (2.19)$$

and

$$N(\phi) = 2\lambda^2 [\sinh(V + \phi) - \sinh V - \cosh V \phi]. \quad (2.20)$$

We claim that V is a good approximation for a solution of (2.15) under the assumption that we choose the parameters μ_j to be given by the relation

$$\log 2\mu_j = H(\xi_j, \xi_j) + \sum_{l \neq j} (-1)^{l+j} G(\xi_j, \xi_l). \quad (2.21)$$

This is the content of estimate (2.22) contained in the following Lemma

Lemma 3. *Assume (2.21) holds true. Then, for any $\alpha \in (0, 1)$, there exists a positive constant C independent of λ such that, for any $y \in \Omega_\lambda$,*

$$|R(y)| \leq C\lambda^\alpha \sum_{j=1}^m \frac{1}{1 + |y - \xi'_j|}, \quad \forall y \in \Omega_\lambda, \quad (2.22)$$

and

$$W(y) = \sum_{j=1}^m \frac{2\mu_j}{|y - \xi'_j - \mu_j v'_j|^2} (1 + \theta_\lambda(y)), \quad (2.23)$$

with

$$|\theta_\lambda(y)| \leq C\lambda^\alpha + C\lambda \sum_{j=1}^m |y - \xi'_j|. \quad (2.24)$$

Proof. First, we observe that a direct consequence of condition (2.21) is that for $|y - \xi'_j| \leq \frac{\delta}{\lambda}$, the following expansion holds true

$$\begin{aligned}
 & (-1)^{j-1} H_j^\lambda(\lambda y) + \sum_{i \neq j} (-1)^{i-1} \left(\log \frac{2\mu_i}{\lambda^2 |y - \xi'_i - \mu_i v(\xi'_i)|^2} + H_i^\lambda(\lambda y) \right) \\
 &= (-1)^{j-1} (H(\xi_j, \xi_j) - \log 2\mu_j) \\
 &\quad + \sum_{i \neq j} (-1)^{i-1} G(\xi_i, \xi_j) + O(\lambda^\alpha) + O(\lambda |y - \xi'_j|) \\
 &= O(\lambda^\alpha) + O(\lambda |y - \xi'_j|). \tag{2.25}
 \end{aligned}$$

We prove (2.22). By definition

$$\begin{aligned}
 -R &= \sum_{j=1}^m (-1)^{j-1} \frac{2\mu_j}{|y - \xi'_j - \mu_j v(\xi'_j)|^2} \\
 &\quad - 2\lambda^2 \sinh V - \frac{\lambda^2}{|\partial\Omega|} \sum_{j=1}^m (-1)^{j-1} \int_{\partial\Omega} e^{u_j^\lambda}.
 \end{aligned}$$

The last term in R can be controlled by $O(\lambda^2)$. Indeed, the following fact holds true

$$\lambda \sum_{j=1}^m (-1)^{j-1} \int_{\partial\Omega} e^{u_j^\lambda} = O \left(\lambda \sum_{j \neq i} |\mu_j - \mu_i| \right) \tag{2.26}$$

as a direct consequence of (2.13).

On the other hand, if $|y - \xi'_j| \leq \frac{\delta}{\lambda}$,

$$\begin{aligned}
 & 2\lambda^2 \sinh V \\
 &= \lambda^2 \left[\exp \left(\sum_{j=1}^m (-1)^{j-1} (u_j^\lambda(\lambda y) + H_j^\lambda(\lambda y)) \right) \right. \\
 &\quad \left. - \exp \left(\sum_{j=1}^m (-1)^j (u_j^\lambda(\lambda y) + H_j^\lambda(\lambda y)) \right) \right] \\
 &= \lambda^2 \left(\frac{2\mu_j}{\lambda^2 |y - \xi'_j - \mu_j v(\xi'_j)|^2} \right)^{(-1)^{j-1}} \\
 &\quad \times \exp \left[(-1)^{j-1} H_j^\lambda(\lambda y) + \sum_{i \neq j} (-1)^{i-1} \left(\log \frac{2\mu_i}{\lambda^2 |y - \xi'_i - \mu_i v(\xi'_i)|^2} + H_i^\lambda(\lambda y) \right) \right] \\
 &\quad - \lambda^2 \left(\frac{2\mu_j}{\lambda^2 |y - \xi'_j - \mu_j v(\xi'_j)|^2} \right)^{(-1)^j}
 \end{aligned}$$

$$\begin{aligned}
 & \times \exp \left[(-1)^{j-1} H_j^\lambda(\lambda y) + \sum_{i \neq j} (-1)^{i-1} \left(\log \frac{2\mu_i}{\lambda^2 |y - \xi'_i - \mu_i v(\xi'_i)|^2} + H_i^\lambda(\lambda y) \right) \right] \\
 & \text{(using (2.25))} \\
 & = \lambda^2 \left[\left(\frac{2\mu_j}{\lambda^2 |y - \xi'_j - \mu_j v(\xi'_j)|^2} \right)^{(-1)^{j-1}} \right. \\
 & \quad \left. - \left(\frac{2\mu_j}{\lambda^2 |y - \xi'_j - \mu_j v(\xi'_j)|^2} \right)^{(-1)^j} \right] e^{O(\lambda^\alpha) + O(\lambda |y - \xi'_j|)} \\
 & = (-1)^{j-1} \frac{2\mu_j}{|y - \xi'_j - \mu_j v(\xi'_j)|^2} \left(1 + O(\lambda^\alpha) + O(\lambda |y - \xi'_j|) \right) + O(\lambda^4).
 \end{aligned} \tag{2.27}$$

Hence, for $|y - \xi'_j| \leq \frac{\delta}{\lambda}$, we get

$$R = \sum_{j=1}^m (-1)^{j-1} \frac{2\mu_j}{|y - \xi'_j - \mu_j v(\xi'_j)|^2} \left(O(\lambda^\alpha) + O(\lambda |y - \xi'_j|) \right).$$

If we are far away from the points, namely if $|y - \xi'_j| > \frac{\delta}{\lambda}$ for all j , then $R = O(\lambda^2)$. This implies (2.22).

Estimates (2.23) and (2.24) follow from the same arguments used to obtain estimate (2.27). \square

3. Analysis of the linearized equation

In this section we study the linear problem

$$\begin{cases} -\Delta \phi = f & \text{in } \Omega_\lambda \\ \frac{\partial \phi}{\partial \nu} = W\phi + h & \text{on } \partial\Omega_\lambda \end{cases} \tag{3.1}$$

together with appropriate orthogonality conditions, where W is a function that satisfies (2.23) and (2.24), and f, h are given. Throughout this section we only assume that the numbers μ_j appearing in (2.23) satisfy $\frac{1}{C} \leq \mu_j \leq C$ independently of λ , and that the points $\xi_j \in \partial\Omega$ are uniformly separated

$$|\xi_i - \xi_j| \geq d \quad \forall i \neq j, \tag{3.2}$$

where $d > 0$ is fixed.

The orthogonality conditions mentioned above are related to the kernel of (3.1) when $\lambda \rightarrow 0$. Let us look at (3.1) with $f \equiv 0, h \equiv 0$ as $\lambda \rightarrow 0$ at a fixed distance from one of the points, say ξ'_j , and let us translate and rotate so that $\xi'_j = 0$ and Ω_λ converges to the upper half plane \mathbb{R}_+^2 . Then equation (3.1) approaches (2.3). By Lemma 1 we know that any bounded solution to (2.3) is a linear combination of z_0 and z_1 defined in (2.1), (2.2). We define appropriate versions of z_0 and z_1

in Ω_λ through a diffeomorphism $F_j : B_\rho(\xi_j) \rightarrow \mathcal{N}_0$ where $\rho > 0$ is fixed and \mathcal{N}_0 is an open neighborhood of the origin such that $F_j(\Omega \cap B_\rho(\xi_j)) = \mathbb{R}_+^2 \cap \mathcal{N}_0$, $F_j(\partial\Omega \cap B_\rho(\xi_j)) = \partial\mathbb{R}_+^2 \cap \mathcal{N}_0$, and such that F_j preserves area. We define for $y \in \Omega_\lambda$,

$$F_j^\lambda(y) = \frac{1}{\lambda} F_j(\lambda y) \quad (3.3)$$

and

$$Z_{ij}(y) = z_{ij}(F_j^\lambda(y)) \quad i = 0, 1 \quad j = 1, \dots, m,$$

where z_{ij} denotes the function z_i with parameter $\mu = \mu_j$ ($i = 0, 1 \quad j = 1, \dots, m$):

$$z_{0j} = 1 - 2\mu_j \frac{x_2 + \mu_j}{x_1^2 + (x_2 + \mu_j)^2}, \quad z_{1j} = -2 \frac{x_1}{x_1^2 + (x_2 + \mu_j)^2}.$$

Next, we fix a large constant $R_0 > 0$ and a non-negative smooth function $\bar{\chi} : \mathbb{R} \rightarrow \mathbb{R}$ so that $\bar{\chi}(r) = 1$ for $r \leq R_0$ and $\bar{\chi}(r) = 0$ for $r \geq R_0 + 1$, $0 \leq \bar{\chi} \leq 1$. Then set

$$\chi_j(y) = \bar{\chi}(|F_j^\lambda(y)|). \quad (3.4)$$

Let $0 < b < 1$ and define

$$Z(y) = \begin{cases} \min(1 - \lambda^b, Z_{0j}(y)) & \text{if } |y - \xi'_j| < \frac{\delta}{\lambda}, \\ 1 - \lambda^b & \text{if } |y - \xi'_j| \geq \frac{\delta}{\lambda} \quad \forall j = 1, \dots, m. \end{cases} \quad (3.5)$$

We will establish *a priori* estimates for solutions to (3.1) under the orthogonality conditions

$$\int_{\Omega_\lambda} \chi_j Z_{1j} \phi = 0 \quad \forall j = 1, \dots, m \quad (3.6)$$

and

$$\int_{\Omega_\lambda} \chi Z \phi = 0, \quad (3.7)$$

where

$$\chi = \sum_{j=1}^m \chi_j.$$

Let us introduce the norms

$$\|h\|_* = \sup_{y \in \partial\Omega_\lambda} \frac{|h(y)|}{\sum_{j=1}^m (1 + |y - \xi'_j|)^{-1-\sigma}}$$

and

$$\|f\|_{**} = \sup_{y \in \Omega_\lambda} \frac{|f(y)|}{\sum_{j=1}^m (1 + |y - \xi'_j|)^{-2-\sigma}},$$

where $\sigma > 0$ is a fixed small constant.

Proposition 1. For fixed $d > 0$ there exist $\lambda_0 > 0$, C such that if $0 < \lambda < \lambda_0$, $\xi_j \in \partial\Omega$ ($j = 1, \dots, m$) satisfy (3.2), and $\phi \in L^\infty(\Omega_\lambda)$ is a solution of (3.1) such that (3.6) and (3.7) hold, then

$$\|\phi\|_{L^\infty(\Omega_\lambda)} \leq C \log \frac{1}{\lambda} (\|f\|_{**} + \|h\|_*).$$

We will prove this estimate by contradiction assuming that there exists a sequence $\lambda \rightarrow 0$, points $\xi_j \in \Omega$ satisfying (3.2) (we omit the dependence on λ in the notation) and functions $h \in L^\infty(\partial\Omega_\lambda)$, $f \in L^\infty(\Omega_\lambda)$, $\phi \in L^\infty(\Omega_\lambda)$ such that

$$\begin{aligned} \|\phi\|_{L^\infty(\Omega_\lambda)} &= 1 \\ \log \frac{1}{\lambda} \|h\|_* &= o(1), \quad \log \frac{1}{\lambda} \|f\|_{**} = o(1). \end{aligned} \quad (3.8)$$

Given $0 < \alpha < 1$, fix $0 < \gamma < \beta < \alpha$ and consider the function p given by

$$p(r) = \begin{cases} 1 & \text{if } r < \lambda^{-\gamma}, \\ \frac{\log \lambda^{-\beta} - \log r}{\log \lambda^{-\beta} - \log \lambda^{-\gamma}} & \text{if } \lambda^{-\gamma} < r < \lambda^{-\beta}, \\ 0 & \text{if } r > \lambda^{-\beta}. \end{cases} \quad (3.9)$$

Define

$$\tilde{Z}_{0j}(y) = z_{0j}(F_j^\lambda(y)) p(|F_j^\lambda(y)|) \quad j = 1, \dots, m.$$

Let

$$\tilde{\phi} = \phi - \sum_{j=1}^m d_j \tilde{Z}_{0j},$$

where the numbers d_j are chosen so that $\int_{\Omega_\lambda} \chi_j Z_{0j} \tilde{\phi} = 0$ for any $j = 1, \dots, m$, namely $d_j = \frac{\int_{\Omega_\lambda} \chi_j Z_{0j} \phi}{\int_{\Omega_\lambda} \chi_j Z_{0j}^2}$. Observe that

$$d_j = O(1), \quad \|\tilde{\phi}\|_{L^\infty(\Omega_\lambda)} = O(1).$$

Furthermore, $\tilde{\phi}$ solves the problem

$$\begin{cases} -\Delta \tilde{\phi} = f + \sum_{j=1}^m d_j \Delta \tilde{Z}_{0j} & \text{in } \Omega_\lambda \\ \frac{\partial \tilde{\phi}}{\partial \nu} = W \tilde{\phi} + h + \sum_{j=1}^m d_j \left(W \tilde{Z}_{0j} - \frac{\partial \tilde{Z}_{0j}}{\partial \nu} \right) & \text{on } \partial\Omega_\lambda \end{cases} \quad (3.10)$$

and satisfies

$$\int_{\Omega_\lambda} \chi_j Z_{ij} \tilde{\phi} = 0 \quad \forall i = 0, 1 \quad \forall j = 1, \dots, m. \quad (3.11)$$

To reach a contradiction we will establish the following

Lemma 4.

$$\tilde{\phi} \rightarrow 0 \quad \text{uniformly in } \Omega_\lambda.$$

Lemma 5.

$$d_j \rightarrow 0 \quad \forall j = 1, \dots, m.$$

This will prove Proposition 1.

We delay the proofs of these lemmas and mention first some key steps.

Lemma 6. *For all $j = 1, \dots, m$ and $R > 0$*

$$\tilde{\phi} \rightarrow 0 \quad \text{uniformly in } \Omega_\lambda \cap B_R(\xi'_j).$$

Proof. Assume that for some $R > 0$ and $j = 1, \dots, m$ there is $c > 0$ so that $\sup_{B_R(\xi'_j)} |\tilde{\phi}| \geq c > 0$ for a subsequence $\lambda \rightarrow 0$. Translate and rotate Ω_λ so that $\xi'_j = 0$ and Ω_λ converges to the upper half plane \mathbb{R}_+^2 . By elliptic estimates $\tilde{\phi} \rightarrow \tilde{\phi}_0$ uniformly on compact sets, and $\tilde{\phi}_0$ is a non-trivial solution of (2.3). Applying Proposition 1 we conclude that $\tilde{\phi}_0$ is a linear combination of z_0 and z_1 . On the other hand, consider the limit as $\lambda \rightarrow 0$ of the orthogonality conditions (3.11). After translation and rotation, Z_{ij} converges to z_i implying $\int_{\mathbb{R}_+^2} \tilde{\chi} z_i \tilde{\phi}_0 = 0$ for $i = 0, 1$. This contradicts the fact that $\tilde{\phi}_0 \neq 0$. \square

Lemma 7.

$$\bar{\phi} \equiv \frac{1}{|\partial\Omega_\lambda|} \int_{\partial\Omega_\lambda} \tilde{\phi} \rightarrow 0.$$

Proof. By potential theory

$$\begin{aligned} \tilde{\phi}(y) - \bar{\phi} &= \frac{1}{2\pi} \int_{\partial\Omega_\lambda} G(\lambda y, \lambda z) \left(W\tilde{\phi} + h + \sum_j d_j \left(W\tilde{Z}_{0j} - \frac{\partial\tilde{Z}_{0j}}{\partial v} \right) \right) dz \\ &\quad + \frac{1}{2\pi} \int_{\Omega_\lambda} G(\lambda y, \lambda z) \left(f + \sum_j d_j \Delta\tilde{Z}_{0j} \right) dz, \end{aligned}$$

where G is Green's function defined in (1.3).

Integrating equation (3.10) yields

$$\int_{\partial\Omega_\lambda} W\tilde{\phi} + h + \sum_j d_j \left(W\tilde{Z}_{0j} - \frac{\partial\tilde{Z}_{0j}}{\partial v} \right) dz + \int_{\Omega_\lambda} \left(f + \sum_j d_j \Delta\tilde{Z}_{0j} \right) dz = 0.$$

Taking into account that $G(\lambda y, \lambda z) = \log \frac{1}{\lambda^2} + \log \frac{1}{|y-z|^2} + H(\lambda y, \lambda z)$, where H is the regular part of Green's function H (c.f. (1.4)) we have

$$\begin{aligned}
 \tilde{\phi}(y) - \bar{\phi} &= \frac{1}{2\pi} \int_{\partial\Omega_\lambda} \left(\log \frac{1}{|y-z|^2} + H(\lambda y, \lambda z) \right) \\
 &\quad \times \left(W\tilde{\phi} + h + \sum_j d_j \left(W\tilde{Z}_{0j} - \frac{\partial \tilde{Z}_{0j}}{\partial v} \right) \right) dz \\
 &\quad + \frac{1}{2\pi} \int_{\Omega_\lambda} \left(\log \frac{1}{|y-z|^2} + H(\lambda y, \lambda z) \right) \left(f + \sum_j d_j \Delta \tilde{Z}_{0j} \right) dz.
 \end{aligned} \tag{3.12}$$

Let us sketch now how the proof works and postpone some of the calculations.

Since $\tilde{\phi}(y) \rightarrow 0$ uniformly on sets of the form $|y - \xi'_j| < R$, we can select a sequence $R_\lambda \rightarrow \infty$ such that

$$\tilde{\phi}(y) \rightarrow 0 \quad \text{uniformly for } |y - \xi'_j| < R_\lambda.$$

We can assume $R_\lambda \rightarrow \infty$ as slowly as we need.

For each $l = 1, \dots, m$ select a point $y_l \in \partial\Omega_\lambda$ so that $|y_l - \xi'_l| = R_\lambda$. We claim that when we evaluate (3.12) at y_l all terms on the right-hand side of (3.12) converge to zero except for $\int_{\Omega_\lambda} \log \frac{1}{|y_l - z|^2} \Delta \tilde{Z}_{0j} dz = 2\pi \delta_{lj} + o(1)$ (where δ_{lj} is Kronecker's delta). Thus, we claim that

$$\tilde{\phi}(y_j) - \bar{\phi} = d_j + o(1) \quad \forall j = 1, \dots, m. \tag{3.13}$$

But the orthogonality condition (3.7) implies that

$$\sum_{j=1}^m d_j a_j = 0 \quad \text{where} \quad a_j = \int_{\Omega_\lambda} \chi_j Z_{0j}^2 > 0. \tag{3.14}$$

Multiplying (3.13) by a_j , adding and using (3.14) we find

$$\sum_{j=1}^m a_j \tilde{\phi}(y_j) - a \bar{\phi} = o(1) \quad \text{where} \quad a = \sum_{j=1}^m a_j.$$

Since $\tilde{\phi}(y_j) \rightarrow 0$ and a is bounded away from zero we reach the conclusion

$$\bar{\phi} = o(1).$$

In what follows we will obtain the necessary estimates to prove (3.13). \square

Claim.

$$\int_{\Omega_\lambda} \log \frac{1}{|y_l - z|^2} \Delta \tilde{Z}_{0j} dz = 2\pi \delta_{lj} + o(1) \quad \forall j, l = 1, \dots, m.$$

Proof. Let

$$\tilde{z}_{0j}(x) = \tilde{Z}_{0j}((F_j^\lambda)^{-1}(x)) = z_{0j}(x)p(|x|), \quad (3.15)$$

where p was defined in (3.9). Let us write $y = (F_j^\lambda)^{-1}(x)$. Since $p'(r)$ has a jump at $r = \lambda^{-\gamma}$ and $r = \lambda^{-\beta}$ and is otherwise smooth, we see that $\Delta_x \tilde{z}_{0j}$ is a measure:

$$\begin{aligned} \Delta_x \tilde{z}_{0j} &= 2\nabla z_{0j} \nabla p + z_{0j} [p'(\lambda^{-\gamma})] \mu_{\lambda^{-\gamma}} + z_{0j} [p'(\lambda^{-\beta})] \mu_{\lambda^{-\beta}} \\ &= 2\nabla z_{0j} \nabla p - z_{0j} \frac{\lambda^\gamma}{(\beta - \gamma) \log \frac{1}{\lambda}} \mu_{\lambda^{-\gamma}} + z_{0j} \frac{\lambda^\beta}{(\beta - \gamma) \log \frac{1}{\lambda}} \mu_{\lambda^{-\beta}}, \end{aligned}$$

where $[p'(r)] = p'(r^+) - p'(r^-)$ denotes the jump of p' at r and let μ_r is the one-dimensional measure on the circle of radius r .

Changing the variables yields

$$\begin{aligned} \int_{\Omega_\lambda} \Delta \tilde{z}_{0j} \varphi &= \int_{\lambda^{-\gamma} < |x| < \lambda^{-\beta}} \left(2\nabla p \nabla z_{0j} + O(\lambda|x| |\nabla^2 \tilde{z}_{0j}|) + O(\lambda |\nabla \tilde{z}_{0j}|) \right) \\ &\quad \times \varphi((F_j^\lambda)^{-1}(x)) dx \\ &\quad - \frac{\lambda^\gamma}{(\beta - \gamma) \log \frac{1}{\lambda}} \int_{r=\lambda^{-\gamma}} (1 + O(\lambda|x|)) z_{0j} \varphi((F_j^\lambda)^{-1}(x)) dx \\ &\quad + \frac{\lambda^\beta}{(\beta - \gamma) \log \frac{1}{\lambda}} \int_{r=\lambda^{-\beta}} (1 + O(\lambda|x|)) z_{0j} \varphi((F_j^\lambda)^{-1}(x)) dx \end{aligned} \quad (3.16)$$

for any $\varphi \in C(\overline{\Omega})$.

Let us consider first the case $l = j$:

$$\begin{aligned} \int_{\Omega_\lambda} \log \frac{1}{|y_j - z|^2} \Delta \tilde{z}_{0j} dz &= \int_{\Omega_\lambda} \left(\log \frac{1}{|y_j - z|^2} - \log \frac{1}{|\xi'_j - z|^2} \right) \Delta \tilde{z}_{0j} dz \\ &\quad + \int_{\Omega_\lambda} \log \frac{1}{|\xi'_j - z|^2} \Delta \tilde{z}_{0j} dz. \end{aligned} \quad (3.17)$$

By the previous remarks, using the fact that $z_{0j}(x) = 1 + O(|x|^{-1})$ and the expansion $(F_j^\lambda)^{-1}(x) = \xi'_j + x + O(\lambda|x|)$ (after rotation) we have

$$\begin{aligned} &\frac{\lambda^\gamma}{(\beta - \gamma) \log \frac{1}{\lambda}} \int_{r=\lambda^{-\gamma}} (1 + O(\lambda|x|)) z_{0j} \log \frac{1}{|\xi'_j - (F_j^\lambda)^{-1}(x)|^2} dx \\ &= \frac{\lambda^\gamma}{(\beta - \gamma) \log \frac{1}{\lambda}} (1 + O(\lambda^{1-\gamma})) (1 + O(\lambda^\gamma)) \pi \lambda^{-\gamma} (2 \log \lambda^{-\gamma} + O(\lambda^{1-\gamma})) \\ &= 2\pi \frac{\gamma}{\beta - \gamma} + O(\lambda^\theta), \end{aligned}$$

where we fix

$$0 < \theta < \min(\gamma, 1 - \beta).$$

Similarly,

$$\begin{aligned} & \frac{\lambda^\beta}{(\beta - \gamma) \log \frac{1}{\lambda}} \int_{r=\lambda^{-\beta}} z_{0j} (1 + O(\lambda|x|)) \log \frac{1}{|\xi'_j - (F_j^\lambda)^{-1}(x)|^2} dx \\ &= 2\pi \frac{\beta}{\beta - \gamma} + O(\lambda^\theta), \end{aligned}$$

and a calculation using $\nabla z_{0j} = O(\frac{1}{|x|})$, $\nabla^2 z_{0j} = O(\frac{1}{|x|^2})$ shows that

$$\begin{aligned} & \int_{\lambda^{-\gamma} < |x| < \lambda^{-\beta}} (2\nabla p \nabla z_{0j} + O(\lambda|x| |\nabla^2 \tilde{z}_{0j}|) + (\lambda |\nabla \tilde{z}_{0j}|)) \log \frac{1}{|\xi'_j - (F_j^\lambda)^{-1}(x)|^2} dx \\ &= O(\lambda^\theta). \end{aligned}$$

Therefore,

$$\int_{\Omega_\lambda} \log \frac{1}{|\xi'_j - z|^2} \Delta \tilde{Z}_{0j} dz = 2\pi + O(\lambda^\theta).$$

For the first integral in the right-hand side of (3.17) we can assume $R_\lambda \rightarrow \infty$ slowly enough so that

$$\lambda^\gamma R_\lambda \rightarrow 0.$$

Then

$$\left| \log \frac{1}{|y_j - z|^2} - \log \frac{1}{|\xi'_j - z|^2} \right| \leq C \frac{|y_j - \xi'_j|}{\lambda^{-\gamma}}$$

and it follows that

$$\left| \int_{\Omega_\lambda} \left(\log \frac{1}{|y_j - z|^2} - \log \frac{1}{|\xi'_j - z|^2} \right) \Delta \tilde{Z}_{0j} dz \right| = O(\lambda^\gamma R_\lambda).$$

Next we show that if $l \neq j$, then

$$\int_{\Omega_\lambda} \log \frac{1}{|y_l - z|^2} \Delta \tilde{Z}_{0j} dz = o(1).$$

In fact,

$$\begin{aligned} \int_{\Omega_\lambda} \log \frac{1}{|y_l - z|^2} \Delta \tilde{Z}_{0j} dz &= \int_{\Omega_\lambda} \left(\log \frac{1}{|y_l - z|^2} - \log \frac{1}{|y_l - \xi'_j|^2} \right) \Delta \tilde{Z}_{0j} dz \\ &\quad + \int_{\Omega_\lambda} \log \frac{1}{|y_l - \xi'_j|^2} \Delta \tilde{Z}_{0j} dz. \end{aligned}$$

We can assume that $R_\lambda < \frac{\lambda^{-\gamma}}{2}$ so that

$$\left| \log \frac{1}{|y_l - z|^2} - \log \frac{1}{|y_l - \xi'_j|^2} \right| \leq C\lambda |z - \xi'_j|.$$

Thus,

$$\left| \int_{\Omega_\lambda} \left(\log \frac{1}{|y_l - z|^2} - \log \frac{1}{|y_l - \xi'_j|^2} \right) \Delta \tilde{Z}_{0j} dz \right| = O(\lambda^\theta).$$

Finally,

$$\begin{aligned} \int_{\Omega_\lambda} \Delta \tilde{Z}_{0j} dz &= -(2\pi + O(\lambda^{1-\gamma}))(1 + O(\lambda^\gamma)) \frac{1}{(\beta - \gamma) \log \frac{1}{\lambda}} \\ &\quad + (2\pi + O(\lambda^{1-\beta}))(1 + O(\lambda^\beta)) \frac{1}{(\beta - \gamma) \log \frac{1}{\lambda}} + O(\lambda^\gamma) \\ &= O(\lambda^\theta) \end{aligned} \quad (3.18)$$

so

$$\log \frac{1}{|y_l - \xi'_j|^2} \int_{\Omega_\lambda} \Delta \tilde{Z}_{0j} dz = o(1). \quad \square$$

Claim. For any $0 < \alpha < 1$

$$\begin{aligned} W(y) \tilde{Z}_{0j}(y) - \frac{\partial \tilde{Z}_{0j}}{\partial v}(y) &= O\left(\frac{\lambda^\alpha}{1 + |y - \xi'_j|}\right) + O\left(\frac{\lambda}{\log \frac{1}{\lambda}}\right), \\ &\text{for } |y - \xi'_j| \leq \frac{\delta}{\lambda}. \end{aligned} \quad (3.19)$$

Proof. Set

$$\tilde{W}(x) = W((F_j^\lambda)^{-1}(x))$$

where the map F_j^λ is defined (3.3). Recall that W satisfies (2.23) and (2.24), that is

$$W(y) = \frac{2\mu_j}{|y - \xi'_j - \mu_j v'_j|^2} (1 + O(\lambda^\alpha(1 + |y|))).$$

Since $(F_j^\lambda)^{-1}(x) = \xi'_j + x + O(\lambda|x|)$ we find

$$\begin{aligned} \tilde{W}(x) &= W((F_j^\lambda)^{-1}(x)) = W(\xi'_j + x + O(\lambda|x|)) \\ &= \frac{2\mu_j}{x_1^2 + \mu_j^2} + O\left(\frac{\lambda^\alpha}{1 + |x|}\right) \quad x = (x_1, 0), |x| < \frac{\delta}{\lambda}. \end{aligned} \quad (3.20)$$

On the other hand

$$\frac{\partial \tilde{Z}_{0j}}{\partial v} = -\frac{\partial \tilde{z}_{0j}}{\partial x_2} + O(\lambda|x| |\nabla \tilde{z}_{0j}|),$$

and using the fact that p has a zero normal derivative on $\partial\mathbb{R}_+^2$ we deduce that

$$\begin{aligned} \frac{\partial \tilde{Z}_{0j}}{\partial v} &= -p \frac{\partial z_{0j}}{\partial x_2} + O(\lambda r (|\nabla p|_{z_{0j}} + p |\nabla z_{0j}|)) \\ &= -p \frac{\partial z_{0j}}{\partial x_2} + O\left(\frac{\lambda}{\log \frac{1}{\lambda}}\right) + O\left(\frac{\lambda}{1+r}\right) \quad r < \frac{\delta}{\lambda}, \end{aligned} \quad (3.21)$$

where $r = |y - \xi'_j|$ (observe that $\nabla p = O\left(\frac{1}{r \log \frac{1}{\lambda}}\right)$).

Using (3.20) we find

$$\frac{\partial \tilde{Z}_{0j}}{\partial v}(y) - W(y) \tilde{Z}_{0j}(y) = O\left(\frac{\lambda}{\log \frac{1}{\lambda}}\right) + O\left(\frac{\lambda^\alpha}{1 + |y - \xi'_j|}\right) \quad |y - \xi'_j| < \frac{\delta}{\lambda}.$$

□

Claim. Similarly

$$\int_{\partial\Omega_\lambda} \log \frac{1}{|y - z|^2} \left(W \tilde{Z}_{0j} - \frac{\partial \tilde{Z}_{0j}}{\partial v} \right) dz = O(\lambda^{1-\beta}) = o(1),$$

and uniformly for $y \in \partial\Omega_\lambda$.

Proof. Using (3.19) and the fact that if $|y - \xi'_j| > \frac{\delta}{\lambda}$ for all j we have $W(y) = O(\lambda^2)$ and $\frac{\partial \tilde{Z}_{0j}}{\partial v} = O(\lambda^2)$, we see that

$$\int_{\partial\Omega_\lambda} \left| W \tilde{Z}_{0j} - \frac{\partial \tilde{Z}_{0j}}{\partial v} \right| = O\left(\lambda^{1-\beta} \frac{1}{\log \frac{1}{\lambda}}\right). \quad (3.22)$$

Since $\log \frac{1}{|y-z|^2} = O\left(\log \frac{1}{\lambda}\right)$ for $|y - z| > R$ where $R > 0$ is fixed and

$$\int_{\partial\Omega_\lambda \cap B_R(y)} \left| \log \frac{1}{|y - z|^2} \right| dz \leq C,$$

we conclude the validity of the assertion. □

Claim.

$$\int_{\partial\Omega_\lambda} \log \frac{1}{|y - z|^2} h(z) dz = o(1) \quad (3.23)$$

and

$$\int_{\Omega_\lambda} \log \frac{1}{|y - z|^2} f(z) dz = o(1). \quad (3.24)$$

Proof. We have $\log \frac{1}{|y-z|^2} = O(\log \frac{1}{\lambda})$ for $|y-z| > R$ where $R > 0$ is fixed, and $\int_{\partial\Omega_\lambda \cap B_R(y)} \left| \log \frac{1}{|y-z|^2} \right| dz \leq C$, and therefore

$$\left| \int_{\partial\Omega_\lambda} \log \frac{1}{|y-z|^2} h dz \right| \leq C \log \frac{1}{\lambda} \|h\|_* = o(1)$$

by hypothesis (3.8). The proof of the other assertion is similar. \square

Claim.

$$\int_{\partial\Omega_\lambda} \log \frac{1}{|y-z|^2} W\tilde{\phi} dz = o(1).$$

Proof. Arguing as before, it is sufficient to show that

$$\log \frac{1}{\lambda} \int_{\partial\Omega_\lambda} W\tilde{\phi} = o(1).$$

Integrating equation (3.10) we find

$$\begin{aligned} & \int_{\partial\Omega_\lambda} \left(W\tilde{\phi} + h + \sum_j d_j \left(W\tilde{Z}_{0j} - \frac{\partial\tilde{Z}_{0j}}{\partial v} \right) \right) dz \\ & + \int_{\Omega_\lambda} \left(f + \sum_j d_j \Delta\tilde{Z}_{0j} \right) dz = 0. \end{aligned}$$

The conclusion follows then from (3.18), (3.22), (3.8). \square

Claim.

$$\begin{aligned} A & \equiv \int_{\partial\Omega_\lambda} H(\lambda y, \lambda z) \left(W\tilde{\phi} + h + \sum_j d_j \left(W\tilde{Z}_{0j} - \frac{\partial\tilde{Z}_{0j}}{\partial v} \right) \right) dz \\ & + \int_{\Omega_\lambda} H(\lambda y, \lambda z) \left(f + \sum_j d_j \Delta\tilde{Z}_{0j} \right) dz = o(1), \end{aligned}$$

uniformly for $y \in \partial\Omega_\lambda$.

Proof. Let

$$\zeta(r) = \begin{cases} 1 & \text{if } r < \lambda^{-1/2} \\ \frac{\log \frac{\delta}{\lambda} - \log r}{\log \frac{\delta}{\lambda} - \log \lambda^{-1/2}} & \text{if } \lambda^{-1/2} < r < \frac{\delta}{\lambda} \\ 0 & \text{if } r > \frac{\delta}{\lambda} \end{cases}$$

and set

$$\psi(z) = \sum_{j=1}^m H(\lambda y, \xi_j) \zeta(|z - \xi_j'|).$$

Multiplying (3.10) by ψ and integrating by parts we have

$$\int_{\Omega_\lambda} \left(f + \sum_j d_j \Delta \tilde{Z}_{0j} \right) \psi + \int_{\partial\Omega_\lambda} \left(W\tilde{\phi} + h + \sum_j d_j \left(W\tilde{Z}_{0j} - \frac{\partial\tilde{Z}_{0j}}{\partial\nu} \right) \right) \psi - \int_{\partial\Omega_\lambda} \tilde{\phi} \frac{\partial\psi}{\partial\nu} + \int_{\Omega_\lambda} \tilde{\phi} \Delta\psi = 0.$$

Subtracting this from A we find that

$$\begin{aligned} A &= \int_{\partial\Omega_\lambda} (H(\lambda y, \lambda z) - \psi) \left(W\tilde{\phi} + h + \sum_j d_j \left(W\tilde{Z}_{0j} - \frac{\partial\tilde{Z}_{0j}}{\partial\nu} \right) \right) dz \\ &\quad + \int_{\Omega_\lambda} (H(\lambda y, \lambda z) - \psi) \left(f + \sum_j d_j \Delta\tilde{Z}_{0j} \right) dz \\ &\quad + \int_{\partial\Omega_\lambda} \tilde{\phi} \frac{\partial\psi}{\partial\nu} - \int_{\Omega_\lambda} \tilde{\phi} \Delta\psi. \end{aligned}$$

Since H and ψ are bounded

$$\left| \int_{\partial\Omega_\lambda} (H(\lambda y, \lambda z) - \psi) h dz \right| \leq C \|h\|_* = o(1) \quad (3.25)$$

$$\left| \int_{\Omega_\lambda} (H(\lambda y, \lambda z) - \psi) f dz \right| \leq C \|f\|_{**} = o(1). \quad (3.26)$$

A calculation shows that

$$\int_{\Omega_\lambda} \tilde{\phi} \Delta\psi = O\left(\frac{1}{\log\frac{\delta}{\lambda}}\right) = o(1), \quad \int_{\partial\Omega_\lambda} \tilde{\phi} \frac{\partial\psi}{\partial\nu} = O\left(\frac{1}{\log\frac{\delta}{\lambda}}\right) = o(1). \quad (3.27)$$

For instance, the first integral in (3.27) can be estimated as follows

$$\left| \int_{\Omega_\lambda} \tilde{\phi} \Delta\psi \right| \leq \|\tilde{\phi}\|_{L^\infty(\Omega_\lambda)} \int_{\Omega_\lambda} |\Delta\psi|.$$

But $\Delta\psi$ is a measure with support on the arcs $r = \lambda^{-1/2}$, $r = \frac{\delta}{\lambda}$ ($r = |z - \xi'_j|$) and

$$\int_{\Omega_\lambda} |\Delta\psi| = O\left(\lambda^{-1/2} \frac{1}{\lambda^{-1/2} \log\frac{1}{\lambda}} + \frac{\delta}{\lambda} \frac{1}{\frac{\delta}{\lambda} \log\frac{1}{\lambda}}\right) = O\left(\frac{1}{\log\frac{1}{\lambda}}\right) = o(1).$$

Now, at distance greater than $\frac{\delta}{\lambda}$ from all ξ'_j we have $W = O(\lambda^2)$ and $H, \tilde{\phi}$ are bounded, thus

$$\int_{\partial\Omega_\lambda \setminus (\cup_j B_{\delta/\lambda}(\xi'_j))} (H(\lambda y, \lambda z) - \psi) W\tilde{\phi} = o(1). \quad (3.28)$$

On the other hand, at distance less than $\frac{\delta}{\lambda}$ from ξ'_j we have $H(\lambda y, \lambda z) - H(\lambda y, \xi_j) = O(\lambda|z - \xi'_j|)$ and $W = O(\frac{1}{r^2})$, $r = |z - \xi'_j|$. So

$$\begin{aligned} & \left| \int_{\partial\Omega_\lambda \cap B_{\lambda^{-1/2}}(\xi'_j)} (H(\lambda y, \lambda z) - \psi(z)) W \tilde{\phi} dz \right| \\ &= \left| \int_{\partial\Omega_\lambda \cap B_{\lambda^{-1/2}}(\xi'_j)} (H(\lambda y, \lambda z) - H(\lambda y, \xi_j)) W \tilde{\phi} dz \right| \\ &\leq \lambda \int_1^{\delta/\lambda} \frac{1}{r} dr = O\left(\lambda \log \frac{1}{\lambda}\right) = o(1). \end{aligned} \quad (3.29)$$

In the region $\lambda^{-1/2} < r = |z - \xi'_j| < \frac{\delta}{\lambda}$ we use the fact that H , ζ , $\tilde{\phi}$ are bounded and that $W = O(\frac{1}{r^2})$, so

$$\begin{aligned} \left| \int_{\partial\Omega_\lambda \cap B_{\delta/\lambda}(\xi'_j) \setminus B_{\lambda^{-1/2}}(\xi'_j)} (H(\lambda y, \lambda z) - \psi(z)) W \tilde{\phi} dz \right| &\leq C \int_{\lambda^{-1/2}}^{\delta/\lambda} \frac{1}{r^2} dr \\ &= O(\lambda^{1/2}) \\ &= o(1). \end{aligned} \quad (3.30)$$

Collecting (3.25)–(3.30) and recalling (3.18), (3.19) we obtain the desired conclusion. \square

Proof of Lemma 4. Let $\hat{\phi}(x) = \tilde{\phi}(x/\lambda)$, $x \in \Omega$. Then $\hat{\phi}$ satisfies

$$\begin{cases} -\Delta \hat{\phi} = \frac{1}{\lambda^2} \left(\hat{f} + \sum_{j=1}^m d_j \Delta \hat{Z}_{0j} \right) & \text{in } \Omega \\ \frac{\partial \hat{\phi}}{\partial \nu} = \frac{1}{\lambda} \left(\hat{W} \hat{\phi} + \hat{h} \sum_{j=1}^m d_j \left(\hat{W} \hat{Z}_{0j} - \frac{\partial \hat{Z}_{0j}}{\partial \nu} \right) \right) & \text{on } \partial\Omega, \end{cases}$$

where $\hat{f}(x) = f(x/\lambda)$, $\hat{h}(x) = h(x/\lambda)$, $\hat{W}(x) = W(x/\lambda)$ and $\hat{Z}_{01}(x) = Z_{01}(x/\lambda)$. For a given $\delta > 0$ let $E_\delta = \Omega \setminus \cup_{j=1}^m B_\delta(\xi_j)$. Then $\frac{1}{\lambda^2} \|\hat{f}\|_{L^\infty(E_\delta)} \leq C \|f\|_{**} \rightarrow 0$, $\frac{1}{\lambda} \|\hat{h}\|_{L^\infty(\partial E_\delta)} \leq C \|h\|_* \rightarrow 0$, and $\frac{1}{\lambda} \|\hat{W} \hat{\phi}\|_{L^\infty(E_\delta)} \leq C \lambda$. Furthermore, in E_δ we have that $\hat{Z}_{0j} \equiv 0$. We also know that $\|\hat{\phi}\|_{L^\infty(\Omega)} \leq 1$ and $f_{\partial\Omega} \hat{\phi} \rightarrow 0$. From this it follows that $\hat{\phi} \rightarrow 0$ uniformly in E_δ and this implies

$$\tilde{\phi} \rightarrow 0 \text{ uniformly in } \Omega_\lambda \setminus \cup_{j=1}^m B_{\delta/\lambda}(\xi'_j), \text{ for any } \delta > 0.$$

For a given $R_1 > 0$ let A_j denote the annulus

$$A_j = B_{\delta/\lambda}(\xi'_j) \setminus B_{R_1}(\xi'_j).$$

Given $\lambda > 0$ small enough, there exist $R_1 > 0$ independent of λ and $\psi_j : \Omega_\lambda \cap A_j \rightarrow \mathbb{R}$ smooth and positive so that

$$\begin{aligned} -\Delta \psi_j &\geq \frac{c}{|y - \xi'_j|^{2+\sigma}} && \text{in } \Omega_\lambda \cap A_j \\ \frac{\partial \psi_j}{\partial \nu} - W \psi_j &\geq \frac{c}{|y - \xi'_j|^{1+\sigma}} && \text{on } \partial\Omega_\lambda \cap A_j \\ c &\leq \psi_j \leq C && \text{in } \Omega_\lambda \cap A_j, \end{aligned}$$

where $C, c > 0$ can be made independent of λ . Indeed, the function

$$\psi_j(y) = \frac{(y - \xi'_j) \cdot \nu'_j}{|y - \xi'_j|^{1+\sigma}} + C_0 \left(1 - \frac{1}{|y - \xi'_j|^\sigma} \right)$$

with C_0 a fixed large constant satisfies the requirements, see [4] (Lemma 4.3).

Thanks to the barrier ψ_j we deduce that the following maximum principle holds in $\Omega_\lambda \cap A_j$: if $\phi \in H^1(\Omega_\lambda \cap A_j)$ satisfies

$$\begin{cases} -\Delta\phi & \geq 0 & \text{in } \Omega_\lambda \cap A_j \\ \frac{\partial\phi}{\partial\nu} - W\phi & \geq 0 & \text{on } \partial\Omega_\lambda \cap A_j \\ \phi & \geq 0 & \text{on } \Omega_\lambda \cap A_j, \end{cases}$$

then $\phi \geq 0$ in $\Omega_\lambda \cap A_j$. By the properties of ψ_j and this maximum principle we deduce that there exists a fixed $C > 0$ so that

$$|\phi| \leq C\psi_j \left(\sup_{\Omega_\lambda \cap \partial B_{R_1}(\xi'_j)} |\phi| + \sup_{\Omega_\lambda \cap \partial B_{\delta/\lambda}(\xi'_j)} |\phi| + \|h\|_* + \|f\|_{**} \right) \quad \text{in } \Omega_\lambda \cap A_j.$$

But $\sup_{\Omega_\lambda \cap \partial B_{R_1}(\xi'_j)} |\phi| \rightarrow 0$ by Lemma 6, and $\sup_{\Omega_\lambda \cap \partial B_{\delta/\lambda}(\xi'_j)} |\phi| \rightarrow 0$ as shown above. This proves the result. \square

Proof of Lemma 5. Multiplying (3.10) by \tilde{Z}_{0j} and integrating we obtain

$$\begin{aligned} & d_j \left(\int_{\Omega_\lambda} (-\Delta\tilde{Z}_{0j})\tilde{Z}_{0j} + \int_{\partial\Omega_\lambda} \tilde{Z}_{0j} \left(\frac{\partial\tilde{Z}_{0j}}{\partial\nu} - W\tilde{Z}_{0j} \right) \right) \\ &= - \int_{\partial\Omega_\lambda} \tilde{Z}_{0j}h - \int_{\Omega_\lambda} \tilde{Z}_{0j}f + \int_{\partial\Omega_\lambda} \tilde{\phi} \left(\frac{\partial\tilde{Z}_{0j}}{\partial\nu} - W\tilde{Z}_{0j} \right) + \int_{\Omega_\lambda} \tilde{\phi}(-\Delta\tilde{Z}_{0j}). \end{aligned}$$

We claim that

$$\int_{\Omega_\lambda} (-\Delta\tilde{Z}_{0j})\tilde{Z}_{0j} + \int_{\partial\Omega_\lambda} \tilde{Z}_{0j} \left(\frac{\partial\tilde{Z}_{0j}}{\partial\nu} - W\tilde{Z}_{0j} \right) \geq \frac{c}{\log\frac{1}{\lambda}}, \quad (3.31)$$

for some fixed $c > 0$. Assuming this for a moment, we can prove the lemma since

$$\left| \int_{\partial\Omega_\lambda} \tilde{Z}_{0j}h \right| \leq \|h\|_* \|\tilde{Z}_{0j}\|_{L^\infty(\Omega_\lambda)} \leq C \log\frac{1}{\lambda} \|h\|_* \frac{1}{\log\frac{1}{\lambda}} = o(1) \frac{1}{\log\frac{1}{\lambda}}$$

and

$$\int_{\Omega_\lambda} \tilde{Z}_{0j}f = o(1) \frac{1}{\log\frac{1}{\lambda}}.$$

Similarly, the other terms can be estimated as follows

$$\left| \int_{\partial\Omega_\lambda} \tilde{\phi} \left(\frac{\partial\tilde{Z}_{0j}}{\partial\nu} - W\tilde{Z}_{0j} \right) \right| \leq \|\tilde{\phi}\|_{L^\infty(\Omega_\lambda)} \int_{\partial\Omega_\lambda} \left| \frac{\partial\tilde{Z}_{0j}}{\partial\nu} - W\tilde{Z}_{0j} \right| = O(\lambda^{1-\beta}),$$

using (3.22) and

$$\left| \int_{\Omega_\lambda} \tilde{\phi}(-\Delta \tilde{Z}_{0j}) \right| \leq \|\tilde{\phi}\|_{L^\infty(\Omega_\lambda)} \int_{\Omega_\lambda} |\Delta \tilde{Z}_{0j}| \leq \frac{C}{\log \frac{1}{\lambda}} \|\tilde{\phi}\|_{L^\infty(\Omega_\lambda)} = o(1) \frac{1}{\log \frac{1}{\lambda}}.$$

Let us prove (3.31) using (3.16) to compute $\int_{\Omega_\lambda} \Delta \tilde{Z}_{0j} \tilde{Z}_{0j}$. For the part of $\Delta \tilde{Z}_{0j}$ supported on $r = \lambda^{-\gamma}$ we have

$$\begin{aligned} & \frac{\lambda^\gamma}{(\beta - \gamma) \log \frac{1}{\lambda}} \int_{r=\lambda^{-\gamma}} (1 + O(\lambda|x|)) z_{0j}^2 p \, dx \\ &= \frac{\lambda^\gamma}{(\beta - \gamma) \log \frac{1}{\lambda}} (1 + O(\lambda^{1-\gamma}))(1 + O(\lambda^\gamma))^2 \pi \lambda^{-\gamma} \\ &= \frac{\pi \gamma}{(\beta - \gamma) \log \frac{1}{\lambda}} + O(\lambda^\theta) \quad 0 < \theta < \min(\gamma, 1 - \beta). \end{aligned}$$

Analogously, for the part supported on $r = \lambda^{-\beta}$ we find that

$$\frac{\lambda^\beta}{(\beta - \gamma) \log \frac{1}{\lambda}} \int_{r=\lambda^{-\beta}} (1 + O(\lambda|x|)) z_{0j}^2 p \, dx = 0$$

since $p(\lambda^{-\beta}) = 0$. Also,

$$\int_{\lambda^{-\gamma} < |x| < \lambda^{-\beta}} (2\nabla p \nabla z_{0j} + O(\lambda|x| |\nabla^2 z_{0j}|) + (\lambda |\nabla \tilde{Z}_{0j}|)) z_{0j} p \, dx = O(\lambda^\theta).$$

Thus,

$$\int_{\Omega_\lambda} \Delta \tilde{Z}_{0j} \tilde{Z}_{0j} = -\frac{\pi \gamma}{(\beta - \gamma) \log \frac{1}{\lambda}} + O(\lambda^\theta).$$

Finally, similarly to (3.22)

$$\int_{\partial \Omega_\lambda} \tilde{Z}_{0j} \left(\frac{\partial \tilde{Z}_{0j}}{\partial \nu} - W \tilde{Z}_{0j} \right) = O(\lambda^{1-\beta}),$$

and this proves (3.31). \square

Proposition 2. *Let $d > 0$ and m a positive even integer. Then there exists $\lambda_0 > 0$ such that for any $0 < \lambda < \lambda_0$, any family of points $\xi_1, \dots, \xi_m \in \partial \Omega$ satisfying (3.2) (i.e. $|\xi_i - \xi_j| \geq d \forall i \neq j$), and any $h \in L^\infty(\partial \Omega_\lambda)$, $f \in L^\infty(\Omega_\lambda)$ there is a unique solution $\phi \in L^\infty(\Omega_\lambda)$, $c_0, c_1, \dots, c_m \in \mathbb{R}$ to*

$$\begin{cases} -\Delta \phi &= f \quad \text{in } \Omega_\lambda \\ \frac{\partial \phi}{\partial \nu} - W \phi &= h + \sum_{j=1}^m c_j \chi_j Z_{1j} + c_0 \chi Z \quad \text{on } \partial \Omega_\lambda \\ \int_{\Omega_\lambda} \chi_j Z_{1j} \phi = 0 \quad \forall j = 1, \dots, m, & \int_{\Omega_\lambda} \chi Z \phi = 0. \end{cases} \quad (3.32)$$

Moreover, there is $C > 0$ independent of λ such that

$$\begin{aligned} \|\phi\|_{L^\infty(\Omega_\lambda)} &\leq C \log \frac{1}{\lambda} (\|h\|_* + \|f\|_{**}), \\ \max(|c_0|, \dots, |c_m|) &\leq C (\|h\|_* + \|f\|_{**}). \end{aligned}$$

Proof. We first deal with the following linear problem

$$\begin{cases} -\Delta\phi &= f + \sum_{j=1}^m d_j \chi_j Z_{1j} + d_0 \chi Z & \text{in } \Omega_\lambda \\ \frac{\partial\phi}{\partial\nu} - W\phi &= h & \text{on } \partial\Omega_\lambda \\ \int_{\Omega_\lambda} \chi_j Z_{1j} \phi = 0 & \forall j = 1, \dots, m, & \int_{\Omega_\lambda} \chi Z \phi = 0, \end{cases} \quad (3.33)$$

where $h \in L^\infty(\partial\Omega_\lambda)$, $f \in L^\infty(\Omega_\lambda)$ are given. Let us show that for any $\phi \in L^\infty(\Omega_\lambda)$, d_0, d_1, \dots, d_m solution to (3.33) we have

$$\|\phi\|_{L^\infty(\Omega_\lambda)} \leq C \log \frac{1}{\lambda} (\|h\|_* + \|f\|_{**}) \quad (3.34)$$

$$|d_j| \leq C (\|h\|_* + \|f\|_{**}) \quad \forall j = 0, \dots, m. \quad (3.35)$$

Given that from Proposition 1 we have

$$\|\phi\|_{L^\infty(\Omega_\lambda)} \leq C \log \frac{1}{\lambda} \left(\|h\|_* + \|f\|_{**} + \sum_{j=0}^m |d_j| \right), \quad (3.36)$$

it suffices to prove that (3.35) holds.

Consider a cut-off function $\bar{\eta}$ such that

$$\begin{aligned} \bar{\eta} &\equiv 1 \text{ in } B_{\frac{\delta}{4\lambda}}(0), \bar{\eta} \equiv 0 \text{ in } \mathbb{R}^2 \setminus B_{\frac{\delta}{3\lambda}}(0) \\ 0 &\leq \bar{\eta} \leq 1, |\nabla\bar{\eta}| \leq C\lambda/\delta, |\nabla^2\bar{\eta}| \leq C\lambda^2/\delta^2 \text{ in } \mathbb{R}^2, \end{aligned}$$

and for $j = 1, \dots, m$ set

$$\eta_j(y) = \bar{\eta}(F_j^\lambda(y)),$$

where F_j^λ is defined in (3.3). Multiplying (3.33) by $\eta_i Z_{1i}$, $i = 1, \dots, m$ and integrating by parts we obtain

$$\begin{aligned} d_i \int_{\Omega_\lambda} \chi_i Z_{1i}^2 &= - \int_{\partial\Omega_\lambda} h \eta_i Z_{1i} - \int_{\Omega_\lambda} f \eta_i Z_{1i} + \int_{\partial\Omega_\lambda} \phi \frac{\partial\eta_i}{\partial\nu} Z_{1i} \\ &+ \int_{\partial\Omega_\lambda} \phi \eta_i \left(\frac{\partial Z_{1i}}{\partial\nu} - W Z_{1i} \right) - \int_{\Omega_\lambda} \phi \Delta(\eta_i Z_{1i}). \end{aligned} \quad (3.37)$$

Since $Z_{1i} = O\left(\frac{1}{1+r}\right)$ and $\nabla\bar{\eta} = O(\lambda)$, we have

$$\left| \int_{\partial\Omega_\lambda} \phi \frac{\partial\eta_i}{\partial\nu} Z_{1i} \right| \leq C \|\phi\|_{L^\infty(\Omega_\lambda)} \lambda \log \frac{1}{\lambda}. \quad (3.38)$$

As in the proof of (3.19) we have

$$\frac{\partial Z_{1i}}{\partial\nu}(y) - W(y) Z_{1i}(y) = O\left(\frac{\lambda^\alpha}{1 + |y - \xi'_j|}\right) \quad y \in \partial\Omega_\lambda, |y - \xi'_j| < \frac{\delta}{\lambda},$$

and this implies

$$\int_{\partial\Omega_\lambda} \eta_i \left| \frac{\partial Z_{1i}}{\partial v} - WZ_{1i} \right| = O(\lambda^{\alpha'}), \quad i = 1, \dots, m, \quad (3.39)$$

where $0 < \alpha' < \alpha$. Since $0 < \alpha < 1$ is arbitrary, so is α' , and so from now on we will just write α .

We also compute

$$\begin{aligned} \Delta(\eta_i Z_{1i}) &= \Delta\eta_i Z_{1i} + 2\nabla\eta_i \nabla Z_{1i} + \eta_i \Delta Z_{1i} \\ &= O\left(\frac{\lambda^2}{1+r}\right) + O\left(\frac{\lambda}{1+r^2}\right) + \eta_i \Delta Z_{1i}. \end{aligned}$$

However, $\Delta_y Z_{1i} = \Delta_x z_1 + O(\lambda|x||\nabla^2 z_1|) + O(\lambda|\nabla z_1|)$ (where we have rotated Ω appropriately and $x = F_i^\lambda(y)$). Thus,

$$\Delta Z_{1i} = O\left(\frac{\lambda}{1+r^2}\right) + O\left(\frac{\lambda^2}{1+r}\right)$$

and it follows that

$$\int_{\Omega_\lambda} |\Delta(\eta_i Z_{1i})| = O\left(\lambda \log \frac{1}{\lambda}\right) = O(\lambda^\alpha). \quad (3.40)$$

Combining (3.37), (3.39) and (3.40) we conclude that

$$d_i \int_{\Omega_\lambda} \chi_i Z_{1i}^2 \leq C(\|h\|_* + \|f\|_{**} + \lambda^\alpha \|\phi\|_{L^\infty(\Omega_\lambda)}),$$

and this together with (3.36) yields

$$|d_i| \leq C \left(\|h\|_* + \|f\|_{**} + \lambda^\alpha \sum_{j=0}^m |d_j| \right), \quad i = 1, \dots, m. \quad (3.41)$$

On the other hand, multiplying (3.33) by Z we obtain

$$d_0 \int_{\Omega_\lambda} \chi Z^2 = - \int_{\Omega_\lambda} fZ - \int_{\partial\Omega_\lambda} hZ + \int_{\partial\Omega_\lambda} \phi \left(\frac{\partial Z}{\partial v} - WZ \right) - \int_{\Omega_\lambda} \phi \Delta Z. \quad (3.42)$$

We estimate as before

$$\left| \int_{\partial\Omega_\lambda} \phi \left(\frac{\partial Z}{\partial v} - WZ \right) \right| \leq \|\phi\|_{L^\infty(\Omega_\lambda)} \int_{\partial\Omega_\lambda} \left| \frac{\partial Z}{\partial v} - WZ \right| \leq C\lambda^{b/2} \|\phi\|_{L^\infty(\Omega_\lambda)} \quad (3.43)$$

and

$$\left| \int_{\Omega_\lambda} \phi \Delta Z \right| \leq \|\phi\|_{L^\infty(\Omega_\lambda)} \int_{\Omega_\lambda} |\Delta Z| \leq C\lambda^b \|\phi\|_{L^\infty(\Omega_\lambda)}. \quad (3.44)$$

From (3.42) and (3.36) we see that

$$|d_0| \leq C \left(\|h\|_* + \|f\|_{**} + \lambda^{b/2} \sum_{j=0}^m |d_j| \right).$$

Using this and (3.41) we deduce (3.35) and (3.34).

Now consider the Hilbert space

$$H = \left\{ \phi \in H^1(\Omega_\lambda) : \int_{\Omega_\lambda} \chi Z \phi = 0, \int_{\Omega_\lambda} \chi_j Z_{1j} \phi = 0 \quad \forall j = 1, \dots, m \right\}$$

with the norm $\|\phi\|_{H^1}^2 = \int_{\Omega_\lambda} |\nabla \phi|^2$. Indeed, this is a norm in H since by choosing R_0 large enough (the size of the support of the cut-off functions χ_j , c.f. (3.4)) we have $\int_{\Omega_\lambda} \chi Z \neq 0$. Equation (3.33) can be formulated as to find $\phi \in H$ such that

$$\int_{\Omega_\lambda} \nabla \phi \nabla \psi - \int_{\partial\Omega_\lambda} W \phi \psi = \int_{\Omega_\lambda} f \psi + \int_{\partial\Omega_\lambda} h \psi \quad \forall \psi \in H.$$

By (3.34) this problem has at most one solution, and by Fredholm's alternative we deduce that given f, h there indeed exists a solution.

For convenience of notation in the rest of the proof we write

$$Z_0 = Z, \quad \chi_0 = \chi \quad \text{and} \quad Z_j = Z_{1j} \quad \forall j = 1, \dots, m.$$

Let $Y_i \in L^\infty(\Omega_\lambda), d_{ij} \in \mathbb{R}i, j = 0, \dots, m$ be the solution to (3.33) with $h = \chi_i Z_{1i}$ and $f = 0$, that is

$$\begin{cases} -\Delta Y_i &= \sum_{j=0}^m d_{ij} \chi_j Z_j \quad \text{in } \Omega_\lambda \\ \frac{\partial Y_i}{\partial \nu} - W Y_i &= -\chi_i Z_i \quad \text{on } \partial\Omega_\lambda \\ \int_{\Omega_\lambda} \chi_j Z_j Y_i &= 0 \quad \forall j = 0, \dots, m. \end{cases} \quad (3.45)$$

There exists a unique $Y_i \in L^\infty(\Omega_\lambda)$ solution to this equation and we have the estimates

$$\|Y_i\|_{L^\infty(\Omega_\lambda)} \leq C \log \frac{1}{\lambda}, \quad |d_{ij}| \leq C, \quad (3.46)$$

for some constant C independent of λ . We shall show that $d_{ij} = A \delta_{ij} + O(\lambda^{b/2})$ where $A > 0$ is independent of λ . Indeed, writing $\eta_0 \equiv 1$, let us multiply (3.45) by $\eta_j Z_j$ and integrate by parts

$$\begin{aligned} d_{ij} \int_{\Omega_\lambda} \chi_j Z_j^2 + \delta_{ij} \int_{\partial\Omega_\lambda} \chi_j Z_j^2 &= \int_{\partial\Omega_\lambda} \left(\frac{\partial Z_j}{\partial \nu} - W Z_j \right) \eta_j Y_i + \int_{\partial\Omega_\lambda} \frac{\partial \eta_j}{\partial \nu} Z_j Y_i \\ &\quad - \int_{\Omega_\lambda} Y_i \Delta(\eta_j Z_j) \\ &= O(\lambda^{b/2}). \end{aligned}$$

To estimate the integrals in the right-hand side above for the case $j = 1, \dots, m$, we argue exactly as in (3.38), (3.39), and (3.40). For the case $j = 0$ we use (3.43) and (3.44).

It follows that the matrix D with entries d_{ij} $i, j = 0, \dots, m$ is invertible for small λ and $\|D^{-1}\| \leq C$ uniformly in λ . To prove the solvability of (3.32) let $f \in L^\infty(\Omega_\lambda)$, $h \in L^\infty(\partial\Omega_\lambda)$ be given. We find ϕ_1, d_0, \dots, d_m the solution to (3.33) and define

$$\phi = \phi_1 + \sum_{i=0}^m c_i Y_i,$$

where c_i is such that $\sum_{i=0}^m c_i d_{ij} = -d_j \forall j = 0, \dots, m$. Then ϕ satisfies (3.32) and we have the estimate

$$\begin{aligned} \|\phi\|_{L^\infty(\Omega_\lambda)} &\leq \|\phi_1\|_{L^\infty(\Omega_\lambda)} + \sum_{i=0}^m |c_i| \leq C \log \frac{1}{\lambda} (\|f\|_{**} + \|h\|_*) + C \sum_{i=1}^m |d_i| \\ &\leq C \log \frac{1}{\lambda} (\|f\|_{**} + \|h\|_*), \end{aligned}$$

by (3.35). \square

The previous result implies that the unique solution $\phi = T(h)$ of (3.32) with $f = 0$ defines a continuous linear map from $L^\infty(\partial\Omega_\lambda)$ with the norm $\|\cdot\|_*$ into $L^\infty(\Omega_\lambda)$. For fixed $h \in L^\infty(\partial\Omega)$ let us compute the derivative of $\phi = T(h)$ with respect to ξ'_l . Formally $Y = \partial_{\xi'_l} \phi$ satisfies the equation

$$\Delta Y = 0 \quad \text{in } \Omega_\lambda,$$

and on $\partial\Omega_\lambda$ the boundary condition

$$\frac{\partial Y}{\partial \nu} - WY = \partial_{\xi'_l}(W)\phi + c_l \partial_{\xi'_l}(Z_{1l}\chi_l) + \sum_{j=1}^m d_j Z_{1j}\chi_j + c_0 \partial_{\xi'_l}(\chi Z) + d_0 \chi Z$$

where (still formally) $d_j = \partial_{\xi'_l}(c_j)$, $j = 0, \dots, m$. The orthogonality conditions now become

$$\begin{aligned} \int_{\Omega_\lambda} Z_{1j}\chi_j Y &= 0, \quad \text{if } j \neq l \\ \int_{\Omega_\lambda} Z_{1l}\chi_l Y &= - \int_{\Omega_\lambda} \partial_{\xi'_l}(Z_{1l}\chi_l)\phi. \\ \int_{\Omega_\lambda} \chi Z Y &= - \int_{\Omega_\lambda} \partial_{\xi'_l}(\chi Z)\phi. \end{aligned}$$

Let us write $\tilde{Y} = Y + b_l \chi_l Z_{1l} + b_0 \chi Z$ where b_0, b_l are defined through

$$b_0 \int_{\Omega_\lambda} \chi^2 Z^2 \equiv \int_{\Omega_\lambda} \phi \partial_{\xi'_l}(\chi Z), \quad b_l \int_{\Omega_\lambda} \chi_l^2 Z_{1l}^2 \equiv \int_{\Omega_\lambda} \phi \partial_{\xi'_l}(\chi_l Z_{1l}).$$

Hence, $\int_{\Omega_\lambda} \tilde{Y} \chi_j Z_{1j} = 0$ for all j and $\int_{\Omega_\lambda} \tilde{Y} \chi Z = 0$, \tilde{Y} satisfies the equation

$$\Delta \tilde{Y} = a \quad \text{in } \Omega_\lambda$$

and the boundary condition

$$\frac{\partial \tilde{Y}}{\partial \nu} - W \tilde{Y} = b + \sum_{j=1}^m d_j Z_{1j} \chi_j + d_0 \chi Z,$$

where

$$a = b_l \Delta(\chi_l Z_{1l}) + b_0 \Delta(\chi Z)$$

and

$$b = \partial_{\xi'_l}(W) \phi + c_l \partial_{\xi'_l}(Z_{1l} \chi_l) + c_0 \partial_{\xi'_l}(\chi Z) + b_l \left(\frac{\partial(\chi_l Z_{1l})}{\partial \nu} - W \chi_l Z_{1l} \right) + b_0 \left(\frac{\partial(\chi Z)}{\partial \nu} - W \chi Z \right),$$

with

$$\|b\|_* \leq C \log \frac{1}{\lambda} \|h\|_*, \quad \|a\|_{**} \leq C \log \frac{1}{\lambda} \|h\|_*.$$

Thus,

$$\|\partial_{\xi'_l} \phi\|_{L^\infty(\Omega_\lambda)} \leq C \left(\log \frac{1}{\lambda} \right)^2 \|h\|_*. \quad (3.47)$$

4. The nonlinear problem with constraints

Let τ be a small parameter and consider

$$V_1(y) = V(y) + \tau Z(y) \quad y \in \Omega_\lambda, \quad (4.1)$$

where V is given by (2.16), and Z is the function introduced in (3.5) at the beginning of Section 3.

A function v of the form

$$v(y) = V_1(y) + \tilde{\phi}(y), \quad y \in \Omega_\lambda$$

is a solution for (2.15) if and only if $\tilde{\phi}$ solves

$$\begin{cases} \Delta \tilde{\phi} = 0 & \text{in } \Omega_\lambda, \\ \frac{\partial \tilde{\phi}}{\partial \nu} - W_1 \tilde{\phi} = R_1 + N_1(\tilde{\phi}) & \text{on } \partial \Omega_\lambda, \end{cases} \quad (4.2)$$

where

$$W_1 = 2\lambda^2 \cosh V_1, \quad (4.3)$$

$$R_1 = - \left[\frac{\partial V_1}{\partial \nu} - 2\lambda^2 \sinh V_1 \right] \quad (4.4)$$

and

$$N_1(\tilde{\phi}) = 2\lambda^2 \left[\sinh(V_1 + \tilde{\phi}) - \sinh V_1 - \cosh(V_1)\tilde{\phi} \right]. \quad (4.5)$$

Observe that from the definition of the function Z we see that $Z(y) = O(1)$ all over Ω_λ . This readily implies that

$$V_1(y) = V(y) + O(|\tau|) \quad \forall y \in \Omega_\lambda. \quad (4.6)$$

We consider first the following auxiliary nonlinear problem

$$\begin{cases} \Delta\phi_1 &= 0 & \text{in } \Omega_\lambda, \\ \frac{\partial\phi_1}{\partial\nu} - W_1\phi_1 &= R_1 + N_1(\phi_1) + \sum_{j=1}^m c_j \chi_j Z_{1j} + c_0 \chi Z & \text{on } \partial\Omega_\lambda, \\ \int_{\Omega_\lambda} \chi_j Z_{1j} \phi_1 &= 0 \quad \forall j = 1, \dots, m, & \int_{\Omega_\lambda} \chi Z \phi_1 = 0 \end{cases} \quad (4.7)$$

where W_1 is as in (4.3) and N_1, R_1 are defined in (4.5) and (4.4) respectively.

Lemma 8. *Let $m > 0, d > 0$. Let α be any number in the interval $(0, 1)$ and $\tau = O(\lambda^\theta)$ with $\theta > \frac{\alpha}{2}$. Then there exist $\lambda_0 > 0, C > 0$ such that for $0 < \lambda < \lambda_0$ and for any $\xi_1, \dots, \xi_m \in \partial\Omega$ satisfying (3.2), problem (4.7) admits a unique solution $\phi_1, c_0, c_1, \dots, c_m$ such that*

$$\|\phi_1\|_{L^\infty(\Omega_\lambda)} \leq C\lambda^\alpha. \quad (4.8)$$

Furthermore, the function $(\tau, \xi') \rightarrow \phi_1(\tau, \xi') \in C(\bar{\Omega}_\lambda)$ is C^1 and

$$\|D_{\xi'}\phi_1\|_{L^\infty(\Omega_\lambda)} \leq C\lambda^\alpha, \quad \|D_\tau\phi_1\|_{L^\infty(\Omega_\lambda)} \leq C\lambda^{\theta_1}, \quad \theta_1 < \theta. \quad (4.9)$$

Proof. First we observe that

$$W_1(y) = W(y) + 2\lambda^2 \sinh(V)\tau Z + \tau^2\lambda^2 \cosh(V + \bar{\tau}Z)Z^2,$$

where $|\bar{\tau}| \leq |\tau|$. The equation for ϕ_1 can be written as

$$\begin{cases} \Delta\phi_1 &= 0 & \text{in } \Omega_\lambda, \\ \frac{\partial\phi_1}{\partial\nu} - W\phi_1 &= \tau B\phi_1 + R_1 + N_1(\phi_1) + \sum_{j=1}^m c_j \chi_j Z_{1j} + c_0 \chi Z & \text{on } \partial\Omega_\lambda, \\ \int_{\Omega_\lambda} \chi_j Z_{1j} \phi_1 &= 0 \quad \forall j = 1, \dots, m, & \int_{\Omega_\lambda} \chi Z \phi_1 = 0, \end{cases} \quad (4.10)$$

where $B = 2\lambda^2 \sinh(V)Z + \tau\lambda^2 \cosh(V + \bar{\tau}Z)Z^2$. Remark that from (2.23) and (2.27) we have the estimate $\|B\|_* \leq C$.

Let A be the operator that associates with any $\phi_1 \in L^\infty(\Omega_\lambda)$ the unique solution given by Proposition 2 for $h = \tau B\phi_1 + R_1 + N_1(\phi_1)$ and $f = 0$. In terms of the operator A , equation (4.10) is equivalent to the fixed point problem

$$\phi_1 = A(\phi_1). \quad (4.11)$$

Let us consider the set

$$\mathcal{F} \equiv \{\phi \in C(\bar{\Omega}_\lambda) : \|\phi\|_{L^\infty(\Omega_\lambda)} \leq \lambda^\alpha\}.$$

From Proposition 2 we get

$$\|A(\phi_1)\|_{L^\infty(\Omega_\lambda)} \leq C |\log \lambda| \left[|\tau| \|B\phi_1\|_* + \|N_1(\phi_1)\|_* + \|R_1\|_* \right].$$

Let us estimate $\|R_1\|_*$. We have

$$\begin{aligned} R_1 &= - \left[\frac{\partial V}{\partial v} + \tau \frac{\partial Z}{\partial v} - 2\lambda^2 \sinh(V + \tau Z) \right] \\ &= R(y) - \tau \frac{\partial Z}{\partial v} + 2\lambda^2 \sinh(V + \tau Z) - 2\lambda^2 \sinh V \\ &= R(y) - \tau \left[\frac{\partial Z}{\partial v} - WZ \right] + \lambda^2 \tau^2 \sinh(V + \bar{\tau} Z) Z^2, \end{aligned}$$

where $|\bar{\tau}| \leq |\tau|$. For $|y - \xi'_j| \leq \frac{\delta}{\lambda}$, we have $\frac{\partial Z}{\partial v} - WZ = O\left(\frac{\lambda^a}{1+|y-\xi'_j|}\right)$ where $0 < a < 1$ will be fixed shortly, while for $|y - \xi'_j| > \frac{\delta}{\lambda}$ we have $\frac{\partial Z}{\partial v} - WZ = O(\lambda^2)$, thus $\|\frac{\partial Z}{\partial v} - WZ\|_* \leq C\lambda^{a-\sigma}$. Similarly $\|R\|_* \leq C\lambda^{a-\sigma}$. On the other hand $\|\lambda^2 \sinh(V + \bar{\tau} Z) Z^2\|_* \leq C$, and hence

$$\|R_1\|_* \leq \|R\|_* + |\tau| \lambda^{a-\sigma} + C\tau^2 \leq C(\lambda^{a-\sigma} + \lambda^{2\theta})$$

since $\tau = O(\lambda^\theta)$. We choose $0 < a < 1$ and $\sigma > 0$ small so that $a - \sigma > \alpha$ (σ is the number that appears in the definition of the norms $\|\cdot\|_*$ and $\|\cdot\|_{**}$).

Furthermore, $\|N_1(\phi_1)\|_* \leq C \|\phi_1\|_{L^\infty(\Omega_\lambda)}^2$ as a direct consequence of (4.5) and

$$|\tau| \|B\phi_1\|_* \leq |\tau| \|\phi_1\|_{L^\infty(\Omega_\lambda)} \|B\|_* \leq C\lambda^{\alpha+\theta}.$$

For any $\psi_1, \psi_2, \psi \in \mathcal{F}$, we get the existence of a positive constant C , such that

$$\begin{aligned} \|A(\psi)\|_{L^\infty(\Omega_\lambda)} &\leq C |\log \lambda| \left[\lambda^{\alpha+\theta} + \lambda^{2\alpha} + \lambda^{a-\sigma} + \lambda^{2\theta} \right], \\ \|A(\psi_1) - A(\psi_2)\|_{L^\infty(\Omega_\lambda)} &\leq C |\log \lambda| (\lambda^\theta + \lambda^\alpha) \|\psi_1 - \psi_2\|_{L^\infty(\Omega_\lambda)}. \end{aligned}$$

It follows that for all λ sufficiently small, A is a contraction mapping of \mathcal{F} , and therefore a unique fixed point of A exists in \mathcal{F} .

Let us now discuss the differentiability of ϕ_1 . Since R_1 depends continuously (in the $*$ -norm) on

$$(\tau, \xi') = (\tau, \xi'_1, \dots, \xi'_m),$$

using the fixed point characterization (4.11) we deduce that the map $(\tau, \xi') \mapsto \phi_1$ is also continuous. Then, formally, for $\beta = \xi'_k$ or $\beta = \tau$,

$$\begin{aligned} -\partial_\beta N_1(\phi_1) &= 2\lambda^2 \left[\left(\cosh(V_1 + \phi_1) - \cosh V_1 - \sinh V_1 \phi_1 \right) \partial_\beta V_1 \right. \\ &\quad \left. + \left(\sinh(V_1 + \phi_1) - \cosh V_1 \right) \partial_\beta \phi_1 \right]. \end{aligned}$$

It can be checked that $\|\partial_\beta V_1\|_*$ is uniformly bounded, for both $\beta = \xi'_k$ and $\beta = \tau$, so we conclude that

$$\begin{aligned} \|\partial_\beta N_1(\phi_1)\|_* &\leq C \left[\|\phi_1\|_{L^\infty(\Omega_\lambda)} + \|\partial_\beta \phi_1\|_{L^\infty(\Omega_\lambda)} \right] \|\phi_1\|_{L^\infty(\Omega_\lambda)} \\ &\leq C \left[\lambda^\alpha + \|\partial_\beta \phi_1\|_{L^\infty(\Omega_\lambda)} \right] \lambda^\alpha. \end{aligned}$$

Using the notation $T(h)$ for the operator that $h \in L^\infty(\Omega_\lambda)$ associates the solution of the linear problem (3.32) with $f = 0$, we may write for $\beta = \xi'_k$,

$$\partial_{\xi'_k} \phi_1 = (\partial_{\xi'_k} T) \left(\tau B \phi_1 + N_1(\phi_1) + R_1 \right) + T \left(\partial_{\xi'_k} \left[\tau B \phi_1 + N_1(\phi_1) + R_1 \right] \right),$$

while for $\beta = \tau$,

$$\partial_\tau \phi_1 = T(B \phi_1 + \tau \partial_\tau(B \phi_1) + \partial_\tau N_1(\phi_1) + \partial_\tau R_1).$$

Thus, from Proposition 2 and (3.47) we deduce for $\beta = \xi'_k$

$$\begin{aligned} \|\partial_{\xi'_k} \phi_1\|_{L^\infty(\Omega_\lambda)} &\leq C |\log \lambda|^2 \|(N_1(\phi_1) + R_1)\|_* + |\log \lambda| \|\partial_{\xi'_k} N_1(\phi_1)\|_* \\ &\quad + \|\partial_{\xi'_k} R_1\|_* \\ &\leq C \lambda^{a-\sigma} |\log \lambda|^2 \leq C \lambda^\alpha, \end{aligned}$$

since it can be seen that $\|\partial_{\xi'_k} R_1\|_* \leq C \lambda^\alpha$. For $\beta = \tau$ we get

$$\|\partial_\tau \phi_1\|_{L^\infty(\Omega_\lambda)} \leq C |\log \lambda| (\|\phi_1\|_{L^\infty(\Omega_\lambda)} + \lambda^{1-\sigma} + \lambda^\theta) \leq C \lambda^{\theta_1}, \quad \theta_1 < \theta$$

since

$$\|\partial_\tau R_1\|_* \leq C \left(\frac{\partial Z}{\partial v} - WZ \right)_* + \tau \|\lambda^2 \sinh(V + \bar{\tau}Z) Z^2\|_* \leq C \lambda^\theta.$$

The above computation can be made rigorous by using the implicit function theorem and the fixed point representation (4.11) which guarantees C^1 regularity in τ and ξ' . \square

Remark. It is possible to verify that given $\tau_1, \tau_2 = O(\lambda^\theta)$ with $\theta > \frac{\alpha}{2}$, the unique solutions ϕ_1, ϕ_2 of Lemma 8 satisfy

$$\|\phi_1 - \phi_2\|_{L^\infty(\Omega_\lambda)} \leq C \lambda^\theta |\tau_1 - \tau_2|. \quad (4.12)$$

This follows from the fixed point characterization (4.11) of these solutions. Indeed, let $A(\tau, \phi)$ be the nonlinear operator introduced in this lemma, i.e. the one that $\phi \in L^\infty(\Omega_\lambda)$ associates the unique solution given by Proposition 2 for $h = \tau B \phi + R_1(\tau) + N_1(\phi_1)$ and $f = 0$ to. Using Proposition 2 we see that

$$\|A(\tau_1, \phi) - A(\tau_2, \phi)\|_{L^\infty(\Omega_\lambda)} \leq C \lambda^\theta |\tau_1 - \tau_2|.$$

Lemma 9. *Let $m > 0$, $d > 0$. For any $0 < \alpha < 1$, there exist $\lambda_0 > 0$, $C > 0$ such that for $0 < \lambda < \lambda_0$, any $\xi_1, \dots, \xi_m \in \partial\Omega$ satisfying (3.2), there exists a unique τ with $|\tau| < C\lambda^{\alpha-b/2}$, such that problem (4.7) admits a unique solution ϕ , c_0, c_1, \dots, c_m with $c_0 = 0$, and such that*

$$\|\phi\|_{L^\infty(\Omega_\lambda)} \leq C\lambda^\alpha. \quad (4.13)$$

Furthermore, the function $\xi' \rightarrow \phi(\xi')$ is C^1 and

$$\|D_{\xi'}\phi\|_{L^\infty(\Omega_\lambda)} \leq C\lambda^\alpha.$$

Proof. Given ξ_1, \dots, ξ_m in $\partial\Omega$ such that $|\xi_i - \xi_j| > d$ and $\tau = O(\lambda^\theta)$ with $\frac{\alpha}{2} < \theta < \alpha$, let $\phi_1, c_0, c_1, \dots, c_m$ be solutions to (4.7). Multiplying (4.7) against Z and integrating by parts, we get

$$\begin{aligned} c_0 \int_{\partial\Omega_\lambda} \chi Z^2 &= - \int_{\partial\Omega_\lambda} \phi_1 \left(W_1 Z - \frac{\partial Z}{\partial v} \right) - \int_{\partial\Omega_\lambda} R_1 Z - \int_{\partial\Omega_\lambda} N_1(\phi_1) Z \\ &\quad - \int_{\Omega_\lambda} \phi_1 \Delta Z - \sum_{j=1}^m c_j \int_{\partial\Omega_\lambda} \chi_j \chi Z_{1j} Z. \end{aligned} \quad (4.14)$$

Now we have:

$$\begin{aligned} \left| \int_{\Omega_\lambda} \phi_1 \Delta Z \right| &\leq C \|\phi_1\|_{L^\infty(\Omega_\lambda)} \lambda^b \leq C\lambda^{\alpha+b}; \\ \left| \int_{\partial\Omega_\lambda} N_1(\phi_1) Z \right| &\leq C \|\phi_1\|_{L^\infty(\Omega_\lambda)}^2 \leq C\lambda^{2\alpha}; \\ \left| \int_{\partial\Omega_\lambda} \phi_1 \left(W_1 Z - \frac{\partial Z}{\partial v} \right) \right| &\leq C\lambda^\alpha \|\phi_1\|_{L^\infty(\Omega_\lambda)} \leq C\lambda^{2\alpha} \\ \left| \int_{\partial\Omega_\lambda} \chi_j \chi Z_{1j} Z \right| &\leq C\lambda \end{aligned}$$

$$\int_{\partial\Omega_\lambda} R_1 Z = \int_{\partial\Omega_\lambda} R Z + \tau \int_{\partial\Omega_\lambda} \left[-\frac{\partial Z}{\partial v} + W Z \right] Z + \tau^2 \lambda^2 \int_{\partial\Omega_\lambda} \sinh(V + \bar{\tau} Z) Z^3$$

Let us estimate the second integral in the right-hand side. Observe that in the regions $\{Z < 1 - \lambda^b\}$ (which are of size $|y - \xi'_j| < \mu_j \lambda^{-b/2}$) we have by a calculation similar to (3.19)

$$\int_{|y - \xi'_j| < \mu_j \lambda^{-b/2}, y \in \partial\Omega_\lambda} \left[-\frac{\partial Z}{\partial v} + W Z \right] Z = O(\lambda^\alpha).$$

For the rest at distance $\mu_j \lambda^{-b/2} < |y - \xi'_j| < \frac{\delta}{\lambda}$ we have that Z is constant, so for a given $j = 1, \dots, m$

$$\begin{aligned} \int_{\mu_j \lambda^{-b/2} < |y - \xi'_j| < \frac{\delta}{\lambda}, y \in \partial\Omega_\lambda} \left[-\frac{\partial Z}{\partial v} + W Z \right] Z &= \int_{\mu_j \lambda^{-b/2} < |y - \xi'_j| < \frac{\delta}{\lambda}, y \in \partial\Omega_\lambda} W Z^2 \\ &= 4\lambda^{b/2} + o(\lambda^{b/2}). \end{aligned}$$

Hence,

$$\int_{\partial\Omega_\lambda} \left[-\frac{\partial Z}{\partial \nu} + WZ \right] Z = 4m\lambda^{b/2} + o(\lambda^{b/2}).$$

We also have

$$\left| \int_{\partial\Omega_\lambda} RZ \right| \leq C\lambda^\alpha$$

and from the expansion (2.27)

$$\lambda^2 \int_{\partial\Omega_\lambda} \sinh(V + \bar{\tau}Z)Z^3 = O(\lambda^{b/2}).$$

This shows that it is possible to find $\tau = O(\lambda^{\alpha-b/2})$ so that $c_0 = 0$. The uniqueness of τ can be seen also from the previous estimates. Indeed, suppose we have $\tau, \bar{\tau} = O(\lambda^\theta)$ and solutions $\phi, \tilde{\phi}$ such that for the corresponding coefficients we have $c_0 = \tilde{c}_0 = 0$. From equation (4.14) and the estimates that follow we obtain

$$\lambda^{b/2} |\tau - \bar{\tau}| \leq C\lambda^\alpha \|\phi - \tilde{\phi}\|_{L^\infty(\Omega_\lambda)} + C|\tau - \bar{\tau}|(|\tau| + |\bar{\tau}|)\lambda^{b/2}$$

and using (4.12) we deduce $\tau = \bar{\tau}$.

Let us now discuss the differentiability of ϕ with respect to ξ' . We have

$$\phi(\xi') = \phi_1(\tau(\xi'), \xi')$$

where ϕ_1 is the solution to problem (4.7) given by Lemma 8 while $\tau(\xi')$ is the unique positive number so that in problem (4.7) we have $c_0 = 0$.

Hence,

$$D_{\xi'_k} \phi(\xi') = D_\tau \phi_1(\tau(\xi'), \xi') D_{\xi'_k} \tau(\xi') + D_{\xi'_k} \phi_1(\tau(\xi'), \xi').$$

Since from (4.14) with $c_0 = 0$ we can deduce that $|D_{\xi'_k} \tau(\xi')| \leq C\lambda^\theta$, from (4.8) and (4.9) we conclude that

$$\|D_{\xi'_k} \phi(\xi')\|_{L^\infty(\Omega_\lambda)} \leq C\lambda^\alpha. \quad \square$$

5. Variational reduction

In view of Lemmas 8 and 9, given $\xi = (\xi_1, \dots, \xi_m) \in \partial\Omega^m$ satisfying $|\xi_i - \xi_j| \geq d\forall i \neq j$, we define $\phi(\xi)$ and $c_j(\xi)$ to be the unique solution to (4.7) with $c_0 = 0$ satisfying the bounds (4.8) and (4.9).

Let

$$J_\lambda(u) = \frac{1}{2} \int_{\Omega_\lambda} |Du|^2 - 2\lambda^2 \int_{\partial\Omega_\lambda} \cosh u \, dx.$$

Given $\xi = (\xi_1, \dots, \xi_m) \in \partial\Omega^m$, define

$$F_\lambda(\xi) = J_\lambda(V_1(\xi) + \phi(\xi)), \tag{5.1}$$

where $V_1(\xi) = V(\xi) + \tau(\xi)Z(\xi)$ with $\tau(\xi)$ given by Lemma 9.

Lemma 10. *If $\xi = (\xi_1, \dots, \xi_m) \in (\partial\Omega)^m$ satisfying (3.2) is a critical point of F_λ then $v = V_1(\xi) + \phi(\xi)$ is a critical point of J_λ , that is, a solution to (2.15).*

Proof. Let $\xi' = \xi/\lambda$. Therefore,

$$\frac{\partial F_\lambda}{\partial \xi_k} = \frac{1}{\lambda} \frac{\partial I_\lambda(V_1(\xi') + \phi(\xi'))}{\partial \xi'_k} = \frac{1}{\lambda} D I_\lambda(V_1(\xi') + \phi(\xi')) \left[\frac{\partial V_1(\xi')}{\partial \xi'_k} + \frac{\partial \phi(\xi')}{\partial \xi'_k} \right].$$

Since $v' = V_1(\xi') + \phi(\xi')$ solves (4.7) with $c_0 = 0$

$$\frac{\partial F_\lambda}{\partial \xi_k} = \frac{1}{\lambda} \sum_{i=1}^m c_i \int_{\partial\Omega_\lambda} \chi_i Z_{1i} \left[\frac{\partial V_1(\xi')}{\partial \xi'_k} + \frac{\partial \phi(\xi')}{\partial \xi'_k} \right].$$

Let us assume that $D F_\lambda(\xi) = 0$. From the previous equation we conclude that

$$\sum_{i=1}^m c_i \int_{\partial\Omega_\lambda} \chi_i Z_{1i} \left[\frac{\partial V_1(\xi')}{\partial \xi'_k} + \frac{\partial \phi(\xi')}{\partial \xi'_k} \right] = 0 \quad \forall k = 1, \dots, m.$$

Since $\left\| \frac{\partial \phi(\xi')}{\partial \xi'_k} \right\|_{L^\infty(\Omega_\lambda)} \leq C\lambda^\alpha$ and $\frac{\partial V_1(\xi')}{\partial \xi'_k} = \pm Z_{1k} + o(1)$ where $o(1)$ is in the L^∞ norm as a direct consequence of (4.6), it follows that

$$\sum_{i=1}^m c_i \int_{\partial\Omega_\lambda} \chi_i Z_{1i} (\pm Z_{1k} + o(1)) = 0 \quad \forall k = 1, \dots, m,$$

which is a strictly diagonal dominant system. This implies that $c_i = 0$ $\forall i = 1, \dots, m$. \square

In order to solve for critical points of the function F_λ , a key step is its expected closeness to the function $J_\lambda(V_1(\xi))$, which we will analyze in the next section.

Lemma 11. *Assume $\alpha \in (\frac{1}{2}, 1)$. The following expansion holds*

$$F_\lambda(\xi) = J_\lambda(V) + \theta_\lambda(\xi),$$

where

$$|\theta_\lambda| \rightarrow 0,$$

uniformly on points satisfying the constraints (3.2).

Proof. We write

$$\begin{aligned} J_\lambda(V_1 + \phi) - J_\lambda(V) &= [J_\lambda(V_1 + \phi) - J_\lambda(V_1)] + [J_\lambda(V_1) - J_\lambda(V)] \\ &= A + B. \end{aligned}$$

Let us estimate A first. Taking into account that $DJ_\lambda(V_1 + \phi)[\phi] = 0$, a Taylor expansion and an integration by parts gives

$$\begin{aligned} A &= \int_0^1 D^2 J_\lambda(V_1 + t\phi)[\phi]^2 (1-t) dt \\ &= \int_0^1 \left(2\lambda^2 \int_{\partial\Omega_\lambda} [\cosh(V_1 + t\phi) - \cosh(V_1 + \phi)] \phi^2 \right. \\ &\quad \left. + \int_{\partial\Omega_\lambda} [N_1(\phi) + R_1] \phi \right) (1-t) dt, \end{aligned} \quad (5.2)$$

so we get

$$J_\lambda(V_1 + \phi) - J_\lambda(V_1) = O(\lambda^{2\alpha})$$

taking into account that $\|\phi\|_{L^\infty(\Omega_\lambda)} \leq C\lambda^\alpha$.

On the other hand,

$$B = \tau J'_\lambda(V + \bar{\tau}Z)[Z]$$

for $|\bar{\tau}| \leq |\tau|$, and since Z is almost an element of the kernel of $J'_\lambda(V)$, we get

$$J_\lambda(V_1) - J_\lambda(V) = o(1)\tau \rightarrow 0.$$

Hence, $|\theta_\lambda(\xi)| = o(1)$ uniformly on points satisfying (3.2).

The continuity in ξ of all these expressions is inherited from that of ϕ and its derivatives in ξ in the L^∞ norm. \square

6. Energy computations and proof of the theorem

In this section we compute the expansion of the energy functional J_λ evaluated at V and we give the proof of Theorem 1.

We have

Lemma 12. *Let $m > 0$, $d > 0$. Let μ_j be given by (2.21) and let V be the function defined in (2.16). Then for any $0 < \alpha < 1$ the following expansion holds true*

$$\begin{aligned} J_\lambda(V) &= 2m\pi \log \frac{1}{\lambda} + m(\beta_0 - 2\pi + 2\pi \log 2) \\ &\quad - \pi \varphi_m(\xi) + O(\lambda^\alpha) \end{aligned} \quad (6.1)$$

uniformly on points $\xi = (\xi_1, \dots, \xi_m) \in (\partial\Omega)^m$ such that $|\xi_i - \xi_j| > d$ for all $i \neq j$. In the previous formula, $\varphi_m(\xi)$ is the function introduced in (1.5), namely

$$\varphi_m(\xi) = \varphi_m(\xi_1, \dots, \xi_m) = \left[\sum_{l=1}^m H(\xi_l, \xi_l) + \sum_{j \neq l} (-1)^{l+j} G(\xi_j, \xi_l) \right]$$

while β_0 is the constant defined by

$$\beta_0 = \int_R \frac{1}{1+x^2} \log \frac{1}{1+x^2}.$$

Proof. Since $V(y) = \sum_{j=1}^m (-1)^{j-1} (u_j^\lambda(\lambda y) + H_j^\lambda(\lambda y))$ satisfies $\Delta V = 0$ in Ω_λ , we write

$$J_\lambda(V) = \frac{1}{2} \int_{\partial\Omega_\lambda} V \frac{\partial V}{\partial \nu} - 2\lambda^2 \int_{\partial\Omega_\lambda} \cosh(V).$$

We compute the second term first:

$$2\lambda^2 \int_{\partial\Omega_\lambda} \cosh(V) = \lambda^2 \sum_{l=1}^m \int_{\partial\Omega_\lambda \cap B_{\delta/\lambda}(\xi'_l)} (e^V + e^{-V}) + O(\lambda^\alpha)$$

Suppose l is odd first. Then, recalling the notation introduced in Section 2, $v_l(y) = u_j^\lambda(y) + 2 \log \lambda$, we get

$$\begin{aligned} \lambda^2 \int_{\partial\Omega_\lambda \cap B_{\delta/\lambda}(\xi'_l)} (e^V + e^{-V}) &= \lambda^2 \int_{\partial\Omega_\lambda \cap B_{\delta/\lambda}(\xi'_l)} e^V + O(\lambda^\alpha) \\ &= \int_{\partial\Omega_\lambda \cap B_{\delta/\lambda}(\xi'_l)} e^{v_l} e^{(-1)^{l-1} H_l^\lambda + \sum_{j \neq l} (-1)^{j-1} (u_j^\lambda + H_j^\lambda)} \\ &\quad + O(\lambda^\alpha) \\ &= 2\pi + O(\lambda^\alpha). \end{aligned}$$

Thus,

$$\lambda^2 \int_{\partial\Omega_\lambda \cap B_{\delta/\lambda}(\xi'_l)} (e^V + e^{-V}) = 2\pi + O(\lambda^\alpha). \quad (6.2)$$

Similarly for l even, we also have (6.2), so we obtain

$$2\lambda^2 \int_{\partial\Omega_\lambda} \cosh(V) = 2m\pi + O(\lambda^\alpha). \quad (6.3)$$

It remains to compute $\int_{\partial\Omega_\lambda} V \frac{\partial V}{\partial \nu}$:

$$\begin{aligned} \int_{\partial\Omega_\lambda} V \frac{\partial V}{\partial \nu} &= \int_{\partial\Omega_\lambda} \left(\sum_{j=1}^m (-1)^{j-1} (u_j^\lambda + H_j^\lambda) \right) \\ &\quad \times \left(\sum_{i=1}^m (-1)^{i-1} e^{v_i} - \frac{1}{|\partial\Omega_\lambda|} \sum_{i=1}^m (-1)^{i-1} \int_{\partial\Omega_\lambda} e^{v_i} \right) \\ &= \int_{\partial\Omega_\lambda} \left(\sum_{j=1}^m (-1)^{j-1} (u_j^\lambda + H_j^\lambda) \right) \left(\sum_{i=1}^m (-1)^{i-1} e^{v_i} \right) \\ &\quad - \frac{1}{|\partial\Omega_\lambda|} \left(\sum_{i=1}^m (-1)^{i-1} \int_{\partial\Omega_\lambda} e^{v_i} \right) \int_{\partial\Omega_\lambda} \left(\sum_{j=1}^m (-1)^{j-1} (u_j^\lambda + H_j^\lambda) \right) \\ &= \sum_{i,j=1}^m (-1)^{i+j} \int_{\partial\Omega_\lambda} e^{v_i} (u_j^\lambda + H_j^\lambda) + O(\lambda^\alpha), \end{aligned}$$

since by (2.26)

$$\sum_{j=1}^m (-1)^{j-1} \int_{\partial\Omega_\lambda} e^{v_j} = O(\lambda).$$

For $j \neq i$, we have

$$\int_{\partial\Omega_\lambda} e^{v_i} (u_j^\lambda + H_j^\lambda) = 2\pi G(\xi_j, \xi_i) + O(\lambda^\alpha). \quad (6.4)$$

For $j = i$, we have using (2.21),

$$\begin{aligned} & \int_{\partial\Omega_\lambda} e^{v_i} (u_i^\lambda + H_i^\lambda) \\ &= \int_{\partial\Omega_\lambda} \frac{2\mu_j}{|y - \mu_j v'_j|^2} \left(\left(\log \frac{1}{\lambda^2} \right) + \log \frac{2\mu_j}{|y - \mu_j v'_j|^2} + H(\xi_j, \xi_j + \lambda y) - \log(2\mu_j) \right) \\ & \quad + O(\lambda^\alpha) \\ &= 2\pi \log \frac{1}{\lambda^2} + 2\pi (H(\xi_j, \xi_j) - \log 2\mu_j) + 2\beta_0 + O(\lambda^\alpha). \end{aligned}$$

Thus,

$$\begin{aligned} \int_{\partial\Omega_\lambda} e^{v_i} (u_i^\lambda + H_i^\lambda) &= 2\pi \log \frac{1}{\lambda^2} + 2\beta_0 + 4\pi \log 2 \\ & \quad + 2\pi \left(-H(\xi_j, \xi_j) - 2 \sum_{i \neq j} (-1)^{i+j} G(\xi_j, \xi_i) \right) + O(\lambda^\alpha) \end{aligned} \quad (6.5)$$

Combining (6.4) and (6.5), we obtain that

$$\begin{aligned} \int_{\partial\Omega_\lambda} V \frac{\partial V}{\partial v} &= \sum_{i,j=1}^m (-1)^{i+j} \int_{\partial\Omega_\lambda} e^{v_i} (u_j^\lambda + H_j^\lambda) + O(\lambda^\alpha) \\ &= 2m\pi \log \frac{1}{\lambda^2} + 2m\beta_0 + 4m\pi \log 2 \\ & \quad + 2\pi \left(-\sum_{i=1}^m H(\xi_i, \xi_i) - \sum_{j \neq i} (-1)^{i+j} G(\xi_j, \xi_i) \right) + O(\lambda^\alpha) \end{aligned} \quad (6.6)$$

Summing up equations (6.3) and (6.6), we finally arrive at

$$\begin{aligned} J_\lambda(V) &= 2m\pi \log \frac{1}{\lambda} + m(\beta_0 - 2\pi + 2\pi \log 2) \\ & \quad - \pi \left[\sum_{l=1}^m H(\xi_l, \xi_l) + \sum_{j \neq l} (-1)^{l+j} G(\xi_j, \xi_l) \right] + O(\lambda^\alpha). \quad \square \end{aligned}$$

We now have all ingredients to give the proof of Theorem 1.

Proof of Theorem 1. Define for $\xi = (\xi_1, \dots, \xi_m) \in (\partial\Omega)^m$ with $|\xi_i - \xi_j| \geq d$, the function

$$v(y) = V_1(\xi)(y) + \phi(\xi)(y) \quad y \in \Omega_\lambda,$$

where $V_1(\xi)$ is given by (4.1) and $\phi(\xi)$ is the unique solution to problem (4.7) with $c_0 = 0$, whose existence and properties are established in Lemma 9. Then, according to Lemma 8, v is solution to (2.15) provided that ξ is a critical point of the function $F_\lambda(\xi)$ defined in (5.1), or equivalently, ξ is a critical point of

$$\tilde{F}_\lambda(\xi) = \frac{1}{\pi} \left(2m\pi \log \frac{1}{\lambda} + m(\beta_0 - 2\pi + 2\pi \log 2) - F_\lambda(\xi) \right).$$

Let $m = 2k$ and $\tilde{\Omega}_m$ be the set of points $\xi = (\xi_1, \dots, \xi_m) \in (\partial\Omega)^m$ ordered clockwise along a given connected component of $\partial\Omega$, such that $|\xi_i - \xi_j| \geq d$ for all $i \neq j$ for some $d > 0$ sufficiently small, so that all the previous results hold true. Namely, if we denote by $p : [0, 2\pi] \rightarrow \partial\Omega$ a continuous parametrization of this connected component of $\partial\Omega$, we can then write

$$\tilde{\Omega}_m = \{ \xi = (p(\theta_1), \dots, p(\theta_m)) \in (\partial\Omega)^m : |p(\theta_i) - p(\theta_j)| \geq d \text{ if } i \neq j \}.$$

It is not restrictive to assume that $0 \in \partial\Omega$. Lemmas 11 and 12 guarantee that for $\xi \in \tilde{\Omega}_m$,

$$\tilde{F}_\lambda(\xi) = \varphi_m(\xi) + \lambda^\alpha \Theta_\lambda(\xi), \quad (6.7)$$

where Θ_λ is uniformly bounded in the considered region as $\lambda \rightarrow 0$. We will show that \tilde{F}_λ has at least two distinct critical points in this region—a fact that when used will prove our result. The function φ_m is C^1 , bounded from above in $\tilde{\Omega}_m$, and if two consecutive points get closer it becomes unbounded from below, implying that

$$\varphi_m(\xi_1, \dots, \xi_m) \rightarrow -\infty \text{ as } |\xi_i - \xi_j| \rightarrow 0 \text{ for some } i \neq j.$$

Hence, since d is arbitrarily small, φ_m has an absolute maximum M_0 in $\tilde{\Omega}_m$, and so does \tilde{F}_λ whenever λ is sufficiently small. Let us call M_λ this value, so that $M_\lambda = M_0 + o(1)$ as $\lambda \rightarrow 0$. On the other hand, Ljusternik–Schnirelmann theory is applicable in our setting, so we can estimate the number of critical points of φ_m in $\tilde{\Omega}_m$ by $\text{cat}(\tilde{\Omega}_m)$, the Ljusternik–Schnirelmann category of $\tilde{\Omega}_m$ relative to $\tilde{\Omega}_m$. We claim that $\text{cat}(\tilde{\Omega}_m) > 1$. Indeed, by contradiction, assume that $\text{cat}(\tilde{\Omega}_m) = 1$. This means that $\tilde{\Omega}_m$ is contractible in itself, namely there exist a point $\xi^0 \in \tilde{\Omega}_m$ and a continuous function $\Gamma : [0, 1] \times \tilde{\Omega}_m \rightarrow \tilde{\Omega}_m$ such that for all $\xi \in \tilde{\Omega}_m$,

$$\Gamma(0, \xi) = \xi, \quad \Gamma(1, \xi) = \xi^0.$$

Let $f : S^1 \rightarrow \tilde{\Omega}_m$ be the continuous function defined by

$$f(x) = \left(p(\theta), p\left(\theta + 2\pi \frac{1}{m}\right), \dots, p\left(\theta + 2\pi \frac{m-1}{m}\right) \right), \\ x = e^{i\theta}, \theta \in [0, 2\pi].$$

Let $\eta : [0, 1] \times S^1 \rightarrow S^1$ be the well-defined continuous map given by

$$\eta(t, x) = \frac{\pi_1 \circ \Gamma(t, f(x))}{\|\pi_1 \circ \Gamma(t, f(x))\|},$$

where π_1 denotes the projection on the first component. The function η is a contraction of S^1 to a point and this gives a contradiction. Thus we conclude that

$$c_0 = \sup_{C \in \Xi} \inf_{\tilde{\xi} \in C} \varphi_m(\tilde{\xi}), \quad (6.8)$$

where

$$\Xi = \{C \subset \tilde{\Omega}_m : C \text{ closed and } \text{cat}(C) \geq 2\}$$

is a finite number, and a critical level for φ_m . Call c_λ the number (6.8) with φ_m replaced by \tilde{F}_λ , so that $c_\lambda = c_0 + o(1)$. If $c_\lambda \neq M_\lambda$, we conclude that there are at least two distinct critical points for \tilde{F}_λ (distinct up to cyclic permutations) in $\tilde{\Omega}_m$. If $c_\lambda = M_\lambda$, we get that there must be a set C , with $\text{cat}(C) \geq 2$, where the function \tilde{F}_λ reaches its absolute maximum. In this case we conclude that there are infinitely many critical points for \tilde{F}_λ in $\tilde{\Omega}_m$. Since cyclic permutations are only finite in number, the result is thus proven. \square

Acknowledgments. This work has been partly supported by grants Fondecyt 1030840, 1040936, 1020815 and FONDAP, Chile, Progetto Nazionale ex 40% *Metodi variazionali e topologici nello studio di fenomeni non lineari*, Italy, and an Earmarked grant CUHK 402304 from RGC, Hong Kong.

References

1. BARAKET, S., PACARD, F.: Construction of singular limits for a semilinear elliptic equation in dimension 2. *Calc. Var. Partial Differential Equations* **6**, 1–38 (1998)
2. BREZIS, H., MERLE, F.: Uniform estimates and blow-up behavior for solutions of $-\Delta u = V(x)e^u$ in two dimensions. *Comm. Partial Differential Equations* **16**, 1223–1253 (1991)
3. BRYAN, K., VOGELIUS, M.: Singular solutions to a nonlinear elliptic boundary value problem originating from corrosion modeling. *Quart. Appl. Math.* **60**, 675–694 (2002)
4. DÁVILA, J., DEL PINO, M., MUSSO, M.: Concentrating solutions in a two-dimensional elliptic problem with exponential Neumann data. To appear in *J. Funct. Anal.* (2005)
5. DECONINCK, J.: Current distributions and electrode shape changes in electrochemical systems. *Lecture Notes in Engineering*, vol. 75. Springer-Verlag, Berlin, 1992
6. DEL PINO, M., KOWALCZYK, M., MUSSO, M.: Singular limits in Liouville-type equations. *Calc. Var. Partial Differential Equations* **24**, 47–81 (2005)
7. ESPOSITO, P., GROSSI, M., PISTOIA, A.: On the existence of blowing-up solutions for a mean field equation. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **22**, 227–257 (2005)
8. KAVIAN, O., VOGELIUS, M.: On the existence and “blow-up” of solutions to a two-dimensional nonlinear boundary-value problem arising in corrosion modelling. *Proc. Roy. Soc. Edinburgh Sect. A* **133**, 119–149 (2003)
9. LI, Y., SHAFRIR, I.: Blow-up analysis for solutions of $-\Delta u = Ve^u$ in dimension two. *Indiana Univ. Math. J.* **43**, 1255–1270 (1994)
10. MEDVILLE, K., VOGELIUS, M.: Blow up behaviour of planar harmonic functions satisfying a certain exponential Neumann boundary condition. *SIAM J. Math. Anal.* **36**, 1772–1806 (2005)

Singular Limits of a Two-Dimensional Boundary Value Problem

11. MEDVILLE, K., VOGELIUS, M.: Existence and blow up of solutions to certain classes of two dimensional nonlinear Neumann problems. To appear in *Ann. Inst. H. Poincaré Anal. Non Linéaire*. (2005)
12. NAGASAKI, K., SUZUKI, Y.: Asymptotic analysis for two-dimensional elliptic eigenvalue problems with exponentially dominated nonlinearities. *Asymptot. Anal.* **3**, 173–188 (1990)
13. VOGELIUS, M., XU, J.M.: A nonlinear elliptic boundary value problem related to corrosion modeling. *Quart. Appl. Math.* **56**, 479–505 (1998)

Departamento de Ingeniería Matemática and CMM,
Universidad de Chile,
Casilla 170 Correo 3,
Santiago, Chile.
e-mail: jdavila@dim.uchile.cl
e-mail: delpino@dim.uchile.cl

and

Dipartimento di Matematica,
Politecnico di Torino,
Corso Duca degli Abruzzi 24,
10129 Torino,
Italy

and

Departamento de Matemática,
Pontificia Universidad Católica de Chile,
Vicuña Mackenna 4860,
Macul, Chile.
e-mail: mmusso@mat.puc.cl

and

Department of Mathematics,
The Chinese University of Hong-Kong,
Shatin, Hong Kong.
e-mail: wei@math.cuhk.edu.hk