

1 OBES SURG  
2 DOI 10.1007/s11695-008-9638-0

4 **Changes in Bone Mineral Density, Body Composition**  
5 **and Adiponectin Levels in Morbid Obese Patients**  
6 **after Bariatric Surgery**

7 **Fernando Carrasco · Manuel Ruz · Pamela Rojas ·**  
8 **Attila Csendes · Annabella Rebolledo · Juana Codoceo ·**  
9 **Jorge Inostroza · Karen Basfi-fer · Karin Papapietro ·**  
10 **Jorge Rojas · Fernando Pizarro · Manuel Olivares**

11 Received: 14 May 2008 / Accepted: 7 July 2008  
12 © Springer Science + Business Media, LLC 2008

15 **Abstract**

16 *Introduction* Gastric bypass surgery (GBP) is increasingly  
17 used as a treatment option in morbid obesity. Little is  
18 known about the effects of this surgery on bone mineral  
19 density (BMD) and the underlying mechanisms. To  
20 evaluate changes on BMD after GBP and its relation with  
21 changes in body composition and serum adiponectin, a  
22 longitudinal study in morbid obese subjects was conducted.  
23 *Methods* Forty-two women (BMI 45.0±4.3 kg/m<sup>2</sup>; 37.7±  
24 9.6 years) were studied before surgery and 6 and 12 months  
25 after GBP. Percentage of body fat (%BF), fat-free mass  
26 (FFM), and BMD were measured by dual-energy X-ray  
27 absorptiometry and serum adiponectin levels by RIA.  
28 *Results* Twelve months after, GBP weight was decreased  
29 by 34.4±6.5% and excess weight loss was 68.2±12.8%.  
30 Significant reduction ( $p<0.001$ ) in total BMD (−3.0±  
31 2.1%), spine BMD (−7.4±6.8%) and hip BMD (−10.5±  
32 5.6%) were observed. Adiponectin concentration increased  
33 from 11.4±0.7 mg/L before surgery to 15.7±0.7 and 19.8±

1.0 at the sixth and twelfth month after GBP, respectively 34  
( $p<0.001$ ). Thirty-seven percent of the variation in total 35  
BMD could be explained by baseline weight, initial BMD, 36  
BF reduction, and adiponectin at the twelfth month ( $r^2=$  37  
0.373;  $p<0.001$ ). Adiponectin at the twelfth month had a 38  
significant and positive correlation with the reduction of 39  
BMD, unrelated to baseline and variation in body compo- 40  
sition parameters (adjusted correlation coefficient:  $r=0.36$ ). 41  
*Conclusion* GBP induces a significant BMD loss related 42  
with changes in body composition, although some meta- 43  
bolic mediators, such as adiponectin increase, may have an 44  
independent action on BMD which deserves further study. 45

**Keywords** Bone mineral density · Gastric bypass · 46  
Adiponectin · Body composition 47

**Introduction** 48

Actually, overweight and obesity represents one of the 49  
major health hazards in the worldwide affecting nearly 1.1 50  
billion subjects [1]. According to the data in the first 51  
national health survey in 2003, the prevalence of over- 52  
weight, obesity, and morbid obesity in Chilean adults were 53  
38%, 22%, and 1.3%, respectively [2]. Obesity is associ- 54  
ated to increased mortality [3] and to a higher risk of type 2 55  
diabetes, hyperlipidemia, sleep apnea, gallbladder disease, 56  
coronary disease, hypertension, musculoskeletal disorders, 57  
several types of cancer and psychosocial disturbances [4, 58  
5]. Besides the greater severity of comorbidities, severe and 59  
morbid obesity (BMI between 35 and 39.9 kg/m<sup>2</sup> and 60  
≥40 kg/m<sup>2</sup>, respectively) sharply reduce life expectancy, 61  
especially in young adults [6]. 62

F. Carrasco (✉) · M. Ruz · P. Rojas · A. Rebolledo · J. Codoceo ·  
J. Inostroza · K. Basfi-fer  
Department of Nutrition, Faculty of Medicine,  
University of Chile,  
Independencia 1027,  
Postal Code 838-0453 Santiago, Chile  
e-mail: fcarrasc@med.uchile.cl

A. Csendes · K. Papapietro · J. Rojas  
Department of Surgery, Clinical Hospital, University of Chile,  
Santiago, Chile

F. Pizarro · M. Olivares  
Institute of Nutrition and Food Technology, University of Chile,  
Santiago, Chile

In these patients, bariatric surgery is considered the most effective method to induce significant weight loss (~70% of excess weight) which is maintained in the long term. This observation is confirmed in two recent meta-analysis showing the efficacy of different types of bariatric surgery, in terms of both weight loss and comorbidities improvement [7, 8]. Therefore, Roux-en-Y gastric bypass (GBP), the most recommended technique currently used, has become the “gold-standard” technique [9–11]. Despite the beneficial results of GBP, some of the warning issues are mortality, and early and late complications. Among the most frequently observed nutritional complications are micronutrient deficiencies such as vitamin B12, folate, zinc, iron, vitamin D and calcium, due to a combination of poor food intake, maldigestion and/or malabsorption [12]. Iron deficiency anemia is the most frequent complication after GBP (~50%), especially in premenopausal women. Emerging concern is being raised related to metabolic bone disease and osteoporosis. In subjects who have undergone bariatric surgery, previous studies have shown increased risks of metabolic bone disease, reduction in bone mineral density (BMD), bone mineral content, and fracture [13, 14]. A number of factors seems to be involved in the reduction of BMD after GBP, such as: elevated levels of serum parathyroid hormone secondary to inadequate calcium intake, intestinal Ca and vitamin D malabsorption [15], reduced biomechanical stress along the weight loss [16, 17], reduced levels of estrogens in women [16], reduction in plasma levels of leptin [18, 19] and ghrelin [20], and increased serum concentrations of adiponectin [21]. The aim of this prospective study was to assess the magnitude of the effect of changes in food intake, micronutrients supplementation, body composition, and serum levels of adiponectin on the reduction in bone mass after gastric bypass. Our hypothesis is that the main contributing factor will be the amount of calcium and vitamin D consumed by supplements.

## Materials and Methods

### Subjects

Forty-two adult women (age  $37.7 \pm 9.6$  years; age range 19–55 years), with BMI  $\geq 40$  kg/m<sup>2</sup> or BMI  $\geq 35$  kg/m<sup>2</sup> with comorbidities such as type-2 diabetes, hypertension, dyslipidemia, sleep apnea, osteoarthritis, among other conditions, who underwent RYGBP in the Department of Surgery of the University of Chile Clinical Hospital were prospectively studied. The mean BMI was  $45.0 \pm 4.3$  kg/m<sup>2</sup>: five patients had a BMI lower than 40 (range 39.3–39.8 kg/m<sup>2</sup>), 31 patients had a BMI between 40 and 49.9, and six patients presented a BMI  $\geq 50$  kg/m<sup>2</sup> (range 50.1–59.2). All patients

who agreed to enter the study signed an authorized consent form. The study was approved by the Ethics Committee for Human Investigation of the University Clinical Hospital. The surgical technique consisted of a 95% distal gastrectomy, with resection of the bypassed stomach, leaving a small gastric pouch of 15–20 ml. [22]. Then, an end-to-side gastrojejunostomy with circular stapler No. 25 was performed. The length of the Roux-loop was 125–150 cm. All patients underwent complete preoperative evaluation.

Before and 6 and 12 months after surgery, the following evaluations were carried out in all patients: anthropometry, food survey, resting energy expenditure (REE), body composition analysis, bone mineral density measurement, and serum adiponectin concentration. In all patients, calcium (640–1,000 mg) and vitamin D (400–800 U) were prescribed as an oral supplementation for the entire period of observation.

### Anthropometric, Food Intake, and Supplements Evaluations

Weight was measured to the nearest 0.1 kg on a digital scale (Seca®, Vogel & Halke GMBH & Co, Germany), and height (m) was measured to the nearest 0.1 cm with a scale-mounted stadiometer. BMI (kg/m<sup>2</sup>) was calculated. Waist (WC) and hip (HC) circumferences (cm) were registered according to the procedures recommended by the Anthropometric Standardization Reference Manual [23]. At baseline, and at the sixth and twelfth month of the postoperative period, patients were interviewed by a dietitian and were enquired to fill a 3-day record of food intake. The recorded data were analyzed using a computer program (Food Processor II®, ESHA Research, Salem, OR, USA) to calculate energy, protein, and calcium intake. A strict record of all types of vitamin and mineral supplements were kept throughout the study.

### Body Composition

Fat-free mass (FFM, kg), Body fat (BF, kg), bone mineral density (BMD, g/cm<sup>2</sup>), and bone mineral content (g) were measured by dual-energy X-ray absorptiometry (DXA) using a Lunar DPX-L densitometer (Lunar, Madison, WI; version 1.30 software) [24, 25]. BMD and BMC were examined in total body, lumbar spine, and total hip. BF was analyzed in the total body, trunk, and in the ROI (Region of Interest). ROI is a rectangular area defined by horizontal lines placed inferiorly on the top of the iliac crests and superiorly at three pixels above the previous line [26].

### Adiponectin Analysis

In serum obtained from morning-fasted blood samples, adiponectin was analyzed by radioimmunoassay using a kit from Linco Research Inc. (St. Charles, MO, USA).

t1.1 **Table 1** Baseline demographics, anthropometrics, body composition and energy metabolism characteristics of patients before bariatric surgery (*n*=42)

t1.2	Mean±SD	Range
t1.3 Age (years)	37.7±9.6	19–55
t1.4 Body weight (kg)	113.9±14.6	88.4–145.5
t1.5 Excess of weight (kg)	58.0±12.0	40.4–83.4
t1.6 BMI (kg/m <sup>2</sup> )	45.0±4.3	39.3–59.2
t1.7 Waist (cm)	119.4±10.2	99–141
t1.8 Hip (cm)	134.5±11.8	107–162
t1.9 Body fat (%)	47.8±5.1	34.5–57.1
t1.10 Trunk body fat (%)	45.4±4.8	33.6–56.9
t1.11 ROI body fat (%)	49.2±4.3	41.1–58.6
t1.12 Total BMD (g/cm <sup>2</sup> )	1.23±0.05	1.14–1.34
t1.13 Total <i>T</i> -score (SD)	1.21±0.74	−0.30–2.73
t1.14 Spine BMD (g/cm <sup>2</sup> )	1.49±0.11	1.25–1.69
t1.15 Spine <i>T</i> -score (SD)	2.44±0.77	0.80–3.80
t1.16 Pelvis BMD (g/cm <sup>2</sup> )	1.28±0.09	1.08–1.44
t1.17 Pelvis <i>T</i> -score (SD)	1.58±0.98	−0.40–3.30

t1.18 *BMI* Body mass index, *ROI* region of interest located in abdominal area 1 cm above iliac crest, *BMD* bone mineral density evaluated by dual X-ray absorptiometry, *T-score* difference between measure and mean value of young adult women in standard deviations

159 Statistical Analysis

160 The design of the study considered obtaining results at  
161 different periods after GBP in the same subjects. Repeated  
162 measures of ANOVA model were used to evaluate the  
163 effects of this surgery on variables such as body composi-  
164 tion, BMD, and adiponectin concentration. The Bonferroni  
165 test was applied for multiple comparisons. To evaluate the

association among variables, Spearman’s correlation coef- 166  
ficient analyses were performed [27]. Regression analysis 167  
models were performed to evaluate the association between 168  
the reduction in BMD, as the dependent variable, and 169  
changes in body composition, nutritional intake, vitamin 170  
and mineral supplementation, and twelfth-month adiponec- 171  
tin levels, as independent variables. To adjust for the 172  
potential effects of baseline body composition parameters 173  
and for changes in body weight and total and regional body 174  
fat, partial correlation coefficients between reduction in 175  
BMD and twelfth-month adiponectin levels were comput- 176  
ed. SPSS 10.0® statistical software was used for all the 177  
analysis (SPSS Inc, Chicago IL, USA). Statistical signifi- 178  
cance was considered at *p*<0.05. 179

Results 180

Anthropometrics, body composition and bone mineral 181  
density characteristics of the forty-two patients before GBP 182  
are showed in Table 1. At baseline, no patient presented a 183  
BMD at a risk level of osteoporosis (*T*-score ≤−1 standard 184  
deviation) in the analyzed regions. 185

Changes in body composition, BMD, and serum 186  
adiponectin levels at the sixth and twelfth-month post 187  
surgery are shown in Table 2. Twelve months after GBP 188  
weight loss ranged between 21.8% and 49.6% of initial 189  
weight, BF between 23.0% and 79.0%, and FFM between 190  
5.9 y 29.4%. Excess weight loss 12 months after GBP was 191  
68.2±12.8% (range, 36.8–95.8). All changes in body 192  
composition were statistically significant (*p*<0.001) among 193  
the three points of evaluation (baseline, sixth, and twelfth 194

t2.1 **Table 2** Changes in weight, body composition, and adiponectin

t2.2 Variable	Baseline	Sixth month	Twelfth month	Change (%) <sup>a</sup>
t2.3 Weight (kg)	113.9±14.6*	82.1±11.1*	74.5±9.9*	−34.4±6.5
t2.4 BMI (kg/m <sup>2</sup> )	45.0±4.3*	32.5±3.9*	29.5±3.9*	−34.4±6.5
t2.5 Body fat (kg)	54.9±11.6*	33.9±9.5*	26.3±8.5*	−52.1±12.4
t2.6 Fat-free mass (kg)	58.9±5.8*	48.2±3.8*	48.2±3.6**	−18.0±5.1
t2.7 Waist (cm)	119.4±10.2*	95.6±8.8*	89.8±8.4*	−24.7±5.1
t2.8 Hip (cm)	134.5±11.8*	113.2±10.1*	106.9±9.5*	−20.3±5.4
t2.9 Trunk body fat (kg)	24.2±6.0*	17.2±6.6*	12.8±4.4*	−46.2±16.5
t2.10 ROI body fat (kg)	1.57±0.41*	0.88±0.27*	0.68±0.25*	−55.8±13.9
t2.11 Total BMD (g/cm <sup>2</sup> )	1.23±0.53*	1.21±0.64*	1.19±0.63*	−3.0±2.1
t2.12 Spine BMD (g/cm <sup>2</sup> )	1.49±0.11*	1.42±0.14*	1.38±0.14****	−7.4±6.8
t2.13 Pelvis BMD (g/cm <sup>2</sup> )	1.28±0.09*	1.19±0.09*	1.14±0.09*	−10.5±5.6
t2.14 Adiponectin	11.4±4.3*	15.7±4.8*	19.8±6.6*	+97.0±99.9

t2.15 *BMI* Body mass index, *ROI* region of interest, *BMD* bone mineral density evaluated by dual X-ray absorptiometry

<sup>a</sup> Percent variation between baseline and final value.  
\**p*<0.001, significant difference between all measures  
\*\**p*<0.001 significant difference with basal measures  
\*\*\**p*<0.05, significant difference with previous measure

month), except FFM which showed no change between month 6 and 12 post GBP.

After GBP, significant reductions in total BMD, spine BMD and pelvis BMD were observed ( $p<0.001$ ; Fig. 1). The change in BMD at the twelfth month showed a significant difference among total BMD ( $-3.0\pm2.1\%$ ), spine BMD ( $-7.4\pm6.8\%$ ) and pelvis BMD ( $-10.5\pm5.6\%$ ;  $p<0.001$ ). Considering total and spine BMD no patient fell to a level considered as osteopenia ( $T$ -score  $\leq -1$  and  $>2.5$  standard deviations) or osteoporosis ( $T$ -score  $\leq -2.5$  standard deviations). Osteopenia however, was present in 4 patients according to pelvis BMD measured 12 months after GBP ( $T$ -score range,  $-1$  to  $-1.7$  standard deviations). Total BMD decrease was positively correlated ( $p<0.05$ ) with baseline weight ( $r=0.36$ ), initial FFM ( $r=0.33$ ), percentage of weight loss ( $r=0.33$ ), adiponectin change ( $r=0.34$ ), and absolute adiponectin at the twelfth-month post GBP ( $r=0.35$ ). After multiple regression analysis, 37% of the reduction of BMD could be explained by the following variables: baseline weight, baseline BMD, BF reduction and adiponectin at the twelfth-month post GBP ( $r^2=0.373$ ;  $p<0.001$ ). After adjusting for covariates (baseline weight, initial BF, initial BMD, weight loss, total, and regional BF loss), adiponectin concentrations at the twelfth month had a positive correlation with BMD decrease (adjusted correlation coefficient:  $r=0.35$ ;  $p<0.05$ ).

Serum adiponectin concentrations increased by 48% at sixth month and by 97% at the twelfth-month post GBP ( $p<0.001$ ). The increase in adiponectin at the twelfth month was negatively correlated with baseline adiponectin levels ( $r=-0.61$ ;  $p<0.001$ ) and positively correlated with the percentage of initial weight loss ( $r=0.41$ ;  $p<0.01$ ), percentage of initial body fat loss ( $r=0.48$ ;  $p<0.005$ ), percentage of initial trunk fat mass loss ( $r=0.42$ ;  $p<0.01$ ), and percentage of initial ROI fat mass loss ( $r=0.47$ ;  $p<0.005$ ).

In relation to nutrient intakes, a significant reduction was observed in energy intake at 6 months after GBP ( $14.6\pm5.9$  vs.  $10.9\pm3.8$  kcal/kg/day;  $p<0.001$ ), although energy

intake at the twelfth month showed no difference with baseline intake ( $14.3\pm2.8$  kcal/kg/day). Protein intake (g/kg/day) showed no significant difference at sixth and twelfth-month post GBP ( $0.56\pm0.22$  vs.  $0.69\pm0.17$  g/kg/day, respectively) compared to baseline values ( $0.61\pm0.19$  g/kg/day).

Dietary calcium intake was no different between initial estimation ( $486\pm222$  mg/day; 95% Confidence Interval (CI): 158–832 mg/day) and the records obtained at sixth ( $393\pm230$  mg/day; 95% CI: 116–838 mg/day) and twelfth-month post GBP ( $468\pm289$  mg/day; 95% CI: 102–1069 mg/day). Calcium intake from supplements was lower in the first 6 months post surgery ( $393\pm230$  mg/day; 95% CI: 319–815 mg/day) compared to the sixth to twelfth-month period ( $468\pm289$  mg/day; 95% CI: 349–973 mg/day;  $p<0.001$ ). Vitamin D intake from supplements was also lower at the first period ( $322\pm191$  U/day; 95% CI: 143–652 U/day) compared to the second period ( $412\pm236$  U/day; 95% CI: 185–762 U/day). No correlation was observed between dietary calcium intake or supplemental Ca plus vitamin D and the changes of total BMD, spine BMD and pelvis BMD.


## Discussion

Beneficial effects of bariatric surgery in morbid obese patients are widely recognized, especially those related to correction of comorbidities [7, 10]. Some potential side-effects however, such as decreased bone mineral density (BMD), are only partially understood. The most dramatic effects of bariatric surgery on bone mineral density have been noted mainly, but not exclusively, in malabsorptive surgery [28]. Roux-en-Y gastric bypass (GBP), although having a malabsorption component, is predominantly a restrictive technique. The extent and clinical implications of potential adverse effects of GBP on bone metabolism are matter of debate [13, 29–31].

In the present study, we observed a 3% decrease of total BMD after 1 year of GBP, with column and pelvis (7.4% and 10.5%, respectively) being their main contributors, which is in agreement with previous reports [13, 29, 30].

The site-dependent differential changes of BMD observed suggest that the effect of reduced body weight accompanied with less mechanic load is highly relevant in determining BMD. Nevertheless, a similar effect could also be induced by secondary hyperparathyroidism [32–34]. This condition is more frequent in individuals older than the subjects studied here. We did not determine parathyroid-related parameters in our subjects; therefore, we cannot disregard the presence of such effect.

Unlike the femur neck and pelvis having predominantly cortical bone, the column has a higher proportion of



**Fig. 1** Bone mineral density after gastric bypass: changes with respect to baseline. \*Significant change from previous value with  $p<0.001$ , \*\*significant change from previous value with  $p<0.05$



trabecular bone. The latter is metabolically more active and, in consequence, it may be more sensitive to humoral factors [21].

Although the literature mentions decreased calcium and vitamin D intake/absorption, secondary hyperparathyroidism and reduced mechanic load on the skeleton as the main factors involved in reduced BMD after GBP, recent evidence suggests that a number of hormonal factors may also participate in this process [28]. The knowledge about the extent of involvement and the mechanisms underlying such effects is limited. Leptin concentration is markedly affected by drastic weight modifications as those observed after GBP. Its association with BMD is rather weak however [19], possibly because this hormone has a dual role. On one hand, it shows an indirect pro-osteoclastic activity by activating the sympathetic system and stimulating of  $\beta_2$  adrenergic receptors. On the other hand, it inhibits osteoclastic differentiation by participating in the cocaine–amphetamine-regulated transcript pathway (CART) [28]. Adiponectin has a clear anti-osteogenic activity, demonstrated by *in vitro* studies, probably through its role as ligand of necrosis factor kappa  $\beta$  (NFK $\beta$ ) receptor [35, 36]. Berner et al. [36] have demonstrated that adiponectin and its receptors (AdipoR1 and AdipoR2) are expressed in human osteoblasts, and also that the supplementation of culture medium with recombinant adiponectin enhances the proliferation of osteoblasts. Adiponectin also is negatively associated with BMD in humans, independently of their body fat mass [21]. In our study, as expected, adiponectin was dramatically increased after GBP. Up to our knowledge, this is the first study reporting a significant association between serum adiponectin and reduction of BMD in non-menopausal women after 1 year of GBP. Since both variables were also associated to initial body fat mass, weight change and fat mass change after 1 year of GBP, correlation analyses were conducted adjusting these variables. Such correction did not affect the described association.

In terms of the effects of dietary changes as determinants of the modifications of BMD in this type of patients, it has been suggested that low intakes of calcium and vitamin D associated to low consumption of dairy products after GBP may be responsible for the increase of PTH and stimulation of bone resorption, especially at the cortical bone. As a result, supplementation with these nutrients seems to be widely justified. Nevertheless, there is no agreement on the optimal amount to be provided after GBP. In the study by Goode et al., secondary hyperparathyroidism post BPG could not be corrected with the administration of 1,200 mg of calcium and 8  $\mu$ g of vitamin D, although the period of observation was only up to 6 months [29]. In our study, the prescribed daily calcium and vitamin D supplements ranged between 640 and 1,000 mg for calcium and between 400 and 800 IU for vitamin D. Those amounts of supplemen-

tation are in agreement with the recommendations of Johnson et al. [30]. As expected, the actual amount consumed varied according the compliance of the individuals. In our study, this reached 83% of the planned amount on average. Finally, the reduction of BMD was not correlated to dietary calcium intakes or to the consumption of calcium plus vitamin D from supplements.

The lack of association between BMD change and calcium and vitamin D total intakes may be explained by the anatomical alterations produced by the GBP. This modification would reduce calcium absorption, which takes place mainly at the duodenum and proximal jejunum. It would also favor fat and fat-soluble vitamins malabsorption as consequence of impaired combination with bile salts.

Since all variables significantly associated to BMD change (initial weight, weight loss, fat mass loss, final adiponectin) only explain 37% of the variance, it is possible that calcium and vitamin-D-impaired absorption is actually playing a major role. In our study, calcium absorption determinations were not carried out. Among the clinical implications of the results, it could be speculated that calcium and vitamin D supplementation, unless they are provided in very high amounts, will have a modest effect preventing BMD decrease.

BMD decrease seems to be highly associated to weight loss in both surgical and non-surgical interventions [16, 37]. This, along with the concomitant adiponectin increase, would condition a loss of protective effect of obesity against osteoporosis [30, 38]. According to Johnson et al. [30, 39], BMD decrease is made evident mainly during the first year, which is also the period when the major weight changes are observed. Beyond that point, both body weight and BMD tend to remain stable.

In conclusion, in our group of non-menopausal women studied after 1 year of gastric bypass, BMD was significantly reduced, mainly at the pelvis. BMD decrease was more important in the patients with greater values of initial body weight, initial BMD, fat mass loss and final serum adiponectin. Adiponectin involvement in the process of bone modeling may explain, at least partially, the decrease of BMD observed after GBP. In order to achieve a better understanding of these observations and their implications, further studies including a wide range of calcium and vitamin D supplements and calcium absorption measurements are needed.

**Acknowledgments** Supported by the National Fund for Science and Technology, Fondecyt grant 1040765. The authors are indebted to Dr Guillermo Watkins, Juan C Díaz, Fernando Maluenda and Dr. Italo Braguetto, from the Department of Surgery, University Clinical Hospital who performed a significant number of the surgical procedures and to the Nutritionists Ms Emma Díaz and Ms Andrea Rizzo for their invaluable collaboration in the dietary control of the patients.

The total supply of calcium–vitamin D supplements used in the study was donated by Andromaco Laboratories (Santiago, Chile). Their contribution allowed for a constant and controlled provision of these products. The multi-vitamin mineral prescriptions were filled by Farmacias Ahumada (Santiago, Chile) at a reduced price. Their contribution is greatly appreciated.

## References

- Haslam DW, James WPT. Obesity. *Lancet* 2005;366:1197–209.
- Ministerio de Salud. Encuesta Nacional de Salud, 2003. [http://epi.minsal.cl/epi/html/invest/ENS/ENS\\_mayo2004.pdf](http://epi.minsal.cl/epi/html/invest/ENS/ENS_mayo2004.pdf). Minsal, Chile.
- Bray GA. Overweight is risking fate: definition, classification, prevalence, and risks. *Ann N Y Acad Sci.* 1987;499:14–28.
- WHO. Obesity. Preventing and management the global epidemic Report of a WHO Consultation on Obesity. Geneva: WHO; 1997.
- Rosmond R, Lapidus L, Marin P, et al. Mental distress, obesity and body fat distribution in middle-aged men. *Obes Res.* 1996;4:245–52.
- Fontaine KR, Redden DT, Wang C, et al. Years of life lost due to obesity. *JAMA* 2003;289:187–93.
- Buchwald H, Avidor Y, Braunwald E, et al. Bariatric surgery: a systematic review and meta-analysis. *JAMA* 2004;292:1724–37.
- Maggard MA, Shugarman LR, Suttrop M, et al. Meta-analysis: surgical treatment of obesity. *Ann Intern Med.* 2005;142:547–59.
- Crookes PF. Surgical treatment of morbid obesity. *Annu Rev Med.* 2006;57:243–64.
- Shah M, Simha V, Garg A. Long-term impact of bariatric surgery on body weight, comorbidities, and nutritional status. *J Clin Endocrinol Metab.* 2006;91:4223–31.
- DeMaria EJ. Bariatric surgery for morbid obesity. *N Engl J Med.* 2007;356:2176–83.
- Bloomberg R, Fleishman A, Nalle J, et al. Nutritional deficiencies following bariatric surgery: what have we learned? *Obes Surg.* 2005;15:145–54.
- Ott MT, Fanti P, Malluche HH, et al. Biochemical evidence of metabolic bone disease in women following roux-Y gastric bypass for morbid obesity. *Obes Surg.* 1992;2:341–8.
- Goldner WS, O'Dorisio TM, Dillon JS, et al. Severe metabolic bone disease as a long-term complication of obesity surgery. *Obes Surg.* 2002;12:685–92.
- Shaker JL, Norton AJ, Woods MF, et al. Secondary hyperparathyroidism and osteopenia in women following gastric exclusion surgery for obesity. *Osteoporos Int.* 1991;1:177–81.
- Ricci TA, Heymsfield SB, Pierson RN, et al. Moderate energy restriction increases bone resorption in obese postmenopausal women. *Am J Clin Nutr.* 2001;73:347–52.
- Edelstein SL, Barrett-Connor E. Relation between body size and bone mineral density in elderly men and women. *Am J Epidemiol.* 1993;138:160–9.
- Baumgartner RN, Ross RR, Waters DL, et al. Serum leptin in elderly people: associations with sex hormones, insulin, and adipose tissue volumes. *Obes Res.* 1999;7:141–9.
- Thomas T, Burguera B. Is leptin the link between fat and bone mass. *J Bone Miner Res.* 2002;17:1563–9.
- Maccarinelli G, Sibilia V, Torsello A, et al. Ghrelin regulates proliferation and differentiation of osteoblastic cells. *J Endocrinol.* 2005;184:249–56.
- Lenchik L, Register TC, Hsu F-C, et al. Adiponectin as a novel determinant of bone mineral density and visceral fat. *Bone* 2003;33:646–51.
- Csendes A, Burdiles P, Papapietro K, et al. Results of gastric bypass plus resection of the distal excluded gastric segment in patients with morbid obesity. *J Gastrointest Surg.* 2005;9:121–31.
- Lohman TG. Anthropometry and body composition. In: Lohman TG, Roche AF, Martorell R, editors. *Anthropometric Standardization Reference Manual*. Champaign, IL: Human Kinetics; 1988. p. 125–9.
- Jebb SA. Measurement of soft tissue composition by dual X-ray absorptiometry. *Br J Nutr.* 1997;77:151–63.
- Clasey JL, Bouchard C, Teates CD, et al. The use of anthropometric and dual-energy X-ray absorptiometry (DXA) measures to estimate total abdominal and abdominal visceral fat in men and women. *Obes Res.* 1999;7:256–64.
- Snijder MB, Visser M, Dekker JM, et al. The prediction of visceral fat by dual-energy X-ray absorptiometry in the elderly: a comparison with computed tomography and anthropometry. *Int J Obes.* 2002;26:984–93.
- Snedecor GW, Cochran WG. *Statistical methods*, 8th edition. Ames, Iowa: The Iowa State University Press; 1989.
- Wucher H, Ciangura C, Poitou C, et al. Effects of weight loss on bone status after bariatric surgery: association between adipokines and bone markers. *Obes Surg.* 2008;18:58–65.
- Goode LR, Brodin RE, Chowdhury HA, et al. Bone and gastric bypass surgery: effects of dietary calcium and vitamin D. *Obes Res.* 2004;12:40–7.
- Johnson JM, Maher JW, Samuel I, et al. Effects of gastric bypass procedures on bone mineral density, calcium, parathyroid hormone, and vitamin D. *J Gastrointest Surg.* 2005;9:1106–10.
- von Mach MA, Stoeckli R, Bilz S, et al. Changes in bone mineral content after surgical treatment of morbid obesity. *Metabolism* 2004;53:918–21.
- El-Kadri LJ, Savassi PR, de Almeida AC, et al. Calcium metabolism in pre- and postmenopausal morbidly obese women at baseline and after laparoscopic roux-en-Y gastric bypass. *Obes Surg.* 2004;14:1062–6.
- Silverberg SJ, Shane E, de la Cruz L, et al. Skeletal disease in primary hyperparathyroidism. *J Bone Miner Res.* 1989;4:283–91.
- Dempster DW, Parisien M, Silverberg SJ, et al. On the mechanism of cancellous bone preservation in postmenopausal women with mild primary hyperparathyroidism. *J Clin Endocrinol Metab.* 1999;84:1562–6.
- Luo XH, Guo LJ, Xie H, et al. Adiponectin stimulates RANKL and inhibits OPG expression in human osteoblasts through the MAPK signaling pathway. *J Bone Miner Res.* 2006;21:1648–56.
- Berner HS, Lyngstadaas SP, Spahr A, et al. Adiponectin and its receptors are expressed in bone-forming cells. *Bone* 2004;35:842–9.
- Andersen RE, Wadden TA, Herzog RJ. Changes in bone mineral content in obese dieting women. *Metabolism* 1997;46:857–61.
- Barrera G, Bunout D, Gattas V, et al. A high body mass index protects against femoral neck osteoporosis in healthy elderly subjects. *Nutrition* 2004;20:769–71.
- Johnson JM, Maher JW, DeMaria EJ, et al. The long-term effects of gastric bypass on vitamin D metabolism. *Ann Surg.* 2006;243:701–4.