

Metabolic profile of the different phenotypes of polycystic ovary syndrome in two Latin American populations

Amanda Ladrón de Guevara, M.D.,^a Carolina Fux-Otta, M.D., Ph.D.,^b Nicolás Crisosto, M.D., Ph.D.,^a Paula Szafryk de Mereshian, M.D., Ph.D.,^b Bárbara Echiburú, Ph.D.,^a Gabriel Iraci, M.D.,^c Francisco Perez-Bravo, Ph.D.,^d and Teresa Sir-Petermann, M.D., Ph.D.^a

^a Endocrinology and Metabolism Laboratory, West Division, School of Medicine, University of Chile, Santiago, Chile; ^b Endocrinology and Diabetes Department, Maternity and Neonatology University Hospital, and ^c Applied Pharmacology Cathedra, National Córdoba University, Córdoba, Argentina; and ^d Laboratory of Nutritional Genomics, Department of Nutrition, Faculty of Medicine, University of Chile, Santiago, Chile

Objective: To evaluate the metabolic profile of Chilean and Argentinian women with polycystic ovary syndrome (PCOS) according to the Rotterdam criteria.

Design: Observational cross-sectional study.

Setting: Academic centers.

Patient(s): Women with PCOS, aged 18–39 years: 220 Chilean (PCOS_{Ch}) and 206 Argentinian (PCOS_{Ar}).

Intervention(s): Physical examination, fasting blood samples for androgens, gonadotropins, metabolic parameters, and a transvaginal ultrasound.

Main Outcome Measure(s): Comparative analysis of the metabolic profile in both populations divided into four phenotypes.

Result(s): The distribution of the different phenotypes was different in both populations. PCOS_{Ch} women showed a higher body mass index and a higher percentage of metabolic syndrome in all phenotypes compared with the PCOS_{Ar} women. The PCOS_{Ar} women exhibited a statistically significantly higher diastolic blood pressure in phenotypes A, B, and C and a higher percentage of hypertension in phenotypes A and D compared with the PCOS_{Ch} women.

Conclusion(s): The data show differences in the metabolic profile of both populations. PCOS_{Ch} women presented with greater metabolic alterations such as dysglycemia and dyslipidemia and a higher prevalence of metabolic syndrome, independent of the phenotype. The PCOS_{Ar} patients showed more elevated blood pressure. Ethnic diversity associated with environmental factors are fundamental elements in the analysis of the PCOS phenotypes. (Fertil Steril® 2014;101:1732–9. ©2014 by American Society for Reproductive Medicine.)

Key Words: Argentina, Chile, metabolic syndrome, polycystic ovary syndrome, Rotterdam phenotypes

Discuss: You can discuss this article with its authors and with other ASRM members at <http://fertstertforum.com/deguevaraa-metabolic-polycystic-ovary-syndrome-latin-america/>



Use your smartphone to scan this QR code and connect to the discussion forum for this article now.*

* Download a free QR code scanner by searching for "QR scanner" in your smartphone's app store or app marketplace.

Received October 11, 2013; revised February 7, 2014; accepted February 11, 2014; published online March 22, 2014.

A.L.d.G. has nothing to disclose. C.F.-O. has nothing to disclose. N.C. has nothing to disclose. P.S.d.M. has nothing to disclose. B.E. has nothing to disclose. G.I. has nothing to disclose. F.P.-B. has nothing to disclose. T.S.-P. has nothing to disclose.

Supported by Fondo Nacional de Desarrollo Científico y Tecnológico (National Fund for Scientific and Technological Research; Fondecyt; Grant 1110864) SOCHED (Chilean Society of Endocrinology and Diabetes; Grant 2009-48 05) and by the Alexander von Humboldt and Florencio Fiorini foundations.

Reprint requests: Teresa Sir-Petermann, M.D., Ph.D., Laboratory of Endocrinology, Department of Medicine W. Division, School of Medicine, Las Palmeras 299, Interior Quinta Normal, Casilla 33052, Correo 33 (Zip Code: 8320000), Santiago, Chile (E-mail: tsir@med.uchile.cl).

Fertility and Sterility® Vol. 101, No. 6, June 2014 0015-0282/\$36.00
Copyright ©2014 American Society for Reproductive Medicine, Published by Elsevier Inc.
<http://dx.doi.org/10.1016/j.fertnstert.2014.02.020>

Polycystic ovary syndrome (PCOS) is a hyperandrogenic disorder associated with chronic oligo-ovulation and polycystic ovary morphology. It is the most common cause of hyperandrogenism, with a prevalence reaching up to 15% when using the Rotterdam criteria described in 2003 (1). Hyperandrogenism and chronic anovulation are the main elements of this definition; the ultrasound pattern of polycystic ovaries is the third

element incorporated in the 2003 consensus (1). According to this definition, the diagnosis of PCOS may be established with two of the three criteria described, generating four different phenotypes:

Phenotype A: hyperandrogenic women with oligoanovulation and transvaginal ultrasound with polycystic ovarian morphology (H/O/PCOM).

Phenotype B: women with hyperandrogenism, oligoanovulation, and without PCOM (H/O).

Phenotype C: hyperandrogenic women with ovulatory cycles and PCOM (H/PCOM).

Phenotype D: normoandrogenic women with oligoanovulation and PCOM (O/PCOM).

Recently, it has been observed that each PCOS phenotype is associated with a different metabolic risk. Most studies have shown that hyperandrogenism is the main factor for the development of metabolic and cardiovascular alterations, with phenotypes A and B having a higher metabolic risk than phenotype D, whose risk is comparable to that of control women (2, 3). However, these phenotypes may vary in individuals and may be modified by changes in weight, treatment, and/or healthy lifestyle factors (4).

It has been proposed that the development of PCOS and its phenotypic expressions may be influenced by genetic and environmental factors that can affect a woman from her intrauterine life until the reproductive stage (4, 5). In this regard, racial and ethnic characteristics are a very important genetic aspect to be considered in PCOS and its phenotypic expressions (6). This background may influence the clinical manifestations of PCOS. For instance, Middle Eastern women have a greater prevalence of hirsutism (2) whereas among East Asian women hirsutism is less common (7, 8). In Indian women, acne has been described as the most prevalent clinical expression of hyperandrogenism (9).

The presence of the metabolic syndrome has been associated with PCOS with variable frequency; its prevalence is lower in East Asian women (7, 8) and higher in women of African, South Asian, and Hispanic descent (2, 10). A northern California study of women with PCOS found that, compared with white patients, black and Hispanic patients were more likely and Asian patients less likely to be obese, the Asian and Hispanic women were more likely to have diabetes, and the black women were more likely and Hispanic women less likely to have hypertension (10).

The phenotypic distribution of PCOS in Latin American populations is only partially known. Thus, our study compares the metabolic profile of women with PCOS from two neighboring countries with different ethnic compositions. Argentina, and especially the city of Córdoba, has been influenced by a large, mainly European immigration wave composed primarily of Italians and secondarily of Spaniards (11). In contrast, in Chile over 65% of the population is of Spanish descent, with the phenotypic characteristics of a mostly white population: the average proportion is 60% Hispanic and 40% Amerindian (12, 13). Our study compares the metabolic profile, classified into the four phenotypes

described according to the Rotterdam criteria, of 220 women with PCOS from Santiago de Chile (PCOS_{Ch}) and 206 women with PCOS from the province of Córdoba, Argentina (PCOS_{Ar}), who have a different ethnic background.

MATERIALS AND METHODS

Patients

We studied 220 PCOS_{Ch} women who were treated at the polyclinic of the Department of Endocrinology and Metabolism of Universidad de Chile in Santiago (Chile) and 206 PCOS_{Ar} women who were treated at the Department of Endocrinology and Diabetes of Universidad de Córdoba (Argentina), both academic health centers serving patients from a socioeconomically middle-class area. All the women were between 18 and 39 years old and had a body mass index (BMI) between 18 and 35. The study was approved by the institutional review boards of both academic centers, and all participants gave their written informed consent.

An analysis of the ethnic background in both countries was conducted. All women belonged to the white race. The PCOS_{Ch} group was composed of women who were 90% Hispanic-Amerindian mixture, 7% Amerindian, and 3% other races. The PCOS_{Ar} group was composed of 87% of European, predominantly Italian, descent; 10% Hispanic-Amerindian mixture; and 3% other ethnicities. The differences in the distribution pattern of ethnicities between the two groups was statistically significant ($P < .001$). Comparisons were made only between the Chilean and Argentinean country groups. We did not make comparisons between the various ethnic groups because it would have meant combining only 10% of the Argentinean group with 90% of the Chilean group; this would have left other two groups too small for analysis and ignored the potential lifestyle differences between the country groups.

The diagnosis of PCOS was established according to the Rotterdam criteria (1), including at least two of the following elements:

Hyperandrogenism (H): modified Ferriman-Gallwey score ≥ 8 and/or total serum testosterone (T) ≥ 80 ng/dL (≥ 2.77 nmol/L).

Ovulatory dysfunction (O): oligomenorrhea (cycles >35 days) or amenorrhea (no menses in the last 6 months), negative pregnancy test, and progesterone level <4 ng/mL (12.72 nmol/L) before beginning of the study.

Polycystic ovaries at transvaginal ultrasound (PCOM): ≥ 12 follicles measuring 2–9 mm in diameter and/or increased ovarian volume (>10 mL) in at least one ovary.

The exclusion criteria were other causes of hyperandrogenism (Cushing syndrome, congenital adrenal hyperplasia, or androgen-secreting tumors), a previous diagnosis of type 2 diabetes mellitus and/or treatment with insulin-sensitizing drugs, or use of contraceptives, antiandrogens, or glucocorticoids 6 months before the beginning of the study. Moreover, all participants had normal thyroid function, normal prolactin levels, and a follicle-stimulating hormone (FSH) level in the

premenopausal range. All patients signed a written informed consent before entering the study, which was conducted in accordance with the Declaration of Helsinki and approved by the institutional ethics committee of each health center.

Study Protocol

A complete clinical history was obtained for all participants, including their family and ethnic background. The estimation of their ethnic background was based on the two surnames of each family member. After the interview, a complete physical examination was performed with anthropometric measurements, including weight, height, waist circumference, waist-to-hip ratio, waist-to-height ratio, body mass index (BMI), systolic blood pressure, and diastolic blood pressure. The degree of hirsutism was determined according to the modified Ferriman-Gallwey score (14).

The studies were conducted during the early follicular phase (days 3 to 7) of a spontaneous cycle. Patients were admitted at 8:00 AM to the corresponding health center after a 12-hour fast, at which point an oral glucose tolerance test was performed (75-g glucose in 250 mL water), with glucose and insulin measurements in the basal sample and 120 minutes after the glucose load. Additionally in the basal sample, concentrations were determined of luteinizing hormone (LH), FSH, T, androstenedione (Δ 4), dehydroepiandrosterone sulphate (DHEAS), 17-hydroxyprogesterone (17-OHP), and sex hormone-binding globulin (SHBG) as well as a lipid profile: total cholesterol (TC), triglycerides, and high-density lipoprotein cholesterol (HDL-C).

Glucose tolerance was evaluated using the criteria of the American Diabetes Association (ADA) (15). Impaired fasting glucose was defined as fasting glycemia >100 mg/dL and <126 mg/dL, and impaired glucose tolerance as a 2-hour glucose, after the glucose load, between 140 and 200 mg/dL. Insulin resistance was calculated through the homeostatic model assessment for insulin resistance (HOMA-IR) (16).

A transvaginal ultrasound examination was performed with a transducer of 3.8–7.5 MHz (SSD-4000; Aloka) in the PCOS_{Ch} group, and with a transvaginal transducer of 5–9 MHz (Voluson 730 Expert; General Electric) in the PCOS_{Ar} women. We identified PCOM according to the Rotterdam consensus as the presence of either ≥ 12 follicles measuring 2–9 mm in diameter and/or an ovarian volume >10 mL in one or more ovaries (1). In each group, the same operator performed all the ultrasound examinations. For comparisons between groups regarding ovarian volume, the average volume was considered.

Assays

Serum glucose was determined by the glucose oxidase method. The intra-assay coefficient of variation (CV) was $<2.0\%$. Lipid profile was determined by standard colorimetric assay (Photometric Instrument 4010; Roche). For both groups, the intra-assay and interassay CVs were less than 10%. The low-density lipoprotein cholesterol (LDL-C) concentration was calculated by Friedewald's formula: $LDL-C = TC - (HDL + Tg/5)$.

The insulin concentration was measured by radioimmunoassay (RIA) (Diagnostic Systems Laboratories) in the PCOS_{Ch} patients (intra-assay and interassay CV: 5% and 8%, respectively), and by electrochemiluminescence in PCOS_{Ar} patients (intra-assay and interassay CV: $<10\%$). The HOMA-IR was calculated by the formula: $(\text{Basal glycemia} \times \text{Basal insulin})/405$. The presence of the metabolic syndrome was defined according to the criteria described in the National Cholesterol Education Program Adult Treatment Panel III guidelines (ATP III) (17).

In both populations, LH, FSH, and DHEAS were determined by electrochemiluminescence assays (Roche). Serum Δ 4 and 17-OHP were analyzed by RIA. We determined SHBG by radioimmunometric assay (Diagnostic Products Corp.) in the PCOS_{Ch} patients and by electrochemiluminescence in the PCOS_{Ar} patients. Serum T was determined by RIA (Diagnostic Systems Laboratories) in the PCOS_{Ch} patients and by electrochemiluminescence in PCOS_{Ar} patients. All assays conducted in both PCOS populations had an intra-assay and interassay CV of $<10\%$. The free androgen index (FAI) was calculated using the following formula: $FAI = [T \text{ (nmol/L)} / SHBG \text{ (nmol/L)}] \times 100$ (18).

Statistical Analysis

Distribution was assessed using the Shapiro-Wilk test. Results are expressed as median and range. The Kruskal-Wallis test was used for comparisons between groups. The chi-square test was used for percentage comparisons. To establish relationships between dichotomous variables, the odds ratio (OR) with a 95% confidence interval (CI) was calculated by logistic regression analysis. All data were analyzed in the STATA 10.0 program (StataCorp). $P < .05$ was considered statistically significant.


RESULTS

Both the PCOS_{Ch} and PCOS_{Ar} groups were classified into the following four phenotypes: A: PCOS_{Ch} $n = 158$, and PCOS_{Ar} $n = 104$; B: PCOS_{Ch} $n = 23$, and PCOS_{Ar} $n = 40$; C: PCOS_{Ch} $n = 36$, and PCOS_{Ar} $n = 41$; D: PCOS_{Ch} $n = 3$, and PCOS_{Ar} $n = 21$. The percentage of distribution of the different PCOS phenotypes was as follows: for PCOS_{Ch}, A 72%; B 10.5%; C 16.5%, and D 1%; and for PCOS_{Ar}, A 50.4%; B 19.4%; C 20%, and D 10% (chi-square = 14.5, $P < .002$).

Regarding the anthropometric variables, BMI was statistically significantly higher in the total PCOS_{Ch} group, but the waist circumference, waist-to-hip ratio, and waist-to-height ratio were comparable between both groups. Diastolic blood pressure was statistically significantly higher in the PCOS_{Ar} group. Family background was established considering both the paternal and maternal first- and second-degree relatives. Regarding family background, 64.5% of PCOS_{Ch} women had a family history of diabetes, and 30% a family history of hypertension. In PCOS_{Ar} women, 27% had a family background of diabetes and 46.3% of hypertension (Supplemental Table 1, available online).

Age was comparable between the PCOS_{Ch} and PCOS_{Ar} women in each phenotype. Figure 1 shows the clinical parameters of the statistically significant differences between the

FIGURE 1



Clinical features in Chilean and Argentinian women with polycystic ovary syndrome according to the different phenotypes. * $P < .05$.

Ladrón de Guevara. PCOS phenotypes in Chile and Argentina. *Fertil Steril* 2014.

PCOS_{Ch} and PCOS_{Ar} women in each of the four phenotypes. The PCOS_{Ch} women showed a statistically significantly higher BMI in all of the phenotypes and a statistically significantly higher Ferriman-Gallwey score in phenotypes A, B, and C, than the PCOS_{Ar} women.

Among the PCOS_{Ar} women with phenotype A, a statistically significantly higher systolic blood pressure was found than in PCOS_{Ch} women. For diastolic blood pressure, PCOS_{Ar} women exhibited statistically significantly higher blood pressure in phenotypes A, B, and C compared with the PCOS_{Ch} women.

Table 1 shows the hormone profile and ovarian volume in the PCOS_{Ch} and PCOS_{Ar} groups according to the different phenotypes. The concentrations of LH and FSH were similar in the A, B, and C phenotypes for both populations, and the LH/FSH ratio was comparable in the four phenotypes between both populations. Different assays were used to assess T and SHBG, so these measurements and the FAI calculation were not compared between the PCOS_{Ch} and PCOS_{Ar} groups. When analyzing the two populations separately, both exhibited a statistically significant difference in T concentration between phenotypes A, B, and C with respect to D. The same occurred with the FAI. Nevertheless, this information has to be taken carefully in the PCOS_{Ch} group as the size of the D phenotype was very small.

Regarding SHBG concentrations, in the PCOS_{Ar} patients there was a statistically significant difference between phenotypes A and D, whereas in the PCOS_{Ch} patients there was no statistically significant difference when comparing the four phenotypes. The $\Delta 4$ concentration was higher in the PCOS_{Ar} group and reached statistical significance in phenotypes B and C in comparison with the PCOS_{Ch} group. The DHEAS con-

centration tended to be higher in the PCOS_{Ar} group, reaching statistical significance only in phenotype C.

The 17-OHP concentration and the average ovarian volumes were discretely higher in phenotype A for PCOS_{Ar}. This difference reached statistical significance.

Metabolic Profile

Table 2 shows the metabolic profile of both populations divided into phenotypes. The 2-hour glucose levels tended to be higher in the PCOS_{Ch} group, and were statistically significantly higher in phenotypes C and D with respect to PCOS_{Ar} (this latter comparison has to be taken carefully as the PCOS_{Ch} D group was very small).

In the PCOS_{Ch} patients, insulin concentration and HOMA-IR were comparable between the four phenotypes. In the PCOS_{Ar} patients, insulin concentration and HOMA-IR were statistically significantly higher in phenotype A compared with C. The finding of type 2 diabetes was generally very low and comparable between both PCOS populations.

The triglyceride concentration was higher in the PCOS_{Ch} group for all phenotypes, and was statistically significant in phenotypes A and B. The same occurred with LDL-C, which was statistically significantly higher in the PCOS_{Ch} group for the A, B, and C phenotypes compared with PCOS_{Ar}. The HDL-C level was lower in all the phenotypes of PCOS_{Ch}, which was statistically significant for the A, B, and C phenotypes compared with the PCOS_{Ar} group. The triglycerides concentration was statistically significantly higher in the PCOS_{Ch} group for all phenotypes compared with PCOS_{Ar}.

Supplemental Table 2 (available online) shows the prevalence of the metabolic syndrome for both total groups

TABLE 1

Hormone profile and ovarian volume in Chilean and Argentinian women with PCOS according to the different phenotypes.

	A		B		C		D	
	PCOS _{Ch} (n = 158)	PCOS _{Ar} (n = 104)	PCOS _{Ch} (n = 23)	PCOS _{Ar} (n = 40)	PCOS _{Ch} (n = 36)	PCOS _{Ar} (n = 41)	PCOS _{Ch} (n = 3)	PCOS _{Ar} (n = 21)
LH (IU/L)	8.7 (1.31-14)	9.1 (3.1-24)	8 (2.2-16.4)	8.6 (3.2-12.8)	3.9 (1-14.7)	7 (4.7-12.7)	6 (5.3-8.1) ^a	9.8 (6.1-19.5)
FSH (IU/L)	5.4 (0.9-14)	5.4 (3-11)	4.6 (1.8-8.3)	5.2 (2.1-9.9)	4.8 (1.6-13.7)	5.1 (3.2-7.3)	4.1 (3.2-4.3) ^a	6.8 (3.2-9.3)
LH/FSH	1.7 (0.3-6.2)	1.6 (0.7-3.5)	1.4 (0.4-3.9)	1.7 (0.6-3.6)	0.99 (0.3-8.3)	1.5 (0.6-2.4)	1.9 (1.2-1.9)	1.7 (0.9-2.5)
T (ng/mL)	0.94 (0.24-2.05)	0.89 (0.76-1.76)	0.93 (0.38-1.41)	0.9 (0.8-1.8)	0.85 (0.34-2.27)	0.87 (0.8-1.3)	0.5 (0.44-0.52) ^b	0.29 (0.21-0.65) ^b
SHBG (nmol/L)	25.4 (9.8-128.6)	39.6 (16-161)	27 (6.61-76)	46.7 (16-100)	37.7 (4.6-137.4)	51.1 (14.3-100)	24 (17-27)	59.2 (29.2-105.1) ^c
FAI	12.17 (1.41-48.82)	8.72 (1.72-25.04)	10.4 (2.6-38.29)	7.3 (3.3-21.8)	8.6 (3.1-49.3)	6.1 (2.8-19.0)	3.7 (3.3-6.7) ^b	1.98 (0.78-5.22) ^b
Δ4 (ng/mL)	3.1 (0.8-7.1)	4.6 (2.3-8)	2.9 (1-5.4) ^a	4.2 (2.2-6.8)	3.2 (0.9-7.4) ^a	4.1 (2.7-7)	2 (1.7-2.3)	2.5 (1-7.1)
DHEAS (μg/dL)	247.1 (194-1,970)	343 (30-607)	248 (167-1,800)	375 (178-557)	272 (181-580) ^a	350 (170-570)	186.6 (172-201)	206.1 (124-301)
17-OHP (ng/mL)	1.5 (0.2-4.6) ^a	1.8 (0.8-3.6)	1.4 (0.1-3.3)	1.7 (0.8-2.3)	1.5 (0.5-3.5)	1.7 (0.9-2.7)	1.2 (1-1.3)	0.89 (0.5-1.7)
Ovarian volume (mL) ^d	10 (2.5-32.9) ^a	11.5 (4.3-25)	6.4 (2.6-8.7)	5.5 (3.6-8.3)	9.8 (4.9-26.3)	10.9 (6-19.5)	8.7 (8.3-15.6)	10.5 (8-30.9)

Note: Comparisons regarding phenotype D of the Chilean PCOS (PCOS_{Ch}) group have to be taken carefully as the size of the group is very small. Testosterone and SHBG levels were measured by radioimmunoassay in PCOS_{Ch} patients and by electrochemiluminescence in Argentinian PCOS (PCOS_{Ar}) patients. Thus, they cannot be directly compared. Δ4 = androstenedione; DHEAS = dehydroepiandrosterone sulfate; FAI = free androgen index; FSH = follicle-stimulating hormone; LH = luteinizing hormone; 17-OHP = 17-hydroxiprogesterone; PCOS = polycystic ovary syndrome; SHBG = sex hormone-binding globulin; T = total testosterone.

^a P < .05 compared with A, B, and C from the same phenotype.

^b P < .05 compared with A, B, and C from the same PCOS group.

^c P < .05 compared with A from the same group.

^d Average ovarian volume.

Ladrón de Guevara. PCOS phenotypes in Chile and Argentina. *Fertil Steril* 2014.

(PCOS_{Ch} = 52.9 % vs. PCOS_{Ar} = 36.5 %; P < .05). Analyzing the metabolic syndrome by each phenotype, the PCOS_{Ch} women showed a higher prevalence of the metabolic syndrome in all phenotypes; this was statistically significant in the B and D phenotypes with respect to PCOS_{Ar} (Table 3).

The percentage of 2-hour glucose > 140 mg/dL in PCOS_{Ch} was statistically significantly higher in phenotypes B, C, and D compared with PCOS_{Ar}. For the percentage of women with increased waist circumference (>88 cm), both populations were comparable between the different phenotypes, except for phenotype D in whom it was higher in PCOS_{Ch} compared with PCOS_{Ar}. The percentage of elevated blood pressure (≥ 130/85) was higher in the PCOS_{Ar} women in the A and D phenotypes. For lipid alterations, the PCOS_{Ch} group showed a statistically significantly higher percentage of HDL-C < 50 mg/dL in the A, B, and C phenotypes; similarly, the percentage of triglycerides > 150 mg/dL was statistically significantly higher in the B, C, and D phenotypes in comparison with the PCOS_{Ar} group.

Multivariate Analysis

In each PCOS group, we evaluated the odds ratio (OR) for the metabolic syndrome according to the phenotype. Multivariate analysis by logistic regression showed that in the PCOS_{Ar} group there is a statistically significant reduction in the OR for the metabolic syndrome in phenotypes B (OR 0.36; 95% CI, 0.14-0.91; P = .03), C (OR 0.24; 95% CI, 0.08-0.66; P = .006), and D (OR 0.28; 95% CI, 0.08-1.04; P = .06) when compared with phenotype A. In the PCOS_{Ch} group, the OR for the metabolic syndrome was not different between phenotype A versus B (OR 1.77; 95% CI, 0.5-6; P = .36), C (OR 0.56; 95% CI, 0.5-1.98; P = .37), or D (OR 2.5; 95% CI, 0.21-29.1; P = .45).

Note that all comparisons regarding the PCOS_{Ch} D group, although statistically significant, are susceptible to bias: this group was composed of only three patients.

DISCUSSION

In the present study, we analyzed two groups of women with PCOS from two academic centers in the cities of Santiago de Chile and Córdoba, Argentina. It should be highlighted that this is the first study comparing two groups of Latin American women in whom the same criteria for the diagnosis of PCOS were used, whose ages were similar but ethnic conformation was different. The distribution of the PCOS phenotypes was different in the two groups studied (Chilean and Argentinian). Both groups showed a higher prevalence of the classic phenotypes (A and B), as has been described in many studies of Western women with PCOS (2). The D phenotype was more prevalent in PCOS_{Ar} (10%), similar to what has been described in the women with PCOS of southern Europe (19). Classification of PCOS into different Rotterdam phenotypes has allowed the identification of the groups who have a greater cardiometabolic risk (20-23). The presence of hyperandrogenism seems to be strongly related to the development of metabolic alterations, with phenotypes A and B having the highest cardiovascular risk whereas

TABLE 2

Metabolic profile in Chilean and Argentinian women with PCOS according to the different phenotypes.

	A		B		C		D	
	PCOS _{Ch} (n = 158)	PCOS _{Ar} (n = 104)	PCOS _{Ch} (n = 23)	PCOS _{Ar} (n = 40)	PCOS _{Ch} (n = 36)	PCOS _{Ar} (n = 41)	PCOS _{Ch} (n = 3)	PCOS _{Ar} (n = 21)
Fasting glucose (mg/dL)	83 (57–134)	87 (64–121)	84 (67–107)	83 (70–110)	83 (67–126)	82 (64–103)	78 (66–99)	82 (70–106)
2-h glucose (mg/dL)	98 (42–210)	89 (50–196)	92 (45–148)	88 (60–170)	94 (60–229) ^a	83 (63–119)	128 ^a (86–149)	81 (68–140)
Fasting insulin (μIU/mL)	16 (2.5–99)	12 (3–34)	14 (4.7–78)	8.5 (3.2–30)	15.7 (3.7–34)	7.7 (2.1–16) ^b	17 (10–21)	10 (3–22)
2-h insulin (μIU/mL)	77.4 (8.2–868)	42.4 (7.3–136)	129.9 (24.7–159)	41.7 (17.6–127)	71.9 (18.6–300)	ND	51 (20–91)	ND
HOMA-IR	3.5 (1.5–21.2)	2.6 (0.5–9.5)	2.4 (0.98–20.6)	1.8 (0.55–7.4)	3 (0.7–8.7)	1.5 (0.3–4) ^b	4.1 (1.6–4.16)	2.1 (0.5–5.6)
Type 2 diabetes (%)	1.3	0	4.3	0	0	0	0	0
Cholesterol (mg/dL)	195 (118–316) ^a	179 (108–261)	202 (8,149–256) ^a	175 (117–260)	185 (130–273)	172 (96–217)	177 (141–235)	171 (110–216)
LDL-C (mg/dL)	130.6 (57.1–218) ^a	106 (34–185)	130 (44–194) ^a	103 (48–176)	111 (63–186) ^a	99 (47–155)	108 (65–136)	100 (60–145)
Triglycerides (mg/dL)	135 (55–388) ^a	109.5 (45–324)	131 (64–230) ^a	80 (44–247)	108 (62–283) ^a	77 (36–170)	167 (74–215) ^a	79 (42–200)
HDL-C (mg/dL)	37.6 (11–74) ^a	50 (26–77)	40.6 (28.7–56.7) ^a	55.5 (30–81)	38 (30–76) ^a	54 (26–86)	43 (36–54)	50 (30–82)

Note: Comparisons regarding phenotype D of the Chilean PCOS (PCOS_{Ch}) group have to be taken carefully as the size of the group is very small. Insulin levels were measured by radioimmunoassay in the PCOS_{Ch} patients and by electrochemoluminescence in Argentinian PCOS (PCOS_{Ar}) patients. Thus, they cannot be directly compared. ND = no data; PCOS = polycystic ovary syndrome.
^a P < .05 compared with PCOS_{Ar} from the same phenotype.
^b P < .05 compared with A from the same PCOS group.

Ladrón de Guevara. PCOS phenotypes in Chile and Argentina. *Fertil Steril* 2014.

phenotype D (normoandrogenic) is comparable to the control women in terms of metabolic profile (3, 23).

Previous studies in which different ethnic groups have been assessed have shown evidence of the profound impact that ethnicity may have on the metabolic parameters of the PCOS phenotypes (9, 23–28). In our study, PCOS_{Ch} women had a higher BMI in all phenotypes compared with the PCOS_{Ar} women; the latter had higher blood pressure with a higher prevalence of hypertension accordingly compared with the PCOS_{Ch} women.

In the PCOS_{Ar} group, the surrogate parameters of insulin resistance and dysglycemia improved from phenotypes A to C. In contrast, this phenomenon was not observed in the PCOS_{Ch} group, in whom these disturbances were found throughout the phenotypes. When comparing the lipid profile between both groups, we observed that the PCOS_{Ch} women showed greater alterations in LDL-C, triglycerides, and HDL-C in all phenotypes, constituting a more atherogenic profile than found in the PCOS_{Ar} women. All these findings described in the PCOS_{Ch} group could constitute a worse metabolic profile independent of the degree of hyperandrogenism.

Regarding type 2 diabetes, its prevalence was generally low, probably because the women in the groups analyzed were under 40 years old. Additionally, after patients who had previously received diagnoses of diabetes or glucose intolerance were specifically excluded from the study, only in the PCOS_{Ch} patients were cases of type 2 diabetes diagnosed through the OGTT. These early alterations in glucose metabolism are comparable with what has been described in other groups, such as the Hispanic and South Asian PCOS patients living in California (10).

When comparing the metabolic syndrome in the PCOS_{Ar} group, we found that the A phenotype exhibited a higher prevalence than the B, C, and D phenotypes. In these last three phenotypes, there was a significant decrease in the OR for the metabolic syndrome, comparable with the group of healthy Argentinian women (14%) (29). In PCOS_{Ch}, the prevalence of the metabolic syndrome in all the phenotypes was slightly higher than that described in healthy women of the same age (41.7%) (30). We found that the OR for the metabolic syndrome was independent of the phenotype evaluated. This observation differs from the findings of Wiltgen and Spritzer (31) in Brazilian women with PCOS.

The characteristics of the D phenotype in the PCOS_{Ch} group are difficult to interpret because the sample was very small (n = 3) and thus susceptible to bias. Even so, the D phenotype of both groups showed a higher BMI and greater metabolic alterations than the D phenotype of other groups who have been studied (2, 3).

To explain the findings regarding the metabolic profile of these two groups of women with PCOS, one must consider the ethnic composition of each group. As previously mentioned, the population of the city of Córdoba has been influenced by a large, mainly Italian and secondarily Spanish immigration wave (11). If we analyze the metabolic characteristics of European women, especially Italians, we can see that they show fewer alterations in glucose metabolism, the metabolic syndrome becomes apparent at older ages, and

TABLE 3

Metabolic syndrome and its components in Chilean and Argentinian women with PCOS according to the different phenotypes.

	A		B		C		D	
	SOP _{Ch} (n = 158)	SOP _{Ar} (n = 104)	SOP _{Ch} (n = 23)	SOP _{Ar} (n = 40)	SOP _{Ch} (n = 36)	SOP _{Ar} (n = 41)	SOP _{Ch} (n = 3)	SOP _{Ar} (n = 21)
MS (%) ^a	44	36.2	58 ^b	15	30	12.2	66 ^b	14.2
Fasting glucose >100 mg/dL (%)	12	19	5.2	2.5	8.1 ^b	2.4	66 ^b	9.5
2-h glucose >140 mg/dL (%)	20	19	15.7 ^b	2.5	16.2 ^b	0	66 ^b	4.7
Waist circumference >88 cm (%)	59.4	71.2	50	41.9	50	39	100 ^b	52.1
Hypertension (%) ≥ 130/85 mm Hg	8.8 ^b	25	10	10	5.4	14.6	0 ^b	14.2
HDL 50 mg/dL (%)	82.9 ^b	52.2	85 ^b	32.6	85.7 ^b	29.3	66.6	52.2
Triglycerides ≥ 150 mg/dL (%)	40	31.8	45.5 ^b	21	35.7 ^b	4.9	66.6 ^b	13

Note: Comparisons regarding phenotype D of the PCOS_{Ch} group have to be taken carefully as the size of the group is very small. HDL = high density lipoprotein; MS = metabolic syndrome; PCOS = polycystic ovary syndrome.

^a Metabolic syndrome according to ATP III (14).

^b P < .05 compared to PCOS_{Ar} from the same phenotype.

Ladrón de Guevara. PCOS phenotypes in Chile and Argentina. *Fertil Steril* 2014.

there is a higher percentage of hypertension than found in Hispanic women of comparable ages (2, 6, 19). As mentioned earlier, 65% of the Chilean population is of Spanish descent, with an average proportion of 60% Hispanic and 40% Amerindian. When we analyze the metabolic profile of Spanish women, we can see that they have a higher trend toward obesity with higher risk of the metabolic syndrome at an early age (<40 years old) and type 2 diabetes, which is similar to the metabolic profile of the PCOS_{Ch} group (2, 10).

It is important to mention that in Hispanic patients the antecedent of a first-degree-relative family member with diabetes is a better predictor for the metabolic syndrome than hyperandrogenemia (10). Regarding the Amerindian component of the PCOS_{Ch} group, the most prevalent Amerindian group in Chile are the Mapuches. Some studies have shown that when the Mapuche people migrate to the urban environment they have a higher risk for developing the metabolic syndrome (32).

During the last decades, people from Latin America have undergone a transition to a less healthy lifestyle (33). The prevalence of obesity is 25% in Chile (34) and 19.4 % in Argentina (35). According to the Food and Agriculture Organization (FAO), the food balance sheets of both countries have some differences in terms of calorie consumption. In this regard, in Argentina animal product consumption is 929 kcal/capita/day and in Chile is 717 kcal/capita/day, but consumption of products such as butter and sugar is very similar. The total calorie intake in Chile is 2,920 kcal/capita/day, and in Argentina is 3,000 kcal/capita/day. Nevertheless, the macronutrient distribution is a little different: for Chile, with 61% carbohydrates, 12% protein, and 27% fat; and for Argentina, with 55% carbohydrates, 13% protein, and 32% fat (www.fao.org). There has been no specific study comparing lifestyle in more detail between the two countries, factors that could explain part of the differences we have observed between our patients.

Ethnic diversity associated to the sociocultural habits of different countries are elements of capital importance in the analysis of the PCOS phenotypes (2, 4). Considered to be a multifactorial disorder, PCOS develops through a combination of genetic risk factors and triggering

environmental factors. The importance of knowing the genetic background resides in adopting lifestyles that may delay the appearance of the pathology and its possible complications. For this reason, it is important to know the local phenotypic characteristics of women with PCOS so that a correct therapeutic and preventive approach may be devised toward the prevalent metabolic aspects.

Acknowledgments: The authors thank Gabriel Cavada for assistance with the statistical analysis of this study.

REFERENCES

1. Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group. Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome. *Fertil Steril* 2004;81:19–25.
2. Fauser B, Tarlatzis B, Rebar R, Legro R. Consensus on women's health aspects of polycystic ovary syndrome (PCOS): the Amsterdam ESHRE/ASRM-Sponsored 3rd PCOS Consensus Workshop Group. *Fertil Steril* 2012;97:28–38.
3. Azziz R, Carmina E, Dewailly D, Diamanti-Kandarakis E, Escobar-Morreale HF, Futterweit W, et al. The Androgen Excess and PCOS Society criteria for the polycystic ovary syndrome: the complete task force report. *Fertil Steril* 2009;91:456–88.
4. Diamanti-Kandarakis E, Christakou C, Marinakis E. Phenotypes and environmental factors: their influence in PCOS. *Curr Pharm Des* 2012;18:270–82.
5. Sir-Petermann T, Maliqueo M, Angel B, Lara HE, Perez-Bravo F, Recabarren SE. Maternal serum androgens in pregnant women with polycystic ovarian syndrome: possible implications in prenatal androgenization. *Hum Reprod* 2002;17:2573–9.
6. Welt CK, Arason G, Gudmundsson JA, Adams J, Palsdottir H, Gudlaugsdottir G, et al. Defining constant versus variable phenotypic features of women with polycystic ovary syndrome using different ethnic groups and populations. *J Clin Endocrinol Metab* 2006;91:4361–8.
7. Chae SJ, Kim JJ, Choi YM, Hwang KR, Jee BC, Ku SY, et al. Clinical and biochemical characteristics of polycystic ovary syndrome in Korean women. *Hum Reprod* 2008;23:1924–31.
8. Wang Y, Qu JW, Wu X, Hou LH, Erkkola R, Wang Y. Different phenotypes of polycystic ovary syndrome by Rotterdam criteria are differently steroidogenic but similarly insulin resistant. *Fertil Steril* 2010;93:1362–5.
9. Wijayaratne CN, Seneviratne RA, Dahanayake S, Kumarapeli V, Palipane E, Kuruppu N, et al. Phenotype and metabolic profile of South Asian women with polycystic ovary syndrome (PCOS): results of a large database from a specialist endocrine clinic. *Hum Reprod* 2011;26:202–13.
10. Lo JC, Feigenbaum SL, Yang J, Pressman AR, Selby JV, Go AS. Epidemiology and adverse cardiovascular risk profile of diagnosed polycystic ovary syndrome. *J Clin Endocrinol Metab* 2006;91:1357–63.

11. Oteiza E, Novick S. *Inmigración y discriminación: políticas y discursos*. 2nd ed. Buenos Aires: Trama Editorial/Prometeo Libros; 2000.
12. Valenzuela C. Human sociogenetics. *Biol Res* 2011;44:393–404.
13. Esteva C. *La hispanización del mestizaje cultural en América*. Madrid: Quinto Centenario; 1981:182–3.
14. Hatch R, Rosenfield M, Kim MH, Tredway D. Hirsutism: implications, etiology and management. *Am J Obstet Gynecol* 1981;40:815–30.
15. American Diabetes Association. *Standards of medical care in diabetes 2013*. *Diabetes Care* 2013;36(Suppl 1):S11.
16. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 1985;28:412–9.
17. National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation* 2002;106:3143–421.
18. Morley JE, Patrick P, Perry HM 3rd. Evaluation of assays available to measure free testosterone. *Metabolism* 2002;51:554–9.
19. Carmina E, Napoli N, Longo RA, Rini GB, Lobo RA. Metabolic syndrome in polycystic ovary syndrome (PCOS): lower prevalence in southern Italy than in the USA and the influence of criteria for the diagnosis of PCOS. *Eur J Endocrinol* 2006;154:141–5.
20. Carmina E, Chu MC, Longo RA, Rini GB, Lobo RA. Phenotypic variation in hyperandrogenic women influences the findings of abnormal metabolic and cardiovascular risk parameters. *J Clin Endocrinol Metab* 2005;90:2545–9.
21. Barber TM, Wass JAH, McCarthy MI, Franks S. Metabolic characteristics of women with polycystic ovaries and oligo-amenorrhoea but normal androgen levels: implications for the management of polycystic ovary syndrome. *Clin Endocrinol* 2007;66:513–7.
22. Moran L, Teede H. Metabolic features of the reproductive phenotypes of polycystic ovary syndrome. *Hum Reprod Update* 2009;15:477–88.
23. Moghetti P, Tosi F, Bonin C, Di Sarra D, Fiers T, Kaufman JM, et al. Divergences in insulin resistance between the different phenotypes of the polycystic ovary syndrome. *J Clin Endocrinol Metab* 2013;98:628–37.
24. Chen X, Yang D, Li L, Feng S, Wang L. Abnormal glucose tolerance in Chinese women with polycystic ovary syndrome. *Hum Reprod* 2006;21:2027–32.
25. Norman R, Mahabeer S, Masters S. Ethnic differences in insulin and glucose response to glucose between white and Indian women with polycystic ovary syndrome. *Fertil Steril* 1995;63:58–62.
26. Guo M, Chen Z, Eijkemans M, Goverde A, Fauser B, Macklon N. Comparison of the phenotype of Chinese versus Dutch Caucasian women presenting with polycystic ovary syndrome and oligo/amenorrhoea. *Hum Reprod* 2012;27:1481–8.
27. Essah P, Nestler J, Carmina E. Differences in dyslipidemia between American and Italian women with polycystic ovary syndrome. *J Endocrinol Invest* 2008;31:35–41.
28. Al-Fozan H, Al-Futaisi A, Morris D, Tulandi T. Insulin responses to the oral glucose tolerance test in women of different ethnicity with polycystic ovary syndrome. *J Obstet Gynaecol Can* 2005;27:33–7.
29. Litwak L, Graffigna M, Abdala M, Akel M, Aranda C, Gutt S, et al. Prevalencia de factores de riesgo cardiovascular en sujetos presuntamente sanos. Estudio epidemiológico multicéntrico. *Rev Argent Endocrinol Metabol* 2004;41:206–13.
30. Ministerio de Salud Pública de Chile. *Encuesta Nacional de Salud, 2009–2010*. Available at: <http://epi.minsal.cl/estudios-y-encuestas-poblacionales/encuestas-poblacionales/encuesta-nacional-de-salud/resultados-ens/>. Last accessed March 12, 2014.
31. Wiltgen D, Spritzer P. Variation in metabolic and cardiovascular risk in women with different polycystic ovary syndrome phenotypes. *Fertil Steril* 2010;94:2493–6.
32. Philco PL, Serón P, Muñoz S, Navia P, Lanús F. Risk factors for metabolic syndrome in a case control study in Temuco, Chile. *Rev Med Chile* 2012;140:334–9.
33. Rivera JA, Barquera S, González-Cossío T, Olaiz G, Sepúlveda J. Nutrition transition in Mexico and other Latin American countries. *Nutr Rev* 2004;62:S149–57.
34. Ministerio de Salud (MINSAL). *II Encuesta Nacional de Salud 2009–2010*. Santiago: MINSAL; 2010.
35. Ministerio de Salud. *Presidencia de la Nación 2009, Encuesta Nacional de Nutrición y Salud*. Available at: <http://www.msal.gov.ar>. Last accessed March 12, 2014.

SUPPLEMENTAL TABLE 1

Anthropometric variables and family background in Chilean and Argentinian total PCOS groups.

	Chilean (n = 220)	Argentinian (n = 206)	P value
Age (y)	26 (18–39)	26 (18–39)	NS
BMI (kg/m ²)	29 (18.2–35)	24.7 (18–35)	< .05
Waist circumference (cm)	90 (59–126)	88 (68–120)	NS
WHR	0.86 (0.68–1.1)	0.87 (0.74–1.5)	NS
WHtR	0.56 (0.38–0.75)	0.54 (0.42–0.64)	NS
SBP (mm/Hg)	112 (100–160)	120 (100–150)	NS
DBP (mm/Hg)	70 (50–100)	80 (60–95)	< .05
Family history of diabetes	64.5%	27%	< .05
Family history of hypertension	30%	46.3%	< .05

Note: Values are median and range. Comparisons assessed with Mann-Whitney test for continuous variables and chi-square test for percentages. BMI = body mass index; DBP = diastolic blood pressure; NS = not statistically significant; PCOS = polycystic ovary syndrome; SBP = systolic blood pressure; WHR = waist to hip ratio; WHtR = waist to height ratio.

Ladrón de Guevara. PCOS phenotypes in Chile and Argentina. *Fertil Steril* 2014.

SUPPLEMENTAL TABLE 2

Percentage of metabolic syndrome and its components in the total Chilean and Argentinian PCOS groups.

	Chilean (n = 220)	Argentinian (n = 206)	P value
MS (%) ^a	52.9	36.5	< .05
Fasting glucose >100 mg/dL (%)	11.43	11.65	NS
Glucose 120 min >140 mg/dL (%)	14.35	5.83	< .05
Waist circumference >88 cm (%)	56.8	52.9	NS
Hypertension \geq 130/85(%)	11.28	15.05	NS
HDL <50 mg/dL (%)	81.08	41.26	< .05
Tg >150 mg/dL(%)	41.44	20.87	< .05

Note: Differences assessed using the chi-square test. HDL = high density lipoprotein; MS = metabolic syndrome; PCOS = polycystic ovary syndrome; Tg = triglycerides.

^a Metabolic syndrome according to the National Cholesterol Education Program Adult Treatment Panel III guidelines (ATP III) (17).

Ladrón de Guevara. PCOS phenotypes in Chile and Argentina. *Fertil Steril* 2014.