

# Cellular Mechanisms of Endoplasmic Reticulum Stress Signaling in Health and Disease. 1. An overview

Estefanie Dufey,<sup>1,2</sup> Denisse Sepúlveda,<sup>1,2</sup> Diego Rojas-Rivera,<sup>1,2</sup> and Claudio Hetz<sup>1,2,3,4</sup>

<sup>1</sup>Biomedical Neuroscience Institute, Faculty of Medicine, University of Chile, Santiago, Chile; <sup>2</sup>Program of Cellular and Molecular Biology, Institute of Biomedical Sciences, Center for Molecular Studies of the Cell, University of Chile, Santiago, Chile; <sup>3</sup>Department of Immunology and Infectious Diseases, Harvard School of Public Health, Boston, Massachusetts; and <sup>4</sup>Neurounion Biomedical Foundation, CENPAR, Santiago, Chile

Submitted 15 August 2014; accepted in final form 17 August 2014

**Dufey E, Sepúlveda D, Rojas-Rivera D, Hetz C.** Cellular Mechanisms of Endoplasmic Reticulum Stress Signaling in Health and Disease. 1. An overview. *Am J Physiol Cell Physiol* 307: C582–C594, 2014. First published August 20, 2014; doi:10.1152/ajpcell.00258.2014.—Increased demand on the protein folding capacity of the endoplasmic reticulum (ER) engages an adaptive reaction known as the unfolded protein response (UPR). The UPR regulates protein translation and the expression of numerous target genes that contribute to restore ER homeostasis or induce apoptosis of irreversibly damaged cells. UPR signaling is highly regulated and dynamic and integrates information about the type, intensity, and duration of the stress stimuli, thereby determining cell fate. Recent advances highlight novel physiological outcomes of the UPR beyond specialized secretory cells, particularly in innate immunity, metabolism, and cell differentiation. Here we discuss studies on the fine-tuning of the UPR and its physiological role in diverse organs and diseases.

ER stress; UPR; protein misfolding

MAINTAINING PROTEIN HOMEOSTASIS or proteostasis is essential for sustaining cell function. In eukaryotic cells, secreted and membrane proteins fold and mature in the lumen of the endoplasmic reticulum (ER) and Golgi apparatus. Secretory proteins are synthesized by ribosomes attached to the ER membrane and are then folded and modified by a large spectrum of chaperones and foldases in the ER lumen. Correctly folded proteins exit the ER and traffic through the secretory pathway to their final destination. This process is precisely coordinated by efficient quality control mechanisms to ensure that functionally folded proteins exit the ER (28). Misfolded glycoproteins are retained in the ER through the calnexin-calreticulin cycle and delivered to the cytosol for proteasomal-mediated degradation via the ER-associated degradation (ERAD) system (28).

Different cellular perturbations can alter ER function and lead to the abnormal accumulation of misfolded proteins. These alterations include, for example, the expression of disease-related mutant proteins, high secretory demands in endocrine and exocrine cells, viral infections that overload the ER with viral-encoded proteins, or loss of calcium homeostasis that affects calcium-dependent chaperones. ER stress activates a conserved signaling pathway to cope with protein folding alterations, that is collectively known as the unfolded protein response (UPR). The UPR transmits information about the protein folding status at the ER lumen to the cytosol and nucleus to engage adaptive responses (134). UPR signaling increases the biogenesis of the ER and other

organelles, enhances folding and quality control mechanisms at the level of gene expression, and fine-tunes protein translation. However, unresolved ER stress results in cell death. Thus, UPR stress sensors can integrate information about the duration and intensity of stress stimuli toward determining cell fate either to adapt and survive or to enter into an apoptotic program.

Accumulating evidence has implicated the UPR in important processes that seem to be independent of its traditional role in the protein folding stress response. Components of the UPR can be differentially engaged to regulate various physiological processes such as lipid metabolism, glucose homeostasis, innate immunity, and cell differentiation (136). Moreover, as demonstrated in a variety of studies in preclinical models, failure of the UPR to sustain ER proteostasis contributes to the development of several pathologies, including metabolic, neurodegenerative, and inflammatory diseases. In agreement with this concept, drug discovery efforts have recently validated ER stress as a therapeutic target to treat several disease conditions (51). Thus, it has become increasingly important to develop a precise understanding of the mechanism of signal transduction of the UPR and its impact on distinct pathologies. This article gives a global view of the signaling mechanism behind the UPR and provides a context to understanding the impact of ER stress in human disease. We review mechanistic aspects of signal transduction by specific UPR stress sensors and how the pathway integrates information in the context of the global proteostasis network to determine cell fate. Finally, we briefly discuss novel physiological outputs of the UPR in different cell types and organs and its possible involvement in the development of human diseases.

Address for reprint requests and other correspondence: C. Hetz, Institute of Biomedical Sciences, Univ. of Chile, Independencia 1027, Santiago, Chile (e-mail: chetz@med.uchile.cl).

## UPR Signaling Branches

In mammals, the UPR is mediated by at least three classes of stress sensors expressed at the ER membrane: PERK (PKR-like ER kinase), IRE1 $\alpha$  and  $\beta$  (inositol-requiring transmembrane kinase/endonucleases), and ATF6 $\alpha$  and  $\beta$  (activating transcription factor 6) (158) (Fig. 1). PERK is a type I transmembrane protein containing an NH<sub>2</sub>-terminal domain that detects luminal ER stress and a cytosolic kinase domain. PERK activation involves its dimerization, autotransphosphorylation, and further oligomerization (96). Activated PERK inhibits protein translation through inactivation by phosphorylation of the eukaryotic translation initiation factor eIF2 $\alpha$ , thus reducing protein synthesis and decreasing misfolded protein load (43). However, some mRNAs containing short open reading frames in their 5'-untranslated regions are preferentially translated when eIF2 $\alpha$  is limiting. One of them is ATF4 mRNA, a

transcription factor that positively regulates a cluster of UPR target genes involved in amino acid metabolism, antioxidant response, folding, and the regulation of apoptosis (41, 74, 77). Two important target genes driven by ATF4 are CHOP (C/EBP homologous protein) and GADD34 (growth arrest and DNA damage-inducible 34). GADD34 encodes a regulatory subunit of the phosphatase protein PP1C that counteracts PERK activity by dephosphorylating eIF2 $\alpha$  under prolonged ER stress (101). CHOP promotes transcription of *BIM* and decreases the expression of *BCL-2*, triggering apoptosis (145, 152). Attempts to define the impact of PERK signaling on gene expression in mammalian cells revealed that nearly half of PERK-dependent targets are ATF4 independent (44), suggesting the existence of other PERK downstream effectors that have not yet been explored. In a pathway that is less well understood, PERK signaling also activates by phosphorylation the transcription




Fig. 1. The unfolded protein response (UPR) signaling. The UPR is mediated by three types of ER stress sensors: IRE1, PERK, and ATF6. Activation of PERK decreases the rate of protein synthesis through phosphorylation of the eukaryotic translation initiation factor eIF2 $\alpha$ . As a consequence, eIF2 $\alpha$  phosphorylation increases the selective translation of the mRNA encoding for ATF4, a transcription factor that induces the expression of genes involved in amino acid (AA) metabolism, antioxidant responses, apoptosis, autophagy, and GADD34. GADD34 counteracts PERK activity by dephosphorylating eIF2 $\alpha$  under prolonged ER stress. Under ER stress conditions, IRE1 dimerizes and autophosphorylates, leading to the activation of its endoribonuclease activity in the cytosolic domain. Active IRE1 processes the mRNA encoding XBP1, a transcription factor that upregulates a subset of UPR target genes related to folding, quality control, and ER-associated degradation (ERAD). IRE1 also degrades certain mRNAs through regulated IRE1-dependent decay (RIDD). ATF6 encodes a transcription factor in its cytosolic domain that localizes to the ER in unstressed cells. Upon the induction of ER stress, ATF6 is processed at the Golgi apparatus, releasing its cytosolic domain which then translocates to the nucleus where it increases the expression of some ER chaperones, ERAD-related genes, and XBP1. See text for definitions of abbreviations.

factors NRF2 (nuclear factor erythroid 2-related factor) and NF- $\kappa$ B (nuclear factor kappa-light-chain-enhancer of activated B cells), which may have consequences in regulating redox metabolism and inflammatory processes, respectively (141).

IRE1 is the most conserved branch of the UPR from yeast to higher eukaryotes (63). IRE1 has two enzymatic activities in its cytosolic region, a serine/threonine kinase (18) and an endoribonuclease (RNase) (15). Under ER stress conditions, IRE1 autotransphosphorylates and homodimerizes, inducing a conformational change that activates its RNase domain (3, 11, 81, 88). IRE1 signals through a unique mechanism: it catalyzes the unconventional splicing of the mRNA encoding XBP1 (X-box binding protein 1), removing 26 nucleotide intron that changes the reading frame (15, 87, 166). Spliced XBP1 mRNA encodes a potent transcription factor called XBP1s, which modulates the expression of a subset of UPR-target genes including ER chaperones, glycosylation enzymes, ERAD components, components of the ER translocon, or involved in the synthesis of phospholipids (87, 134). Additionally, other RNAs are targets of IRE1 through a process termed regulated IRE1-dependent decay (RIDD) (39, 59). RIDD is conserved in mammals, yeast, and plants (58, 78, 110), and its substrates are defined in a cell-type-specific manner (24, 31, 108, 127). IRE1 is also involved in the degradation of certain microRNAs that have been linked to the regulation of apoptosis, cell migration, energy metabolism, and inflammation (107, 150). IRE1 can also control distinct signaling events that cross talk with other classical stress pathways, highlighting the MAP kinases. The cytosolic domain of IRE1 binds the adaptor protein TRAF2 (TNF receptor-associated factor 2), promoting the activation of ASK1 (apoptosis-signal-regulating kinase 1) and the JNK (c-Jun NH<sub>2</sub>-terminal kinase) pathway (116, 151). IRE1 also modulates other “alarm pathways” including p38, ERK (114), and NF- $\kappa$ B (60). In general, these signaling events are independent of XBP1 and could have an impact on a wide range of physiological processes ranging from apoptosis/survival, macroautophagy, proliferation, metabolism, and inflammation.

ATF6 is a type II glycoprotein that contains a single transmembrane domain. Under ER stress conditions, ATF6 is transported to the Golgi apparatus where it is processed by the site-1 and site-2 proteases (45, 120, 140). This processing event releases the cytosolic fragment of the protein ATF6f, which acts as a transcription factor and regulates the expression of a subgroup of target genes related to ERAD and protein folding at the ER, in addition to controlling the transcription of XBP1 (1, 45). Furthermore, ATF6f forms heterodimers with XBP1s (166) and thereby, can regulate specific gene expression patterns (143). Moreover, UPR transcription factors can heterodimerize with other transcription factors such as CREB, PCAF, and others (8, 90, 169), resulting in distinct effects on gene expression of a cell-type and stimuli-dependent manner (49), demonstrating the complex and dynamic transcriptional programs activated by the UPR.

### Cell Fate and ER Stress

Under ER stress, it is possible to distinguish two temporal and global phases in the UPR. In cells undergoing ER stress, both PERK and IRE1 control immediate reactions to stress before any transcriptional reaction is engaged. One of the fastest consequences triggered by PERK activation is the repression of global

protein translation through the control of the phosphorylation stage of eIF2 $\alpha$  (41). Similarly, IRE1-RNase activity degrades mRNA coding for secretory proteins that are predicted to be difficult to fold (39, 58, 59, 107). However, this concept is still evolving and there are increasing examples showing that IRE1 also targets mRNAs encoding for proteins localized in the nucleus and cytosol (107), having an immediate impact on RNA stability and as a consequence protein translation.

ER stress also attenuates the translocation of secretory and membrane proteins in a signal sequence-selective manner to reduce ER lumen protein overload, a system termed “preemptive quality control” (69), mediating the cotranslational degradation of diverse ER proteins at the cytosol (124). In addition, early stages of the ER stress response could modulate the transfer of calcium from the ER to mitochondria, which stimulate mitochondrial bioenergetics and ATP production (13, 71).

Under ER stress conditions, autophagy is also activated as a survival pathway. Autophagy is involved in many physiological processes and is essential to maintain metabolic homeostasis. Cells undergoing ER stress activate autophagy to eliminate damaged cellular components and aggregated proteins by the lysosomal pathway. Usually, autophagy is induced as a protective mechanism; however, when autophagy is overactive it can be deleterious to cellular survival (9). ER stress-dependent autophagy is mediated by the binding of IRE1 to the adaptor protein TRAF2, followed by the downstream activation of JNK that modulates Beclin 1 activity (118), an essential autophagy regulator (82). In addition, ATF4 can also induce genes involved in autophagy such as *ATG12*, *ATG5*, and *BECN1* (4). Besides, autophagy-defective cells show upregulation of essential ER chaperones (105), suggesting a close homeostatic balance between the autophagy and UPR pathways. The increase in both autophagy and mitochondrial bioenergetics contributes to the restoration of proteostasis. Together, these immediate responses represent a first barrier to cope with ER stress.

The second phase in the adaptive response controlled by the UPR involves reprogramming of gene expression in part through XBP1s, ATF4, and ATF6f. As mentioned, these factors regulate the expression of a large range of partially overlapping target genes that contribute to an increase in the folding capacity of the ER and an improvement in the efficiency of quality control and protein degradation mechanism (152). Under chronic or irreversible ER stress, apoptosis is triggered as a late event, a process dependent on the canonical intrinsic mitochondrial pathway (Fig. 2). Transcriptional and posttranscriptional mechanisms are activated to regulate proapoptotic members of the BCL-2 family that facilitate cytochrome *c* release from the mitochondria and calcium release from the ER to engage downstream apoptotic signaling events (130). The molecular events that determine how the UPR switches its signaling from an adaptive reaction to activate cell death programs are poorly understood.

Diverse UPR signals emerging from the ER converge into the mitochondrial outer membrane permeabilization (MOMP) to release apoptogenic factors (123, 168). The BCL-2 family of proteins is a group of upstream regulators of MOMP that comprises both anti- and proapoptotic components (22, 167). Antiapoptotic BCL-2 family members are characterized by the presence of four BH domains (BH1–4), and their mechanism of action is the inhibition of the conformational activation of BAX and BAK through direct or indirect mechanisms (147).




Fig. 2. ER stress-mediated apoptosis. The BCL-2 family of protein plays an essential role in the control of apoptosis under chronic ER stress. Activation of the proapoptotic BCL-2 family members BAX and BAK at the mitochondria is a key step in the induction of apoptosis, leading to the release of cytochrome *c* and activation of downstream caspases. Upstream regulators of BAX and BAK are the BH3-only proteins, another subset of proapoptotic members of the BCL-2 family. Activation of the UPR stress sensor PERK induces the transcriptional induction of the transcription factor CHOP, which downregulates the antiapoptotic protein BCL-2 and induces GADD34. In addition, the UPR controls the transcriptional upregulation of BH3-only proteins (i.e., PUMA, BIM, and NOXA) possibly through p53, CHOP, and ATF4. BIM protein levels can be regulated by phosphorylation, ubiquitination, and proteasomal degradation. The BH3-only protein BID also activates apoptosis when it is cleaved by caspase-2. In addition, PERK is required at the mitochondrial-associated ER membranes (MAMs) to modulate cytochrome *c* release and apoptosis by controlling calcium signaling and ROS production. Active IRE1 also binds TRAF2, leading to the activation of the proapoptotic kinases JNK and ASK. Thus, IRE1 degrades several mRNAs through a process known as RIDD. RIDD has a prosurvival function degrading mRNA coding for proteins with a high tendency to misfold and also has a proapoptotic activity by degrading mRNAs coding for key ER chaperones. Also, the endoribonuclease activity of IRE1 can cleave miRNAs that regulate the expression of proapoptotic proteins. At the ER membrane, TMBIM6/BI-1 inhibits Ca<sup>2+</sup> release. Mitochondrial Ca<sup>2+</sup> overload induces opening of the permeability transition pore (PTP), which leads to loss of mitochondrial inner membrane potential ( $\Delta\psi$ ), ionic unbalances, matrix swelling, and mitochondrial inner and outer membrane permeabilization (MOMP). PTP has been associated with cytochrome *c* release. See text for definitions of abbreviations.

Proapoptotic components of the family can be subdivided into “multidomain” members displaying homology in the BH1–3 domains, such as BAX, BAK, and BOK, and the “BH3-only” members, which are characterized by the presence of only one BH3 domain critical for apoptosis activation (72). Chronic ER stress leads to the transcriptional and posttranslational upregulation of proapoptotic BH3-only proteins, such as BIM, PUMA, BID, and NOXA. This engages the activation of downstream proapoptotic proteins BAX and BAK (89). The control of expression of BCL-2 members under chronic ER stress has been attributed in part to the PERK-ATF4-CHOP axis (109). Thus, CHOP represses the expression of BCL-2 and upregulates the transcription of BIM, promoting apoptosis. Interestingly, p53 is also involved in the upregulation of BH3-only proteins such as PUMA and NOXA under ER stress (92). GADD34 expression enhances reactive oxygen species (ROS) production possibly by promoting protein synthesis overloading, which may increase proteotoxicity (80). This event triggers cell death possibly due to augmented ROS production and ATP depletion (40). The BH3-only protein BID

can be posttranscriptionally activated via cleavage mediated by caspase-2 (130, 146). ER stress-dependent apoptosis is also highly controlled by another family of cell death regulators, known as the Bax-inhibitor 1 (BI-1) or TMBIM protein family (133). The TMBIM family has a well-described impact on UPR signaling and ER calcium homeostasis under ER stress. Both TMBIM6/BI-1 (20, 161) and TMBIM3/GRINA have antiapoptotic activity under chronic ER stress, but only the mRNA coding for TMBIM3/GRINA is upregulated by ER stress through the PERK-ATF4 branch (132). Another complementary mechanism involved in cell death induced by prolonged ER stress is ER calcium release (54, 146), which sensitizes mitochondria to undergo MOMP.

Prolonged activation of IRE1 has been associated with the induction of apoptosis, possibly due to the activation of ASK1/JNK (151) and by the occurrence of RIDD of certain mRNAs and miRNAs that encode ER chaperones and regulators of cell death such as caspase-2 (150). Although many distinct mechanisms have been shown to engage apoptosis by ER stress, the effects of these individual events are only partial, suggesting that the apo-

ptosis programs triggered by the UPR are highly complex and dependent on the cross talk between different signaling pathways.

### *Regulation of UPR Stress Sensors*

The first step in the induction of the UPR is the detection of abnormal levels of unfolded proteins in the ER lumen. A few models have been proposed to explain how UPR stress sensors monitor unfolded protein load. The mechanism of activation of IRE1 has been primarily studied in yeast (75). This model suggested that binding of BiP, an ER-resident chaperone, to IRE1 and PERK, or IRE1p, retains the sensors in a monomeric and inactive state, thereby preventing their oligomerization. Under ER stress conditions, BiP preferentially interacts with unfolded proteins, releasing its association with IRE1 and PERK luminal domains, enabling their spontaneous oligomerization (11, 75, 121, 134). In this model, BiP is the actual sensor because of its ability to detect misfolded proteins, whereas PERK and IRE1 operate as signal transducers. Further studies in yeast, however, have shown that IRE1p or BiP mutants that disrupt their binding do not dramatically alter the activation of the UPR (11, 75). An alternative “direct-recognition” model has emerged based on structural and biochemical analysis of yeast IRE1p. The crystal structure of the yeast IRE1 luminal domain (19) and recent biochemical evidence has shown that IRE1p directly binds to unfolded proteins, triggering IRE1p activation and oligomerization (32). In yeast, BiP dissociation and reassociation to IRE1p may help fine-tune its activation but also modulate the inactivation phase that is observed after prolonged ER stress (75).

The luminal event mediated by IRE1 $\beta$  involves direct interaction with unfolded proteins, whereas IRE1 $\alpha$  activation is mediated by association/dissociation with BiP (119). The three-dimensional structure of the mammalian IRE1 $\alpha$  luminal domain is similar to the yeast IRE1; however, the binding of IRE1 $\alpha$  to unfolded proteins is theoretically incompatible because the binding pocket is narrow in the crystal structure reported (172) and this binding has not been recapitulated *in vitro* when compared with the yeast IRE1p luminal domain (75). Besides, in contrast to the yeast UPR, mutations in IRE1 $\alpha$  that reduce its ability to bind BiP enhance the ability of this sensor to be activated, even in the absence of stress (76).

Little is known about how ATF6 responds to ER stress. ATF6 also associates with BiP under resting conditions, and this binding masks a Golgi apparatus localization signal, retaining ATF6 at the ER membrane (142). Upon ER stress, BiP is released, thus allowing the translocation of ATF6 to the Golgi apparatus where it undergoes proteolytic processing. The ATF6 luminal domain also contains intramolecular and intermolecular disulfide bonds that may serve as redox sensors to monitor the ER environment. Interestingly, one study indicated that ATF6 could be selectively activated by the overload of the ER membrane with proteins (100). It remains to be determined whether PERK or ATF6 activation also involves the direct recognition of unfolded proteins. These models require further biochemical analysis to fully understand ER stress-sensing mechanisms.

### *Fine-Tuning ER Stress Signaling: The UPRosome*

The kinetics of activation and signal attenuation between the three UPR stress sensors could differ depending on the nature and duration of the stress stimuli and the cell type analyzed,

suggesting the existence of modulatory mechanisms that fine-tune the UPR beyond the accumulation of misfolded proteins at the ER (49). Interestingly, accumulating evidence suggests that the UPR may be engaged, even, in the absence of stress through signal transduction mechanisms that affect UPR stress sensors, mediated in part by the binding of cofactors or posttranslational modifications. Thus, the mechanisms underlying the activation of the UPR may differ from yeast to mammals and may involve distinct selective modulatory events.

The integration of UPR signaling determines cell fate under ER stress. As mentioned, increasing evidence indicates that the selective modulation of specific UPR branches impacts the balance between adaptation/survival and cell death under ER stress. Although PERK and IRE1 share functionally similar luminal sensing domains (96), depending on the cell type, the signaling kinetics of these sensors is markedly different. For example, in certain conditions, IRE1 signaling is turned off by prolonged ER stress (93), whereas PERK signaling can be sustained until apoptosis occurs (94). Attenuation of IRE1 signaling under chronic ER stress is predicted to reduce the expression of XBPIs and as a consequence its pro-survival effects, whereas sustained PERK signaling favors the upregulation of proapoptotic factors. The differences between the regulation of these sensors has been explained by a mechanism that involves structural changes in their cytosolic domains and the physical association of positive and negative regulators. These regulatory factors could specifically affect their activation and modulate the intensity of downstream signaling outputs. Most of the studies on protein-protein interactions have been performed with IRE1, leading to the definition of a dynamic signaling platform that has been referred to as the “UPRosome” (52). In this section we discuss a few regulatory mechanisms that fine-tune the amplitude and kinetics of individual UPR signaling branches.

Several studies have uncovered an interesting cross talk between IRE1 and the apoptosis machinery (50). An initial discovery from our group indicated that IRE1 signaling is selectively enhanced by the expression of BAX and BAK, involving the formation of a protein complex with the cytosolic domain of IRE1 (50). Similarly, expression of the proapoptotic BH3-only proteins BIM and PUMA at the ER membrane triggers the activation of the JNK pathway in an IRE1- and BAK-dependent manner (79). These BCL-2 family members also modulate the maintenance of IRE1 signaling under prolonged ER stress (131). In this context, a subgroup of proapoptotic BCL-2 family members may have a dual function: They operate as pro-survival factors, instigating early adaptive responses to cope with ER stress via IRE1 and XBPI signaling, and also mediate downstream and late effector functions in apoptosis at the mitochondria. Several other components of the UPRosome complex have been reported (49), including the recent discovery of the nonmuscle myosin IIB, which controls the formation of the IRE1 clusters (46) and the disulfide isomerase PDIA6, which modulates the redox status of IRE1 through its luminal region (27, 37). In addition, indirect evidence suggests that XBPI mRNA splicing and RIDD activity may also be differentially modulated (107), adding an additional layer of complexity to the regulation of IRE1.

As mentioned, under constant ER stress, IRE1 is turned off. Inactivation of IRE1 involves the dissolution of IRE1 clusters and dephosphorylation of IRE1, leading to a decline of XBPI

mRNA splicing (91). Bax-inhibitor 1 (BI-1) is a central mediator of IRE1 attenuation as demonstrated in different studies in cell culture and animal models of ER stress (6, 7, 95, 135). In other experimental systems the opposite behavior has been observed, where XBP1 mRNA splicing is sustained over time while eIF2 $\alpha$  phosphorylation is attenuated through a feedback loop that induces its phosphatase (14, 98, 117). Thus, the UPR is fine-tuned in a dynamic manner through the assembling of distinct factors to the UPRosome, which potentially establishes a stress threshold to engage the UPR.

In addition to the negative regulation of eIF2 $\alpha$  phosphorylation by GADD34 (117), PERK can also be modulated by other components by a physical interaction. For example, under ER stress conditions, p58 IPK expression is upregulated and binds to PERK, reducing its kinase activity (156, 165). Recent evidence has also shown that mitofusin 2 interacts with PERK and that deficiency of mitofusin 2 enhances PERK phosphorylation, impacting apoptosis and autophagy (113). Other components also selectively modulate PERK, in particular Nck1 (164), a splicing variant of BiP (115) and the calcium-dependent phosphatase calcineurin (12). Although less well studied, ATF6f is also modulated through interactions with other factors. The UPR target gene *WFS1* represses ATF6 signaling possibly by inducing its proteasome-dependent degradation (29). ATF6 is also modulated by redox changes of its luminal cysteine and also by glycosylation, where the protein disulfide isomerase PDIA5 modulates ATF6 activation (57).

Systematic studies are needed to define the interactome of UPR stress sensors and to understand how their composition is modulated by ER stress in different contexts. We predict that the identification of distinct UPRosomes will emerge from these studies, where their identity may depend on the cell type and stimuli analyzed. All these examples demonstrate the highly regulated and dynamic nature of the UPR. Assembling of specific UPRosomes may allow the integration of information regarding the type and intensity of the stress stimulus toward reinforcing specific outputs of the UPR according to the cell's needs.

### *UPR in Physiology and Disease*

UPR activation is observed in many physiological processes beyond the homeostatic control of protein folding. Components of the UPR play a central role in the normal development and differentiation of specialized secretory cells, including B cells, pancreatic  $\beta$ -cells and salivary glands. In addition, recent studies have associated the UPR with the control of innate immunity and in the control of energy metabolism and the synthesis of cholesterol and lipids. Furthermore, abnormal ER stress levels are involved in several diseases including diabetes mellitus, neurodegeneration, cancer, and other pathologies (Fig. 3). In this section, we describe some examples that highlight the impact of the UPR to the development of disease and its participation in the homeostatic control of diverse tissues (Table 1).

During the differentiation of B cells to antibody-secreting plasma cells, XBP1s is required to induce cell differentiation in addition to enhancing the folding capacity of the cell (65, 129). Professional secretory immune cells lacking XBP1 display severe abnormalities in their development and function as initially reported in plasma B cells (65) and dendritic cells (66). ATF6 is also activated during plasma cell differentiation (33),

while PERK remains inactive (34). In contrast, pancreatic  $\beta$ -cells require PERK signaling to maintain endocrine function. In fact, genetic targeting of PERK/eIF2 $\alpha$  signaling leads to  $\beta$ -cell deficiency, altering insulin production and glucose metabolism, which results in early-onset diabetes mellitus (42, 139, 171). Similarly, developmental ablation of XBP1 triggers a collapse in the exocrine pancreas and salivary glands, whereas endocrine pancreas in adults is also altered, leading to hypoglycemia (50, 84, 85). XBP1 conditional deletion in pancreatic  $\beta$ -cells causes hyperglycemia, glucose intolerance, and markedly decreases the number of insulin granules, in addition to impaired proinsulin processing (2, 85). Interestingly, insulin mRNA is a RIDD target and its expression is also controlled by XBP1 (39). XBP1-deficient animals present hypoplastic fetal livers, with reduced hematopoiesis triggering early death during development due to anemia (128). Postnatal XBP1 liver-specific knockout mice do not show any spontaneous abnormalities, with no evidence of liver damage; however, a dramatic alteration in lipid synthesis is observed (86). In the liver, IRE1 $\alpha$  controls lipogenesis and lipoprotein metabolism through the RIDD pathway (144).

UPR components have a crucial role in bone development and osteoblast function. PERK-deficient animals develop drastic osteopenia involving a reduction of trabecular bone thickness and volume (138, 159, 171). Interestingly, targeting XBP1 in intestinal epithelial cells triggers spontaneous enteritis, augmenting the susceptibility to colitis and chronic inflammation (70). Moreover, a polymorphism in the *XBP1* gene was found in patients with Crohn's disease (35). In addition, emerging roles of the UPR in innate immunity have been described, involving activities in macrophage and dendritic cell differentiation and function (103, 122). XBP1 is induced downstream of Toll-like receptors, modulating the production of proinflammatory cytokines in macrophages (103). Genetic manipulation of key UPR components has also demonstrated a crucial role of the pathway in gastric zymogenic cell differentiation (61) and intestinal epithelial cell differentiation (47), among many other relevant physiological roles (reviewed in Ref. 17).

In relation to the phenotypes observed in UPR target mouse models, PERK knockout animals are viable, allowing the early characterization of its function in vivo in the field (42). IRE1 $\alpha$  null mice develop dramatic defects during development, leading to lethality at 12.5 days of gestation (170). These mice show a decrease of vascular endothelial growth factor and abnormalities in the blood vessels of the placenta (27). Remarkably, there is a rescue of the embryonic lethality in conditional knockout mice when IRE1 $\alpha$  is reconstituted in the placenta (68). Surprisingly, these animals did not develop hypoplasia in the liver (68) as described for XBP1-deficient mice (68) and, in contrast to the phenotypes described for XBP1-deficient animals, IRE1 $\alpha$  deletion caused only mild abnormalities of exocrine tissues and only slightly altered blood glucose level and serum immunoglobulin levels (67). One possible explanation is that the deletion of XBP1 produces the overactivation of IRE1 $\alpha$  activity that decreases some RIDD target genes, compensating for XBP1 loss-of-function (62). In addition, neither ATF6 $\alpha$  nor ATF6 $\beta$  is essential for embryonic development and single knockout animals do not show evident growth defects and develop normally into adulthood (160, 162). Importantly, ATF6 $\alpha$  and ATF6 $\beta$  double deficiency results in full embryonic lethality, suggesting that these proteins

have functional redundancy, with essential functions during mouse development (162). However, ATF6 $\alpha$  knockout animals are hypersensitive to experimental ER stress as revealed by the unexpected lethality observed after the intraperitoneal injection of the ER stress agent tunicamycin, associated with the generation of acute liver and kidney damage (160, 162). ATF6 $\alpha$  plays an important role in glucose and lipid homeostasis under both pharmacologically induced and physiological ER stress. Moreover, ATF6 $\alpha$ -deficient mice showed augmented hepatic triacylglycerol levels, fat deposits, and increased lipid droplets, which are associated with liver microvesicular steatosis (137, 163). In agreement with this concept, this abnormal hepatic phenotype is observed during the genetic ablation of any of the three UPR branches (137).

The physiological role of the UPR in many organs translates into the development of several diseases when the ER proteostasis is irreversibly damaged. For example, increased levels of ER stress are observed in models of obesity and diabetes, leading to inflammatory reactions and insulin resistance (30). Suppression of insulin signaling pathways occurs through IRE1-dependent hyperactivation of JNK and subsequent phosphorylation of insulin receptor substrate on serine residues (125). Wolcott-Rallison syndrome (childhood diabetes) mutations in PERK have been associated with abnormal function of

pancreatic islets (25), consistent with the phenotypes described in *perk*-deficient animals (43).

One of the major areas of therapeutic development in the UPR field is cancer. Early studies demonstrated that the UPR is essential for the survival and growth of tumor cells into solid tumors, providing an adaptive capacity to the adverse microenvironmental conditions generated by hypoxic conditions (23, 99). In addition, accumulating evidence indicates that the UPR contributes to tumor growth, metastasis, and angiogenesis (5, 16, 26, 111, 153). Importantly, genomic screening of cancer cells has revealed that IRE1 is one of the kinases most frequently mutated in cancer (36, 38, 126). More recently, XBP1-dependent gene signatures were proposed as a predictor of the aggressiveness of different types of cancer (21, 73). Many different drug screening efforts have identified small molecules that selectively inhibit PERK or the RNase activity of IRE1, exhibiting potent antitumor activity in a variety of cancer models in vivo (51).

Another area of active research in the UPR field is neuroscience. Conditional deletion of XBP1 in the central nervous system has been performed to test its contribution to neurodegeneration in many disease models (55). In addition, deletion of PERK in the adult forebrain resulted in reduced eIF2 $\alpha$  phosphorylation and ATF4 expression, leading to altered be-




Fig. 3. Role of the UPR in physiology and diseases. Genetic manipulation of major UPR components has revealed its relevance in the function of diverse organs and cell types, in addition to its contribution to a variety of diseases using preclinical mouse models. Important functions have been reported in brain, bone marrow, heart, liver, pancreas, intestine, and gastric system (blue boxes). Pathologies where abnormal ER stress levels play a relevant role in disease include diabetes, neurodegeneration, ischemia, cancer, and other diseases (red boxes).

Table 1. *UPR mouse models and associated phenotypes*

| UPR Member                  | Animal Model                                | Phenotype  | Reference          |
|-----------------------------|---|--|--------------------|
| IRE1 $\alpha$               | Full knockout                               | Embryonic lethality after 12.5–13 days of gestation.   | (68, 170)          |
| IRE1 $\alpha$               | Conditional embryo-specific knockout        | Hyperglycemia, mild hypoinsulinemia. Abnormal histological structure of the acinar pancreas and salivary serous tissues.                     | (67)               |
| IRE1 $\beta$                | Full knockout                               | Increased susceptibility to experimental colitis.  | (10, 64, 102, 149) |
| XBP1                        | Full knockout                               | Hypoplastic fetal liver, reduced hematopoiesis, and embryonic death from anemia at 12.5 days of gestation.                                   | (128)              |
| XBP1                        | Adult liver-specific knockout               | No gross liver abnormalities. Reduced plasma levels of cholesterol and triglycerides due to decreased hepatic lipid synthesis and secretion. | (86)               |
| XBP1                        | Conditional $\beta$ -cell-specific knockout | Mild hyperglycemia and glucose intolerance due to impaired proinsulin processing and reduced insulin secretion.                              | (85)               |
| PERK                        | Full knockout                               | Hyperglycemia, impaired synthesis of insulin, and digestive enzymes. $\beta$ -Cell loss. Growth retardation and skeletal dysplasia.          | (42, 171)          |
| PERK                        | Brain-specific knockout                     | Cognitive deficits in information processing.  | (148)              |
| ATF4                        | Full knockout                               | Defective eye lens development. Severe anemia due to impaired fetal-liver hematopoiesis.   | (48, 83, 104)      |
| ATF6 $\alpha$               | Full knockout                               | Hypoglycemia, insulin resistance, and liver steatosis in response to pharmacologically induced ER stress.                                    | (154, 163)         |
| ATF6 $\beta$                | Full knockout                               | No obvious phenotype.  | (162)              |
| ATF6 $\alpha$ and - $\beta$ | Double knockout                             | Embryonic lethality.   | (162)              |

UPR, unfolded protein response; IRE1, inositol-requiring transmembrane kinase/endonuclease 1; ER, endoplasmic reticulum; PERK, PKR-like ER kinase; ATF4, activating transcription factor 4.

havior in several tests (148). Interestingly, disruption of PERK improves learning and memory in an Alzheimer's disease mouse model (97). Although the role of the UPR in the physiology of the nervous system remains poorly understood, the occurrence of chronic ER stress is widely described in most models of neurodegeneration and human postmortem studies (55). ER stress is found in the brain of patients affected with diseases such as Alzheimer's, Parkinson's, Huntington's, amyotrophic lateral sclerosis (ALS), and prion-related disorders. These diseases have different clinical manifestations; they all involve the accumulation of protein aggregates containing misfolded proteins. Initially, it was proposed that UPR activation in neurodegenerative diseases represents a pathogenic event; however, the scenario is much more complex because genetic manipulation of distinct UPR signaling components in preclinical models of neurodegeneration have revealed contrasting and sometimes opposing results (55). Activation of the UPR may enhance or prevent neurodegeneration in different diseases, and in some cases may have opposite effects depending on the signaling branch activated. For example, we have reported that targeting XBP1 in ALS (56) or Huntington's disease (157) provides protection through upregulation of autophagy, whereas in spinal cord injury it reduces motor recovery (155). In prion diseases, XBP1 deficiency did not have any effects (53). In contrast, targeting PERK/ATF4 signaling in ALS (106) or prion diseases (112) protects against degeneration; it has no effects in Huntington's disease (157) and exacerbates the effects of spinal cord injury (155). The relevance of the UPR in brain disorders seems to be specific to distinct diseases, therefore, a systematic study of the functional significance of ER stress is needed to understand the contribution of UPR in neurodegeneration. In summary, overall, the UPR is emerging as a relevant homeostatic network that could contribute to the pathogenesis of many common human diseases.

### Perspectives

The adaptive capacity to protein folding stress is fundamental to sustain the physiological function of specialized secretory

cells. The UPR constitutes an extremely dynamic and complex network of signals, whose impact on cell fate depends heavily on a larger cellular context. Mechanistic studies have revealed that the effector consequences of UPR signaling (outputs) are diverse and have effects beyond its classical role as an adjustor of the protein folding status. In addition, understanding the way in which the UPR is fine-tuned is acquiring more relevance because of its fundamental impact to understanding how cells transit from an adaptive to an apoptotic phase. We believe that defining the UPR stress sensor interactome is needed to uncover the composition of the UPRosome and how the pathway is connected to other signaling events.

The UPR has been involved in several diseases with a high prevalence and mortality, and studies continue to identify novel connections between ER stress and disease. Modulating the activity of the UPR on a disease context has been validated using pharmacology and gene therapy approaches in a variety of preclinical models (51). Owing to the impact of ER stress signaling on the sustenance of various organ functions, it is difficult to predict the side effects of targeting the pathway with chronic administration of UPR-targeting small molecules. Future clinical trials will unveil the true potential of this pathway as a therapeutic target. Applications in the area of biomarkers for diagnosis and prediction of disease prognosis are also emerging in the field which we believe will represent an enormous advance to monitor clinical trials in the future as a mirror of the health of the proteome.

### ACKNOWLEDGMENTS

We thank Andrew Foley for critical feedback on this review.

### GRANTS

This work was funded by Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT) no. 1140549, Millennium Institute no. P09-015-F, Ring Initiative ACT1109, Science and Technology Development Fund (FONDEF) Grant D1111007, Comisión Nacional de Investigación Científica y Tecnológica (CONICYT) Grant USA2013-0003, ECOS-CONICYT C13S02, The Michael J. Fox Foundation for Parkinson Research, ALS Therapy Alliance, Muscular Dystrophy Association, Alzheimer's Disease Association, and Foundation COPEC-UC (C. Hetz), FONDECYT 3130365 (D. Rojas-Rivera), and CONICYT PhD fellowships (E. Dufey, D. Sepúlveda).



## DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the author(s).

## AUTHOR CONTRIBUTIONS

E.D. conception and design of research; E.D., D.S., and D.R.-R. prepared figures; E.D., D.S., D.R.-R., and C.H. drafted manuscript; E.D., D.S., D.R.-R., and C.H. edited and revised manuscript; E.D. and C.H. approved final version of manuscript.

## REFERENCES

- Adachi Y, Yamamoto K, Okada T, Yoshida H, Harada A, Mori K. ATF6 is a transcription factor specializing in the regulation of quality control proteins in the endoplasmic reticulum. *Cell Struct Funct* 33: 75–89, 2008.
- Akiyama M, Liew CW, Lu S, Hu J, Martinez R, Hambro B, Kennedy RT, Kulkarni RN. X-box binding protein 1 is essential for insulin regulation of pancreatic alpha-cell function. *Diabetes* 62: 2439–2449, 2013.
- Ali MM, Bagratuni T, Davenport EL, Nowak PR, Silva-Santisteban MC, Hardcastle A, McAndrews C, Rowlands MG, Morgan GJ, Aherne W, Collins I, Davies FE, Pearl LH. Structure of the Ire1 autophosphorylation complex and implications for the unfolded protein response. *EMBO J* 30: 894–905, 2011.
- Ameri K, Harris AL. Activating transcription factor 4. *Int J Biochem Cell Biol* 40: 14–21, 2008.
- Auf G, Jabouille A, Guerit S, Pineau R, Delugin M, Boucheccareilh M, Magnin N, Favereaux A, Maitre M, Gaiser T, von Deimling A, Czabanka M, Vajkoczy P, Chevet E, Bikfalvi A, Moenner M. Inositol-requiring enzyme 1alpha is a key regulator of angiogenesis and invasion in malignant glioma. *Proc Natl Acad Sci USA* 107: 15553–15558, 2010.
- Bailly-Maitre B, Belgardt BF, Jordan SD, Coornaert B, von Freyend MJ, Kleinriders A, Mauer J, Cuddy M, Kress CL, Willmes D, Essig M, Hampel B, Protzer U, Reed JC, Bruning JC. Hepatic Bax inhibitor-1 inhibits IRE1alpha and protects from obesity-associated insulin resistance and glucose intolerance. *J Biol Chem* 285: 6198–6207, 2010.
- Bailly-Maitre B, Fondevila C, Kaldas F, Droin N, Luciano F, Ricci JE, Croxton R, Krajewska M, Zapata JM, Kupiec-Weglinski JW, Farmer D, Reed JC. Cytoprotective gene bi-1 is required for intrinsic protection from endoplasmic reticulum stress and ischemia-reperfusion injury. *Proc Natl Acad Sci USA* 103: 2809–2814, 2006.
- Baumeister P, Luo S, Skarnes WC, Sui G, Seto E, Shi Y, Lee AS. Endoplasmic reticulum stress induction of the Grp78/BiP promoter: activating mechanisms mediated by YY1 and its interactive chromatin modifiers. *Mol Cell Biol* 25: 4529–4540, 2005.
- Bernales S, Schuck S, Walter P. ER-phagy: selective autophagy of the endoplasmic reticulum. *Autophagy* 3: 285–287, 2007.
- Bertolotti A, Wang X, Novoa I, Jungreis R, Schlessinger K, Cho JH, West AB, Ron D. Increased sensitivity to dextran sodium sulfate colitis in IRE1beta-deficient mice. *J Clin Invest* 107: 585–593, 2001.
- Bertolotti A, Zhang Y, Hendershot LM, Harding HP, Ron D. Dynamic interaction of BiP and ER stress transducers in the unfolded-protein response. *Nat Cell Biol* 2: 326–332, 2000.
- Bollo M, Paredes RM, Holstein D, Zhelezova N, Camacho P, Lechleiter JD. Calcineurin interacts with PERK and dephosphorylates calnexin to relieve ER stress in mammals and frogs. *PLoS One* 5: e11925, 2010.
- Bravo R, Gutierrez T, Paredes F, Gatica D, Rodriguez AE, Pedrozo Z, Chiong M, Parra V, Quest AF, Rothermel BA, Lavandro S. Endoplasmic reticulum: ER stress regulates mitochondrial bioenergetics. *Int J Biochem Cell Biol* 44: 16–20, 2012.
- Brush MH, Weiser DC, Shenolikar S. Growth arrest and DNA damage-inducible protein GADD34 targets protein phosphatase 1 alpha to the endoplasmic reticulum and promotes dephosphorylation of the alpha subunit of eukaryotic translation initiation factor 2. *Mol Cell Biol* 23: 1292–1303, 2003.
- Calfon M, Zeng H, Urano F, Till JH, Hubbard SR, Harding HP, Clark SG, Ron D. IRE1 couples endoplasmic reticulum load to secretory capacity by processing the XBP-1 mRNA. *Nature* 415: 92–96, 2002.
- Clarke HJ, Chambers JE, Liniker E, Marciniak SJ. Endoplasmic reticulum stress in malignancy. *Cancer Cell* 25: 563–573, 2014.
- Cornejo VH, Pihan P, Vidal RL, Hetz C. Role of the unfolded protein response in organ physiology: lessons from mouse models. *IUBMB Life* 65: 962–975, 2013.
- Cox JS, Shamu CE, Walter P. Transcriptional induction of genes encoding endoplasmic reticulum resident proteins requires a transmembrane protein kinase. *Cell* 73: 1197–1206, 1993.
- Credle JJ, Finer-Moore JS, Papa FR, Stroud RM, Walter P. On the mechanism of sensing unfolded protein in the endoplasmic reticulum. *Proc Natl Acad Sci USA* 102: 18773–18784, 2005.
- Chae HJ, Kim HR, Xu C, Bailly-Maitre B, Krajewska M, Krajewski S, Banares S, Cui J, Digicaylioglu M, Ke N, Kitada S, Monosov E, Thomas M, Kress CL, Babendure JR, Tsien RY, Lipton SA, Reed JC. Bi-1 regulates an apoptosis pathway linked to endoplasmic reticulum stress. *Mol Cell* 15: 355–366, 2004.
- Chen X, Iliopoulos D, Zhang Q, Tang Q, Greenblatt MB, Hatziaepostolou M, Lim E, Tam WL, Ni M, Chen Y, Mai J, Shen H, Hu DZ, Adoro S, Hu B, Song M, Tan C, Landis MD, Ferrari M, Shin SJ, Brown M, Chang JC, Liu XS, Glimcher LH. XBP1 promotes triple-negative breast cancer by controlling the HIF1alpha pathway. *Nature* 508: 103–107, 2014.
- Danial NN, Korsmeyer SJ. Cell death: critical control points. *Cell* 116: 205–219, 2004.
- Dejeans N, Manie S, Hetz C, Bard F, Hupp T, Agostinis P, Samali A, Chevet E. Addicted to secrete - novel concepts and targets in cancer therapy. *Trends Mol Med* 20: 242–250, 2014.
- Dejeans N, Pluquet O, Lhomond S, Grise F, Boucheccareilh M, Juin A, Meynard-Cadars M, Bidaud-Meynard A, Gentil C, Moreau V, Saltel F, Chevet E. Autocrine control of glioma cells adhesion and migration through IRE1alpha-mediated cleavage of SPARC mRNA. *J Cell Sci* 125: 4278–4287, 2012.
- Delepine M, Nicolino M, Barrett T, Golamaully M, Lathrop GM, Julier C. EIF2AK3, encoding translation initiation factor 2-alpha kinase 3, is mutated in patients with Wolcott-Rallison syndrome. *Nat Genet* 25: 406–409, 2000.
- Drogat B, Auguste P, Nguyen DT, Boucheccareilh M, Pineau R, Nalbantoglu J, Kaufman RJ, Chevet E, Bikfalvi A, Moenner M. IRE1 signaling is essential for ischemia-induced vascular endothelial growth factor-A expression and contributes to angiogenesis and tumor growth in vivo. *Cancer Res* 67: 6700–6707, 2007.
- Eletto D, Dersh D, Gidalevitz T, Argon Y. Protein disulfide isomerase A6 controls the decay of IRE1alpha signaling via disulfide-dependent association. *Mol Cell* 53: 562–576, 2014.
- Ellegaard L, Helenius A. Quality control in the endoplasmic reticulum. *Nat Rev Mol Cell Biol* 4: 181–191, 2003.
- Fonseca SG, Ishigaki S, Osowski CM, Lu S, Lipson KL, Ghosh R, Hayashi E, Ishihara H, Oka Y, Permutt MA, Urano F. Wolfram syndrome 1 gene negatively regulates ER stress signaling in rodent and human cells. *J Clin Invest* 120: 744–755, 2010.
- Fu S, Watkins SM, Hotamisligil GS. The role of endoplasmic reticulum in hepatic lipid homeostasis and stress signaling. *Cell Metab* 15: 623–634, 2012.
- Gaddam D, Stevens N, Hollien J. Comparison of mRNA localization and regulation during endoplasmic reticulum stress in *Drosophila* cells. *Mol Biol Cell* 24: 14–20, 2013.
- Gardner BM, Walter P. Unfolded proteins are Ire1-activating ligands that directly induce the unfolded protein response. *Science* 333: 1891–1894, 2011.
- Gass JN, Gifford NM, Brewer JW. Activation of an unfolded protein response during differentiation of antibody-secreting B cells. *J Biol Chem* 277: 49047–49054, 2002.
- Gass JN, Jiang HY, Wek RC, Brewer JW. The unfolded protein response of B-lymphocytes: PERK-independent development of antibody-secreting cells. *Mol Immunol* 45: 1035–1043, 2008.
- Glas J, Seiderer J, Czamara D, Pasciuto G, Diegelmann J, Wetzke M, Olszak T, Wolf C, Muller-Myhsok B, Balschun T, Achkar JP, Kamboh MI, Franke A, Duerr RH, Brand S. PTGER4 expression-modulating polymorphisms in the 5p13.1 region predispose to Crohn's disease and affect NF-kappaB and XBP1 binding sites. *PLoS One* 7: e52873, 2012.
- Greenman C, Stephens P, Smith R, Dalgleish GL, Hunter C, Bignell G, Davies H, Teague J, Butler A, Stevens C, Edkins S, O'Meara S, Vastrik I, Schmidt EE, Avis T, Barthorpe S, Bhamra G, Buck G, Choudhury B, Clements J, Cole J, Dicks E, Forbes S, Gray K, Halliday K, Harrison R, Hills K, Hinton J, Jenkinson A, Jones D, Menzies A, Mironenko T, Perry J, Raine K, Richardson D, Shepherd

- R, Small A, Tofts C, Varian J, Webb T, West S, Widaa S, Yates A, Cahill DP, Louis DN, Goldstraw P, Nicholson AG, Brasseur F, Looijenga L, Weber BL, Chiew YE, DeFazio A, Greaves MF, Green AR, Campbell P, Birney E, Easton DF, Chenevix-Trench G, Tan MH, Khoo SK, Teh BT, Yuen ST, Leung SY, Wooster R, Futreal PA, Stratton MR. Patterns of somatic mutation in human cancer genomes. *Nature* 446: 153–158, 2007.
37. Groenendyk J, Peng Z, Dudek E, Fan X, Mizianty MJ, Dufey E, Urria H, Sepulveda D, Rojas-Rivera D, Lim Y, Kim do H, Baretta K, Srikanth S, Gwack Y, Ahnn J, Kaufman RJ, Lee SK, Hetz C, Kurgan L, Michalak M. Interplay Between the oxidoreductase PDIA6 and microRNA-322 controls the response to disrupted endoplasmic reticulum calcium homeostasis. *Sci Signal* 7: ra54, 2014.
  38. Guichard C, Amaddeo G, Imbeaud S, Ladeiro Y, Pelletier L, Maad IB, Calderaro J, Bioulac-Sage P, Letexier M, Degos F, Clement B, Balabaud C, Chevet E, Laurent A, Couchy G, Letouze E, Calvo F, Zucman-Rossi J. Integrated analysis of somatic mutations and focal copy-number changes identifies key genes and pathways in hepatocellular carcinoma. *Nat Genet* 44: 694–698, 2012.
  39. Han D, Lerner AG, Vande Walle L, Upton JP, Xu W, Hagen A, Backes BJ, Oakes SA, Papa FR. IRE1alpha kinase activation modes control alternate endoribonuclease outputs to determine divergent cell fates. *Cell* 138: 562–575, 2009.
  40. Han J, Back SH, Hur J, Lin YH, Gildersleeve R, Shan J, Yuan CL, Krokowski D, Wang S, Hatzoglou M, Kilberg MS, Sartor MA, Kaufman RJ. ER-stress-induced transcriptional regulation increases protein synthesis leading to cell death. *Nat Cell Biol* 15: 481–490, 2013.
  41. Harding HP, Novoa I, Zhang Y, Zeng H, Wek R, Schapira M, Ron D. Regulated translation initiation controls stress-induced gene expression in mammalian cells. *Mol Cell* 6: 1099–1108, 2000.
  42. Harding HP, Zeng H, Zhang Y, Jungries R, Chung P, Plesken H, Sabatini DD, Ron D. Diabetes mellitus and exocrine pancreatic dysfunction in *perk<sup>-/-</sup>* mice reveals a role for translational control in secretory cell survival. *Mol Cell* 7: 1153–1163, 2001.
  43. Harding HP, Zhang Y, Bertolotti A, Zeng H, Ron D. Perk is essential for translational regulation and cell survival during the unfolded protein response. *Mol Cell* 5: 897–904, 2000.
  44. Harding HP, Zhang Y, Zeng H, Novoa I, Lu PD, Calfon M, Sadri N, Yun C, Popko B, Paules R, Stojdl DF, Bell JC, Hettmann T, Leiden JM, Ron D. An integrated stress response regulates amino acid metabolism and resistance to oxidative stress. *Mol Cell* 11: 619–633, 2003.
  45. Haze K, Yoshida H, Yanagi H, Yura T, Mori K. Mammalian transcription factor ATF6 is synthesized as a transmembrane protein and activated by proteolysis in response to endoplasmic reticulum stress. *Mol Biol Cell* 10: 3787–3799, 1999.
  46. He Y, Beatty A, Han X, Ji Y, Ma X, Adelstein RS, Yates JR 3rd, Kempthues K, Qi L. Nonmuscle myosin IIB links cytoskeleton to IRE1alpha signaling during ER stress. *Dev Cell* 23: 1141–1152, 2012.
  47. Heijmans J, van Lidth de Jeude JF, Koo BK, Rosekrans SL, Wieleenga MC, van de Wetering M, Ferrante M, Lee AS, Onderwater JJ, Paton JC, Paton AW, Mommaas AM, Kodach LL, Hardwick JC, Hommes DW, Clevers H, Muncan V, van den Brink GR. ER stress causes rapid loss of intestinal epithelial stemness through activation of the unfolded protein response. *Cell Rep* 3: 1128–1139, 2013.
  48. Hettmann T, Barton K, Leiden JM. Microphthalmia due to p53-mediated apoptosis of anterior lens epithelial cells in mice lacking the CREB-2 transcription factor. *Dev Biol* 222: 110–123, 2000.
  49. Hetz C. The unfolded protein response: controlling cell fate decisions under ER stress and beyond. *Nat Rev Mol Cell Biol* 13: 89–102, 2012.
  50. Hetz C, Bernasconi P, Fisher J, Lee AH, Bassik MC, Antonsson B, Brandt GS, Iwakoshi NN, Schinzel A, Glimcher LH, Korsmeyer SJ. Proapoptotic BAX and BAK modulate the unfolded protein response by a direct interaction with IRE1alpha. *Science* 312: 572–576, 2006.
  51. Hetz C, Chevet E, Harding HP. Targeting the unfolded protein response in disease. *Nat Rev Drug Discov* 12: 703–719, 2013.
  52. Hetz C, Glimcher LH. Fine-tuning of the unfolded protein response: assembling the IRE1alpha interactome. *Mol Cell* 35: 551–561, 2009.
  53. Hetz C, Lee AH, Gonzalez-Romero D, Thielen P, Castilla J, Soto C, Glimcher LH. Unfolded protein response transcription factor XBP-1 does not influence prion replication or pathogenesis. *Proc Natl Acad Sci USA* 105: 757–762, 2008.
  54. Hetz C, Martinon F, Rodriguez D, Glimcher LH. The unfolded protein response: integrating stress signals through the stress sensor IRE1alpha. *Physiol Rev* 91: 1219–1243, 2011.
  55. Hetz C, Mollereau B. Disturbance of endoplasmic reticulum proteostasis in neurodegenerative diseases. *Nat Rev Neurosci* 15: 233–249, 2014.
  56. Hetz C, Thielen P, Matus S, Nassif M, Court F, Kiffin R, Martinez G, Cuervo AM, Brown RH, Glimcher LH. XBP-1 deficiency in the nervous system protects against amyotrophic lateral sclerosis by increasing autophagy. *Genes Dev* 23: 2294–2306, 2009.
  57. Higa A, Taouji S, Lhomond S, Jensen D, Fernandez-Zapico ME, Simpson JC, Pasquet JM, Schekman R, Chevet E. Endoplasmic reticulum stress-activated transcription factor ATF6alpha requires the disulfide isomerase PDIA5 to modulate chemoresistance. *Mol Cell Biol* 34: 1839–1849, 2014.
  58. Hollien J, Lin JH, Li H, Stevens N, Walter P, Weissman JS. Regulated Ire1-dependent decay of messenger RNAs in mammalian cells. *J Cell Biol* 186: 323–331, 2009.
  59. Hollien J, Weissman JS. Decay of endoplasmic reticulum-localized mRNAs during the unfolded protein response. *Science* 313: 104–107, 2006.
  60. Hu P, Han Z, Couvillon AD, Kaufman RJ, Exton JH. Autocrine tumor necrosis factor alpha links endoplasmic reticulum stress to the membrane death receptor pathway through IRE1alpha-mediated NF-kappaB activation and down-regulation of TRAF2 expression. *Mol Cell Biol* 26: 3071–3084, 2006.
  61. Huh WJ, Esen E, Geahlen JH, Bredemeyer AJ, Lee AH, Shi G, Konecny SF, Glimcher LH, Mills JC. XBP1 controls maturation of gastric zymogenic cells by induction of MIST1 and expansion of the rough endoplasmic reticulum. *Gastroenterology* 139: 2038–2049, 2010.
  62. Hur KY, So JS, Ruda V, Frank-Kamenetsky M, Fitzgerald K, Kotliansky V, Iwakaki T, Glimcher LH, Lee AH. IRE1alpha activation protects mice against acetaminophen-induced hepatotoxicity. *J Exp Med* 209: 307–318, 2012.
  63. Imagawa Y, Hosoda A, Sasaka S, Tsuru A, Kohno K. RNase domains determine the functional difference between IRE1alpha and IRE1beta. *FEBS Lett* 582: 656–660, 2008.
  64. Iqbal J, Dai K, Seimon T, Jungreis R, Oyadomari M, Kuriakose G, Ron D, Tabas I, Hussain MM. IRE1beta inhibits chylomicron production by selectively degrading MTP mRNA. *Cell Metab* 7: 445–455, 2008.
  65. Iwakoshi NN, Lee AH, Glimcher LH. The X-box binding protein-1 transcription factor is required for plasma cell differentiation and the unfolded protein response. *Immunity* 19: 29–38, 2003.
  66. Iwakoshi NN, Pypaert M, Glimcher LH. The transcription factor XBP-1 is essential for the development and survival of dendritic cells. *J Exp Med* 204: 2267–2275, 2007.
  67. Iwakaki T, Akai R, Kohno K. IRE1alpha disruption causes histological abnormality of exocrine tissues, increase of blood glucose level, and decrease of serum immunoglobulin level. *PLoS One* 5: e13052, 2010.
  68. Iwakaki T, Akai R, Yamanaka S, Kohno K. Function of IRE1 alpha in the placenta is essential for placental development and embryonic viability. *Proc Natl Acad Sci USA* 106: 16657–16662, 2009.
  69. Kang SW, Rane NS, Kim SJ, Garrison JL, Taunton J, Hegde RS. Substrate-specific translocational attenuation during ER stress defines a pre-emptive quality control pathway. *Cell* 127: 999–1013, 2006.
  70. Kaser A, Lee AH, Franke A, Glickman JN, Zeissig S, Tilg H, Nieuwenhuis EE, Higgins DE, Schreiber S, Glimcher LH, Blumberg RS. XBP1 links ER stress to intestinal inflammation and confers genetic risk for human inflammatory bowel disease. *Cell* 134: 743–756, 2008.
  71. Kaufman RJ, Malhotra JD. Calcium trafficking integrates endoplasmic reticulum function with mitochondrial bioenergetics. *Biochim Biophys Acta* 1843: 2233–2239, 2014.
  72. Kelekar A, Thompson CB. Bcl-2-family proteins: the role of the BH3 domain in apoptosis. *Trends Cell Biol* 8: 324–330, 1998.
  73. Kharabi Masouleh B, Geng H, Hertz C, Chan LN, Logan AC, Chang MS, Huang C, Swaminathan S, Sun H, Paietta E, Melnick AM, Koeffler P, Muschen M. Mechanistic rationale for targeting the unfolded protein response in pre-B acute lymphoblastic leukemia. *Proc Natl Acad Sci USA* 111: E2219–E2228, 2014.
  74. Kim R, Emi M, Tanabe K, Murakami S. Role of the unfolded protein response in cell death. *Apoptosis* 11: 5–13, 2006.
  75. Kimata Y, Kohno K. Endoplasmic reticulum stress-sensing mechanisms in yeast and mammalian cells. *Curr Opin Cell Biol* 23: 135–142, 2011.
  76. Kimata Y, Oikawa D, Shimizu Y, Ishiwata-Kimata Y, Kohno K. A role for BiP as an adaptor for the endoplasmic reticulum stress-sensing protein Ire1. *J Cell Biol* 167: 445–456, 2004.
  77. Kimball SR, Clemens MJ, Tilleray VJ, Wek RC, Horvitz RL, Jefferson LS. The double-stranded RNA-activated protein kinase PKR is

- dispensable for regulation of translation initiation in response to either calcium mobilization from the endoplasmic reticulum or essential amino acid starvation. *Biochem Biophys Res Commun* 280: 293–300, 2001.
78. **Kimmig P, Diaz M, Zheng J, Williams CC, Lang A, Aragon T, Li H, Walter P.** The unfolded protein response in fission yeast modulates stability of select mRNAs to maintain protein homeostasis. *eLife* 1: e00048, 2012.
  79. **Klee M, Pallauf K, Alcalá S, Fleischer A, Pimentel-Muinos FX.** Mitochondrial apoptosis induced by BH3-only molecules in the exclusive presence of endoplasmic reticular Bak. *EMBO J* 28: 1757–1768, 2009.
  80. **Kojima E, Takeuchi A, Haneda M, Yagi A, Hasegawa T, Yamaki K, Takeda K, Akira S, Shimokata K, Isobe K.** The function of GADD34 is a recovery from a shutoff of protein synthesis induced by ER stress: elucidation by GADD34-deficient mice. *FASEB J* 17: 1573–1575, 2003.
  81. **Korennykh AV, Egea PF, Korostelev AA, Finer-Moore J, Zhang C, Shokat KM, Stroud RM, Walter P.** The unfolded protein response signals through high-order assembly of Ire1. *Nature* 457: 687–693, 2009.
  82. **Kroemer G, Marino G, Levine B.** Autophagy and the integrated stress response. *Mol Cell* 40: 280–293, 2010.
  83. **Lange PS, Chavez JC, Pinto JT, Coppola G, Sun CW, Townes TM, Geschwind DH, Ratan RR.** ATF4 is an oxidative stress-inducible, prodeath transcription factor in neurons in vitro and in vivo. *J Exp Med* 205: 1227–1242, 2008.
  84. **Lee AH, Chu GC, Iwakoshi NN, Glimcher LH.** XBP-1 is required for biogenesis of cellular secretory machinery of exocrine glands. *EMBO J* 24: 4368–4380, 2005.
  85. **Lee AH, Heidtman K, Hotamisligil GS, Glimcher LH.** Dual and opposing roles of the unfolded protein response regulated by IRE1alpha and XBP1 in proinsulin processing and insulin secretion. *Proc Natl Acad Sci USA* 108: 8885–8890, 2011.
  86. **Lee AH, Scapa EF, Cohen DE, Glimcher LH.** Regulation of hepatic lipogenesis by the transcription factor XBP1. *Science* 320: 1492–1496, 2008.
  87. **Lee K, Tirasophon W, Shen X, Michalak M, Prywes R, Okada T, Yoshida H, Mori K, Kaufman RJ.** IRE1-mediated unconventional mRNA splicing and S2P-mediated ATF6 cleavage merge to regulate XBP1 in signaling the unfolded protein response. *Genes Dev* 16: 452–466, 2002.
  88. **Lee KP, Dey M, Neculai D, Cao C, Dever TE, Sicheri F.** Structure of the dual enzyme Ire1 reveals the basis for catalysis and regulation in nonconventional RNA splicing. *Cell* 132: 89–100, 2008.
  89. **Letai A, Bassik MC, Walensky LD, Sorcinelli MD, Weiler S, Korsmeyer SJ.** Distinct BH3 domains either sensitize or activate mitochondrial apoptosis, serving as prototype cancer therapeutics. *Cancer Cell* 2: 183–192, 2002.
  90. **Lew QJ, Chu KL, Lee J, Koh PL, Rajasegaran V, Teo JY, Chao SH.** PCAF interacts with XBP-1S and mediates XBP-1S-dependent transcription. *Nucleic Acids Res* 39: 429–439, 2011.
  91. **Li H, Korennykh AV, Behrman SL, Walter P.** Mammalian endoplasmic reticulum stress sensor IRE1 signals by dynamic clustering. *Proc Natl Acad Sci USA* 107: 16113–16118, 2010.
  92. **Li J, Lee B, Lee AS.** Endoplasmic reticulum stress-induced apoptosis: multiple pathways and activation of p53-up-regulated modulator of apoptosis (PUMA) and NOXA by p53. *J Biol Chem* 281: 7260–7270, 2006.
  93. **Lin JH, Li H, Yasumura D, Cohen HR, Zhang C, Panning B, Shokat KM, Lavail MM, Walter P.** IRE1 signaling affects cell fate during the unfolded protein response. *Science* 318: 944–949, 2007.
  94. **Lin JH, Li H, Zhang Y, Ron D, Walter P.** Divergent effects of PERK and IRE1 signaling on cell viability. *PLoS One* 4: e4170, 2009.
  95. **Lisbona F, Rojas-Rivera D, Thielen P, Zamorano S, Todd D, Martinon F, Glavic A, Kress C, Lin JH, Walter P, Reed JC, Glimcher LH, Hetz C.** BAX inhibitor-1 is a negative regulator of the ER stress sensor IRE1alpha. *Mol Cell* 33: 679–691, 2009.
  96. **Liu CY, Schroder M, Kaufman RJ.** Ligand-independent dimerization activates the stress response kinases IRE1 and PERK in the lumen of the endoplasmic reticulum. *J Biol Chem* 275: 24881–24885, 2000.
  97. **Ma T, Trinh MA, Wexler AJ, Bourbon C, Gatti E, Pierre P, Cavener DR, Klann E.** Suppression of eIF2alpha kinases alleviates Alzheimer's disease-related plasticity and memory deficits. *Nat Neurosci* 16: 1299–1305, 2013.
  98. **Ma Y, Hendershot LM.** Delineation of a negative feedback regulatory loop that controls protein translation during endoplasmic reticulum stress. *J Biol Chem* 278: 34864–34873, 2003.
  99. **Ma Y, Hendershot LM.** The role of the unfolded protein response in tumour development: friend or foe? *Nat Rev Cancer* 4: 966–977, 2004.
  100. **Maiuolo J, Bulotta S, Verderio C, Benfante R, Borgese N.** Selective activation of the transcription factor ATF6 mediates endoplasmic reticulum proliferation triggered by a membrane protein. *Proc Natl Acad Sci USA* 108: 7832–7837, 2011.
  101. **Marciniak SJ, Yun CY, Oyadomari S, Novoa I, Zhang Y, Jungreis R, Nagata K, Harding HP, Ron D.** CHOP induces death by promoting protein synthesis and oxidation in the stressed endoplasmic reticulum. *Genes Dev* 18: 3066–3077, 2004.
  102. **Martino MB, Jones L, Brighton B, Ehre C, Abdulah L, Davis CW, Ron D, O'Neal WK, Ribeiro CM.** The ER stress transducer IRE1beta is required for airway epithelial mucin production. *Mucosal Immunol* 6: 639–654, 2013.
  103. **Martinon F, Chen X, Lee AH, Glimcher LH.** TLR activation of the transcription factor XBP1 regulates innate immune responses in macrophages. *Nat Immunol* 11: 411–418, 2010.
  104. **Masuoka HC, Townes TM.** Targeted disruption of the activating transcription factor 4 gene results in severe fetal anemia in mice. *Blood* 99: 736–745, 2002.
  105. **Mathew R, Karp CM, Beaudoin B, Vuong N, Chen G, Chen HY, Bray K, Reddy A, Bhanot G, Gelinas C, Dipaola RS, Karantzawa-Wadsworth V, White E.** Autophagy suppresses tumorigenesis through elimination of p62. *Cell* 137: 1062–1075, 2009.
  106. **Matus S, Lopez E, Valenzuela V, Nassif M, Hetz C.** Functional contribution of the transcription factor ATF4 to the pathogenesis of amyotrophic lateral sclerosis. *PLoS One* 8: e66672, 2013.
  107. **Maurel M, Chevet E, Tavernier J, Gerlo S.** Getting RIDD of RNA: IRE1 in cell fate regulation. *Trends Biochem Sci* 39: 245–254, 2014.
  108. **Maurel M, Dejeans N, Taouji S, Chevet E, Grosset CF.** MicroRNA-1291-mediated silencing of IRE1alpha enhances Glypican-3 expression. *RNA* 19: 778–788, 2013.
  109. **McCullough KD, Martindale JL, Klotz LO, Aw TY, Holbrook NJ.** Gadd153 sensitizes cells to endoplasmic reticulum stress by down-regulating Bcl2 and perturbing the cellular redox state. *Mol Cell Biol* 21: 1249–1259, 2001.
  110. **Mishiba K, Nagashima Y, Suzuki E, Hayashi N, Ogata Y, Shimada Y, Koizumi N.** Defects in IRE1 enhance cell death and fail to degrade mRNAs encoding secretory pathway proteins in the *Arabidopsis* unfolded protein response. *Proc Natl Acad Sci USA* 110: 5713–5718, 2013.
  111. **Moenner M, Pluquet O, Boucheccareilh M, Chevet E.** Integrated endoplasmic reticulum stress responses in cancer. *Cancer Res* 67: 10631–10634, 2007.
  112. **Moreno JA, Halliday M, Molloy C, Radford H, Verity N, Axten JM, Ortori CA, Willis AE, Fischer PM, Barrett DA, Mallucci GR.** Oral treatment targeting the unfolded protein response prevents neurodegeneration and clinical disease in prion-infected mice. *Sci Transl Med* 5: 206ra138, 2013.
  113. **Munoz JP, Ivanova S, Sanchez-Wandelmer J, Martinez-Cristobal P, Noguera E, Sancho A, Diaz-Ramos A, Hernandez-Alvarez MI, Sebastian D, Mauvezin C, Palacin M, Zorzano A.** Mfn2 modulates the UPR and mitochondrial function via repression of PERK. *EMBO J* 32: 2348–2361, 2013.
  114. **Nguyen DT, Kebache S, Fazel A, Wong HN, Jenna S, Emadali A, Lee EH, Bergeron JJ, Kaufman RJ, Larose L, Chevet E.** Nck-dependent activation of extracellular signal-regulated kinase-1 and regulation of cell survival during endoplasmic reticulum stress. *Mol Biol Cell* 15: 4248–4260, 2004.
  115. **Ni M, Zhou H, Wey S, Baumeister P, Lee AS.** Regulation of PERK signaling and leukemic cell survival by a novel cytosolic isoform of the UPR regulator GRP78/BiP. *PLoS One* 4: e6868, 2009.
  116. **Nishitoh H, Matsuzawa A, Tobiume K, Saegusa K, Takeda K, Inoue K, Hori S, Kakizuka A, Ichijo H.** ASK1 is essential for endoplasmic reticulum stress-induced neuronal cell death triggered by expanded polyglutamine repeats. *Genes Dev* 16: 1345–1355, 2002.
  117. **Novoa I, Zeng H, Harding HP, Ron D.** Feedback inhibition of the unfolded protein response by GADD34-mediated dephosphorylation of eIF2alpha. *J Cell Biol* 153: 1011–1022, 2001.
  118. **Ogata M, Hino S, Saito A, Morikawa K, Kondo S, Kanemoto S, Murakami T, Taniguchi M, Tani I, Yoshinaga K, Shiosaka S, Hammarback JA, Urano F, Imaizumi K.** Autophagy is activated for

- cell survival after endoplasmic reticulum stress. *Mol Cell Biol* 26: 9220–9231, 2006.
119. **Oikawa D, Kitamura A, Kinjo M, Iwawaki T.** Direct association of unfolded proteins with mammalian ER stress sensor, IRE1beta. *PLoS One* 7: e51290, 2012.
  120. **Okada T, Haze K, Nadanaka S, Yoshida H, Seidah NG, Hirano Y, Sato R, Negishi M, Mori K.** A serine protease inhibitor prevents endoplasmic reticulum stress-induced cleavage but not transport of the membrane-bound transcription factor ATF6. *J Biol Chem* 278: 31024–31032, 2003.
  121. **Okamura K, Kimata Y, Higashio H, Tsuru A, Kohno K.** Dissociation of Kar2p/BiP from an ER sensory molecule, Ire1p, triggers the unfolded protein response in yeast. *Biochem Biophys Res Commun* 279: 445–450, 2000.
  122. **Osorio F, Tavernier SJ, Hoffmann E, Saeys Y, Martens L, Veters J, Delrue I, De Rycke R, Parthoens E, Pouliot P, Iwawaki T, Janssens S, Lambrecht BN.** The unfolded-protein-response sensor IRE-1alpha regulates the function of CD8alpha+ dendritic cells. *Nat Immunol* 15: 248–257, 2014.
  123. **Ow YP, Green DR, Hao Z, Mak TW.** Cytochrome *c*: functions beyond respiration. *Nat Rev Mol Cell Biol* 9: 532–542, 2008.
  124. **Oyadomari S, Yun C, Fisher EA, Kreglinger N, Kreibich G, Oyadomari M, Harding HP, Goodman AG, Harant H, Garrison JL, Taunton J, Katze MG, Ron D.** Cotranslocational degradation protects the stressed endoplasmic reticulum from protein overload. *Cell* 126: 727–739, 2006.
  125. **Ozcan U, Cao Q, Yilmaz E, Lee AH, Iwakoshi NN, Ozdelen E, Tuncman G, Gorgun C, Glimcher LH, Hotamisligil GS.** Endoplasmic reticulum stress links obesity, insulin action, and type 2 diabetes. *Science* 306: 457–461, 2004.
  126. **Parsons DW, Jones S, Zhang X, Lin JC, Leary RJ, Angenendt P, Mankoo P, Carter H, Siu IM, Gallia GL, Olivari A, McLendon R, Rasheed BA, Keir S, Nikolskaya T, Nikolsky Y, Busam DA, Tekleab H, Diaz LA Jr, Hartigan J, Smith DR, Strausberg RL, Marie SK, Shinjo SM, Yan H, Riggins GJ, Bigner DD, Karchin R, Papadopoulos N, Parmigiani G, Vogelstein B, Velculescu VE, and Kinzler KW.** An integrated genomic analysis of human glioblastoma multiforme. *Science* 321: 1807–1812, 2008.
  127. **Pluquet O, Dejeans N, Bouche-careilh M, Lhomond S, Pineau R, Higa A, Delugin M, Combe C, Loriot S, Cubel G, Dugot-Senant N, Vital A, Loiseau H, Gosline SJ, Taouji S, Hallett M, Sarkaria JN, Anderson K, Wu W, Rodriguez FJ, Rosenbaum J, Saltel E, Fernandez-Zapico ME, Chevret E.** Posttranscriptional regulation of PER1 underlies the oncogenic function of IRE1alpha. *Cancer Res* 73: 4732–4743, 2013.
  128. **Reimold AM, Etkin A, Clauss I, Perkins A, Friend DS, Zhang J, Horton HF, Scott A, Orkin SH, Byrne MC, Grusby MJ, Glimcher LH.** An essential role in liver development for transcription factor XBP-1. *Genes Dev* 14: 152–157, 2000.
  129. **Reimold AM, Iwakoshi NN, Manis J, Vallabhajosyula P, Szomolanyi-Tsuda E, Gravallesse EM, Friend D, Grusby MJ, Alt F, Glimcher LH.** Plasma cell differentiation requires the transcription factor XBP-1. *Nature* 412: 300–307, 2001.
  130. **Rodriguez D, Rojas-Rivera D, Hetz C.** Integrating stress signals at the endoplasmic reticulum: The BCL-2 protein family rheostat. *Biochim Biophys Acta* 1813: 564–574, 2011.
  131. **Rodriguez DA, Zamorano S, Lisbona F, Rojas-Rivera D, Urra H, Cubillos-Ruiz JR, Armisen R, Henriquez DR, Cheng EH, Letek M, Vaisar T, Irrazabal T, Gonzalez-Billault C, Letai A, Pimentel-Muinos FX, Kroemer G, Hetz C.** BH3-only proteins are part of a regulatory network that control the sustained signalling of the unfolded protein response sensor IRE1alpha. *EMBO J* 31: 2322–2335, 2012.
  132. **Rojas-Rivera D, Armisen R, Colombo A, Martinez G, Eguiguren AL, Diaz A, Kiviluoto S, Rodriguez D, Patron M, Rizzuto R, Bultynck G, Concha ML, Sierralta J, Stutzin A, Hetz C.** TMBIM3/GRINA is a novel unfolded protein response (UPR) target gene that controls apoptosis through the modulation of ER calcium homeostasis. *Cell Death Differ* 19: 1013–1026, 2012.
  133. **Rojas-Rivera D, Hetz C.** TMBIM protein family: ancestral regulators of cell death. *Oncogene*. doi: 10.1038/nc.2014.6. [Epub ahead of print].
  134. **Ron D, Walter P.** Signal integration in the endoplasmic reticulum unfolded protein response. *Nat Rev Mol Cell Biol* 8: 519–529, 2007.
  135. **Rong J, Chen L, Toth JI, Tcherpakov M, Petroski MD, Reed JC.** Bifunctional apoptosis regulator (BAR), an endoplasmic reticulum (ER)-associated E3 ubiquitin ligase, modulates BI-1 protein stability and function in ER Stress. *J Biol Chem* 286: 1453–1463, 2011.
  136. **Rutkowski DT, Hegde RS.** Regulation of basal cellular physiology by the homeostatic unfolded protein response. *J Cell Biol* 189: 783–794, 2010.
  137. **Rutkowski DT, Wu J, Back SH, Callaghan MU, Ferris SP, Iqbal J, Clark R, Miao H, Hassler JR, Fornek J, Katze MG, Hussain MM, Song B, Swathirajan J, Wang J, Yau GD, Kaufman RJ.** UPR pathways combine to prevent hepatic steatosis caused by ER stress-mediated suppression of transcriptional master regulators. *Dev Cell* 15: 829–840, 2008.
  138. **Saito A, Ochiai K, Kondo S, Tsumagari K, Murakami T, Cavener DR, Imaizumi K.** Endoplasmic reticulum stress response mediated by the PERK-eIF2(alpha)-ATF4 pathway is involved in osteoblast differentiation induced by BMP2. *J Biol Chem* 286: 4809–4818, 2011.
  139. **Scheuner D, Song B, McEwen E, Liu C, Laybutt R, Gillespie P, Saunders T, Bonner-Weir S, Kaufman RJ.** Translational control is required for the unfolded protein response and in vivo glucose homeostasis. *Mol Cell* 7: 1165–1176, 2001.
  140. **Schindler AJ, Schekman R.** In vitro reconstitution of ER-stress induced ATF6 transport in COPII vesicles. *Proc Natl Acad Sci USA* 106: 17775–17780, 2009.
  141. **Schroder M, Kaufman RJ.** The mammalian unfolded protein response. *Annu Rev Biochem* 74: 739–789, 2005.
  142. **Shen J, Chen X, Hendershot L, Prywes R.** ER stress regulation of ATF6 localization by dissociation of BiP/GRP78 binding and unmasking of Golgi localization signals. *Dev Cell* 3: 99–111, 2002.
  143. **Shoulders MD, Ryno LM, Genereux JC, Moresco JJ, Tu PG, Wu C, Yates JR 3rd, Su AI, Kelly JW, Wiseman RL.** Stress-independent activation of XBP1s and/or ATF6 reveals three functionally diverse ER proteostasis environments. *Cell Rep* 3: 1279–1292, 2013.
  144. **So JS, Hur KY, Tarrío M, Ruda V, Frank-Kamenetsky M, Fitzgerald K, Koteliensky V, Lichtman AH, Iwawaki T, Glimcher LH, Lee AH.** Silencing of lipid metabolism genes through IRE1alpha-mediated mRNA decay lowers plasma lipids in mice. *Cell Metab* 16: 487–499, 2012.
  145. **Szegezdi E, Logue SE, Gorman AM, Samali A.** Mediators of endoplasmic reticulum stress-induced apoptosis. *EMBO Rep* 7: 880–885, 2006.
  146. **Tabas I, Ron D.** Integrating the mechanisms of apoptosis induced by endoplasmic reticulum stress. *Nat Cell Biol* 13: 184–190, 2011.
  147. **Tait SW, Green DR.** Mitochondria and cell death: outer membrane permeabilization and beyond. *Nat Rev Mol Cell Biol* 11: 621–632, 2010.
  148. **Trinh MA, Kaphzan H, Wek RC, Pierre P, Cavener DR, Klann E.** Brain-specific disruption of the eIF2alpha kinase PERK decreases ATF4 expression and impairs behavioral flexibility. *Cell Rep* 1: 676–688, 2012.
  149. **Tsuru A, Fujimoto N, Takahashi S, Saito M, Nakamura D, Iwano M, Iwawaki T, Kadokura H, Ron D, Kohno K.** Negative feedback by IRE1beta optimizes mucin production in goblet cells. *Proc Natl Acad Sci USA* 110: 2864–2869, 2013.
  150. **Upton JP, Wang L, Han D, Wang ES, Huskey NE, Lim L, Truitt M, McManus MT, Ruggero D, Goga A, Papa FR, Oakes SA.** IRE1alpha cleaves select microRNAs during ER stress to derepress translation of proapoptotic Caspase-2. *Science* 338: 818–822, 2012.
  151. **Urra F, Wang X, Bertolotti A, Zhang Y, Chung P, Harding HP, Ron D.** Coupling of stress in the ER to activation of JNK protein kinases by transmembrane protein kinase IRE1. *Science* 287: 664–666, 2000.
  152. **Urra H, Dufey E, Lisbona F, Rojas-Rivera D, Hetz C.** When ER stress reaches a dead end. *Biochim Biophys Acta* 1833: 3507–3517, 2013.
  153. **Urra H, Hetz C.** A novel ER stress-independent function of the UPR in angiogenesis. *Mol Cell* 54: 542–544, 2014.
  154. **Usui M, Yamaguchi S, Tanji Y, Tominaga R, Ishigaki Y, Fukumoto M, Katagiri H, Mori K, Oka Y, Ishihara H.** Atf6alpha-null mice are glucose intolerant due to pancreatic beta-cell failure on a high-fat diet but partially resistant to diet-induced insulin resistance. *Metabolism* 61: 1118–1128, 2012.
  155. **Valenzuela V, Collyer E, Armentano D, Parsons GB, Court FA, Hetz C.** Activation of the unfolded protein response enhances motor recovery after spinal cord injury. *Cell Death Dis* 3: e272, 2012.
  156. **van Huizen R, Martindale JL, Gorospe M, Holbrook NJ.** P58<sup>IPK</sup>, a novel endoplasmic reticulum stress-inducible protein and potential negative regulator of eIF2alpha signaling. *J Biol Chem* 278: 15558–15564, 2003.
  157. **Vidal RL, Figueroa A, Court FA, Thielen P, Molina C, Wirth C, Caballero B, Kiffin R, Segura-Aguilar J, Cuervo AM, Glimcher LH, Hetz C.** Targeting the UPR transcription factor XBP1 protects against

- Huntington's disease through the regulation of FoxO1 and autophagy. *Hum Mol Genet* 21: 2245–2262, 2012.
158. **Walter P, Ron D.** The unfolded protein response: from stress pathway to homeostatic regulation. *Science* 334: 1081–1086, 2011.
159. **Wei J, Sheng X, Feng D, McGrath B, Cavener DR.** PERK is essential for neonatal skeletal development to regulate osteoblast proliferation and differentiation. *J Cell Physiol* 217: 693–707, 2008.
160. **Wu J, Rutkowski DT, Dubois M, Swathirajan J, Saunders T, Wang J, Song B, Yau GD, Kaufman RJ.** ATF6alpha optimizes long-term endoplasmic reticulum function to protect cells from chronic stress. *Dev Cell* 13: 351–364, 2007.
161. **Xu Q, Reed JC.** Bax inhibitor-1, a mammalian apoptosis suppressor identified by functional screening in yeast. *Mol Cell* 1: 337–346, 1998.
162. **Yamamoto K, Sato T, Matsui T, Sato M, Okada T, Yoshida H, Harada A, Mori K.** Transcriptional induction of mammalian ER quality control proteins is mediated by single or combined action of ATF6alpha and XBP1. *Dev Cell* 13: 365–376, 2007.
163. **Yamamoto K, Takahara K, Oyadomari S, Okada T, Sato T, Harada A, Mori K.** Induction of liver steatosis and lipid droplet formation in ATF6alpha-knockout mice burdened with pharmacological endoplasmic reticulum stress. *Mol Biol Cell* 21: 2975–2986, 2010.
164. **Yamani L, Latreille M, Larose L.** Interaction of Nck1 and PERK phosphorylated at Y(5)(6)(1) negatively modulates PERK activity and PERK regulation of pancreatic beta-cell proinsulin content. *Mol Biol Cell* 25: 702–711, 2014.
165. **Yan W, Frank CL, Korth MJ, Sopher BL, Novoa I, Ron D, Katze MG.** Control of PERK eIF2alpha kinase activity by the endoplasmic reticulum stress-induced molecular chaperone P58IPK. *Proc Natl Acad Sci USA* 99: 15920–15925, 2002.
166. **Yoshida H, Matsui T, Yamamoto A, Okada T, Mori K.** XBP1 mRNA is induced by ATF6 and spliced by IRE1 in response to ER stress to produce a highly active transcription factor. *Cell* 107: 881–891, 2001.
167. **Youle RJ, Strasser A.** The BCL-2 protein family: opposing activities that mediate cell death. *Nat Rev Mol Cell Biol* 9: 47–59, 2008.
168. **Zamzami N, Marchetti P, Castedo M, Zanin C, Vayssiere JL, Petit PX, Kroemer G.** Reduction in mitochondrial potential constitutes an early irreversible step of programmed lymphocyte death in vivo. *J Exp Med* 181: 1661–1672, 1995.
169. **Zeng L, Liu YP, Sha H, Chen H, Qi L, Smith JA.** XBP-1 couples endoplasmic reticulum stress to augmented IFN-beta induction via a cis-acting enhancer in macrophages. *J Immunol* 185: 2324–2330, 2010.
170. **Zhang K, Wong HN, Song B, Miller CN, Scheuner D, Kaufman RJ.** The unfolded protein response sensor IRE1alpha is required at 2 distinct steps in B cell lymphopoiesis. *J Clin Invest* 115: 268–281, 2005.
171. **Zhang P, McGrath B, Li S, Frank A, Zambito F, Reinert J, Gannon M, Ma K, McNaughton K, Cavener DR.** The PERK eukaryotic initiation factor 2 alpha kinase is required for the development of the skeletal system, postnatal growth, and the function and viability of the pancreas. *Mol Cell Biol* 22: 3864–3874, 2002.
172. **Zhou J, Liu CY, Back SH, Clark RL, Peisach D, Xu Z, Kaufman RJ.** The crystal structure of human IRE1 luminal domain reveals a conserved dimerization interface required for activation of the unfolded protein response. *Proc Natl Acad Sci USA* 103: 14343–14348, 2006.

