

Intermediate reduction method and infinitely many positive solutions of nonlinear Schrödinger equations with non-symmetric potentials

Manuel del Pino · Juncheng Wei · Wei Yao

Received: 27 March 2014 / Accepted: 22 May 2014 / Published online: 29 June 2014
© Springer-Verlag Berlin Heidelberg 2014

Abstract We consider the standing-wave problem for a nonlinear Schrödinger equation, corresponding to the semilinear elliptic problem

$$-\Delta u + V(x)u = |u|^{p-1}u, \quad u \in H^1(\mathbb{R}^2),$$

where $V(x)$ is a uniformly positive potential and $p > 1$. Assuming that

$$V(x) = V_\infty + \frac{a}{|x|^m} + O\left(\frac{1}{|x|^{m+\sigma}}\right), \quad \text{as } |x| \rightarrow +\infty,$$

for instance if $p > 2$, $m > 2$ and $\sigma > 1$ we prove the existence of infinitely many positive solutions. If $V(x)$ is radially symmetric, this result was proved in [43]. The proof without symmetries is much more difficult, and for that we develop a new *intermediate Lyapunov–Schmidt reduction method*, which is a compromise between the finite and infinite dimensional versions of it.

Mathematics Subject Classification 35J15 · 35J61 · 35B08 · 35B09 · 35Q55

Communicated by P. Rabinowitz.

M. del Pino (✉) · W. Yao
Departamento de Ingeniería Matemática and Centro de Modelamiento Matemático (UMI 2807 CNRS),
Universidad de Chile, Casilla 170 Correo 3, Santiago, Chile
e-mail: delpino@dim.uchile.cl

W. Yao
e-mail: wyao.cn@gmail.com

J. Wei
Department of Mathematics, University of British Columbia, Vancouver, BC V6T 1Z2, Canada

J. Wei
Department of Mathematics, The Chinese University of Hong Kong, Shatin, NT, Hong Kong
e-mail: jcwei@math.ubc.ca

Contents

1 Introduction and statement of the main result 474
 2 Description of the construction 478
 3 Preliminaries 484
 4 The Lyapunov–Schmidt reduction 490
 4.1 Linear analysis 491
 4.2 Nonlinear analysis 495
 5 A further reduction process 500
 5.1 Projections 500
 5.2 The invertibility of T 507
 5.3 Reduction to one dimension 507
 6 Proof of Theorem 1.1: variational reduction 509
 7 Generalizations and discussion 510
 7.1 More general nonlinearities 510
 7.2 Sign-changing solutions 510
 7.3 Remarks on condition (1.14) 511
 7.4 The anisotropic case 512
 7.5 Optimal condition on the decay 512
 7.6 Higher dimensions 512
 7.7 Higher dimensional concentration phenomena 513
 Appendix A: Circulant matrices and proof of Lemma 5.4 513
 Appendix B: Energy expansion 520
 References 522

1 Introduction and statement of the main result

In this paper we consider the problem of finding positive solutions of the classical semilinear elliptic problem

$$-\Delta u + V(x)u = |u|^{p-1}u \text{ in } \mathbb{R}^N, \tag{1.1}$$

where $\Delta = \sum_{j=1}^N \frac{\partial^2}{\partial x_j^2}$ stands for the Laplace operator in \mathbb{R}^N , $V(x)$ is a non-negative potential, and $p > 1$.

Equation (1.1) arises in various branches of applied mathematics and physics (cf. [12] and references therein). For instance, in condensed matter physics one simulates the interaction effect among many particles to obtain a focusing nonlinear Schrödinger equation of the form

$$i\hbar \frac{\partial \psi}{\partial t} = -\hbar^2 \Delta \psi + W(x)\psi - |\psi|^{p-1}\psi \text{ in } [0, \infty) \times \mathbb{R}^N, \tag{1.2}$$

where i is the imaginary unit, \hbar the Planck constant and $W(x)$ a given potential. *Standing wave solutions* of (1.2) are those of the form

$$\psi(t, x) = e^{-i\lambda t/\hbar} u(\hbar^{-1}x),$$

where $u(x)$ is a real-valued function. Then (1.2) reduces to Eq. (1.1) for u , where $V(x) = W(\hbar x) - \lambda$.

In what follows, we shall only consider positive, finite energy solutions of (1.1). Namely, we are concerned with the problem:

$$\begin{cases} -\Delta u + V(x)u - u^p = 0 & \text{in } \mathbb{R}^N, \\ u > 0 \text{ in } \mathbb{R}^N, u \in H^1(\mathbb{R}^N). \end{cases} \tag{1.3}$$

Associated to (1.3) is the energy functional

$$\mathcal{E}(u) = \frac{1}{2} \int_{\mathbb{R}^N} \{|\nabla u|^2 + V(x)u^2\} dx - \frac{1}{p+1} \int_{\mathbb{R}^N} u_+^{p+1} dx, \tag{1.4}$$

where $u_+ = \max\{u, 0\}$. In all what follows, we make the following structure assumptions on V and p :

$$V \text{ is locally H\"older continuous, } V \in L^\infty(\mathbb{R}^N) \text{ and } V_0 = \inf_{x \in \mathbb{R}^N} V(x) > 0; \tag{1.5}$$

$$1 < p < \infty \text{ for } N = 2 \text{ and } 1 < p < \frac{N+2}{N-2} \text{ for } N \geq 3. \tag{1.6}$$

Under these hypotheses, it is standard that classical solutions of (1.3) correspond precisely to non-trivial critical points of \mathcal{E} in $H^1(\mathbb{R}^N)$.

Let us denote the set of solutions of problem (1.3) by \mathcal{S}_V . A natural question is whether or not $\mathcal{S}_V \neq \emptyset$. When V is radially symmetric, the answer is yes (cf. Theorem 4.6 in [24]). But the answer is no when the potential is increasing along a direction (cf. Theorem 1.1 in [15]).

If we further assume that

$$\lim_{|x| \rightarrow \infty} V(x) = V_\infty > 0, \tag{1.7}$$

the existence of a positive solution of (1.3) has been widely investigated. For example, if we further suppose that

$$\inf_{x \in \mathbb{R}^N} V(x) < V_\infty, \tag{1.8}$$

then one can show that (1.3) has a least energy (ground state) solution by using the concentration compactness principle (cf. [24, 31, 32, 39]). But if (1.8) does not hold, problem (1.3) may not have a least energy solution and solutions have to be sought for at higher energy levels. Results in this direction are contained in [6, 7, 9], where a positive solution has been found by variational methods under a suitable decay condition on V at infinity.

The structure of the solution set \mathcal{S}_V may be quite rich and interesting. Let us consider for instance the semi-classical limit case:

$$\begin{cases} -\varepsilon^2 \Delta u + W(x)u - u^p = 0 & \text{in } \mathbb{R}^N, \\ u > 0 & \text{in } \mathbb{R}^N, u \in H^1(\mathbb{R}^N), \end{cases} \tag{1.9}$$

where $\varepsilon > 0$ is a small parameter. Naturally, problem (1.9) is equivalent to problem (1.3) for $V(x) = W(\varepsilon x)$. It is known that as ε goes to zero, highly concentrated solutions near critical points of the potential W can be found, see [1, 10, 11, 17–20, 25, 27, 37, 42], or near higher dimensional stationary sets of other auxiliary potentials [3, 21, 33, 41]. The number of solutions of (1.9) may depend on the number or type of the critical points of $W(x)$. It is rather difficult task to understand the structure of \mathcal{S}_V for an arbitrary potential V . For instance, a conspicuously unanswered question is whether or not $\mathcal{S}_V \neq \emptyset$ for any potential V satisfying (1.7).

Summing up, the above-mentioned work concerns the existence of positive solutions, i.e., $\mathcal{S}_V \neq \emptyset$. There is less work on the multiplicity of positive solutions of (1.3), namely on estimating $\#\mathcal{S}_V$. A seminal result in this direction was given in Coti-Zelati and Rabinowitz [16] where $V(x)$ is spatially periodic. In that situation they prove the existence of infinitely many positive solutions, distinct up to periodic translations, via variational methods.

Recently, by assuming that $V = V(|x|)$ is radially symmetric, the second author and Yan [43] proved that problem (1.3) has infinitely many positive non-radial solutions if there are constants $V_\infty > 0, a > 0, m > 1$, and $\sigma > 0$, such that

$$V(r) = V_\infty + \frac{a}{r^m} + O\left(\frac{1}{r^{m+\sigma}}\right), \quad \text{as } r \rightarrow +\infty. \tag{1.10}$$

An alternative proof through min-max methods was given by Devillanova and Solimini [23].

The proof in [43] uses in essential way the radial symmetry of the potential V . On the other hand, it is conjectured there that the result should remain true when the symmetry requirement is lifted:

Conjecture *Problem (1.3) has infinitely many positive solutions if there are constants $V_\infty > 0, a > 0, m > 1,$ and $\sigma > 0,$ such that the potential $V(x)$ satisfies*

$$V(x) = V_\infty + \frac{a}{|x|^m} + O\left(\frac{1}{|x|^{m+\sigma}}\right), \quad \text{as } |x| \rightarrow +\infty. \tag{1.11}$$

Results in this direction with non-symmetric potentials, as far as we know, there are only perturbative results (cf. [4, 14]). For instance, if $V(x)$ tends to V_∞ from above with a suitable rate:

$$V(x) \geq V_0 > 0, \quad \lim_{|x| \rightarrow \infty} (V(x) - V_\infty) e^{\bar{\eta}|x|} = +\infty, \quad \text{for some } \bar{\eta} \in (0, \sqrt{V_\infty}), \tag{1.12}$$

and V satisfies a global condition:

$$\sup_{y \in \mathbb{R}^N} \|V(x) - V_\infty\|_{L^{N/2}(B_1(y))} < \mathcal{V}, \tag{1.13}$$

where \mathcal{V} is a sufficiently small positive constant (with no explicit expression), Cerami et al. [14] proved that problem (1.3) has infinitely many positive solutions by purely variational methods. (In [4], Ao and Wei gave a new proof of this result, using localized energy method. The new techniques also allow them to deal with more general nonlinearity).

The main purpose of this paper is to prove the above conjecture under some additional assumptions. In [43], the fact that V is radially symmetric allows to build a k -bump solution for an arbitrary $k \geq 1$ with a k -dyadic symmetry, reducing the problem to just adjusting one parameter representing the location of a single bump along a given ray. A finite dimensional Lyapunov–Schmidt reduction method is used.

When V is non-symmetric, we cannot constrain the bump configuration to any symmetry class. We are thus forced to deal with a large number of bumps and therefore with a huge number of parameters which need to be adjusted. This poses a tremendous difficulty in the construction comparatively to [43]. In Lyapunov–Schmidt reduction for problems like (1.3) the situation of adjusting a finite number of points (*finite dimensional Lyapunov–Schmidt reduction method*), and that of adjusting a higher dimensional object such a geodesic in a suitable metric as limiting concentration sets (*infinite dimensional Lyapunov–Schmidt reduction method*) have been treated. In this work we develop an *intermediate Lyapunov–Schmidt reduction method*, which consists of the finite dimensional procedure for large number of reduced equations, which in the limit become an ODE system of limiting Jacobi-type operators (see (8.9) below). Treating the discrete problem needs a method, technically delicate, which we interpret as an intermediate procedure between the finite and the infinite dimensional one (see for instance [21, 22, 38] and references therein for the latter). The main difference between the intermediate and infinite dimensional reduction, is that in the latter procedure only the variations in the normal direction are needed so the usual Jacobi operator for a curve appears. In the former procedure we also need to take into account variations in the tangential direction of points, which in the limit may be interpreted as a reparametrization of the curve. This seems to be a new procedure, with potentially many interesting applications.

Our main result removes the symmetry assumption on V when $N = 2$.

Theorem 1.1 *Let $N = 2$. Suppose that $V(x)$ satisfies (1.5) and (1.11) for some constants $V_\infty > 0, a > 0$, and*

$$\min \left\{ 1, \frac{p-1}{2} \right\} m > 2, \quad \sigma > 2. \tag{1.14}$$

Then problem (1.3) has infinitely many non-radial positive solutions, whose energy can be made arbitrarily large.

If (1.11) holds in the C^1 sense, then “ $\sigma > 2$ ” in (1.14) can be improved to be “ $\sigma > 1$ ”. The condition on p can be further relaxed if we assume more regularity of the condition (1.11) or if p is an integer.

The results in [4, 14] make us think that condition (1.11) could be improved. In fact we believe that the optimal condition should be (1.12). We stress that our result does not require a global perturbative assumption such as (1.13) on V . In addition, it is worth pointing out that the results on the existence of positive solutions in [6, 7, 9] do not include the polynomial decay case (1.11).

Finally, we remark that for $N \geq 3$, Theorem 1.1 holds if we assume the following additional symmetry assumption on V : after suitably rotating the coordinate system,

$$V(x) = V(x', x'') = V(x', -x''), \tag{1.15}$$

where $x = (x', x'') \in \mathbb{R}^2 \times \mathbb{R}^{N-2}$. An open question is whether or not the same result holds when $N \geq 3$ with no extra assumption made.

Throughout the paper, we shall use the following notation and conventions:

- For quantities A_K and B_K , we write $A_K \sim B_K$ to denote that there exists a positive constant C such that $1/C \leq A_K/B_K \leq C$ for K sufficiently large; $A_K = O(B_K)$ means that $|A_K/B_K|$ is uniformly bounded as K tends to infinity; $A_K = o(B_K)$ denotes that $|A_K/B_K| \rightarrow 0$ as $K \rightarrow \infty$.
- For simplicity, the letter C denotes various generic constant which is independent of K . It is allowed to vary from line to line, and also within the same formula.
- We will use the same $|y| = \|y\|_2$ for the Euclidean norm in various Euclidean spaces \mathbb{R}^N when no confusion can arise and we always denote the inner product of a and b in \mathbb{R}^N by $a \cdot b$.
- For the index $j \in \{1, 2, \dots, K\}$, we shall always use the convention that $j - 1 = K$ if $j = 1$ and $j + 1 = 1$ if $j = K$.
- The cardinality of a finite set E will be denoted by $\#E$; The Lebesgue measure of a set $E \subset \mathbb{R}^N$ will be denoted by $|E|$.
- The transpose of a matrix A will be denoted by A^T .
- For each function $w(x)$ defined in \mathbb{R}^N , if w is radially symmetric, then there is a real function $\tilde{w}(r)$ such that $w(x) = \tilde{w}(|x|)$. With slight abuse of notation, we will simply write $w(r)$ instead of $\tilde{w}(r)$.

In the next section, we will describe the procedure of our construction and give the main ideas of each step.

2 Description of the construction

We will prove the slightly more general version of the Theorem 1.1, for $N \geq 3$ where we assume even symmetry in $N - 2$ the remaining variables in the sense that

$$V(x) = V(x', x'') = V(x', -x''), \tag{2.1}$$

where $x = (x', x'') \in \mathbb{R}^2 \times \mathbb{R}^{N-2}$. We assume this henceforth.

We shall briefly describe the solutions to be constructed later and will give the main ideas in the procedure of the construction. In particular, we shall introduce the *intermediate reduction method*, which we believe can be very useful in other contexts.

Firstly, without loss of generality, we can assume that $V_\infty = 1$ by suitable scaling. As developed in [43], we will use the loss of compactness to build up solutions. More precisely, we will construct solutions with large number of spikes whose inter-distances and distances from the origin are sufficiently large.

By the asymptotic behaviour of V at infinity, the basic building block is the ground state (radial) solution w of the limit problem at infinity:

$$\begin{cases} -\Delta w + w - w^p = 0, & w > 0 \text{ in } \mathbb{R}^N, \\ w = w(|x|), & w \in H^1(\mathbb{R}^N). \end{cases} \tag{2.2}$$

The solutions we construct will be small perturbations of the sum of copies of w , centered at some carefully chosen points on $\mathbb{R}^2 \times \{0\} \subset \mathbb{R}^N$, where 0 is the zero vector in \mathbb{R}^{N-2} .

Let $K \in \mathbb{N}_+$ be the number of spikes, whose locations are given by $Q_j \in \mathbb{R}^N, j = 1, \dots, K$. We define

$$w_{Q_j}(x) = w(x - Q_j) \text{ and } U(x) = \sum_{j=1}^K w_{Q_j}(x), \text{ for } x \in \mathbb{R}^N. \tag{2.3}$$

A natural and central question is how to choose Q_j 's such that a small perturbation of U will be a genuine solution.

Assuming that

$$\inf_{1 \leq j \leq K} |Q_j| \rightarrow +\infty \text{ and } \inf_{j \neq l} |Q_j - Q_l| \rightarrow +\infty,$$

by the asymptotic behaviour of V at infinity and the property of w , one can get (at least formally) the following energy expansion

$$\mathcal{E}(U) = \underbrace{K I_0 + a_0 \sum_{j=1}^K |Q_j|^{-m} - \frac{1}{2} \gamma_0 \sum_{j \neq l} w(|Q_j - Q_l|)}_{J(Q_1, \dots, Q_K)} + \text{other terms}, \tag{2.4}$$

where I_0, a_0 and γ_0 are positive constants. Here $\mathcal{E}(U)$ is the energy functional defined at (1.4) and we denote the leading order expansion as $J(Q_1, \dots, Q_K)$.

Observe that for any rotation R_θ around the origin in \mathbb{R}^N , there holds

$$J(R_\theta Q_1, \dots, R_\theta Q_K) = J(Q_1, \dots, Q_K).$$

Hence any critical point of $J(Q_1, \dots, Q_K)$ is *degenerate*. Therefore, except in the symmetric class, it is not easy to find critical points of small perturbations of $J(Q_1, \dots, Q_K)$. This means that it is not easy to apply the localized energy method directly. However, this observation

gives us some enlightenment in the non-symmetric setting. Actually, when we restrict Q_j 's on a plane, this suggests us to introduce one more parameter to deal with the degeneracy due to rotations as we will see in Sect. 5.

Under the condition (1.15), there is no essential difference between $N = 2$ and $N \geq 3$. Hence from now on we will restrict Q_j 's on the plane $\mathbb{R}^2 \times \{0\} \subset \mathbb{R}^N$. To describe further the configuration space of Q_j 's, we define

$$Q_j^0 = (R \cos \theta_j, R \sin \theta_j, 0) \in \mathbb{R}^2 \times \{0\}, \quad \text{for } j = 1, \dots, K,$$

where

$$\theta_j = \alpha + (j - 1) \frac{2\pi}{K} \in \mathbb{R}.$$

Here α is the parameter representing the degeneracy due to rotations, and R is a positive constant to be determined later. Observe that each point Q_j^0 depends on α . Thus we write $Q_j^0 = Q_j^0(\alpha)$. When $\alpha = 0$, the Q_j^0 's are the points used in [43]. If $V(x)$ is radially symmetric, it is obvious that the parameter α plays no role in the construction in [43]. But it is very important in our construction as we will see in Sect. 6.

For the constant R , we introduce the so-called balancing condition:

$$a_0 m R^{-m-1} = 2 \sin \frac{\pi}{K} \Psi \left(2R \sin \frac{\pi}{K} \right), \tag{2.5}$$

where $a_0 = \frac{a}{2} \int_{\mathbb{R}^N} w^2 dx > 0$, and Ψ is the interaction function defined by

$$\Psi(s) = - \int_{\mathbb{R}^N} w(x - s\vec{e}) \operatorname{div} (w^p(x)\vec{e}) dx. \tag{2.6}$$

Here \vec{e} can be any unit vector in \mathbb{R}^N (cf. [34,35]). The balancing condition (2.5) can either be understood as a consequence of a conservation law or can be seen as a condition such that the approximation U is very close to a genuine solution (cf. Appendix in [35]). Assuming that

$$d = 2R \sin \frac{\pi}{K} \rightarrow +\infty, \quad \text{as } K \rightarrow +\infty,$$

we will see that (cf. Lemma 3.3)

$$|Q_j^0| = R \sim \frac{m}{2\pi} K \ln K, \quad \text{and } \min_{j \neq l} \left\{ |Q_j^0 - Q_l^0| \right\} = d \sim m \ln K.$$

Next we define a small neighbourhood of $Q^0 = (Q_1^0, \dots, Q_K^0)$ on $(\mathbb{R}^2 \times \{0\})^K$ in a suitable norm to be made precise and introduce another parameter. Let $f_j, g_j \in \mathbb{R}, j = 1, \dots, K$, we define

$$Q_j = Q_j^0 + f_j \vec{n}_j + g_j \vec{t}_j = (R + f_j) \vec{n}_j + g_j \vec{t}_j, \tag{2.7}$$

where

$$\vec{n}_j = (\cos \theta_j, \sin \theta_j, 0), \quad \text{and } \vec{t}_j = (-\sin \theta_j, \cos \theta_j, 0).$$

Keep in mind that f_j and g_j measure the displacement in the normal and tangential directions, respectively.

Writing $Q_j = Q_j(\alpha), \vec{n}_j = \vec{n}_j(\alpha)$ and $\vec{t}_j = \vec{t}_j(\alpha)$, we note the following trivial but important fact:

$$Q_j(\alpha + 2\pi) = Q_j(\alpha), \quad \forall \alpha \in \mathbb{R}, \quad \text{and } \forall j = 1, \dots, K. \tag{2.8}$$

We can now introduce another parameter \mathbf{q} and define a suitable norm. Denote

$$\mathbf{q} = (f_1, \dots, f_K, g_1, \dots, g_K)^T \in \mathbb{R}^{2K},$$

that is, $q_j = f_j$ and $q_{K+j} = g_j$ for $j = 1, \dots, K$. We define

$$\dot{\mathbf{q}} = (\dot{f}_1, \dots, \dot{f}_K, \dot{g}_1, \dots, \dot{g}_K)^T, \quad \text{and} \quad \ddot{\mathbf{q}} = (\ddot{f}_1, \dots, \ddot{f}_K, \ddot{g}_1, \dots, \ddot{g}_K)^T,$$

where for $j = 1, \dots, K$,

$$\begin{aligned} \dot{f}_j &= (f_{j+1} - f_j) \frac{K}{2\pi}, & \ddot{f}_j &= (f_{j+1} - 2f_j + f_{j-1}) \frac{K^2}{4\pi^2}, \\ \dot{g}_j &= (g_{j+1} - g_j) \frac{K}{2\pi}, & \ddot{g}_j &= (g_{j+1} - 2g_j + g_{j-1}) \frac{K^2}{4\pi^2}, \\ f_{K+1} &= f_1, & f_0 &= f_K, & g_{K+1} &= g_1, & g_0 &= g_K. \end{aligned}$$

Observe that if $f_j = f(\theta_j)$ for some 2π periodic smooth function f , then \dot{f}_j is the forward difference of f and \ddot{f}_j is the 2nd order central difference of f .

With these notation, we can define the configuration space of Q_j 's by

$$\Lambda_K = \left\{ (Q_1, \dots, Q_K) \in (\mathbb{R}^2 \times \{0\})^K \mid Q_j \text{ is defined by (2.7) and } \|\mathbf{q}\|_* \leq 1 \right\},$$

where $\|\mathbf{q}\|_* = \|\mathbf{q}\|_\infty + \|\dot{\mathbf{q}}\|_\infty + \|\ddot{\mathbf{q}}\|_\infty$ is a norm on \mathbb{R}^{2K} . In the following, we assume that Q_j is defined by (2.7), the parameter $\alpha \in \mathbb{R}$ and the parameter \mathbf{q} satisfy

$$\|\mathbf{q}\|_* = \|\mathbf{q}\|_\infty + \|\dot{\mathbf{q}}\|_\infty + \|\ddot{\mathbf{q}}\|_\infty \leq 1. \tag{2.9}$$

For any $(Q_1, \dots, Q_K) \in \Lambda_K$, an easy computation shows that for $j = 1, \dots, K$,

$$|Q_j| = R + f_j + O(R^{-1}),$$

and

$$|Q_{j+1} - Q_j| = d + 2(f_j + \dot{g}_j) \frac{\pi}{K} + O(K^{-2}).$$

Define $\rho = \min_{j \neq l} \{|Q_j - Q_l|\}$, it follows that

$$\rho = d + O(K^{-1}), \quad \text{and} \quad \min_{j=1, \dots, K} \{|Q_j|\} = R + O(1). \tag{2.10}$$

We will prove Theorem 1.1 by showing the following result.

Theorem 2.1 *Under the assumption of Theorem 1.1, there is a positive integer K_0 such that: for all integer $K \geq K_0$, there exist $\alpha \in [0, 2\pi)$ and $(Q_1, \dots, Q_K) \in \Lambda_K$ such that problem (1.3) has two solutions of the form*

$$u(x) = \sum_{j=1}^K w(x - Q_j) + \phi(x), \tag{2.11}$$

where $\|\phi\|_{H^1(\mathbb{R}^N)} + \|\phi\|_{L^\infty(\mathbb{R}^N)} \rightarrow 0$ as $K \rightarrow +\infty$. Moreover, the energy of u is given by

$$\mathcal{E}(u) = K \left(\frac{1}{2} - \frac{1}{p+1} \right) \int_{\mathbb{R}^N} w^{p+1} dx + o(1). \tag{2.12}$$

Remark 1 It is worth pointing out that the solutions constructed in this paper are different from those found in [4, 14]. The reason is simply that the inter-distances and distances from the origin of the spikes of the solutions given in (2.11) tend to infinity uniformly as K goes to infinity, but those of the solutions found in [4, 14] do not.

Remark 2 The fact that we can find at least **two** solutions of the form (2.11) is nontrivial. This is due to the fact that we need to choose the **first** starting point Q_1^0 . It turns out that there are at least two such points to choose (see Sect. 6).

Remark 3 As $K \rightarrow +\infty$, (f_j, g_j) is the discretization of two second order ordinary differential equations (8.9).

To prove Theorem 2.1, it is sufficient to show that for K sufficiently large there are parameters α and \mathbf{q} such that $U + \phi$ is a genuine solution for a small perturbation ϕ . To achieve this goal, we will adopt the techniques in the singularly perturbed problem. Unlike problem (1.9), there is no apparent parameter in (1.3). As stated in Theorem 2.1, we use the number of the spikes as the ε -type parameter. This idea comes directly from [43] and goes back at least as far as to [30].

Before we sketch the procedure of our proof, we briefly introduce the abstract set-up of the Lyapunov–Schmidt reduction (although it is always used in a framework that occurs often in bifurcation theory).

Let X, Y be Banach spaces and $S(u)$ is a C^1 map from X to Y . To study the equation $S(u) = 0$, a natural way is to find approximations first and then to look for genuine solutions as (small) perturbations of approximations. Assume that U_λ are the approximations, where $\lambda \in \Lambda$ is the parameter (we think of Λ as the configuration space). Writing $u = U_\lambda + \phi$, then solving $S(u) = 0$ amounts to solve

$$L[\phi] + E + N(\phi) = 0, \tag{2.13}$$

where

$$L[\phi] = S'(U_\lambda)[\phi], \quad E = S(U_\lambda), \quad \text{and} \quad N(\phi) = S(U_\lambda + \phi) - S(U_\lambda) - S'(U_\lambda)[\phi].$$

Here $S'(U_\lambda)$ is the Fréchet derivative of S at U_λ , E denotes the error of approximation, and $N(\phi)$ denotes the nonlinear term. In order to solve (2.13), we try to invert the linear operator L so that we can rephrase the problem as a fixed point problem. That is, when L has a uniformly bounded inverse in a suitable space, one can rewrite Eq. (2.13) as

$$\phi = -L^{-1}[E + N(\phi)] = \mathcal{A}(\phi).$$

What is left is to use fixed point theorems such as contraction mapping theorem.

The Lyapunov–Schmidt reduction deals with the situation when the linear operator L is Fredholm and its eigenfunction space associated to small eigenvalues has finite dimension. Assuming that $\{\mathcal{Z}_1, \dots, \mathcal{Z}_n\}$ is a basis of the eigenfunction space associated to small eigenvalues of L , we can divide the procedure of solving (2.13) into two steps:

- (i) solving the projected problem for any $\lambda \in \Lambda$,

$$\begin{cases} L[\phi] + E + N(\phi) = \sum_{j=1}^n c_j \mathcal{Z}_j, \\ \langle \phi, \mathcal{Z}_j \rangle = 0, \quad \forall j = 1, \dots, n, \end{cases}$$

where c_j may be constant or function depending on the form of $\langle \phi, \mathcal{Z}_j \rangle$;

(ii) solving the reduced problem

$$c_j(\lambda) = 0, \quad \forall j = 1, \dots, n,$$

by adjusting λ .

Let us now turn to our problem (1.3). In this case,

$$\begin{aligned} S(u) &= -\Delta u + V(x)u - u_+^p, \\ L[\phi] &= -\Delta\phi + V(x)\phi - pU^{p-1}\phi, \\ E &= -\Delta U + V(x)U - U^p, \\ N(\phi) &= -(U + \phi)_+^p + U^p + pU^{p-1}\phi. \end{aligned}$$

Observe that all of these quantities depend implicitly on α and \mathbf{q} even though this is not apparent in the notation.

By the Lyapunov–Schmidt reduction, the procedure of construction is made up of several steps which we explain next and postpone the proofs of major facts in later sections.

Step 1: Solving the projected problem

Let $\alpha \in \mathbb{R}$ and \mathbf{q} satisfy (2.9). We look for a function ϕ and some multiplier $\widehat{\beta} \in \mathbb{R}^{2K}$ such that

$$\begin{cases} L[\phi] + E + N(\phi) = \widehat{\beta} \cdot \frac{\partial U}{\partial \mathbf{q}}, \\ \int_{\mathbb{R}^N} \phi \mathcal{Z}_{Q_j} dx = 0, \quad \forall j = 1, \dots, K, \end{cases} \tag{2.14}$$

where the vector field \mathcal{Z}_{Q_j} is defined by

$$\mathcal{Z}_{Q_j}(x) = \nabla w(x - Q_j). \tag{2.15}$$

By direct computation, we have

$$\frac{\partial U}{\partial \mathbf{q}} = -(\mathcal{Z}_{Q_1} \cdot \vec{n}_1, \dots, \mathcal{Z}_{Q_K} \cdot \vec{n}_K, \mathcal{Z}_{Q_1} \cdot \vec{t}_1, \dots, \mathcal{Z}_{Q_K} \cdot \vec{t}_K)^T.$$

This is the first step in the Lyapunov–Schmidt reduction. It is done in Sect. 4 through some a priori estimates and contraction mapping theorem. A required element in this step is the non-degeneracy of w (cf. Lemma 3.1). It is worth pointing out that the function ϕ and the multiplier $\widehat{\beta}$ found in Step 1 depend on the parameters α and \mathbf{q} . Hence we write $\phi = \phi(x; \alpha, \mathbf{q})$ and $\widehat{\beta} = \widehat{\beta}(\alpha, \mathbf{q})$.

Step 2: Solving the reduced problem

By Step 1, it is known that $\widehat{\beta}$ is small. But it is not easy to solve $\widehat{\beta}(\alpha, \mathbf{q}) = 0$ directly since the linear part of the expansion of $\widehat{\beta}$ in \mathbf{q} is degenerate (due to the invariance of $J(Q_1, \dots, Q_K)$ under rotations).

More precisely, let us write

$$\widehat{\beta}(\alpha, \mathbf{q}) = \widetilde{T}\mathbf{q} + \Phi(\alpha, \mathbf{q}),$$

where $\widetilde{T}\mathbf{q}$ is the linear part and $\Phi(\alpha, \mathbf{q})$ denotes the remaining term. As we will see in Sect. 5, $\widetilde{T}\mathbf{q}$ does not depend on α and there is a unique vector (up to a scalar)

$$\mathbf{q}_0 = \underbrace{(0, \dots, 0)}_K, \underbrace{(1, \dots, 1)}_K)^T \in \mathbb{R}^{2K}$$

such that $\widetilde{T}\mathbf{q}_0 = 0$.

By the Lyapunov–Schmidt reduction again (called the secondary Lyapunov–Schmidt reduction), the step of solving the reduced problem $\widehat{\beta}(\alpha, \mathbf{q}) = 0$ can be divided into two steps. To write the projected problem of $\widehat{\beta} = 0$ in a proper form, note that

$$\frac{\partial U}{\partial \alpha} = R \sum_{j=1}^K \frac{\partial U}{\partial g_j} + \sum_{j=1}^K \left(f_j \frac{\partial U}{\partial g_j} - g_j \frac{\partial U}{\partial f_j} \right) = (R\mathbf{q}_0 + \mathbf{q}^\perp) \cdot \frac{\partial U}{\partial \mathbf{q}},$$

where $\mathbf{q}^\perp = (-\vec{g}, \vec{f})$ for $\mathbf{q} = (\vec{f}, \vec{g})$. Hence we define

$$\vec{\beta} = \widehat{\beta} - \gamma(R\mathbf{q}_0 + \mathbf{q}^\perp), \quad \text{for every } \gamma \in \mathbb{R}. \tag{2.16}$$

With this notation,

$$\widehat{\beta} \cdot \frac{\partial U}{\partial \mathbf{q}} = \vec{\beta} \cdot \frac{\partial U}{\partial \mathbf{q}} + \gamma \frac{\partial U}{\partial \alpha}.$$

Obviously the new multiplier $\vec{\beta}$ depends on the parameters α, \mathbf{q} and γ . Thus we write $\vec{\beta} = \vec{\beta}(\alpha, \mathbf{q}, \gamma)$.

Step 2.A: Solving $\vec{\beta}(\alpha, \mathbf{q}, \gamma) = 0$ by adjusting γ and \mathbf{q}

In this step, for each $\alpha \in \mathbb{R}$, we are going to find parameters (γ, \mathbf{q}) such that

$$\vec{\beta}(\alpha, \mathbf{q}, \gamma) = 0, \quad \text{and } \mathbf{q} \perp \mathbf{q}_0. \tag{2.17}$$

It can be seen as the step of solving the projected problem in the secondary Lyapunov–Schmidt reduction. To achieve it, we will use the condition (1.14). This step is done in Sect. 5 by using various integral estimates and contraction mapping theorem. A key element in this step is the invertibility of an $2K \times 2K$ matrix whose proof is given in Appendix A.

When Step 2.A is done, we denote the unique solution of (2.17) by $(\gamma(\alpha), \mathbf{q}(\alpha))$. Then the original problem (1.3) is reduced to the problem $\gamma(\alpha) = 0$ of one dimension.

Step 2.B: Solving $\gamma(\alpha) = 0$ by choosing α

At the last step, we want to prove that there exists an α such that $\gamma(\alpha) = 0$. As a result, the function $u = U + \phi$ is a genuine solution of (1.3).

This step is the second step of solving the reduced problem in the secondary Lyapunov–Schmidt reduction. To achieve this step, by Step 2.A, the function $\phi = \phi(x; \alpha, \mathbf{q}(\alpha))$ found in Step 1 solves the following problem:

$$\begin{cases} L[\phi] + E + N(\phi) = \gamma(\alpha) \frac{\partial U}{\partial \alpha}, \\ \int_{\mathbb{R}^N} \phi \mathcal{Z}_{Q_j} dx = 0, \quad \forall j = 1, \dots, K, \end{cases} \tag{2.18}$$

where all of the quantities depending implicitly on (α, \mathbf{q}) are taken values at $(\alpha, \mathbf{q}(\alpha))$. To solve $\gamma(\alpha) = 0$, we first apply the so-called variational reduction (often used in the localized energy method) to show that equation $\gamma(\alpha) = 0$ has a solution if the reduced energy function $F(\alpha) = \mathcal{E}(U + \phi)$ has a critical point. Secondly, by using (2.8), it is easy to check that $F(\alpha)$ is 2π periodic in α . Hence it has at least **two** critical points. More details of this step will be given in Sect. 6.

Finally, this paper is organized as follows. Some preliminary facts and estimates are explained in Sect. 3. In Sect. 4 we apply the standard Lyapunov–Schmidt reduction for Step 1. Sect. 5 contains a further reduction process for Step 2.A which reduces the original problem to a one-dimensional one. In Sect. 6 we carry out Step 2.B and then complete the proof of Theorem 2.1. At the last, we discuss some possible extensions in Sect. 7.

3 Preliminaries

In this section we present some preliminary facts and some useful estimates.

First we recall some basic and useful properties of the standard spike solution w defined by (2.2) and those of the interaction function Ψ defined in (2.6).

Lemma 3.1 *If $1 < p < \infty$ for $N = 2$ and $1 < p < \frac{N+2}{N-2}$ for $N \geq 3$, then every positive solution of the problem:*

$$\begin{cases} -\Delta u + u - u^p = 0 & \text{in } \mathbb{R}^N, \\ u > 0 & \text{in } \mathbb{R}^N, \quad u \in H^1(\mathbb{R}^N), \end{cases} \tag{3.1}$$

has the form $w(\cdot - Q)$ for some $Q \in \mathbb{R}^N$, where $w(x) = w(|x|) \in C^\infty(\mathbb{R}^N)$ is the unique positive radial solution which satisfies

$$\lim_{r \rightarrow +\infty} r^{\frac{N-1}{2}} e^r w(r) = c_{N,p}, \quad \lim_{r \rightarrow +\infty} \frac{w'(r)}{w(r)} = -1. \tag{3.2}$$

Here $c_{N,p}$ is a positive constant depending only on N and p . Furthermore, the Morse index of w is one and w is nondegenerate in the sense that

$$\text{Ker}(-\Delta + 1 - pw^{p-1}) \cap L^\infty(\mathbb{R}^N) = \text{Span}\{\partial_{x_1} w, \dots, \partial_{x_N} w\}.$$

Proof This result is well known. For the proof we refer the reader to [12,40] for the existence, [26] for the symmetry, [29] for the uniqueness, Appendix C in [36] for the nondegeneracy, and [8] for the Morse index.

Lemma 3.2 *For s sufficiently large,*

$$\Psi(s) = c_{N,p} s^{-\frac{N-1}{2}} e^{-s} (1 + O(s^{-1})), \tag{3.3}$$

where $c_{N,p} > 0$ is a constant depending only on N and p .

Proof This lemma follows from Taylor’s theorem and the Lebesgue dominated convergence theorem. We omit it here and refer to [27,34] for details.

Next we study the balancing condition (2.5). Assuming that

$$d = 2R \sin \frac{\pi}{K} \rightarrow +\infty, \quad \text{as } K \rightarrow +\infty,$$

by the expansion (3.3), both positive numbers R and d are uniquely determined by K . Moreover, we have the following expansions.

Lemma 3.3 *For K sufficiently large,*

$$d = m \ln K + \left(m - \frac{N-3}{2}\right) \ln(m \ln K) + O(1), \tag{3.4}$$

$$R = \frac{m}{2\pi} K \ln K + \frac{1}{2\pi} \left(m - \frac{N-3}{2}\right) K \ln(m \ln K) + O(K).$$

Proof From the balancing condition (2.5), the number d satisfies the equation

$$d^{m+1} \Psi(d) = a_0 m \left(2 \sin \frac{\pi}{K}\right)^m. \tag{3.5}$$

By using (3.3), for K sufficiently large, Eq. (3.5) becomes

$$c_{N,p} d^{m+1-\frac{N-1}{2}} e^{-d} (1 + O(d^{-1})) = a_0 m (2\pi)^m K^{-m} (1 + O(K^{-2})),$$

from which we obtain

$$d^{m+1-\frac{N-1}{2}} e^{-d} \sim K^{-m}.$$

Let $d = m \ln K + d_1$ with $d_1 = o(\ln K)$, we have

$$c_{N,p} (m \ln K)^{m+1-\frac{N-1}{2}} e^{-d_1} (1 + o(1)) = a_0 m (2\pi)^m (1 + O(K^{-2})).$$

It follows that

$$e^{d_1} \sim (m \ln K)^{m+1-\frac{N-1}{2}}.$$

Therefore,

$$d_1 = \left(m - \frac{N-3}{2} \right) \ln(m \ln K) + O(1),$$

from which we get the expansions of d and R . □

In the next section, we will apply the Lyapunov–Schmidt reduction. After refinements by many authors working on the subject or on closely related problems, this type of argument is rather standard now. However technical difficulties arise when the number of spikes goes to infinity or the number of spikes is infinity (cf. [34]). To deal with these difficulties, the following lemmas are useful.

Lemma 3.4 *There exists a constant C_N depending only on N such that for any $K \in \mathbb{N}_+$ and any $\mathbf{Q} = (Q_1, \dots, Q_K) \in (\mathbb{R}^N)^K$,*

$$\#\{Q_j \mid \ell\rho/2 \leq |Q_j - x| < (\ell + 1)\rho/2\} \leq C_N (\ell + 1)^{N-1}, \tag{3.6}$$

for all $x \in \mathbb{R}^N$ and all $\ell \in \mathbb{N}$, where $\rho = \min_{j \neq l} \{|Q_j - Q_l|\}$. In particular, if $\mathbf{Q} = (Q_1, \dots, Q_K) \in (\mathbb{R}^2 \times \{0\})^K$, then for all $x \in \mathbb{R}^N$ and all $\ell \in \mathbb{N}$,

$$\#\{Q_j \mid \ell\rho/2 \leq |Q_j - x| < (\ell + 1)\rho/2\} \leq 6(\ell + 1). \tag{3.7}$$

Proof When $\rho = 0$, the result is trivial. It remains to consider the case $\rho > 0$. For $\ell = 0$, it suffices to take $C_N \geq 1$. For $\ell \geq 1$, let Q_{j_k} 's ($k = 1, \dots, n$) be the points satisfying

$$\ell\rho/2 \leq |Q_{j_k} - x| < (\ell + 1)\rho/2.$$

By the triangle inequality, we have

$$(\ell - 1)\rho/2 \leq |y - x| < (\ell + 2)\rho/2, \quad \forall y \in B_{\rho/2}(Q_{j_k}).$$

Hence for all $k = 1, \dots, n$,

$$B_{\rho/2}(Q_{j_k}) \subset B_{(\ell+2)\rho/2}(x) \setminus B_{(\ell-1)\rho/2}(x).$$

Since $B_{\rho/2}(Q_{j_l}) \cap B_{\rho/2}(Q_{j_k}) = \emptyset$ for $l \neq k$, we conclude that

$$\sum_{k=1}^n |B_{\rho/2}(Q_{j_k})| \leq |B_{(\ell+2)\rho/2}(x) \setminus B_{(\ell-1)\rho/2}(x)|.$$

Therefore, taking $C_N = \sup_{\ell \in \mathbb{N}_+} \frac{(\ell+2)^N - (\ell-1)^N}{(\ell+1)^{N-1}}$, we have

$$n \leq (\ell + 2)^N - (\ell - 1)^N \leq C_N(\ell + 1)^{N-1},$$

which implies (3.6).

If $\mathbf{Q} = (Q_1, \dots, Q_K) \in (\mathbb{R}^2 \times \{0\})^K$, the above argument implies that

$$\sum_{k=1}^n |B_{\rho/2}(Q_{jk}) \cap \mathbb{R}^2 \times \{0\}| \leq |B_{(\ell+2)\rho/2}(x) \setminus B_{(\ell-1)\rho/2}(x) \cap \mathbb{R}^2 \times \{0\}|,$$

which implies that

$$n \leq (\ell + 2)^2 - (\ell - 1)^2 \leq 6(\ell + 1).$$

Therefore, we get the estimate (3.7) if we restrict $\mathbf{Q} = (Q_1, \dots, Q_K)$ on $(\mathbb{R}^2 \times \{0\})^K$. \square

Given $\mathbf{Q} = (Q_1, \dots, Q_K) \in (\mathbb{R}^N)^K$ with $\rho = \min_{j \neq l} \{|Q_j - Q_l|\} > 0$, for any $\ell \in \mathbb{N}$, we divide \mathbb{R}^N into $K + 1$ parts:

$$\Omega_j^\ell = \left\{ x \in \mathbb{R}^N \mid |x - Q_j| = \min_{1 \leq l \leq K} |x - Q_l| \leq \ell\rho/2 \right\}, \quad \forall j = 1, \dots, K,$$

and $\Omega_{K+1}^\ell = \mathbb{R}^N \setminus \cup_{j=1}^K \Omega_j^\ell$. Then the interior of $\Omega_j^\ell \cap \Omega_l^\ell$ is an empty set for $j \neq l$.

Lemma 3.5 *Suppose that $\Gamma(r)$ is a positive decreasing function defined on $[0, \infty)$ such that for some $b \in \mathbb{R}$ and $\eta > 0$,*

$$\Gamma(r) \sim r^b e^{-\eta r} \quad \text{as } r \rightarrow +\infty. \tag{3.8}$$

Then there exist positive constants ρ_0 and C (independent of K) such that

- (i) *for all $K, \ell \in \mathbb{N}_+$, all $(Q_1, \dots, Q_K) \in (\mathbb{R}^N)^K$ with $\rho \geq \rho_0$, and all $x \in \Omega_{j_0}^\ell$ ($j_0 = 1, \dots, K$), we have*

$$\sum_{j=1}^K \Gamma(|x - Q_j|) \leq C \ell^{N-1} \Gamma(|x - Q_{j_0}|). \tag{3.9}$$

In particular, if $(Q_1, \dots, Q_K) \in (\mathbb{R}^2 \times \{0\})^K$, then

$$\sum_{j=1}^K \Gamma(|x - Q_j|) \leq C \ell \Gamma(|x - Q_{j_0}|).$$

- (ii) *for all $(Q_1, \dots, Q_K) \in (\mathbb{R}^N)^K$ with $\rho \geq \rho_0$ and all $j_0 \in \{1, \dots, K\}$,*

$$\sum_{j \neq j_0} \Gamma(|Q_{j_0} - Q_j|) \leq C \Gamma(\rho). \tag{3.10}$$

Remark 4 *A similar result holds when $\Gamma(r)$ has polynomial decay. For example, if for some integer $n \in \mathbb{N}_+$,*

$$\Gamma(r) \sim r^b \text{ as } r \rightarrow +\infty, \quad \text{where } b < -n,$$

then there are positive constants ρ_0 and C (independent of K) such that for all $K \in \mathbb{N}_+$, all $(Q_1, \dots, Q_K) \in (\mathbb{R}^n \times \{0\})^K$ with $\rho \geq \rho_0$, and all $j_0 \in \{1, \dots, K\}$,

$$\sum_{j \neq j_0} \Gamma(|Q_j - Q_{j_0}|) \leq C \Gamma(\rho).$$

It seems that this kind of property will be useful and important in the construction of infinitely many solutions of problem with critical growth.

Proof Given $x \in \Omega_{j_0}^\ell$, by definition we have

$$|x - Q_{j_0}| \leq \ell\rho/2 \quad \text{and} \quad |x - Q_j| \leq |x - Q_{j_0}| + \rho, \quad \forall j = 1, \dots, K.$$

Thus there is an integer $0 \leq \ell_0 \leq \ell$ such that

$$\ell_0\rho/2 \leq |x - Q_{j_0}| < (\ell_0 + 1)\rho/2.$$

By the property of $\Gamma(r)$ and Lemma 3.4, for ρ sufficiently large, we have

$$\begin{aligned} \sum_{j=1}^K \Gamma(|x - Q_j|) &\leq C_N(\ell_0 + 1)^{N-1} \Gamma(|x - Q_{j_0}|) + C_N \sum_{s=\ell_0+1}^{+\infty} (s + 1)^{N-1} \Gamma(s\rho/2) \\ &\leq C_N(\ell_0 + 1)^{N-1} \Gamma(|x - Q_{j_0}|) + C(\ell_0 + 2)^{N-1} \Gamma((\ell_0 + 1)\rho/2) \\ &\leq C\ell^{N-1} \Gamma(|x - Q_{j_0}|), \end{aligned}$$

where in the second inequality we use the following inequality:

$$\sum_{s=\ell_0+1}^{+\infty} \frac{(s + 1)^{N-1}}{(\ell_0 + 2)^{N-1}} \frac{\Gamma(s\rho/2)}{\Gamma((\ell_0 + 1)\rho/2)} \leq C.$$

To prove it, for ρ sufficiently large, by (3.8), we have

$$\frac{\Gamma(s\rho/2)}{\Gamma((\ell_0 + 1)\rho/2)} \leq C \left(\frac{s}{\ell_0 + 1} \right)^b e^{-\eta(s-\ell_0-1)\rho/2}.$$

Hence

$$\begin{aligned} \sum_{s=\ell_0+1}^{+\infty} \frac{(s + 1)^{N-1}}{(\ell_0 + 2)^{N-1}} \frac{\Gamma(s\rho/2)}{\Gamma((\ell_0 + 1)\rho/2)} &\leq C \sum_{s=\ell_0+1}^{+\infty} \left(\frac{s}{\ell_0 + 1} \right)^{N-1+b} e^{-\eta(s-\ell_0-1)\rho/2} \\ &\leq C \sum_{s=\ell_0+1}^{+\infty} e^{-\eta(s-\ell_0-1)\rho/4} \\ &\leq C \int_0^{+\infty} e^{-\eta t\rho/4} dt \leq C. \end{aligned}$$

In particular, if $(Q_1, \dots, Q_K) \in (\mathbb{R}^2 \times \{0\})^K$, then by (3.7), we can take $N = 2$ in the above argument.

To deduce (3.10) from (3.9), denote

$$\widehat{Q} = (Q_1, \dots, Q_{j_0-1}, Q_{j_0+1}, \dots, Q_K) \in (\mathbb{R}^N)^{K-1},$$

and

$$\widehat{\rho} = \min_{j \neq l} \{|Q_j - Q_l| \mid j \neq j_0, l \neq j_0\} \geq \rho.$$

Take $j_1 \in \{1, \dots, K\}$ such that

$$|Q_{j_0} - Q_{j_1}| = \min_{l \neq j_0} \{|Q_{j_0} - Q_l|\},$$

and choose $\ell \in \mathbb{N}_+$ satisfying

$$(\ell - 1)\widehat{\rho}/2 < |Q_{j_0} - Q_{j_1}| \leq \ell\widehat{\rho}/2.$$

Then by (3.9), we have

$$\begin{aligned} \sum_{j \neq j_0} \Gamma(|Q_{j_0} - Q_j|) &\leq C \left(1 + \frac{2|Q_{j_0} - Q_{j_1}|}{\widehat{\rho}}\right)^{N-1} \Gamma(|Q_{j_0} - Q_{j_1}|) \\ &\leq C\Gamma(|Q_{j_0} - Q_{j_1}|) + C\Gamma(\widehat{\rho}) \leq C\Gamma(\rho), \end{aligned}$$

where in the second inequality we use the following inequality:

$$|Q_{j_0} - Q_{j_1}|^{N-1} \Gamma(|Q_{j_0} - Q_{j_1}|) \leq C\widehat{\rho}^{N-1} \Gamma(\widehat{\rho}), \quad \text{for } |Q_{j_0} - Q_{j_1}| \geq \widehat{\rho} \geq \rho_0.$$

To prove it, we only need to apply (3.8). □

A simple corollary is the following result which is useful in our construction.

Corollary 3.6 *There are positive constants ρ_0 and C (independent of K) such that for all $K, \ell \in \mathbb{N}_+$, all $(Q_1, \dots, Q_K) \in (\mathbb{R}^2 \times \{0\})^K$ with $\rho \geq \rho_0$, and all $x \in \Omega_{j_0}^\ell$ ($j_0 = 1, \dots, K$), we have*

$$\sum_{j=1}^K w_{Q_j}(x) \leq C\ell w_{Q_{j_0}}(x), \tag{3.11}$$

and

$$\sum_{j \neq j_0} e^{-\eta|Q_j - Q_{j_0}|} \leq C e^{-\eta\rho}. \tag{3.12}$$

To analyze the interactions between spikes, we prove some estimates concerning convolution of functions with suitable exponential decays.

Lemma 3.7 *Given Γ_1, Γ_2 two positive continuous radial functions on \mathbb{R}^N with the following property:*

$$\Gamma_1(r) \sim r^{b_1} e^{-\eta_1 r}, \quad \text{and} \quad \Gamma_2(r) \sim r^{b_2} e^{-\eta_2 r}, \quad \text{as } r \rightarrow +\infty,$$

where $b_1, b_2 \in \mathbb{R}, \eta_1 > 0, \eta_2 > 0$. Let $\xi \in \mathbb{R}^N$ tends to infinity. Then, the following asymptotic estimates hold:

(i) *If $\eta_1 < \eta_2$, then*

$$\int_{\mathbb{R}^N} \Gamma_1(x - \xi) \Gamma_2(x) dx \sim |\xi|^{b_1} e^{-\eta_1 |\xi|}.$$

Clearly, if $\eta_1 > \eta_2$, a similar expression holds, by replacing b_1 and η_1 with b_2 and η_2 .

(ii) *If $\eta_1 = \eta_2$, suppose that $b_1 \geq b_2$ for simplicity. Then*

$$\int_{\mathbb{R}^N} \Gamma_1(x - \xi) \Gamma_2(x) dx \sim \begin{cases} |\xi|^{b_1+b_2+\frac{N+1}{2}} e^{-\eta_1 |\xi|}, & \text{if } b_2 > -\frac{N+1}{2}, \\ (|\xi|^{b_1} \ln |\xi|) e^{-\eta_1 |\xi|}, & \text{if } b_2 = -\frac{N+1}{2}, \\ |\xi|^{b_1} e^{-\eta_1 |\xi|}, & \text{if } b_2 < -\frac{N+1}{2}. \end{cases}$$

Proof This result follows from the Lebesgue dominated convergence theorem. The argument is standard and is omitted here, we refer the interested reader to Lemma 3.7 in [2] for details. \square

By the property of w , as a corollary of Lemma 3.7, we have the following integral estimates.

Lemma 3.8 *Suppose that $|Q_j - Q_k|$ is sufficiently large, then the following estimates hold:*

(i) *for every $p > 1$,*

$$\int_{\mathbb{R}^N} w_{Q_j} w_{Q_k}^p dx = (\gamma_0 + o(1))w(|Q_j - Q_k|),$$

where $\gamma_0 = \int_{\mathbb{R}^N} w^p(x)e^{-x_1} dx > 0$ is a constant;

(ii)

$$\int_{\mathbb{R}^N} w_{Q_j} w_{Q_k} dx = O(e^{-|Q_k - Q_j|} |Q_k - Q_j|^{-(N-3)/2});$$

(iii) *let $\Omega_k = \{x \in \mathbb{R}^N \mid |x - Q_k| = \min_{1 \leq j \leq K} |x - Q_j|\}$, then*

$$\int_{\Omega_k} w_{Q_j}^p w_{Q_k} dx = O\left(e^{-\frac{p+1}{2}|Q_j - Q_k|} |Q_j - Q_k|^{-\frac{N-3}{2}}\right),$$

and

$$\int_{\Omega_k} w_{Q_j}^2 w_{Q_k}^{p-1} dx = O\left(e^{-\min\{2, \frac{p+1}{2}\}|Q_j - Q_k|} |Q_j - Q_k|^{-\frac{N-3}{2}}\right).$$

Proof Since the argument of proof is somewhat standard, we give only the main ideas of the proof.

- (i) It follows from Lemma 3.1 and Lebesgue’s dominated convergence theorem (see e.g. the arguments used in the proof of Lemma 2.5, in [30]).
- (ii) By Lemma 3.1 and a simple computation, we get the estimate from Lemma 3.7.
- (iii) By the definition, for all $x \in \Omega_k$, $w_{Q_j}(x) \leq w_{Q_k}(x)$ for every $1 \leq j \leq K$. Hence by Lemmas 3.1 and 3.7, we have

$$\int_{\Omega_k} w_{Q_j}^p w_{Q_k} dx \leq \int_{\Omega_k} w_{Q_j}^{\frac{p+1}{2}} w_{Q_k}^{\frac{p+1}{2}} dx \leq C e^{-\frac{p+1}{2}|Q_j - Q_k|} |Q_j - Q_k|^{-\frac{N-3}{2}},$$

and

$$\begin{aligned} \int_{\Omega_k} w_{Q_j}^2 w_{Q_k}^{p-1} dx &\leq C \int_{\Omega_k} w_{Q_j}^{\min\{2, \frac{p+1}{2}\}} w_{Q_k}^{\min\{2, \frac{p+1}{2}\}} dx \\ &\leq C e^{-\min\{2, \frac{p+1}{2}\}|Q_j - Q_k|} |Q_j - Q_k|^{-\frac{N-3}{2}}. \end{aligned}$$

\square

Using above integral estimates, we can get the expansion of the energy of approximate solution.

Lemma 3.9 For K sufficiently large, for any $\alpha \in \mathbb{R}$ and \mathbf{q} satisfying (2.9) we have

$$\begin{aligned} \mathcal{E}(U) &= KI_0 + (a_0 + o(1)) \sum_{j=1}^K |Q_j|^{-m} - \frac{1}{2} \sum_{i \neq j} (\gamma_0 + o(1)) w(|Q_i - Q_j|) \\ &\quad + O(KR^{-2m}) + O\left(Ke^{-\min\{2, \frac{p+1}{2}\}d} d^{-\frac{N-3}{2}}\right), \end{aligned}$$

where $\gamma_0 = \int_{\mathbb{R}^N} w^p(x)e^{-x_1} dx$ is a positive constant given in Lemma 3.8,

$$I_0 = \left(\frac{1}{2} - \frac{1}{p+1}\right) \int_{\mathbb{R}^N} w^{p+1} dx, \text{ and } a_0 = \frac{a}{2} \int_{\mathbb{R}^N} w^2 dx.$$

Proof The proof is delayed to Appendix B. □

4 The Lyapunov–Schmidt reduction

The aim of this section is to achieve Step 1 in the procedure of our construction described in Sect. 2.

Before stating the main result, we first introduce some notation. Let $\eta \in (0, 1)$ be a constant chosen later, we define the weighted norm:

$$\|h\|_{**} = \sup_{x \in \mathbb{R}^N} \left| \left(\sum_{j=1}^K e^{-\eta|x-Q_j|} \right)^{-1} h(x) \right|, \tag{4.1}$$

where Q_j is defined in (2.7). In what follows, we assume that $(Q_1, \dots, Q_K) \in \Lambda_K$, i.e., the parameter \mathbf{q} satisfies (2.9).

We first claim that

$$\|h\|_{L^\infty(\mathbb{R}^N)} \leq C \|h\|_{**} \text{ and } \|h\|_{L^q(\mathbb{R}^N)} \leq CK \|h\|_{**} \text{ for } 1 \leq q < \infty. \tag{4.2}$$

Indeed, the second inequality in (4.2) follows directly from

$$|h(x)| \leq \|h\|_{**} \sum_{j=1}^K e^{-\eta|x-Q_j|}, \quad \forall x \in \mathbb{R}^N.$$

To prove the first inequality in (4.2), it suffices to show that $\sum_{j=1}^K e^{-\eta|x-Q_j|} \leq C$. Indeed, for any $x \in \mathbb{R}^N$, we can choose $\ell \in \mathbb{N}_+$ and $j_0 \in \{1, \dots, K\}$ such that $x \in \Omega_{j_0}^\ell \setminus \Omega_{j_0}^{\ell-1}$. Hence by Lemma 3.5,

$$0 < \sum_{j=1}^K e^{-\eta|x-Q_j|} \leq C\ell^{N-1} e^{-\eta(\ell-1)\rho/2} \leq C. \tag{4.3}$$

Denote

$$\mathcal{B}_{**} = \left\{ h \in L^\infty(\mathbb{R}^N) \mid \|h\|_{**} < \infty \right\}.$$

Then \mathcal{B}_{**} is a Banach space with the norm $\|h\|_{**}$. To show the completeness, suppose that $\{h_n\}$ is a Cauchy sequence in \mathcal{B}_{**} . By (4.2), $\{h_n\}$ is also a Cauchy sequence in $L^\infty(\mathbb{R}^N)$. Hence h_n converges to a function h_∞ in $L^\infty(\mathbb{R}^N)$. By the definition of Cauchy sequence, for any $\varepsilon > 0$, there is $n_0 \in \mathbb{N}$ such that

$$|h_n(x) - h_k(x)| \left(\sum_{j=1}^K e^{-\eta|x-Q_j|} \right)^{-1} \leq \|h_n - h_k\|_{**} < \varepsilon, \quad \forall x \in \mathbb{R}^N, \quad \text{if } n, k \geq n_0.$$

Letting $k \rightarrow +\infty$, we get

$$|h_n(x) - h_\infty(x)| \left(\sum_{j=1}^K e^{-\eta|x-Q_j|} \right)^{-1} < \varepsilon, \quad \forall x \in \mathbb{R}^N, \quad \text{if } n \geq n_0,$$

which implies that $\|h_n - h_\infty\|_{**} \rightarrow 0$ as $n \rightarrow +\infty$.

Now we can state our main result in this section.

Proposition 4.1 *Suppose that $V(x)$ satisfies (1.11) for constants $V_\infty > 0, a \in \mathbb{R}, m \geq 1$ and $\sigma > 0$. If $N \geq 3$, we further assume (1.15). Then there is a positive integer K_0 such that: for all $K \geq K_0$, every $\alpha \in \mathbb{R}$, and \mathbf{q} satisfies (2.9), there exists a unique function $\phi \in W^{2,2}(\mathbb{R}^N) \cap \mathcal{B}_K$ and a unique multiplier $\widehat{\beta} \in \mathbb{R}^{2K}$ such that*

$$\begin{cases} L[\phi] + E + N(\phi) = \widehat{\beta} \cdot \frac{\partial U}{\partial \mathbf{q}}, \\ \int_{\mathbb{R}^N} \phi \mathcal{Z}_{Q_j} dx = 0, \quad \forall j = 1, \dots, K, \end{cases} \tag{4.4}$$

where

$$\mathcal{B}_K = \left\{ \phi \in L^\infty(\mathbb{R}^N) \mid \|\phi\|_{**} \leq C_0 K^{-\min\{1, \frac{p-\eta}{2}\}m} (\ln K)^{-\frac{1}{2}} \right\}.$$

Here C_0 is a positive constant independent of K . Moreover, $(\alpha, \mathbf{q}) \mapsto \phi(x; \alpha, \mathbf{q})$ is of class C^1 , and

$$R^{-1} \left\| \frac{\partial \phi}{\partial \alpha} \right\|_{**} + \left\| \frac{\partial \phi}{\partial \mathbf{q}} \right\|_{**} \leq C \left(K^{-\min\{1, \frac{p-\eta}{2}\}m} (\ln K)^{-\frac{1}{2}} \right)^{\min\{p-1, 1\}}.$$

The proof of Proposition 4.1 is somewhat standard and can be divided into two steps:

- (i) studying the invertibility of the linear operator;
- (ii) applying fixed point theorems.

4.1 Linear analysis

Let M denotes an $2K \times 2K$ matrix defined by

$$M_{jk} = \int_{\mathbb{R}^N} \frac{\partial U}{\partial q_j} \frac{\partial U}{\partial q_k} dx, \quad \forall j, k = 1, \dots, 2K. \tag{4.5}$$

Lemma 4.2 *Assume that $m \geq 1$. Then for K sufficiently large, given any vector $\vec{b} \in \mathbb{R}^{2K}$, there exists a unique vector $\widehat{\beta} \in \mathbb{R}^{2K}$ such that $M\widehat{\beta} = \vec{b}$. Moreover,*

$$\|\widehat{\beta}\|_\infty \leq C \|\vec{b}\|_\infty, \tag{4.6}$$

for some constant C independent of K .

Proof To prove the existence, it is sufficient to prove the a priori estimate (4.6). Suppose that $|\widehat{\beta}_j| = \|\widehat{\beta}\|_\infty$, by the definition, we have

$$\sum_{k=1}^K M_{jk} \widehat{\beta}_k = b_j. \tag{4.7}$$

For the entries M_{jk} , by Lemmas 3.3 and 3.8, we get

$$|M_{jk}| \leq C e^{-d} d^{-\frac{N-3}{2}} \leq CK^{-m} (m \ln K)^{-m}, \quad \forall k \neq j, \tag{4.8}$$

and

$$M_{jj} = \int_{\mathbb{R}^N} (\partial_{x_1} w)^2 dx = c_0 > 0, \quad \forall j = 1, \dots, 2K. \tag{4.9}$$

Hence by (4.7)–(4.9), for K sufficiently large, we have

$$c_0 \|\widehat{\beta}\|_\infty \leq c_0 |\widehat{\beta}_j| \leq \sum_{k \neq j} |M_{jk}| |\widehat{\beta}_k| + |b_j| \leq \frac{c_0}{2} \|\widehat{\beta}\|_\infty + \|\vec{b}\|_\infty,$$

from which the desired result follows. □

We can now formulate our main result in this subsection.

Lemma 4.3 *Under the assumption of Proposition 4.1, there is a positive integer K_0 such that: for all $K \geq K_0$, every $\alpha \in \mathbb{R}$, and \mathbf{q} satisfies (2.9), and for all $h \in \mathcal{B}_{**}$, there exists a unique function $\phi \in W^{2,2}(\mathbb{R}^N) \cap \mathcal{B}_{**}$ and a unique multiplier $\widehat{\beta} \in \mathbb{R}^{2K}$ such that*

$$\begin{cases} L[\phi] = h + \widehat{\beta} \cdot \frac{\partial U}{\partial \mathbf{q}}, \\ \int_{\mathbb{R}^N} \phi \mathcal{Z}_{Q_j} dx = 0, \quad \forall j = 1, \dots, K. \end{cases} \tag{4.10}$$

Moreover, we have

$$\|\phi\|_{**} + \|\widehat{\beta}\|_\infty \leq C \|h\|_{**}, \tag{4.11}$$

for some positive constant C independent of K .

Proof To solve (4.10), we first consider weak solutions. Define

$$\mathcal{H} = \left\{ u \in H^1(\mathbb{R}^N) \mid (u, (-\Delta + 1)^{-1} \mathcal{Z}_{Q_j}) = 0, \quad \forall j = 1, \dots, K \right\}.$$

Then \mathcal{H} is a Hilbert space with the standard inner product:

$$(u, v) = \int_{\mathbb{R}^N} (\nabla u \nabla v + uv) dx.$$

Since the vector function \mathcal{Z}_{Q_j} decays exponentially at infinity, by integration by parts, it is not hard to show that for $\phi \in H^1(\mathbb{R}^N)$, $\phi \in \mathcal{H}$ is equivalent to

$$\int_{\mathbb{R}^N} \phi \mathcal{Z}_{Q_j} dx = 0, \quad \forall j = 1, \dots, K.$$

As usual, $\phi \in \mathcal{H}$ is a weak solution of (4.10) if and only if it satisfies the following equation:

$$\int_{\mathbb{R}^N} \{ \nabla \phi \nabla \varphi + V(x) \phi \varphi - p U^{p-1} \phi \varphi \} dx = \int_{\mathbb{R}^N} h \varphi dx, \quad \forall \varphi \in \mathcal{H}.$$

By the Riesz representation theorem, the last equation can be written as

$$\phi + \mathcal{K}[\phi] = \widehat{h},$$

where \widehat{h} is defined by duality and \mathcal{K} is a linear compact operator due to the exponential decay of U and $|V(x) - 1| \leq C|x|^{-m}$ for $|x|$ large. Using the Fredholm alternative, showing that Eq. (4.10) has a unique weak solution is equivalent to showing that it has a unique solution for $h = 0$. Moreover, by (4.2), $h \in L^q(\mathbb{R}^N)$ for all $1 < q < \infty$. By the standard elliptic regularity results, $\phi \in W^{2,q}(\mathbb{R}^N)$. Hence ϕ is a strong solution and $\phi \in L^\infty(\mathbb{R}^N)$ by the Sobolev imbedding theorem. Therefore, to prove Lemma 4.3, it is sufficient to prove the a priori estimate (4.11).

To prove (4.11), we first multiply Eq. (4.10) by $\frac{\partial U}{\partial \mathbf{q}}$ and integrate over \mathbb{R}^N to obtain

$$M\widehat{\beta} = \int_{\mathbb{R}^N} L[\phi] \frac{\partial U}{\partial \mathbf{q}} dx - \int_{\mathbb{R}^N} h \frac{\partial U}{\partial \mathbf{q}} dx, \tag{4.12}$$

where M is an $2K \times 2K$ matrix defined in (4.5).

By the integration by parts,

$$\int_{\mathbb{R}^N} L[\phi] \mathcal{Z}_{Q_k} dx = \int_{\mathbb{R}^N} \phi L[\mathcal{Z}_{Q_k}] dx.$$

Observe that

$$L[\mathcal{Z}_{Q_k}] = (V(x) - 1) \nabla w_{Q_k} - p \left(U^{p-1} - w_{Q_k}^{p-1} \right) \nabla w_{Q_k}.$$

We claim that

$$\left| \int_{\mathbb{R}^N} L[\phi] \mathcal{Z}_{Q_k} dx \right| \leq Cde^{-\min\{1, \frac{p}{2}\}d} \|\phi\|_\infty. \tag{4.13}$$

Indeed, on one hand, by the assumption (1.11) and $\phi \in \mathcal{H}$, we have

$$\left| \int_{\mathbb{R}^N} (V(x) - 1) \nabla w_{Q_k} \phi dx \right| \leq C(R^{-m-1} \ln K + R^{-m-\sigma}) \|\phi\|_\infty.$$

On the other hand, by mean value theorem and (3.11), for $|x - Q_k| < 2m \ln K$, we have

$$|U^{p-1} - w_{Q_k}^{p-1}| \leq Cw_{Q_k}^{p-2} \sum_{j \neq k} w_{Q_j}.$$

Thus by Lemma 3.8,

$$\left| \int_{\mathbb{R}^N} -p \left(U^{p-1} - w_{Q_k}^{p-1} \right) \nabla w_{Q_k} \phi dx \right| \leq Cde^{-\min\{1, \frac{p}{2}\}d} \|\phi\|_\infty.$$

Combining the above estimates we get (4.13).

Since w decays exponentially at infinity, we have

$$\left| \int_{\mathbb{R}^N} h \mathcal{Z}_{Q_k} dx \right| \leq C \|h\|_{**}. \tag{4.14}$$

Combining the above estimates (4.13) and (4.14), by Lemma 4.2, we get

$$\|\widehat{\beta}\|_\infty \leq C \left(d e^{-\min\{1, \frac{\rho}{2}\}d} \|\phi\|_\infty + \|h\|_{**} \right). \tag{4.15}$$

Now we prove the a priori estimate (4.11). First we show that $\|\phi\|_{**} < \infty$. To prove it, by the maximum principle, we prove that there exist constants τ and C (both independent of K) such that for all $x \in \mathbb{R}^N \setminus \cup_{j=1}^K B(Q_j, \tau)$,

$$|\phi(x)| \leq C \left(\|L[\phi]\|_{**} + \sup_{1 \leq j \leq K} \|\phi\|_{L^\infty(B(Q_j, \tau))} \right) \sum_{j=1}^K e^{-\eta|x-Q_j|}. \tag{4.16}$$

To prove the above pointwise estimate, we first show the independence of τ on K , for $x \in \mathbb{R}^N \setminus \cup_{j=1}^K B(Q_j, \tau)$, by Lemma 3.4, we have

$$\begin{aligned} U(x) &\leq \sum_{|Q_j-x| < \rho/2} w(x-Q_j) + \sum_{\ell=1}^\infty \sum_{\ell\rho/2 \leq |Q_j-x| < (\ell+1)\rho/2} w(x-Q_j) \\ &\leq w(\tau) + C \sum_{\ell=1}^\infty \ell^{N-1} e^{-\ell\rho/2} \leq Cw(\tau). \end{aligned}$$

Thus we can take τ sufficiently large but independent of K such that

$$pU^{p-1}(x) \leq (V_0 - \eta^2)/4, \quad \forall x \in \mathbb{R}^N \setminus \cup_{j=1}^K B(Q_j, \tau). \tag{4.17}$$

Now we claim that for τ sufficiently large (independent of K), in $\mathbb{R}^N \setminus \cup_{j=1}^K B(Q_j, \tau)$,

$$L[W_-] \geq c_0 W_-, \quad \text{and} \quad L[W_+] \geq c_0 W_+$$

where $W_\pm(x) = \sum_{j=1}^K e^{\pm\eta|x-Q_j|}$ and $c_0 > 0$ is a constant independent of K . Indeed, for $x \in \mathbb{R}^N \setminus \cup_{j=1}^K B(Q_j, \tau)$,

$$L[W_\pm] = \sum_{j=1}^K \left\{ V(x) - \eta^2 \mp \frac{N-1}{|x-Q_j|} \eta - pU^{p-1} \right\} e^{\pm\eta|x-Q_j|} \geq \frac{V_0 - \eta^2}{2} W_\pm,$$

by the assumption (V1) and inequality (4.17).

The remaining part in the proof of (4.16) is to apply the maximum principle for the linear operator L in $\mathbb{R}^N \setminus \cup_{j=1}^K B(Q_j, \tau)$ to obtain

$$|\phi(x)| \leq C \left(\|L[\phi]\|_{**} + \sup_{1 \leq j \leq K} \|\phi\|_{L^\infty(B(Q_j, \tau))} \right) \sum_{j=1}^K e^{-\eta|x-Q_j|} + \delta \sum_{j=1}^K e^{\eta|x-Q_j|}$$

for any $\delta > 0$, where C is a constant independent of K and δ . Letting $\delta \rightarrow 0$, we get the desired estimate (4.16). Hence

$$\|\phi\|_{**} \leq C \left(\|L[\phi]\|_{**} + \sup_{1 \leq j \leq K} \|\phi\|_{L^\infty(B(Q_j, \tau))} \right) < \infty. \tag{4.18}$$

Now we can prove the a priori estimate (4.11). Arguing by contradiction, assume that there is a sequence of $(\phi^{(K)}, h^{(K)})$ satisfying (4.10) such that

$$\|\phi^{(K)}\|_{**} = 1, \quad \text{and} \quad \|h^{(K)}\|_{**} = o(1), \quad \text{as } K \rightarrow \infty.$$

(For simplicity, in the following we will drop “(K)” in the superscript.) As a consequence of (4.15),

$$\left| \widehat{\beta} \cdot \frac{\partial U}{\partial \mathbf{q}}(x) \right| \leq C \left(d e^{-\min\{1, \frac{p}{2}\}d} \|\phi\|_\infty + \|h\|_{**} \right) \sum_{j=1}^K e^{-\eta|x-Q_j|}.$$

Since $\|\phi\|_\infty \leq C\|\phi\|_{**}$ and $\|h\|_{**} = o(1)$, we get $\|L[\phi]\|_{**} = o(1)$. Hence (4.18) implies that there exists a subsequence of Q_j such that

$$\|\phi\|_{L^\infty(B(Q_j, \tau))} \geq C > 0 \tag{4.19}$$

for some fixed constant C (independent of K). Since $\|\phi\|_\infty \leq 1$, by elliptic regularity estimates, we get $\|\phi\|_{C^1(\mathbb{R}^N)} \leq C$. Applying Ascoli–Arzela’s theorem, one can find a subsequence of Q_j such that $\phi(x + Q_j)$ converge (on compact sets) to ϕ_∞ . It is not hard to show that ϕ_∞ is a bounded (weak and then strong) solution (actually bound by $e^{-\eta|x|}$) of

$$-\Delta\phi_\infty + \phi_\infty - p w^{p-1} \phi_\infty = 0.$$

Furthermore, since ϕ satisfies the orthogonality condition $\int_{\mathbb{R}^N} \phi \mathcal{Z}_{Q_j} dx = 0$, the limit function ϕ_∞ satisfies $\int_{\mathbb{R}^N} \phi_\infty \nabla w = 0$. By the non-degeneracy of w , one has $\phi_\infty \equiv 0$, which is in contradiction with (4.19). This completes the proof of Lemma 4.3. \square

Remark 5 If $V(x)$ is a bounded measurable function such that there is no nontrivial solution of

$$-\Delta\phi + V(x)\phi = 0, \quad |\phi(x)| \leq C e^{-\eta|x|} \quad \text{in } \mathbb{R}^N, \tag{4.20}$$

the above argument still work by adding 0 to the points Q_j ’s.

Remark 6 Since the Morse index of w is finite, using a similar argument in the proof of Lemma 4.3 (cf. [4]), one can show that

$$\|\phi\|_{H^1(\mathbb{R}^N)} \leq C \|h\|_{L^2(\mathbb{R}^N)} \tag{4.21}$$

for some positive constant C independent of K . Indeed, since τ is independent of K , one can first prove that

$$C \|\phi\|_{H^1(\mathbb{R}^N)}^2 \leq \int_{\mathbb{R}^N} \{ |\nabla\phi|^2 + V(x)\phi^2 - p U^{p-1} \phi^2 \} dx.$$

4.2 Nonlinear analysis

Summarizing, for any $h \in \mathcal{B}_{**}$, by Lemma 4.3, there is a unique function $\phi \in \mathcal{H} \cap W^{2,2}(\mathbb{R}^N) \cap \mathcal{B}_{**}$ satisfying (4.10). Hence we can define a linear operator from \mathcal{B}_{**} to $\mathcal{H} \cap W^{2,2}(\mathbb{R}^N) \cap \mathcal{B}_{**}$ and denote it by L^{-1} . Then Eq. (4.4) is equivalent to

$$\phi = -L^{-1} [E + N(\phi)].$$

Before we give the complete proof of Proposition 4.1, we first show the estimate of the error.

Lemma 4.4 *Given $(Q_1, \dots, Q_K) \in \Lambda_K$, then for any fixed $0 < \eta < 1$ and K sufficiently large, there is a constant C (independent of K) such that*

$$\|E\|_{**} \leq C K^{-\min\{1, \frac{p-\eta}{2}\}m} (\ln K)^{-\frac{1}{2}}. \tag{4.22}$$

Proof By the definition, we have

$$E = \underbrace{\sum_{j=1}^K (V(x) - 1) w_{Q_j}}_{E_1} - \underbrace{\left\{ \left(\sum_{j=1}^K w_{Q_j} \right)^p - \sum_{j=1}^K w_{Q_j}^p \right\}}_{E_2}.$$

Claim 1: There exists a constant C (independent of K) such that

$$\|E_1\|_{**} \leq CK^{-m} \leq CK^{-m} (\ln K)^{-m}. \tag{4.23}$$

Claim 2: There exists a constant C (independent of K) such that

$$\|E_2\|_{**} \leq Cd^{-\frac{N-1}{2}} e^{-\min\{1, \frac{p-\eta}{2}\}d} \leq CK^{-\min\{1, \frac{p-\eta}{2}\}m} (\ln K)^{-\min\{\frac{N-1}{2}, m+1\}}. \tag{4.24}$$

If both Claim 1 and Claim 2 are true, the desired estimate (4.22) follows.

Proof of Claim 1: Note that for $|x| < R/3$, by the triangle inequality, we have

$$|x - Q_j| \geq |Q_j| - |x| \geq R/2.$$

Hence for all $|x| < R/3$, by $V \in L^\infty(\mathbb{R}^N)$ and Lemma 3.1, we get

$$|E_1(x)| \leq C \sum_{j=1}^K w(x - Q_j) \leq Ce^{-(1-\eta)R/2} \sum_{j=1}^K e^{-\eta|x-Q_j|} \leq CK^{-m-3} \sum_{j=1}^K e^{-\eta|x-Q_j|}.$$

For $|x| \geq R/3$, by the assumption (1.11), we have $|V(x) - 1| \leq CK^{-m}$. Hence for all $|x| \geq R/3$,

$$|E_1(x)| \leq CK^{-m} \sum_{j=1}^K w(x - Q_j) \leq CK^{-m} \sum_{j=1}^K e^{-\eta|x-Q_j|}.$$

Combining these estimates, Claim 1 follows.

Proof of Claim 2: For $x \in \Omega_{K+1}^\ell$, where $\ell \in \mathbb{N}_+$ is chosen later, we have

$$\begin{aligned} |E_2(x)| &\leq K^{p-1} \sum_{j=1}^K w_{Q_j}^p(x) + \sum_{j=1}^K w_{Q_j}^p(x) \\ &\leq CK^{p-1} \sum_{j=1}^K e^{-p|x-Q_j|} \leq CK^{p-1} e^{-(p-\eta)\ell\rho/2} \sum_{j=1}^K e^{-\eta|x-Q_j|}. \end{aligned}$$

Since $\rho > \frac{m}{2} \ln K$, by choosing $\ell > \frac{4(p+m+2)}{m(p-1)}$ (independent of K), we have

$$K^{p-1} e^{-(p-\eta)\ell\rho/2} \leq K^{p-1} K^{-(p-\eta)\ell m/4} \leq CK^{-m-3}.$$

Thus for all $x \in \Omega_{K+1}^\ell$,

$$|E_2(x)| \leq CK^{-m-3} \sum_{j=1}^K e^{-\eta|x-Q_j|}.$$

By the definition of $\Omega_j^\ell, j = 1, \dots, K, |x - Q_j| \leq |x - Q_k|$ for all $x \in \Omega_j^\ell$ and $1 \leq k \leq K$. Thus $|x - Q_k| \geq \frac{\rho}{2}$ for $k \neq j$. Hence by mean value theorem and (3.11), for all $x \in \Omega_j^\ell$, we have

$$\begin{aligned} |E_2(x)| &\leq \left| \left(\sum_{k=1}^K w_{Q_k} \right)^p - w_{Q_j}^p \right| + \sum_{i \neq j} w_{Q_i}^p \leq p \left(\sum_{k=1}^K w_{Q_k} \right)^{p-1} \sum_{k \neq j} w_{Q_k} + \sum_{k \neq j} w_{Q_k}^p \\ &\leq C \ell^{(N-1)(p-1)} w_{Q_j}^{p-1} \sum_{k \neq j} w_{Q_k}. \end{aligned}$$

Since ℓ is independent of K , for all $x \in \Omega_j^\ell$, by Theorem 3.1 and (3.12) we have

$$\begin{aligned} |E_2(x)| &\leq C e^{-(p-1)|x-Q_j|} \sum_{k \neq j} \rho^{-\frac{N-1}{2}} e^{-|x-Q_k|} \\ &\leq C \rho^{-\frac{N-1}{2}} e^{-\eta|x-Q_j|} \sum_{k \neq j} e^{-\min\{1, \frac{p-\eta}{2}\}|Q_j-Q_k|} \\ &\leq C d^{-\frac{N-1}{2}} e^{-\min\{1, \frac{p-\eta}{2}\}d} e^{-\eta|x-Q_j|}. \end{aligned}$$

Combining these estimates, Claim 2 follows. □

Now we are in the position to give the proof of Proposition 4.1.

Proof of Proposition 4.1 Let C_0 be a positive number to be determined later, we define

$$\mathcal{B}_K = \left\{ \phi \in L^\infty(\mathbb{R}^N) : \|\phi\|_{**} \leq C_0 K^{-\min\{1, \frac{p-\eta}{2}\}m} (\ln K)^{-\frac{1}{2}} \right\}.$$

Then \mathcal{B}_K is a non-empty closed set in \mathcal{B}_{**} . Now we define a map $\mathcal{A} : \mathcal{B}_K \mapsto \mathcal{H} \cap W^{2,2}(\mathbb{R}^N) \cap \mathcal{B}_{**}$ by

$$\mathcal{A}(\phi) = -L^{-1} [E + N(\phi)].$$

Now solving Eq. (4.4) is equivalent to finding a fixed point for the map \mathcal{A} .

Since ϕ is uniformly bounded for $\phi \in \mathcal{B}_K$, by the mean value theorem, there is a positive constant C such that for all $\phi \in \mathcal{B}_K$,

$$|N(\phi)| \leq C |\phi|^{\min\{p,2\}},$$

and for all $\phi_1, \phi_2 \in \mathcal{B}_K$,

$$|N(\phi_1) - N(\phi_2)| \leq C (|\phi_1|^{\min\{p-1,1\}} + |\phi_2|^{\min\{p-1,1\}}) |\phi_1 - \phi_2|.$$

Thus by (4.3), one has

$$\|N(\phi)\|_{**} \leq C \|\phi\|_{**}^{\min\{p,2\}},$$

and we have that

$$\|N(\phi_1) - N(\phi_2)\|_{**} \leq C (\|\phi_1\|_{**}^{\min\{p-1,1\}} + \|\phi_2\|_{**}^{\min\{p-1,1\}}) \|\phi_1 - \phi_2\|_{**}.$$

Hence by Lemmas 4.3 and 4.4, for K sufficiently large and C_0 large we have

$$\|\mathcal{A}(\phi)\|_{**} \leq C (\|E\|_{**} + \|N(\phi)\|_{**}) \leq C_0 K^{-\min\{1, \frac{p-\eta}{2}\}m} (\ln K)^{-\frac{1}{2}},$$

and

$$\|\mathcal{A}(\phi_1) - \mathcal{A}(\phi_2)\|_{**} \leq C \|N(\phi_1) - N(\phi_2)\|_{**} \leq \frac{1}{2} \|\phi_1 - \phi_2\|_{**},$$

which shows that \mathcal{A} is a contraction mapping on \mathcal{B}_K . Hence there is a unique $\phi \in \mathcal{B}_K$ such that (4.4) holds.

Now we come to the differentiability of $\phi(x; \alpha, \mathbf{q})$ as function of (α, \mathbf{q}) . Consider the following map $\mathcal{T} : \mathbb{R} \times \mathbb{R}^{2K} \times \mathcal{B} \times \mathbb{R}^{2K} \rightarrow \mathcal{B} \times \mathbb{R}^{2K}$ of class C^1 :

$$\mathcal{T}(\alpha, \mathbf{q}, \phi, \widehat{\beta}) = \begin{pmatrix} (-\Delta + 1)^{-1} S(U + \phi) - \widehat{\beta} \cdot (-\Delta + 1)^{-1} \frac{\partial U}{\partial \mathbf{q}} \\ \int_{\mathbb{R}^N} \phi \mathcal{Z}_{Q_1} dx \\ \vdots \\ \int_{\mathbb{R}^N} \phi \mathcal{Z}_{Q_K} dx \end{pmatrix},$$

where $\mathcal{B} = W^{2,2}(\mathbb{R}^N) \cap \mathcal{B}_{**}$. Equation (4.4) is equivalent to $\mathcal{T}(\alpha, \mathbf{q}, \phi, \widehat{\beta}) = 0$. By the above argument, we know that, given $\alpha \in \mathbb{R}$ and \mathbf{q} satisfying (2.9), there is a unique local solution $(\phi(\alpha, \mathbf{q}), \widehat{\beta}(\alpha, \mathbf{q}))$. For simplicity, in the following, we write $(\phi, \widehat{\beta}) = (\phi(\alpha, \mathbf{q}), \widehat{\beta}(\alpha, \mathbf{q}))$. We claim that the linear operator

$$\frac{\partial \mathcal{T}(\alpha, \mathbf{q}, \phi, \widehat{\beta})}{\partial(\phi, \widehat{\beta})} \Big|_{(\alpha, \mathbf{q}, \phi, \widehat{\beta})} : \mathcal{B} \times \mathbb{R}^{2K} \rightarrow \mathcal{B} \times \mathbb{R}^{2K}$$

is invertible for K large. Then the C^1 -regularity of $(\alpha, \mathbf{q}) \mapsto (\phi, \widehat{\beta})$ follows from the Implicit Function Theorem. Indeed we have

$$\frac{\partial \mathcal{T}(\alpha, \mathbf{q}, \phi, \widehat{\beta})}{\partial(\phi, \widehat{\beta})} \Big|_{(\alpha, \mathbf{q}, \phi, \widehat{\beta})} [\varphi, \vec{\zeta}] = \begin{pmatrix} (-\Delta + 1)^{-1} S'(U + \phi)[\varphi] - \vec{\zeta} \cdot (-\Delta + 1)^{-1} \frac{\partial U}{\partial \mathbf{q}} \\ \int_{\mathbb{R}^N} \varphi \mathcal{Z}_{Q_1} dx \\ \vdots \\ \int_{\mathbb{R}^N} \varphi \mathcal{Z}_{Q_K} dx \end{pmatrix}.$$

Since $\|\phi\|_{**} \leq C_0 K^{-\min\{1, \frac{n-2}{2}\}m} (\ln K)^{-\frac{1}{2}}$, by Lemma 4.2, the argument in the proof of Lemma 4.3 shows that $\frac{\partial \mathcal{T}(\alpha, \mathbf{q}, \phi, \widehat{\beta})}{\partial(\phi, \widehat{\beta})} \Big|_{(\alpha, \mathbf{q}, \phi, \widehat{\beta})}$ is invertible for K sufficiently large. This concludes the proof of Proposition 4.1.

Next we study the dependence of ϕ on (α, \mathbf{q}) . Assume that we have two solutions corresponding to two sets of parameters. One of them denoted by

$$L[\phi] + E + N(\phi) = \widehat{\beta} \cdot \nabla_{\mathbf{q}} U,$$

corresponds to the parameters α and \mathbf{q} ; the other denoted by

$$\mathring{L}[\mathring{\phi}] + \mathring{E} + \mathring{N}(\mathring{\phi}) = \mathring{\beta} \cdot \mathring{\nabla}_{\mathbf{q}} U,$$

corresponds to the parameters $\mathring{\alpha}$ and $\mathring{\mathbf{q}}$. Observe that ϕ is L^2 -orthogonal to $\nabla_{\mathbf{q}} U$ while $\mathring{\phi}$ is L^2 -orthogonal to $\mathring{\nabla}_{\mathbf{q}} U$. To compare $\mathring{\phi}$ with ϕ , we first choose a vector $\vec{\omega}$ so that

$$\mathring{\phi}_{\vec{\omega}} = \mathring{\phi} + \vec{\omega} \cdot \nabla_{\mathbf{q}} U$$

satisfies the same orthogonality condition as ϕ . Moreover, by the equation of $\mathring{\phi}$, the function $\mathring{\phi}_{\vec{\omega}}$ satisfies the equation

$$L[\mathring{\phi}_{\vec{\omega}}] + (\mathring{L} - L)[\mathring{\phi}] - \vec{\omega} \cdot L[\nabla_{\mathbf{q}} U] + \mathring{E} + \mathring{N}(\mathring{\phi}) + \mathring{\beta} \cdot (\nabla_{\mathbf{q}} U - \mathring{\nabla}_{\mathbf{q}} U) = \mathring{\beta} \cdot \nabla_{\mathbf{q}} U.$$

Taking the difference with the equation satisfied by ϕ , we get

$$L[\hat{\phi}_\omega - \phi] = (L - \hat{L})[\hat{\phi}] + \bar{\omega} \cdot L[\nabla_{\mathbf{q}}U] + (E - \hat{E}) + (N(\phi) - \hat{N}(\hat{\phi})) - \hat{\beta} \cdot (\nabla_{\mathbf{q}}U - \hat{\nabla}_{\mathbf{q}}U) + (\hat{\beta} - \beta) \cdot \nabla_{\mathbf{q}}U.$$

Note that by (2.7), for $j = 1, \dots, K$, we have

$$|\hat{Q}_j - Q_j| \leq C(R|\hat{\alpha} - \alpha| + \|\hat{\mathbf{q}} - \mathbf{q}\|_\infty).$$

Assume that $(R|\hat{\alpha} - \alpha| + \|\hat{\mathbf{q}} - \mathbf{q}\|_\infty) \leq 1/2$, then we have

$$\|(L - \hat{L})[\hat{\phi}]\|_{**} \leq CK^{-\min\{1, \frac{p-\eta}{2}\}m} (\ln K)^{-\frac{1}{2}} (R|\hat{\alpha} - \alpha| + \|\hat{\mathbf{q}} - \mathbf{q}\|_\infty),$$

$$\|\bar{\omega} \cdot L[\nabla_{\mathbf{q}}U]\|_{**} \leq CK^{-\min\{1, \frac{p-\eta}{2}\}m} (\ln K)^{-\frac{1}{2}} \|\bar{\omega}\|_\infty,$$

$$\|E - \hat{E}\|_{**} \leq CK^{-\min\{1, \frac{p-\eta}{2}\}m} (\ln K)^{-\frac{1}{2}} (R|\hat{\alpha} - \alpha| + \|\hat{\mathbf{q}} - \mathbf{q}\|_\infty),$$

$$\begin{aligned} \|N(\phi) - \hat{N}(\hat{\phi})\|_{**} &\leq C(\|\phi\|_{**}^{p-1} + \|\hat{\phi}\|_{**}^{p-1})\|\phi - \hat{\phi}\|_{**} + C\|\phi\|_{**}^{\min\{p-1, 1\}} \\ &\quad (R|\hat{\alpha} - \alpha| + \|\hat{\mathbf{q}} - \mathbf{q}\|_\infty) \\ &\leq CK^{-\min\{1, \frac{p-\eta}{2}\}m(p-1)} (\ln K)^{-\frac{p-1}{2}} \|\phi - \hat{\phi}\|_{**} \\ &\quad + C \left(K^{-\min\{1, \frac{p-\eta}{2}\}m} (\ln K)^{-\frac{1}{2}} \right)^{\min\{p-1, 1\}} (R|\hat{\alpha} - \alpha| + \|\hat{\mathbf{q}} - \mathbf{q}\|_\infty), \end{aligned}$$

$$\begin{aligned} \|\hat{\beta} \cdot (\nabla_{\mathbf{q}}U - \hat{\nabla}_{\mathbf{q}}U)\|_{**} &\leq C\|\hat{\beta}\|_\infty (R|\hat{\alpha} - \alpha| + \|\hat{\mathbf{q}} - \mathbf{q}\|_\infty) \\ &\leq CK^{-\min\{1, \frac{p-\eta}{2}\}m} (\ln K)^{-\frac{1}{2}} (R|\hat{\alpha} - \alpha| + \|\hat{\mathbf{q}} - \mathbf{q}\|_\infty). \end{aligned}$$

Hence by Lemma 4.3,

$$\begin{aligned} \|\hat{\phi}_\omega - \phi\|_{**} + \|\hat{\beta} - \beta\|_\infty &\leq C \left(K^{-\min\{1, \frac{p-\eta}{2}\}m} (\ln K)^{-\frac{1}{2}} \right)^{\min\{p-1, 1\}} \\ &\quad (R|\hat{\alpha} - \alpha|_\infty + \|\hat{\mathbf{q}} - \mathbf{q}\|_\infty) \\ &\quad + CK^{-\min\{1, \frac{p-\eta}{2}\}m} (\ln K)^{-\frac{1}{2}} \|\bar{\omega}\|_\infty \\ &\quad + CK^{-\min\{1, \frac{p-\eta}{2}\}m(p-1)} (\ln K)^{-\frac{p-1}{2}} \|\hat{\phi} - \phi\|_{**}. \end{aligned}$$

On the other hand, by the definition of $\hat{\phi}_\omega$, we have

$$\begin{aligned} \|\bar{\omega}\|_\infty &\leq C\|\hat{\phi}\|_{**} (R|\hat{\alpha} - \alpha|_\infty + \|\hat{\mathbf{q}} - \mathbf{q}\|_\infty) \\ &\leq CK^{-\min\{1, \frac{p-\eta}{2}\}m} (\ln K)^{-\frac{1}{2}} (R|\hat{\alpha} - \alpha|_\infty + \|\hat{\mathbf{q}} - \mathbf{q}\|_\infty). \end{aligned}$$

Hence

$$\|\hat{\phi} - \phi\|_{**} + \|\hat{\beta} - \beta\|_\infty \leq C \left(K^{-\min\{1, \frac{p-\eta}{2}\}m} (\ln K)^{-\frac{1}{2}} \right)^{\min\{p-1, 1\}} (R|\hat{\alpha} - \alpha|_\infty + \|\hat{\mathbf{q}} - \mathbf{q}\|_\infty).$$

Therefore, we conclude that

$$R^{-1} \left\| \frac{\partial \phi}{\partial \alpha} \right\|_{**} + \left\| \frac{\partial \phi}{\partial \mathbf{q}} \right\|_{**} \leq C \left(K^{-\min\{1, \frac{p-\eta}{2}\}m} (\ln K)^{-\frac{1}{2}} \right)^{\min\{p-1, 1\}}.$$

□

5 A further reduction process

The main purpose of this section is to achieve Step 2.A. As explained in Sect. 2, we define

$$\vec{\beta} = \widehat{\beta} - \gamma(R\mathbf{q}_0 + \mathbf{q}^\perp), \text{ for every } \gamma \in \mathbb{R}. \tag{5.1}$$

Then Eq. (4.4) becomes

$$L[\phi] + E + N(\phi) = \vec{\beta} \cdot \frac{\partial U}{\partial \mathbf{q}} + \gamma \frac{\partial U}{\partial \alpha}. \tag{5.2}$$

Note that ϕ does not depend on γ , but $\vec{\beta}$ depends on the parameters α , \mathbf{q} and γ and we write $\vec{\beta} = \vec{\beta}(\alpha, \mathbf{q}, \gamma)$.

In this section we are going to solve $\vec{\beta}(\alpha, \mathbf{q}, \gamma) = 0$ for each $\alpha \in \mathbb{R}$ by adjusting γ and \mathbf{q} . To this end, we multiply (5.2) by $\frac{\partial U}{\partial \mathbf{q}}$ and integrate over \mathbb{R}^N to conclude that

$$\int_{\mathbb{R}^N} (E + L[\phi] + N(\phi)) \frac{\partial U}{\partial \mathbf{q}} dx = M\vec{\beta} + \gamma \int_{\mathbb{R}^N} \frac{\partial U}{\partial \alpha} \frac{\partial U}{\partial \mathbf{q}} dx.$$

By Lemma 4.2, solving $\vec{\beta}(\alpha, \mathbf{q}, \gamma) = 0$ amounts to solve

$$\int_{\mathbb{R}^N} (E + L[\phi] + N(\phi)) \frac{\partial U}{\partial \mathbf{q}} dx = \gamma \int_{\mathbb{R}^N} \frac{\partial U}{\partial \alpha} \frac{\partial U}{\partial \mathbf{q}} dx. \tag{5.3}$$

For this purpose, in the next subsection, we will evaluate the projection of the error and the projections of the terms involving ϕ .

5.1 Projections

We first compute $\int_{\mathbb{R}^N} E \frac{\partial U}{\partial \mathbf{q}} dx$. Recall that

$$\frac{\partial U}{\partial \mathbf{q}} = -(\mathcal{Z}_{Q_1} \cdot \vec{n}_1, \dots, \mathcal{Z}_{Q_K} \cdot \vec{n}_K, \mathcal{Z}_{Q_1} \cdot \vec{t}_1, \dots, \mathcal{Z}_{Q_K} \cdot \vec{t}_K)^T.$$

Lemma 5.1 *Under the assumption of Proposition 4.1, for sufficiently large K , the following expansion holds:*

$$\begin{aligned} \int_{\mathbb{R}^N} E \mathcal{Z}_{Q_k} dx &= a_0 m |Q_k|^{-m-1} \frac{Q_k}{|Q_k|} + \sum_{j \neq k} \Psi(|Q_j - Q_k|) \frac{(Q_j - Q_k)}{|Q_j - Q_k|} \\ &+ R^{-m-\sigma} \Pi_{k,1}(\alpha, \mathbf{q}) + R^{-m-3} \Pi_{k,2}(\alpha, \mathbf{q}) + R^{-2m} \Pi_{k,3}(\alpha, \mathbf{q}) \\ &+ R^{-\min\{2-\eta, \frac{p+1-\eta}{2}\}m} \Pi_{k,4}(\alpha, \mathbf{q}) \end{aligned}$$

where η is a small positive constant chosen later, $a_0 = \frac{a}{2} \int_{\mathbb{R}^N} w^2(x) dx$, and $\Pi_{k,l}(\alpha, \mathbf{q})$'s are smooth vector valued functions, which are uniformly bounded as $K \rightarrow \infty$.

Proof By the definition,

$$\int_{\mathbb{R}^N} E \mathcal{Z}_{Q_k} dx = \underbrace{\sum_{j=1}^K \int_{\mathbb{R}^N} (V(x) - 1) w_{Q_j} \nabla w_{Q_k} dx}_{I_1} - \underbrace{\int_{\mathbb{R}^N} \left\{ \left(\sum_{j=1}^K w_{Q_j} \right)^p - \sum_{j=1}^K w_{Q_j}^p \right\} \nabla w_{Q_k} dx}_{I_2}.$$

Claim 1: There are smooth vector valued functions $\Pi_{k,1}, \Pi_{k,2}$ and $\Pi_{k,3}$ of α and \mathbf{q} such that

$$I_1 = a_0 m |\mathcal{Q}_k|^{-m-1} \frac{\mathcal{Q}_k}{|\mathcal{Q}_k|} + R^{-m-\sigma} \Pi_{k,1}(\alpha, \mathbf{q}) + R^{-m-3} \Pi_{k,2}(\alpha, \mathbf{q}) + R^{-2m} \Pi_{k,3}(\alpha, \mathbf{q}).$$

Claim 2: There exists a smooth vector valued function $\Pi_{k,4}$ of α and \mathbf{q} such that

$$I_2 = - \sum_{j \neq k} \Psi(|\mathcal{Q}_j - \mathcal{Q}_k|) \frac{(\mathcal{Q}_j - \mathcal{Q}_k)}{|\mathcal{Q}_j - \mathcal{Q}_k|} - R^{-\min\{2-\eta, \frac{p+1-\eta}{2}\}m} \Pi_{k,4}(\alpha, \mathbf{q}),$$

where η is a small positive constant chosen later.

Combining Claim 1 and Claim 2, we get the desired result. The remainder of this proof will be devoted to the proofs of Claim 1 and Claim 2.

Proof of Claim 1: We divide I_1 into two parts:

$$I_1 = \underbrace{\int_{\mathbb{R}^N} (V(x) - 1) w_{\mathcal{Q}_k} \nabla w_{\mathcal{Q}_k} dx}_{I_{11}} + \sum_{j \neq k} \underbrace{\int_{\mathbb{R}^N} (V(x) - 1) w_{\mathcal{Q}_j} \nabla w_{\mathcal{Q}_k} dx}_{I_{12}}.$$

Write $I_{11} = I_{111} + I_{112}$, where

$$I_{111} = \int_{|x| \leq \frac{R}{3}} (V(x) - 1) w_{\mathcal{Q}_k} \nabla w_{\mathcal{Q}_k} dx, \text{ and } I_{112} = \int_{|x| > \frac{R}{3}} (V(x) - 1) w_{\mathcal{Q}_k} \nabla w_{\mathcal{Q}_k} dx.$$

On one hand, since $V \in L^\infty(\mathbb{R}^N)$ and (2.10), by the triangle inequality and the property of w , we get $|I_{111}| \leq C e^{-R/2} \leq C R^{-m-3}$. On the other hand, by (1.11) and Taylor’s theorem,

$$\begin{aligned} I_{112} &= \int_{|x| > \frac{R}{3}} \frac{a}{|x|^m} w_{\mathcal{Q}_k} \nabla w_{\mathcal{Q}_k} dx + O(R^{-m-\sigma}) \\ &= \int_{|y+\mathcal{Q}_k| > \frac{R}{3}} \frac{a}{|y + \mathcal{Q}_k|^m} w(y) \nabla w(y) dy + O(R^{-m-\sigma}) \\ &= m |\mathcal{Q}_k|^{-m-1} \frac{\mathcal{Q}_k}{|\mathcal{Q}_k|} \frac{a}{2} \int_{\mathbb{R}^N} w^2(x) dx + O(R^{-m-\sigma}) + O(R^{-m-3}), \end{aligned}$$

where in the last equality we use the following identities:

$$\int_{\mathbb{R}^N} w(y) \nabla w(y) dy = 0, \int_{\mathbb{R}^N} y_j y_k w(y) \nabla w(y) dy = 0, \forall j, k = 1, \dots, N,$$

and

$$\int_{\mathbb{R}^N} (y^T \vec{e}) w(y) \nabla w(y) dy = -\frac{1}{2} \int_{\mathbb{R}^N} w^2 dy \vec{e}, \forall \vec{e} \in \mathbb{R}^N.$$

Therefore, we get

$$I_{11} = a_0 m |\mathcal{Q}_k|^{-m-1} \frac{\mathcal{Q}_k}{|\mathcal{Q}_k|} + O(R^{-m-\sigma}) + O(R^{-m-3}).$$

If (1.11) holds in the C^1 sense, i.e.,

$$\nabla V(x) = -\frac{ma}{|x|^{m+1}} \frac{x}{|x|} + O\left(\frac{1}{|x|^{m+1+\sigma}}\right), \text{ as } |x| \rightarrow +\infty,$$

by a similar argument, we get

$$I_{11} = a_0 m |Q_k|^{-m-1} \frac{Q_k}{|Q_k|} + O(R^{-m-1-\sigma}) + O(R^{-m-3}).$$

For the term I_{12} , we claim that there exists a constant C (independent of K) such that

$$|I_{12}| \leq CR^{-m} e^{-d} d^{-\frac{N-3}{2}} \leq CR^{-2m}.$$

Indeed, by Lemmas 3.3, 3.5 and 3.8, we have

$$\begin{aligned} & \left| \sum_{j \neq k} \int_{\mathbb{R}^N} (V(x) - 1) w_{Q_j} \nabla w_{Q_k} dx \right| \\ & \leq \left| \sum_{j \neq k} \int_{|x| < \frac{R}{3}} (V(x) - 1) w_{Q_j} \nabla w_{Q_k} dx \right| + \left| \sum_{j \neq k} \int_{|x| \geq \frac{R}{3}} (V(x) - 1) w_{Q_j} \nabla w_{Q_k} dx \right| \\ & \leq CK e^{-R/2} + C \sum_{j \neq k} R^{-m} e^{-|Q_j - Q_k|} |Q_j - Q_k|^{-\frac{N-3}{2}} \\ & \leq CR^{-m} e^{-d} d^{-\frac{N-3}{2}} \leq CR^{-2m}. \end{aligned}$$

Combining the above estimates, we complete the proof of Claim 1.

Proof of Claim 2: We first divide I_2 into two parts:

$$\begin{aligned} I_2 &= \underbrace{\int_{\mathbb{R}^N} p w_{Q_k}^{p-1} \left(\sum_{j \neq k} w_{Q_j} \right) \nabla w_{Q_k} dx}_{I_{21}} \\ &+ \underbrace{\int_{\mathbb{R}^N} \left\{ \left(\sum_{j=1}^K w_{Q_j} \right)^p - w_{Q_k}^p - p w_{Q_k}^{p-1} \sum_{j \neq k} w_{Q_j} - \sum_{j \neq k} w_{Q_j}^p \right\} \nabla w_{Q_k} dx}_{I_{22}}. \end{aligned}$$

By the definition (2.6) of the interaction function Ψ ,

$$I_{21} = - \sum_{j \neq k} \Psi(|Q_j - Q_k|) \frac{(Q_j - Q_k)}{|Q_j - Q_k|}.$$

For the term I_{22} , we divide the domain of integration into $K + 1$ parts: $\Omega_1^\ell, \dots, \Omega_{K+1}^\ell$. On Ω_{K+1}^ℓ , by taking ℓ large but independent of K , we have

$$\left| \left(\sum_{j=1}^K w_{Q_j} \right)^p - w_{Q_k}^p - p w_{Q_k}^{p-1} \sum_{j \neq k} w_{Q_j} - \sum_{j \neq k} w_{Q_j}^p \right| \leq CK^p e^{-p\ell\rho/2} \leq CK^{-m-3}.$$

On $\Omega_l^\ell, l = 1, \dots, K$, by the definition, we have

$$|x - Q_l| \leq |x - Q_j|, \forall x \in \Omega_l^\ell, \text{ and } \forall 1 \leq j \leq K. \tag{5.4}$$

Since ℓ is independent of K , for $x \in \Omega_k^\ell$, by mean value theorem and (3.11),

$$\begin{aligned} & \left| \left\{ \left(\sum_{j=1}^K w_{Q_j} \right)^p - w_{Q_k}^p - p w_{Q_k}^{p-1} \sum_{j \neq k} w_{Q_j} - \sum_{j \neq k} w_{Q_j}^p \right\} \nabla w_{Q_k} \right| \\ & \leq C \left\{ w_{Q_k}^{p-1} \left(\sum_{j \neq k} w_{Q_j} \right)^2 + w_{Q_k} \sum_{j \neq k} w_{Q_j}^p \right\}. \end{aligned}$$

Note that for $x \in \Omega_k^\ell, |x - Q_j| \geq \rho/2$ for all $j \neq k$. Hence by Lemma 3.1, (5.4) and Lemma 3.5, and by choosing $0 < \eta < 1$ sufficiently small, we have

$$\begin{aligned} & w_{Q_k}^{p-1} \left(\sum_{j \neq k} w_{Q_j} \right)^2 + w_{Q_k} \sum_{j \neq k} w_{Q_j}^p \\ & \leq C \rho^{-\min\{2,p\} \frac{N-1}{2}} \left\{ e^{-(p-1-\eta)|x-Q_k|} \left(\sum_{j \neq k} e^{-|x-Q_j|} \right)^2 + e^{-(1-\eta)|x-Q_k|} \sum_{j \neq k} e^{-p|x-Q_j|} \right\} w_{Q_k}^\eta \\ & \leq C e^{-\min\{2, \frac{p+1-\eta}{2}\} d} d^{-\min\{2,p\} \frac{N-1}{2}} w_{Q_k}^\eta. \end{aligned}$$

Similarly, for all $x \in \Omega_l^\ell (l \neq k)$, by mean value theorem, (3.11), Lemma 3.1, (5.4) and Lemma 3.5, we have

$$\begin{aligned} & \left| \left\{ \left(\sum_{j=1}^K w_{Q_j} \right)^p - w_{Q_k}^p - p w_{Q_k}^{p-1} \sum_{j \neq k} w_{Q_j} - \sum_{j \neq k} w_{Q_j}^p \right\} \nabla w_{Q_k} \right| \\ & \leq C \left\{ w_{Q_k} w_{Q_l}^{p-1} \sum_{j \neq l} w_{Q_j} + w_{Q_k}^p w_{Q_l} \right\}, \end{aligned}$$

and

$$\begin{aligned} & w_{Q_k} w_{Q_l}^{p-1} \sum_{j \neq l} w_{Q_j} + w_{Q_k}^p w_{Q_l} \\ & \leq C \rho^{-\min\{2,p\} \frac{N-1}{2}} \left\{ e^{-(2-\eta)|x-Q_k|} e^{-(p-1)|x-Q_l|} + e^{-(p-\eta)|x-Q_k|} e^{-|x-Q_l|} \right. \\ & \quad \left. + e^{-(1-\eta)|x-Q_k|} e^{-(p-1)|x-Q_l|} \sum_{j \neq k,l} e^{-|x-Q_j|} \right\} e^{-\eta|x-Q_k|} \\ & \leq C d^{-\min\{2,p\} \frac{N-1}{2}} e^{-\min\{2-\eta, \frac{p+1-\eta}{2}, p-\eta\} d} e^{-\eta|x-Q_k|}. \end{aligned}$$

Since $p > 1$, we can choose $\eta > 0$ such that $p - \eta > \frac{p+1-\eta}{2}$. Hence, there is a constant (independent of K) such that for all $x \in \mathbb{R}^N$,

$$\left| \left\{ \left(\sum_{j=1}^K w_{Q_j} \right)^p - w_{Q_k}^p - p w_{Q_k}^{p-1} \sum_{j \neq k} w_{Q_j} - \sum_{j \neq k} w_{Q_j}^p \right\} \nabla w_{Q_k} \right| \leq C d^{-\min\{2,p\} \frac{N-1}{2}} e^{-\min\{2-\eta, \frac{p+1-\eta}{2}\}d} e^{-\eta|x-Q_k|}.$$

Therefore,

$$|I_{22}| \leq C d^{-\min\{2,p\} \frac{N-1}{2}} e^{-\min\{2-\eta, \frac{p+1-\eta}{2}\}d} \leq C R^{-\min\{2-\eta, \frac{p+1-\eta}{2}\}m}.$$

Combining the above estimates, we complete the proof of Claim 2. □

Now let us analyze $\int_{\mathbb{R}^N} E \frac{\partial U}{\partial \mathbf{q}} dx$. Before doing this, we define

$$\widehat{d} = -\frac{\Psi'(d)}{\Psi(d)}d = d + O(1),$$

and for $j = 1, \dots, K$, we denote

$$\bar{f}_j = \dot{f}_j + \dot{f}_{j-1} = (f_{j+1} - f_{j-1}) \frac{K}{2\pi}, \quad \bar{g}_j = \dot{g}_j + \dot{g}_{j-1} = (g_{j+1} - g_{j-1}) \frac{K}{2\pi}.$$

Lemma 5.2 *Under the assumption of Proposition 4.1, for sufficiently large K , the following expansion holds:*

$$\begin{aligned} - \int_{\mathbb{R}^N} E \frac{\partial U}{\partial \mathbf{q}} dx &= a_0 m R^{-m-2} T \mathbf{q} + R^{-m-\sigma} \Pi_1(\alpha, \mathbf{q}) + R^{-m-3} \Pi_2(\alpha, \mathbf{q}) + R^{-2m} \Pi_3(\alpha, \mathbf{q}) \\ &\quad + R^{-\min\{2-\eta, \frac{p+1-\eta}{2}\}m} \Pi_4(\alpha, \mathbf{q}) + R^{-m-3} (\ln K)^2 \Pi_5(\alpha, \mathbf{q}, \dot{\mathbf{q}}, \ddot{\mathbf{q}}), \end{aligned}$$

where $\Pi_1(\alpha, \mathbf{q}), \dots, \Pi_4(\alpha, \mathbf{q}), \Pi_5(\alpha, \mathbf{q}, \dot{\mathbf{q}}, \ddot{\mathbf{q}})$ are uniformly bounded smooth vector valued functions with $\Pi_5(\alpha, 0, 0, 0) = 0$, and T is an $2K \times 2K$ matrix defined by

$$T = \begin{pmatrix} c_1 A_1 + c_4 I & c_2 A_2 \\ -c_2 A_2 & c_3 A_1 \end{pmatrix}, \tag{5.5}$$

Here I is the $K \times K$ identity matrix, both A_1 and A_2 are $K \times K$ circulant matrices given by

$$A_1 = \begin{pmatrix} -2 & 1 & 0 & \dots & 0 & 1 \\ 1 & -2 & 1 & 0 & \dots & 0 \\ 0 & 1 & -2 & 1 & 0 & \dots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & 1 & -2 & 1 \\ 1 & 0 & \dots & 0 & 1 & -2 \end{pmatrix}, \quad A_2 = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 & -1 \\ -1 & 0 & 1 & 0 & \dots & 0 \\ 0 & -1 & 0 & 1 & 0 & \dots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & -1 & 0 & 1 \\ 1 & 0 & \dots & 0 & -1 & 0 \end{pmatrix},$$

and c_1, c_2, c_3, c_4 are constants given by

$$c_1 = \frac{K^2}{4\pi^2}, \quad c_2 = (\widehat{d} - 1) \frac{K}{4\pi}, \quad c_3 = -\widehat{d} \frac{K^2}{4\pi^2}, \quad c_4 = \widehat{d} - m - 1. \tag{5.6}$$

Proof First a simple computation shows that

$$\begin{aligned}
 |Q_k|^{-m-1} \frac{Q_k}{|Q_k|} &= |Q_k^0 + q_k|^{-m-1} \frac{(Q_k^0 + q_k)}{|Q_k^0 + q_k|} \\
 &= R^{-m-1} \vec{n}_k + R^{-m-2} \{g_k \vec{t}_k - (m + 1) f_k \vec{n}_k\} + O(R^{-m-3}).
 \end{aligned}$$

To estimate I_2 , by direct computation, we have

$$\begin{aligned}
 &\frac{Q_{j+1} - Q_j}{|Q_{j+1} - Q_j|} \\
 &= \left\{ -\sin \frac{\pi}{K} + (\dot{f}_j - g_j)R^{-1} - \dot{g}_j R^{-1} \frac{\pi}{K} - (\dot{f}_j - g_j)(f_j + \dot{g}_j)R^{-2} \right\} \vec{n}_j \\
 &\quad + \left\{ \cos \frac{\pi}{K} + (\dot{f}_j - g_j)R^{-1} \frac{\pi}{K} - \frac{1}{2}(\dot{f}_j - g_j)^2 R^{-2} \right\} \vec{t}_j + O(K^{-3}(\ln K)^{-1}),
 \end{aligned}$$

$$\begin{aligned}
 &\frac{Q_{j-1} - Q_j}{|Q_{j-1} - Q_j|} \\
 &= \left\{ -\sin \frac{\pi}{K} + (-\dot{f}_{j-1} + g_j)R^{-1} - \dot{g}_{j-1} R^{-1} \frac{\pi}{K} - (-\dot{f}_{j-1} + g_j)(f_j + \dot{g}_{j-1})R^{-2} \right\} \vec{n}_j \\
 &\quad - \left\{ \cos \frac{\pi}{K} + (-\dot{f}_{j-1} + g_j)R^{-1} \frac{\pi}{K} - \frac{1}{2}(-\dot{f}_{j-1} + g_j)^2 R^{-2} \right\} \vec{t}_j + O(K^{-3}(\ln K)^{-1}),
 \end{aligned}$$

and

$$\begin{aligned}
 &\Psi(|Q_{j+1} - Q_j|) \\
 &= \Psi(d + (2f_j + \bar{g}_j) \frac{\pi}{K}) + \Psi'(d) \left\{ 2\ddot{g}_j \frac{\pi^2}{K^2} + 2\dot{f}_j \frac{\pi^2}{K^2} + (\dot{f}_j - g_j)^2 R^{-1} \frac{\pi}{K} \right\} \\
 &\quad + \Psi(d) O(K^{-3}),
 \end{aligned}$$

$$\begin{aligned}
 &\Psi(|Q_{j-1} - Q_j|) \\
 &= \Psi(d + (2f_j + \bar{g}_j) \frac{\pi}{K}) + \Psi'(d) \left\{ -2\ddot{g}_j \frac{\pi^2}{K^2} - 2\dot{f}_{j-1} \frac{\pi^2}{K^2} + (-\dot{f}_{j-1} + g_j)^2 R^{-1} \frac{\pi}{K} \right\} \\
 &\quad + \Psi(d) O(K^{-3}).
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 &\sum_{j \in \{k-1, k+1\}} \Psi(|Q_j - Q_k|) \frac{(Q_j - Q_k)}{|Q_j - Q_k|} \\
 &= -2 \sin \frac{\pi}{K} \Psi(d) \vec{n}_j + 2\Psi'(d) \frac{\pi^2}{K^2} \left\{ -(2f_j + \bar{g}_j) \vec{n}_j + (\bar{f}_j + 2\ddot{g}_j) \vec{t}_j \right\} \\
 &\quad + \Psi(d) R^{-1} \frac{\pi}{K} \left\{ (2\ddot{f}_j - \bar{g}_j) \vec{n}_j + (\bar{f}_j - 2g_j) \vec{t}_j \right\} + \Psi(d) O(K^{-3}).
 \end{aligned}$$

Combining the above estimates and Lemma 5.1, we get

$$\begin{aligned} \int_{\mathbb{R}^N} E \mathcal{Z}_{Q_k} dx &= a_0 m R^{-m-2} \left\{ -(m+1) f_k + (\ddot{f}_k - \frac{1}{2} \ddot{g}_k) + \widehat{d}(f_k + \frac{1}{2} \ddot{g}_k) \right\} \vec{n}_k \\ &\quad + a_0 m R^{-m-2} \left\{ g_k + (\frac{1}{2} \ddot{f}_k - g_k) - \widehat{d}(\frac{1}{2} \ddot{f}_k + \ddot{g}_k) \right\} \vec{t}_k \\ &\quad + R^{-m-\sigma} \Pi_{k,1}(\alpha, \mathbf{q}) + R^{-m-3} \Pi_{k,2}(\alpha, \mathbf{q}) + R^{-2m} \Pi_{k,3}(\alpha, \mathbf{q}) \\ &\quad + R^{-\min\{2-\eta, \frac{p+1-\eta}{2}\}m} \Pi_{k,4}(\alpha, \mathbf{q}) + R^{-m-3} (\ln K)^2 \Pi_{k,5}(\alpha, \mathbf{q}, \dot{\mathbf{q}}, \ddot{\mathbf{q}}), \end{aligned}$$

where $\Pi_{k,l}(\alpha, \mathbf{q})$'s and $\Pi_{k,5}(\alpha, \mathbf{q}, \dot{\mathbf{q}}, \ddot{\mathbf{q}})$ are smooth vector valued functions, which are uniformly bounded as $K \rightarrow \infty$. Moreover, $\Pi_{k,5}(\alpha, 0, 0, 0) = 0$. The desired result follows. \square

Next we compute $\int_{\mathbb{R}^N} (L[\phi] + N(\phi)) \frac{\partial U}{\partial \mathbf{q}} dx$.

Lemma 5.3 *Under the assumption of Proposition 4.1, for sufficiently large K , the following expansions hold true:*

$$\int_{\mathbb{R}^N} L[\phi] \frac{\partial U}{\partial \mathbf{q}} dx = K^{-\min\{2, \frac{p}{2} + 1 - \frac{\eta}{2}, p - \frac{\eta}{2}\}m} (\ln K)^2 \Pi_6(\alpha, \mathbf{q}),$$

and

$$\int_{\mathbb{R}^N} N(\phi) \frac{\partial U}{\partial \mathbf{q}} dx = K^{-\min\{2, p - \eta\}m} (\ln K)^{-1} \Pi_7(\alpha, \mathbf{q}),$$

where $\Pi_6(\alpha, \mathbf{q}), \Pi_7(\alpha, \mathbf{q})$ are uniformly bounded smooth vector valued functions.

Proof By integration by parts, (4.13) and Proposition 4.1, we have

$$\begin{aligned} \left| \int_{\mathbb{R}^N} L[\phi] \mathcal{Z}_{Q_k} dx \right| &= \left| \int_{\mathbb{R}^N} \phi L[\mathcal{Z}_{Q_k}] dx \right| \leq C d e^{-\min\{1, \frac{p}{2}\}d} \|\phi\|_{L^\infty(\mathbb{R}^N)} \\ &\leq C K^{-\min\{2, \frac{p}{2} + 1 - \frac{\eta}{2}, p - \frac{\eta}{2}\}m} (\ln K)^2. \end{aligned}$$

For the second estimate, since $\|\phi\|_{**} \leq C K^{-\min\{1, \frac{p-\eta}{2}\}m} (\ln K)^{-\frac{1}{2}}$, one has

$$|N(\phi)| \leq \begin{cases} C U^{p-2} |\phi|^2, & \text{for } |\phi| \leq U/2, \\ C |\phi|^p, & \text{for } |\phi| \geq U/2. \end{cases}$$

We claim that

$$\left| \int_{\mathbb{R}^N} N(\phi) \mathcal{Z}_{Q_k} dx \right| \leq C \|\phi\|_{**}^2 \leq C K^{-\min\{2, p-\eta\}m} (\ln K)^{-1}.$$

Indeed, when $p \geq 2$, this follows from $|N(\phi)| \leq C |\phi|^2$. Now we consider the case $p < 2$. In this case, it is not hard to get $|N(\phi)| \leq C U^{p-2} |\phi|^2$. Since $U \geq w_{Q_k}$, we have

$$|N(\phi) \mathcal{Z}_{Q_k}| \leq C U^{p-2} w_{Q_k} |\phi|^2 \leq C w_{Q_k}^{p-1} |\phi|^2, \text{ if } p < 2,$$

from which we get the desired result. \square

5.2 The invertibility of T

In this subsection, we study the linear problem $T\vec{\mathbf{q}} = \mathbf{b}$ and get the following result, whose proof is delayed to Appendix A.

Lemma 5.4 *There is an $K_0 \in \mathbb{N}$ such that for all $K \geq K_0$ and every $\mathbf{b} \in \mathbb{R}^{2K}$, there exists a unique vector $\vec{\mathbf{q}} \in \mathbb{R}^{2K}$ and a unique constant $\gamma \in \mathbb{R}$ such that*

$$T\vec{\mathbf{q}} = \mathbf{b} + \gamma \mathbf{q}_1, \quad \vec{\mathbf{q}} \perp \mathbf{q}_0, \quad \text{where } \mathbf{q}_1 = M(R\mathbf{q}_0 + \mathbf{q}^\perp). \tag{5.7}$$

Moreover, there is a positive constant C which is independent of K such that

$$\|\vec{\mathbf{q}}\|_* \leq C(\ln K)^2 \|\mathbf{b}\|_\infty. \tag{5.8}$$

Denote the inverse of T in the sense of Lemma 5.4 by T^{-1} . Since \mathbf{q}_1 depends on the parameter \mathbf{q} , the matrix T^{-1} depends on \mathbf{q} and thus we write $T^{-1} = T_{\mathbf{q}}^{-1}$.

5.3 Reduction to one dimension

Now we can state the main result in this section.

Proposition 5.5 *Under the assumption of Theorem 1.1, there is an integer $K_0 > 0$ such that: for all integer $K \geq K_0$ and for each $\alpha \in \mathbb{R}$, there exists a unique $(\mathbf{q}, \gamma) = (\mathbf{q}(\alpha), \gamma(\alpha))$ such that $\vec{\beta}(\alpha, \mathbf{q}, \gamma) = 0$. As a result, $\phi(x; \alpha, \mathbf{q}(\alpha))$ and $\gamma(\alpha)$ satisfy the equation:*

$$\begin{cases} L[\phi] + E + N(\phi) = \gamma \frac{\partial U}{\partial \alpha}, \\ \int_{\mathbb{R}^N} \phi \mathcal{Z}_{Q_j} dx = 0, \quad \forall j = 1, \dots, K. \end{cases} \tag{5.9}$$

Moreover, the function $\phi(x; \alpha, \mathbf{q}(\alpha))$ is of class C^1 in α , and we have

$$\|\phi\|_{**} \leq C_0 K^{-\min\{1, \frac{p-2}{2}\}m} (\ln K)^{-\frac{1}{2}}, \quad \|\mathbf{q}\|_* + R^{-1} \|\partial_\alpha \mathbf{q}\|_* \leq C K^{-\mu} (\ln K)^2, \tag{5.10}$$

where C is a positive constant (independent of K), and

$$0 < \mu < \min \left\{ \sigma - 2, \min \left\{ 1 - \eta, \frac{p - 1 - \eta}{2} \right\} m - 2, 1 \right\}. \tag{5.11}$$

To prove Proposition 5.5, it suffices to solve $\vec{\beta}(\alpha, \mathbf{q}, \gamma) = 0$ for each α . By the results in the preceding subsections, we can rewrite this equation in a more explicit form.

Lemma 5.6 *For every $\alpha \in \mathbb{R}$, the equation $\vec{\beta}(\alpha, \mathbf{q}, \gamma) = 0$ is equivalent to*

$$-a_0 m R^{-m-2} T\mathbf{q} + \Phi(\alpha, \mathbf{q}) = \gamma \mathbf{q}_1, \tag{5.12}$$

where T is the $2K \times 2K$ matrix defined in (5.5), Φ denotes the remaining term, and

$$\mathbf{q}_1 = \int_{\mathbb{R}^N} \frac{\partial U}{\partial \alpha} \frac{\partial U}{\partial \mathbf{q}} dx = M(R\mathbf{q}_0 + \mathbf{q}^\perp).$$

By Lemmas 5.2, 5.6 and 5.6, we have the following expansion of $\Phi(\alpha, \mathbf{q})$.

Lemma 5.7 *Under the assumption of Proposition 4.1, for K sufficiently large, the following expansion holds:*

$$\begin{aligned} \Phi(\mathbf{q}) &= R^{-m-\sigma} \Pi_1(\alpha, \mathbf{q}) + R^{-m-3} \Pi_2(\alpha, \mathbf{q}) + R^{-2m} \Pi_3(\alpha, \mathbf{q}) \\ &\quad + R^{-\min\{2-\eta, \frac{p+1-\eta}{2}\}m} \Pi_4(\alpha, \mathbf{q}) + R^{-m-3} (\ln K)^2 \Pi_5(\alpha, \mathbf{q}, \dot{\mathbf{q}}, \ddot{\mathbf{q}}) \\ &\quad + K^{-\min\{2, \frac{p}{2}+1-\frac{\eta}{2}, p-\frac{\eta}{2}\}m} (\ln K)^2 \Pi_6(\alpha, \mathbf{q}) + K^{-\min\{2, p-\eta\}m} (\ln K)^{-1} \Pi_7(\alpha, \mathbf{q}), \end{aligned}$$

where $\Pi_j(\alpha, \mathbf{q})$'s and $\Pi_5(\alpha, \mathbf{q}, \dot{\mathbf{q}}, \ddot{\mathbf{q}})$ are smooth vector valued functions, which are uniformly bounded as $K \rightarrow \infty$. Moreover, $\Pi_5(\alpha, 0, 0, 0) = 0$.

Now we are going to solve (5.12) and then complete the proof of Proposition 5.5.

Proof of Proposition 5.5 By Lemma 5.4, Eq. (5.12) is equivalent to

$$\mathbf{q} = (a_0 m)^{-1} T_{\mathbf{q}}^{-1} [R^{m+2} \Phi(\alpha, \mathbf{q})] = \mathcal{F}(\mathbf{q}).$$

By Lemma 5.7, since $\min\{2 - \eta, \frac{p+1-\eta}{2}\} \leq \min\{2, p - \eta\} \leq \min\{2, \frac{p}{2} + 1 - \frac{\eta}{2}, p - \frac{\eta}{2}\}$ for $0 < \eta < p - 1$, we get

$$(a_0 m)^{-1} R^{m+2} \Phi(\alpha, \mathbf{q}) = K^{-\mu} \tilde{\Pi}(\alpha, \mathbf{q}) + (K^{-1} \ln K) \tilde{\Xi}(\alpha, \mathbf{q}, \dot{\mathbf{q}}, \ddot{\mathbf{q}}),$$

where both $\tilde{\Pi}$ and $\tilde{\Xi}$ are smooth vector valued functions, which are uniformly bounded as K tends to infinity. Moreover, $\tilde{\Xi}(\alpha, 0, 0, 0) = 0$.

Hence by Lemma 5.4, for $\|\mathbf{q}\|_* \leq 1/2$, we have

$$\|\mathcal{F}(\mathbf{q})\|_* \leq C (K^{-\mu} (\ln K)^2 + K^{-1} (\ln K)^3) \leq C K^{-\mu} (\ln K)^2,$$

and

$$\|\mathcal{F}(\mathbf{q}) - \mathcal{F}(\hat{\mathbf{q}})\|_* \leq C (K^{-\mu} (\ln K)^2 + K^{-1} (\ln K)^3) \|\mathbf{q} - \hat{\mathbf{q}}\|_* \leq \frac{1}{2} \|\mathbf{q} - \hat{\mathbf{q}}\|_*,$$

since $\|(T_{\mathbf{q}}^{-1} - T_{\hat{\mathbf{q}}}^{-1})\mathbf{b}\|_* \leq C K R^{-m} (\ln K)^2 \|\mathbf{b}\|_\infty \|\mathbf{q} - \hat{\mathbf{q}}\|_*$. Therefore, \mathcal{F} is a contraction mapping. By the Banach fixed point theorem, the result follows.

To show the differentiability of $\mathbf{q}(\alpha)$. Consider the map $\mathcal{T}(\alpha, \mathbf{q}) = \mathbf{q} - \mathcal{F}(\alpha, \mathbf{q}) : \mathbb{R} \times \mathbb{R}^{2K} \mapsto \mathbb{R}^{2K}$ of class C^1 . Since $\frac{\partial \mathcal{F}}{\partial \mathbf{q}} = O(K^{-\mu-1} (\ln K)^2)$, $\frac{\partial \mathcal{T}}{\partial \mathbf{q}}|_{\alpha, \mathbf{q}(\alpha)} = I - \frac{\partial \mathcal{F}}{\partial \mathbf{q}}(\alpha, \mathbf{q}(\alpha))$ is invertible, we get the differentiability of $\mathbf{q}(\alpha)$.

Next we study the dependence of \mathbf{q} on α . Assume that we have two solutions corresponding to two sets of parameters. One of them denoted by

$$\mathbf{q} = (a_0 m)^{-1} T_{\mathbf{q}}^{-1} [R^{m+2} \Phi(\alpha, \mathbf{q})],$$

corresponds to α ; the other denoted by

$$\hat{\mathbf{q}} = (a_0 m)^{-1} T_{\hat{\mathbf{q}}}^{-1} [R^{m+2} \Phi(\hat{\alpha}, \hat{\mathbf{q}})],$$

corresponds to $\hat{\alpha}$. Assume that $R|\hat{\alpha} - \alpha| \leq 1/2$, by a direct computation and Lemma 5.4, we have

$$\|\mathbf{q} - \hat{\mathbf{q}}\|_* \leq C K^{-\mu} (\ln K)^2 (R|\hat{\alpha} - \alpha|),$$

from which we get the desired result. □

6 Proof of Theorem 1.1: variational reduction

In this section, our purpose is to achieve Step 2.B in the setting up of the problem and then complete the proof of Theorem 2.1.

To solve $\gamma(\alpha) = 0$ in Step 2.B, we will apply the variational reduction. To do this, we first introduce some notation. Let $\alpha \in \mathbb{R}$ and $\phi = \phi(x; \alpha, \mathbf{q}(\alpha))$ be the function given in Proposition 5.5, we define the reduced energy function by

$$F(\alpha) = \mathcal{E}(U + \phi) : \mathbb{R} \rightarrow \mathbb{R}, \tag{6.1}$$

where we write $U = U(x; \alpha, \mathbf{q}(\alpha))$.

By (2.8), both U and ϕ are 2π periodic in α . Hence by Proposition 5.5, the reduced energy function $F(\alpha)$ has the following property.

Lemma 6.1 *The function $F(\alpha)$ is of class C^1 and satisfies $F(\alpha + 2\pi) = F(\alpha)$ for every $\alpha \in \mathbb{R}$.*

With this notation, the next lemma shows that if $F(\alpha)$ has a critical point then $\gamma(\alpha) = 0$ has a solution. In other words, after the Lyapunov–Schmidt reduction, the following lemma concerns the relation between the critical points of $F(\alpha)$ and those of the energy functional $\mathcal{E}(u)$.

Lemma 6.2 *Under the assumption of Proposition 5.5, there exists $K_0 \in \mathbb{N}_+$ such that: for all integer $K \geq K_0$, if α_0 is a critical point of $F(\alpha)$, then $\gamma(\alpha_0) = 0$ and the corresponding function*

$$u(x) = U(x; \alpha_0, \mathbf{q}(\alpha_0)) + \phi(x; \alpha_0, \mathbf{q}(\alpha_0))$$

is a solution of (1.3).

Proof By Proposition 5.5, for K sufficiently large and for every $\alpha \in \mathbb{R}$, $\phi = \phi(x; \alpha, \mathbf{q}(\alpha))$ satisfies the equation

$$S(U + \phi) = \gamma(\alpha) \frac{\partial U}{\partial \alpha}. \tag{6.2}$$

By the definition (6.1), we obtain

$$F'(\alpha) = \int_{\mathbb{R}^N} S(U + \phi)(\partial_\alpha U + \partial_\alpha \phi) \, dx,$$

where $\partial_\alpha U = \frac{\partial U}{\partial \alpha} + \frac{\partial U}{\partial \mathbf{q}} \cdot \partial_\alpha \mathbf{q}$ and $\partial_\alpha \phi = \frac{\partial \phi}{\partial \alpha} + \frac{\partial \phi}{\partial \mathbf{q}} \cdot \partial_\alpha \mathbf{q}$. Thus by using (6.2),

$$F'(\alpha) = \gamma(\alpha) \int_{\mathbb{R}^N} \frac{\partial U}{\partial \alpha} (\partial_\alpha U + \partial_\alpha \phi) \, dx.$$

If α_0 be a critical point of $F(\alpha)$, then $F'(\alpha_0) = 0$. Hence to prove $\gamma(\alpha_0) = 0$, it is sufficient to show that

$$\int_{\mathbb{R}^N} \frac{\partial U}{\partial \alpha} (\partial_\alpha U + \partial_\alpha \phi) \, dx \neq 0. \tag{6.3}$$

In fact, by Propositions 5.5 and 5.5, we have

$$\partial_\alpha U + \partial_\alpha \phi = \frac{\partial U}{\partial \alpha} + \frac{\partial U}{\partial \mathbf{q}} \cdot \partial_\alpha \mathbf{q} + \frac{\partial \phi}{\partial \alpha} + \frac{\partial \phi}{\partial \mathbf{q}} \cdot \partial_\alpha \mathbf{q}. \tag{6.4}$$

Recall that

$$\frac{\partial U}{\partial \alpha} = (R\mathbf{q}_0 + \mathbf{q}^\perp) \cdot \frac{\partial U}{\partial \mathbf{q}},$$

hence by Propositions 4.1 and 5.5, we have

$$K^{-1}R^{-2} \int_{\mathbb{R}^N} \frac{\partial U}{\partial \alpha} (\partial_\alpha U + \partial_\alpha \phi) dx = (1 + o(1)) \int_{\mathbb{R}^N} (\partial_{x_1} w)^2 dx,$$

which implies (6.3) and completes the proof. □

Proof of Theorem 2.1 By Lemma 6.1, $F(\alpha)$ is 2π periodic and of class C^1 . Hence it has at least two critical points (maximum and minimum points) in $[0, 2\pi)$. Therefore, Theorem 2.1 follows from Lemma 6.2. □

7 Generalizations and discussion

In this section we first give some slight extensions of the results proved in the previous sections. Finally we would like to discuss some related questions we do not answer in this paper.

7.1 More general nonlinearities

Unlike the minimization method, we do not use the homogeneous property of the nonlinearity of Eq. (1.3). Therefore, our argument can be applied to construct infinitely many positive solutions for a more general problem:

$$\begin{cases} -\Delta u + V(x)u - f(u) = 0 & \text{in } \mathbb{R}^N, \\ u > 0 & \text{in } \mathbb{R}^N, u \in H^1(\mathbb{R}^N), \end{cases}$$

where $f : \mathbb{R} \rightarrow \mathbb{R}$ is at least $C^{1,\nu}(\mathbb{R})$ for some $\nu \in (0, 1)$, and satisfies the following conditions:

- (f₁) $f(u) = 0$ for $u \leq 0$, $f(0) = f'(0) = 0$;
- (f₂) The equation

$$\begin{cases} -\Delta u + V_\infty u - f(u) = 0 & \text{in } \mathbb{R}^N, \\ u > 0 & \text{in } \mathbb{R}^N, u \in H^1(\mathbb{R}^N), \end{cases}$$

has a nondegenerate solution w , in the sense that

$$\text{Ker}(-\Delta + 1 - f'(w)) \cap L^\infty(\mathbb{R}^N) = \text{Span}\{\partial_{x_1} w, \dots, \partial_{x_N} w\}.$$

Remark 7 It is not hard to see that our argument can be used to deal with the homogeneous Dirichlet boundary condition problem of (1.3) in $\mathbb{R}^N \setminus \overline{\Omega}$, where Ω is a bounded domain in \mathbb{R}^N .

7.2 Sign-changing solutions

Suppose that (1.11) holds for some constants

$$V_\infty > 0, a < 0, \text{ and } \min \left\{ 1, \frac{p-1}{2} \right\} m > 2, \sigma > 2.$$

If $N \geq 3$, we further assume (1.15), using almost the same argument, our method can be applied to construct infinitely many sign-changing solutions of the problem

$$-\Delta u + V(x)u - |u|^{p-1}u = 0 \text{ in } \mathbb{R}^N, u \in H^1(\mathbb{R}^N). \tag{7.1}$$

A similar result can be found in [13] when $V(x)$ tends to V_∞ from below with a suitable rate. We emphasize that our method can be applied to a more general non-even nonlinearity.

7.3 Remarks on condition (1.14)

In this subsection we consider the possible ways to improve the condition (1.14). Recall that the key step in our method is to solve the following equation:

$$-a_0 m R^{-m-2} T \mathbf{q} + \Phi(\alpha, \mathbf{q}) = \gamma \mathbf{q}_1,$$

where the property of T has been described in Lemma 5.4. We pose the condition (1.14) such that $\Phi(\alpha, \mathbf{q})$ is a smaller term comparing to $R^{-m-2} T \mathbf{q}$. Hence, to refine the condition (1.14), the estimate of $\Phi(\alpha, \mathbf{q})$ is the key point. A better estimate on $\Phi(\alpha, \mathbf{q})$ gives a weaker condition on m and σ . For example, for $N = 2$, if we assume that the following asymptotic behaviour of V holds in the C^1 sense:

$$V(x) = V_\infty + \frac{a}{|x|^m} + \frac{a_1(\theta)}{|x|^{m+1}} + O\left(\frac{1}{|x|^{m+1+\sigma_1}}\right), \text{ as } |x| \rightarrow \infty,$$

where $a_1(\theta)$ is an 2π periodic smooth function. Here (r, θ) is the polar coordinate. Then “ $\sigma > 2$ ” in the condition (1.14) can be improved to be “ $\sigma_1 > 0$ ”.

To get a better estimate of $\Phi(\alpha, \mathbf{q})$, an improvement of approximation is needed. Recall that

$$E = \underbrace{\sum_{j=1}^K (V(x) - 1) w_{Q_j}}_{E_1} - \underbrace{\left\{ \left(\sum_{j=1}^K w_{Q_j} \right)^p - \sum_{j=1}^K w_{Q_j}^p \right\}}_{E_2}.$$

The leading term of E_1 is given by $\sum_{j=1}^K (V(Q_j) - 1) w_{Q_j}$, which is $O(R^{-m})$ by (1.11).

Moreover, it is known that $\varphi_0 = -\frac{1}{p-1} w - \frac{1}{2} x \cdot \nabla w$ is the explicit solution of

$$L_0[\varphi_0] = -\Delta \varphi_0 + \varphi_0 - p w^{p-1} \varphi_0 = w \text{ in } \mathbb{R}^N, \text{ and } \int_{\mathbb{R}^N} \varphi_0 \nabla w \, dx = 0.$$

Thanks to the polynomial decay of V , we can improve the approximation and write E_1 into two parts: one is $O(R^{-m})$ having explicit form; and the other one is $O(R^{-m-1})$, which is enough for a better estimate of the part in $\Phi(\alpha, \mathbf{q})$ from E_1 .

However, for the term E_2 , the situation is more difficult. Recall that in the proof of Claim 2 of Lemma 4.4, we show that $E_2 \sim d^{-\frac{N-1}{2}} e^{-\min\{1, \frac{p}{2}\}d}$. Since $w(r) \sim r^{-\frac{N-1}{2}} e^{-r}$, subtracting the terms of order $d^{-\frac{N-1}{2}} e^{-\min\{1, \frac{p}{2}\}d}$, the next term is $O(d^{-\frac{N-1}{2}-1} e^{-\min\{1, \frac{p}{2}\}d})$. It is not enough for a better estimate of the part in $\Phi(\alpha, \mathbf{q})$ from E_2 .

Except for getting a better estimate of $\Phi(\alpha, \mathbf{q})$, another feasible way to improve the condition (1.14) is to apply min-max theorems to study the reduced energy functional $\mathcal{E}(U)$ and its small perturbations. A useful property is that the matrix T has only one zero eigenvalue, $K - 1$ negative eigenvalues, and K positive eigenvalues.

7.4 The anisotropic case

Observe that the leading term in (1.11) is radial. Thus it is interesting to consider the fully anisotropic case of the problem (1.3) for $N = 2$.

Question 1 *Do there still exist infinitely many positive solution of problem (1.3) if there are constants $V_\infty > 0, m > 1, \sigma > 0$ and a positive 2π periodic smooth function $a(\theta)$ such that*

$$V(x) = V_\infty + \frac{a(\theta)}{|x|^m} + O\left(\frac{1}{|x|^{m+\sigma}}\right), \text{ as } |x| \rightarrow \infty. \tag{V3}$$

Here (r, θ) is the polar coordinate.

Inspired by the results in [4, 14], the answer is very likely yes. But in this situation, the idea of uniformly distribution of points on curves does not work. Indeed, let $\Gamma = (r(\theta), \theta)$ be a closed curve in the polar coordinate system. Denote its length by L and its natural parameter by $s(\theta)$. If one puts K points on the stretched curve $R\Gamma$ for some positive constant R . After some computations, we get the balancing condition on Γ and R :

$$\begin{cases} \left(\frac{1}{2} \int_{\mathbb{R}^N} w^2 dy\right) R^{-m-1} = C\Psi\left(\frac{RL}{K}\right) \frac{L}{K}, \\ C \{ma(\theta(s))\gamma(s) - a'(\theta(s))\gamma^\perp(s)\} |\gamma(s)|^{-m-2} + \gamma''(s) = 0, \end{cases} \tag{7.2}$$

where $C > 0$ is a constant, L is the length of γ and $\theta(s)$ is the inverse of $s(\theta)$. However, it can be proved that system (7.2) has a solution if and only if $a(\theta) \equiv \text{constant}$. Therefore, it is to be expected that the spikes cannot be uniformly distributed on a closed curve if $a(\theta)$ is not a constant function.

A feasible way to answer Question 1 is to develop a theory like [5]. But a more accurate reduction procedure would be required since the mutual angles between the adjacent rays connecting spikes and the origin goes to zero as K tends to infinity.

7.5 Optimal condition on the decay

It seems that our argument here can only deal with the case of polynomial decay. Inspired by [4, 14], it is reasonable to believe that there are infinitely many positive solutions when the potential V satisfies the following decay assumption:

$$\exists \bar{\eta} \in (0, \sqrt{V_\infty}) : \lim_{|x| \rightarrow \infty} (V(x) - V_\infty) e^{\bar{\eta}|x|} = +\infty. \tag{V4}$$

Hence it is natural to ask the following question:

Question 2 *Does $S_V \neq \emptyset$ for any potential V satisfying (1.7)? Is the condition (V4) sufficient and necessary for $\#S_V = \infty$, i.e., on the existence of infinitely many positive solution of problem (1.3)?*

7.6 Higher dimensions

For the higher dimension case, i.e., $N \geq 3$, as we have seen, our arguments still work under the weak symmetry condition (1.15). It is natural to ask the following question:

Question 3 Does the weak symmetry condition (1.15) can be dropped when $N \geq 3$? Do the points can be distributed (uniformly) over the surface of a higher dimensional set, such as sphere in \mathbb{R}^3 ?

7.7 Higher dimensional concentration phenomena

Next we turn to the higher dimensional concentration phenomena. Inspired by the results in [3,21,33,41] and [8,34], it is interesting to ask the following question:

Question 4 Does there exist solution of problem (1.3) concentrating on higher dimensional sets, e.g., curves? That is, does the Ambrosetti–Malchiodi–Ni conjecture in [3] still hold without the small parameter ε , even in the radial symmetry case? If the answer is yes, are the solutions constructed in Theorem 2.1 bifurcations sets?

Under the condition (1.7), Question 4 is not easy even in the radial symmetry case. For example, assuming that $N = 2$ and $V(x)$ is radially symmetric, if one try to construct a positive solution concentrating on a circle with radius R , a simple computation gives $V'(R) \sim 1/R$, which is incompatible with $\lim_{|x| \rightarrow \infty} V(x) = V_\infty$.

Acknowledgments The authors are grateful to Frank Pacard for sharing his ideas and for useful discussions. Manuel del Pino is supported by Fondecyt Grant 110181 and Fondo Basal CMM. J. Wei is supported by a NSERC from Canada. The research of W. Yao is supported by Fondecyt Grant 3130543.

Appendix A: Circulant matrices and proof of Lemma 5.4

In this section we will prove Lemma 5.4. To this end, we need some notation. Denote the K -dimensional complex vector space and the ring of $K \times K$ complex matrices by \mathbb{C}^K and \mathbb{M}_K , respectively. Let $\mathbf{b} = (b_1, b_2, \dots, b_K) \in \mathbb{C}^K$, we define a shift operator $S : \mathbb{C}^K \rightarrow \mathbb{C}^K$ by

$$S(b_1, b_2, \dots, b_K) = (b_K, b_1, \dots, b_{K-1}).$$

Definition 8.1 (cf. [28]) The circulant matrix $B = \text{circ} \{\mathbf{b}\}$ associated to the vector $\mathbf{b} = (b_1, b_2, \dots, b_K) \in \mathbb{C}^K$ is the $K \times K$ matrix whose n th row is $S^{n-1}\mathbf{b}$:

$$B = \begin{pmatrix} b_1 & b_2 & \cdots & b_{K-1} & b_K \\ b_K & b_1 & \cdots & b_{K-2} & b_{K-1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ b_3 & b_4 & \cdots & b_1 & b_2 \\ b_2 & b_3 & \cdots & b_K & b_1 \end{pmatrix}.$$

We denote by $\text{Circ}(K) \subset \mathbb{M}_K$ the set of all $K \times K$ complex circulant matrices.

With this notation, both A_1 and A_2 in (5.5) are $K \times K$ circulant matrices. In fact,

$$A_1 = \text{circ} \{(-2, 1, 0, \dots, 0, 1)\} \quad \text{and} \quad A_2 = \text{circ} \{(0, 1, 0, \dots, 0, -1)\}.$$

Let $\epsilon = e^{i\frac{2\pi}{K}}$ be a primitive K -th root of unity, we define

$$X_l = \frac{1}{\sqrt{K}}(1, \epsilon^{l-1}, \epsilon^{2(l-1)}, \dots, \epsilon^{(K-1)(l-1)})^T \in \mathbb{C}^K, \quad \text{for } l = 1, \dots, K,$$

and

$$P_K = \frac{1}{\sqrt{K}} \begin{pmatrix} 1 & 1 & \dots & 1 & 1 \\ 1 & \epsilon & \dots & \epsilon^{K-2} & \epsilon^{K-1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & \epsilon^{K-2} & \dots & \epsilon^{(K-2)^2} & \epsilon^{(K-2)(K-1)} \\ 1 & \epsilon^{K-1} & \dots & \epsilon^{(K-1)(K-2)} & \epsilon^{(K-1)^2} \end{pmatrix} \in \mathbb{M}_K.$$

For the circulant matrix $B = \text{circ} \{ \mathbf{b} \}$, let

$$\lambda_l = b_1 + b_2 \epsilon^{l-1} + \dots + b_K \epsilon^{(K-1)(l-1)}, \quad \text{for } l = 1, \dots, K. \tag{8.1}$$

A simple calculation shows that $BX_l = \lambda_l X_l$. Hence λ_l is an eigenvalue of B with normalized eigenvector X_l . Since $\{X_1, \dots, X_K\}$ is a linearly independent set of vectors in \mathbb{C}^K , all of the eigenvalues of B are given by $\lambda_l, l = 1, \dots, K$.

Lemma 8.2 (cf. [28]) *All circulant matrices have the same ordered set of orthonormal eigenvectors $\{X_l\}$. Moreover, P_K is the diagonalizable matrix.*

Using these notation, we study the invertibility of T .

Lemma 8.3 *There is an $K_0 \in \mathbb{N}$ such that for all $K \geq K_0$ and every $\mathbf{b} \in \mathbb{R}^{2K}$, there exists a unique vector $\mathbf{q} \in \mathbb{R}^{2K}$ and a unique constant $\gamma \in \mathbb{R}$ such that*

$$T\mathbf{q} = \mathbf{b} + \gamma \mathbf{q}_0, \quad \mathbf{q} \perp \mathbf{q}_0. \tag{8.2}$$

Moreover, there is a positive constant C which is independent of K such that

$$\|\mathbf{q}\|_2 \leq C\|\mathbf{b}\|_2, \quad \|\dot{\mathbf{q}}\|_2 \leq C(\ln K)^{1/2}\|\mathbf{b}\|_2, \quad \text{and} \quad \|\ddot{\mathbf{q}}\|_2 \leq C(\ln K)^{3/2}\|\mathbf{b}\|_2. \tag{8.3}$$

Furthermore, the number of zero (negative, positive) eigenvalues of T is 1 ($K - 1, K$), respectively.

Proof Note that (8.3) is the Euclidean norm, hence it suffices to perform the analysis of the eigenvalues. To this end, first by (8.1), the eigenvalues of A_1 are

$$\lambda_{1,l} = -2 + \epsilon^{l-1} + \epsilon^{(K-1)(l-1)} = -4 \sin^2 \frac{(l-1)\pi}{K}, \quad l = 1, \dots, K, \tag{8.4}$$

and the eigenvalues of A_2 are

$$\lambda_{2,l} = \epsilon^{l-1} - \epsilon^{(K-1)(l-1)} = 2i \sin \frac{2(l-1)\pi}{K}, \quad l = 1, \dots, K. \tag{8.5}$$

Write $\text{diag} (c_1, \dots, c_K)$ for a diagonal matrix whose diagonal entries starting in the upper left corner are c_1, \dots, c_K . Denote the diagonal matrix of A_1 and A_2 by

$$D_1 = \text{diag} (\lambda_{1,1}, \lambda_{1,2}, \dots, \lambda_{1,K}) \quad \text{and} \quad D_2 = \text{diag} (\lambda_{2,1}, \lambda_{2,2}, \dots, \lambda_{2,K}), \quad \text{respectively.}$$

Since P_K is the diagonalizable matrix for circulant matrices, we have

$$P^{-1}TP = \begin{pmatrix} P_K^{-1} & 0 \\ 0 & P_K^{-1} \end{pmatrix} T \begin{pmatrix} P_K & 0 \\ 0 & P_K \end{pmatrix} = \begin{pmatrix} c_1 D_1 + c_4 I & c_2 D_2 \\ -c_2 D_2 & c_3 D_1 \end{pmatrix}. \tag{8.6}$$

Since the matrix T is real and symmetric, all its eigenvalues are real and satisfy the equations

$$\Lambda^2 - [(c_1 + c_3)\lambda_{1,l} + c_4] \Lambda + (c_1 \lambda_{1,l} + c_4) (c_3 \lambda_{1,l}) + c_2^2 \lambda_{2,l}^2 = 0, \tag{8.7}$$

for $l = 1, \dots, K$. Let

$$\alpha_l = (c_1 + c_3)\lambda_{1,l} + c_4, \text{ and } \beta_l = (c_1\lambda_{1,l} + c_4)(c_3\lambda_{1,l}) + c_2^2\lambda_{2,l}^2, \forall l = 1, \dots, K.$$

Then by (5.6), (8.4) and (8.5), we have

$$\alpha_l = (\widehat{d} - 1) \frac{K^2}{\pi^2} \sin^2 \frac{(l-1)\pi}{K} + (\widehat{d} - m - 1) > 0,$$

and

$$\beta_l = - \left\{ \left(\widehat{d} \frac{K^2}{\pi^2} - (\widehat{d} - 1)^2 \right) \sin^2 \frac{(l-1)\pi}{K} + (m-1)\widehat{d} + 1 \right\} \frac{K^2}{\pi^2} \sin^2 \frac{(l-1)\pi}{K} \leq 0.$$

Denote the solutions of (8.7) by $\Lambda_{1,l}$ and $\Lambda_{2,l}$ with $\Lambda_{1,l} \leq \Lambda_{2,l}$ for $l = 1, \dots, K$. Then

$$\Lambda_{1,l} = \frac{\alpha_l}{2} \left(-\sqrt{1 - \frac{4\beta_l}{\alpha_l^2}} + 1 \right) \leq 0, \quad \Lambda_{2,l} = \frac{\alpha_l}{2} \left(\sqrt{1 - \frac{4\beta_l}{\alpha_l^2}} + 1 \right) > 0, \quad \forall l = 1, \dots, K.$$

In particular, for $l = 1$, we have

$$\Lambda_{1,1} = 0, \quad \Lambda_{2,1} = \widehat{d} - m - 1. \tag{8.8}$$

For $l = 2, \dots, K$, by Lemma 3.3, we have

$$-\frac{4\beta_l}{\alpha_l^2} \leq \frac{\left(\widehat{d} \frac{K^2}{\pi^2} \sin^2 \frac{(l-1)\pi}{K} + m\widehat{d} \right) \frac{4K^2}{\pi^2} \sin^2 \frac{(l-1)\pi}{K}}{\left(\frac{\widehat{d} K^2}{2\pi^2} \sin^2 \frac{(l-1)\pi}{K} + \frac{\widehat{d}}{2} \right)^2} \leq \frac{C}{\widehat{d}},$$

and

$$-\frac{4\beta_l}{\alpha_l^2} \geq \frac{\left(\frac{\widehat{d} K^2}{2\pi^2} \sin^2 \frac{(l-1)\pi}{K} \right) \frac{4K^2}{\pi^2} \sin^2 \frac{(l-1)\pi}{K}}{\left(\widehat{d} \frac{K^2}{\pi^2} \sin^2 \frac{(l-1)\pi}{K} + \widehat{d} \right)^2} \geq \frac{C}{\widehat{d}}, \quad \forall l = 2, \dots, K,$$

where C is a positive constant. Therefore, for all $l = 2, \dots, K$,

$$-\Lambda_{1,l} \geq \frac{\widehat{d}}{2} \cdot \frac{C}{\widehat{d}} \geq \frac{1}{2}C, \text{ and } \Lambda_{2,l} \geq \alpha_l \geq \frac{1}{2}\widehat{d}, \text{ for some constant } C > 0,$$

from which we get $\|\mathbf{q}\|_2 \leq C\|\mathbf{b}\|_2$.

Define $\widehat{f}_j = f_{j+1} - f_j$ and $\widehat{g}_j = g_{j+1} - g_j$. Then

$$\begin{cases} c_1(\widehat{f}_j - \widehat{f}_{j-1}) + c_2(\widehat{g}_j + \widehat{g}_{j-1}) = \phi_j - c_4f_j, \\ -c_2(\widehat{f}_j + \widehat{f}_{j-1}) + c_3(\widehat{g}_j - \widehat{g}_{j-1}) = \varphi_j, \end{cases}$$

where $\phi_j = \mathbf{b}_j$ and $\varphi_j = \mathbf{b}_{K+j}$ for $j = 1, \dots, K$. By a similar argument in the proof of $\|\mathbf{q}\|_2 \leq C\|\mathbf{b}\|_2$, we can get

$$\|\dot{\mathbf{q}}\|_2 \leq C(\ln K)^{1/2}\|\mathbf{b}\|_2.$$

When this is done, let $\widetilde{f}_j = f_{j+1} - 2f_j + f_{j-1}$ and $\widetilde{g}_j = g_{j+1} - 2g_j + g_{j-1}$. Then

$$\begin{cases} c_1\widetilde{f}_j = \phi_j - c_4f_j - c_2(\widehat{g}_j + \widehat{g}_{j-1}), \\ c_3\widetilde{g}_j = \varphi_j + c_2(\widehat{f}_j + \widehat{f}_{j-1}). \end{cases}$$

Using a similar argument, by the definition of c_j 's, we get

$$\|\ddot{\mathbf{q}}\|_2 \leq C(\ln K)^{3/2} \|\mathbf{b}\|_2.$$

□

Now we are going to prove Lemma 5.4. An important observation is that the system $T\mathbf{q} = \mathbf{b}$ can be seen as the discretization of the following continuous system:

$$\begin{cases} -(m + 1)f(\theta) + (f'' + f)(\theta) + (\widehat{d} - 1)(f + g')(\theta) = \phi(\theta), & \theta \in (0, 2\pi), \\ f'(\theta) - \widehat{d}(f' + g'')(\theta) = \varphi(\theta), & \theta \in (0, 2\pi), \\ f(0) = f(2\pi), f'(0) = f'(2\pi), g(0) = g(2\pi), g'(0) = g'(2\pi), \end{cases} \tag{8.9}$$

where $f'' + f$ is the Jacobi operator on S^1 .

Lemma 8.4 *For K sufficiently large, given ϕ, φ satisfying $\int_0^{2\pi} \varphi = 0$, the system (8.9) has a unique solution (f, g) satisfying $\int_0^{2\pi} g = 0$. Moreover, there exists a constant $C > 0$ such that*

$$\|f\|_{C^2([0, 2\pi])} + \|g\|_{C^2([0, 2\pi])} \leq C (\|\phi\|_{C^0([0, 2\pi])} + \|\varphi\|_{C^0([0, 2\pi])}). \tag{8.10}$$

Proof Let $h = \widehat{d}(f + g')$, then system (8.9) becomes

$$\begin{cases} f'' - mf + \frac{\widehat{d}-1}{\widehat{d}}h = \phi, \\ f' - h' = \varphi. \\ f, h \text{ are } 2\pi \text{ periodic.} \end{cases} \tag{8.11}$$

Since $\int_0^{2\pi} \varphi = 0$, from the second equation we get

$$h(\theta) = f(\theta) - \int_0^\theta \varphi - c_h. \tag{8.12}$$

Here we take $c_h = \frac{1-\widehat{d}}{2\pi} \int_0^{2\pi} f - \frac{1}{2\pi} \int_0^{2\pi} \int_0^\theta \varphi$ such that $-\int_0^{2\pi} f + \widehat{d}^{-1} \int_0^{2\pi} h = 0$. Hence g can be solved by

$$g(\theta) = - \int_0^\theta f + \widehat{d}^{-1} \int_0^\theta h + c_g,$$

where we take $c_g = \frac{1}{2\pi} \int_0^{2\pi} \int_0^\theta f - \widehat{d}^{-1} \frac{1}{2\pi} \int_0^{2\pi} \int_0^\theta h$ such that $\int_0^{2\pi} g = 0$.

By (8.12) the first equation in (8.11) becomes

$$f'' - (m - 1 + \frac{1}{\widehat{d}})f = \phi + \frac{\widehat{d} - 1}{\widehat{d}} \left[\int_0^\theta \varphi - \frac{1}{2\pi} \int_0^{2\pi} \int_0^\theta \varphi \right] - \frac{(\widehat{d} - 1)^2}{2\pi \widehat{d}} \int_0^{2\pi} f.$$

To solve the above equation, we first integrate it over $[0, 2\pi]$ to get $\int_0^{2\pi} f = \frac{1}{\widehat{d}-1-m} \int_0^{2\pi} \phi$. Thus

$$f'' - (m - 1 + \frac{1}{\widehat{d}})f = \left[\phi - \frac{(\widehat{d} - 1)^2}{\widehat{d}(\widehat{d} - 1 - m)} \frac{1}{2\pi} \int_0^{2\pi} \phi \right] + \frac{\widehat{d} - 1}{\widehat{d}} \left[\int_0^\theta \varphi - \frac{1}{2\pi} \int_0^{2\pi} \int_0^\theta \varphi \right].$$

Note that $m - 1 + \frac{1}{\widehat{d}} > 0$, by the boundary condition, f is uniquely given and satisfies $\|f\|_{C^2([0,2\pi])} \leq C (\|\phi\|_{C^0([0,2\pi])} + \|\varphi\|_{C^0([0,2\pi])})$. Therefore,

$$g(\theta) = - \int_0^\theta f + \widehat{d}^{-1} \int_0^\theta h + c_g,$$

satisfies the same inequality, where

$$h(\theta) = f(\theta) + \frac{(\widehat{d} - 1)}{2\pi(\widehat{d} - 1 - m)} \int_0^{2\pi} \phi - \left[\int_0^\theta \varphi - \frac{1}{2\pi} \int_0^{2\pi} \int_0^\theta \varphi \right].$$

□

Remark 8 Let $c = \sqrt{m - 1 + \frac{1}{\widehat{d}}}$, by the equation of f , we can get

$$f(\theta) = \int_0^{2\pi} \left[\frac{(\widehat{d} - 1)^2}{\widehat{d}(\widehat{d} - m - 1)} \frac{1}{2\pi c^2} - G_0(\theta, s) \right] \phi(s) ds + \int_0^{2\pi} \frac{\widehat{d} - 1}{\widehat{d}} \left[\int_0^t G_0(\theta, t) - \frac{s}{2\pi c^2} \right] \varphi(s) ds,$$

where

$$G_0(\theta, s) = \begin{cases} \frac{1}{2c(e^{2\pi c} - 1)} [e^{2\pi c} e^{c(\theta-s)} + e^{-c(\theta-s)}], & \text{if } \theta \leq s, \\ \frac{1}{2c(e^{2\pi c} - 1)} [e^{c(\theta-s)} + e^{2\pi c} e^{-c(\theta-s)}], & \text{if } \theta > s. \end{cases}$$

Moreover, there is a Green’s matrix

$$G(\theta, t) = \begin{pmatrix} G_{11}(\theta, s) & G_{12}(\theta, s) \\ G_{21}(\theta, s) & G_{22}(\theta, s) \end{pmatrix}$$

such that

$$f(\theta) = \int_0^{2\pi} G_{11}(\theta, s) \phi(s) ds + \int_0^{2\pi} G_{12}(\theta, s) \varphi(s) ds,$$

and

$$g(\theta) = \int_0^{2\pi} G_{21}(\theta, s) \phi(s) ds + \int_0^{2\pi} G_{22}(\theta, s) \varphi(s) ds.$$

Actually,

$$G_{11}(\theta, s) = \frac{(\widehat{d} - 1)^2}{\widehat{d}(\widehat{d} - m - 1)} \frac{1}{2\pi c^2} - G_0(\theta, s),$$

$$G_{12}(\theta, s) = \frac{\widehat{d} - 1}{\widehat{d}} \left[\int_0^s G_0(\theta, t) dt - \frac{s}{2\pi c^2} \right],$$

$$G_{21}(\theta, s) = \frac{(\widehat{d} - 1)^3}{\widehat{d}^2(\widehat{d} - m - 1)} \left(\frac{1}{2c^2} - \frac{\theta}{2\pi c^2} \right) + \frac{(\widehat{d} - 1)}{\widehat{d}(\widehat{d} - m - 1)} \left(\frac{\theta}{2\pi} - \frac{1}{2} \right) + \frac{(\widehat{d} - 1)}{\widehat{d}} \left[\int_0^\theta G_0(t, s) dt - \frac{1}{2\pi} \int_0^{2\pi} \int_0^\theta G_0(t, s) dt d\theta \right],$$

and

$$G_{22}(\theta, s) = \frac{(\widehat{d} - 1)}{\widehat{d}} \left(\frac{\theta s}{2\pi c^2} - \frac{s}{2c^2} \right) + \frac{1}{\widehat{d}} \left[\left(\pi - \frac{s}{2} + \frac{s^2}{4\pi} \right) - (\theta - s)_+ - \frac{\theta s}{2\pi} \right] + \frac{(\widehat{d} - 1)}{\widehat{d}} \left[- \int_0^\theta \int_0^s G_0(t, r) dr dt + \frac{1}{2\pi} \int_0^{2\pi} \int_0^\theta \int_0^s G_0(t, r) dr dt d\theta \right].$$

Lemma 8.5 *Under the assumption of Lemma 8.3, there is a positive constant C which is independent of K such that*

$$\|\mathbf{q}\|_* \leq C(\ln K)^2 \|\mathbf{b}\|_\infty. \tag{8.13}$$

Proof **Claim 1:** There is a positive constant C (independent of K) such that

$$\|\mathbf{q}\|_\infty \leq C \|\mathbf{b}\|_\infty. \tag{8.14}$$

To prove it, we only need to consider the case $\mathbf{b} \perp \mathbf{q}_0$. For $j = 1, \dots, K$, we define

$$\mathbf{q}_j = (f_{1,j}, \dots, f_{K,j}, g_{1,j}, \dots, g_{K,j})^T,$$

where

$$f_{l,j} = \frac{2\pi}{K} G_{11}(\theta_l, \theta_j), \quad g_{l,j} = \frac{2\pi}{K} G_{21}(\theta_l, \theta_j).$$

This corresponds to take $\phi = \frac{2\pi}{K} \delta(\theta - \theta_j)$, $\varphi = 0$ for $j = 1, \dots, K$, where δ is the delta function in the distribution theory. For $j = K + 2, \dots, 2K$, we define

$$\mathbf{q}_j = (f_{1,j}, \dots, f_{K,j}, g_{1,j}, \dots, g_{K,j})^T,$$

where

$$f_{l,j} = -\frac{2\pi}{K} G_{12}(\theta_l, \theta_1) + \frac{2\pi}{K} G_{12}(\theta_l, \theta_j), \quad g_{l,j} = -\frac{2\pi}{K} G_{22}(\theta_l, \theta_1) + \frac{2\pi}{K} G_{22}(\theta_l, \theta_j).$$

This corresponds to take $\phi = 0$, $\varphi = -\frac{2\pi}{K} \delta(\theta - \theta_1) + \frac{2\pi}{K} \delta(\theta - \theta_j)$ for $j = K + 2, \dots, 2K$. By the property of Green’s matrix, for $j = 1, \dots, K$, we have

$$T\mathbf{q}_j = \vec{e}_j + \vec{\tau}_j;$$

for $j = K + 2, \dots, 2K$, we have

$$T\mathbf{q}_j = -\vec{e}_{K+1} + \vec{e}_j + \vec{\tau}_j,$$

where $\vec{e}_j = (\delta_{j,1}, \dots, \delta_{j,2K})^T$ is standard orthonormal basis of \mathbb{R}^{2K} and $\vec{\tau}_j = O(K^{-2})$ is the local truncation error for the Green’s matrix in the finite difference method. Since $\{\vec{e}_1, \dots, \vec{e}_K, -\vec{e}_{K+1} + \vec{e}_{K+2}, -\vec{e}_{K+1} + \vec{e}_{2K}\}$ is a basis of $\{\mathbf{b} \in \mathbb{R}^{2K} \mid \mathbf{b} \perp \mathbf{q}_0\}$ and write $\mathbf{b} =$

$\sum_{j=1}^K b_j \vec{e}_j + \sum_{j=K+2}^{2K} b_j (-\vec{e}_{K+1} + \vec{e}_j)$. Then we get $\mathbf{q} = \widehat{\mathbf{q}} + \widetilde{\mathbf{q}}$, where $\widehat{\mathbf{q}} = \sum_{j=1}^K b_j \mathbf{q}_j + \sum_{j=K+2}^{2K} b_j \mathbf{q}_j$ and $T\widetilde{\mathbf{q}} = \sum_{j \neq K+1} b_j \vec{e}_j$.

On one hand, by the property of Green’s matrix, we get $\|\widehat{\mathbf{q}}\|_\infty \leq C\|\mathbf{b}\|_\infty$. On the other hand, by Lemma 8.3, we get

$$\|\widetilde{\mathbf{q}}\|_\infty \leq \|\widetilde{\mathbf{q}}\|_2 \leq C\|\sum_{j \neq K+1} b_j \vec{e}_j\|_2 \leq CK^{1/2}\|\sum_{j \neq K+1} b_j \vec{e}_j\|_\infty \leq CK^{-1/2}\|\mathbf{b}\|_\infty.$$

Combining these two estimates, we get $\|\mathbf{q}\|_\infty \leq C\|\mathbf{b}\|_\infty$.

Claim 2: There is a positive constant C (independent of K) such that

$$\|\mathbf{q}\|_* = \|\mathbf{q}\|_\infty + \|\dot{\mathbf{q}}\|_\infty + \|\ddot{\mathbf{q}}\|_\infty \leq C(\ln K)^2\|\mathbf{b}\|_\infty. \tag{8.15}$$

Proof of Claim 2:

Define $\widehat{f}_j = f_{j+1} - f_j$ and $\widehat{g}_j = g_{j+1} - g_j$. Then

$$\begin{cases} c_1(\widehat{f}_j - \widehat{f}_{j-1}) + c_2(\widehat{g}_j + \widehat{g}_{j-1}) = \phi_j - c_4 f_j, \\ -c_2(\widehat{f}_j + \widehat{f}_{j-1}) + c_3(\widehat{g}_j - \widehat{g}_{j-1}) = \varphi_j. \end{cases}$$

By using a similar argument, one can get

$$\|\dot{\mathbf{q}}\|_\infty \leq C\|\mathbf{b}\|_\infty + (\ln K)\|\mathbf{q}\|_\infty \leq C(\ln K)\|\mathbf{b}\|_\infty. \tag{8.16}$$

Let $\widetilde{f}_j = f_{j+1} - 2f_j + f_{j-1}$ and $\widetilde{g}_j = g_{j+1} - 2g_j + g_{j-1}$. Then

$$\begin{cases} c_1 \widetilde{f}_j = \phi_j - c_4 f_j - c_2(\widehat{g}_j + \widehat{g}_{j-1}), \\ c_3 \widetilde{g}_j = \varphi_j + c_2(\widehat{f}_j + \widehat{f}_{j-1}). \end{cases}$$

Similarly we obtain

$$\|\ddot{\mathbf{q}}\|_\infty \leq C\|\mathbf{b}\|_\infty + (\ln K)\|\mathbf{q}\|_\infty + (\ln K)\|\dot{\mathbf{q}}\|_\infty \leq C(\ln K)^2\|\mathbf{b}\|_\infty. \tag{8.17}$$

The above estimate is enough for our application. Actually if one can show that $\|\widehat{\mathbf{q}}\|_* \leq C\|\mathbf{b}\|_\infty$, then we can get $\|\mathbf{q}\|_* \leq C\|\mathbf{b}\|_\infty$ by Lemma 8.3 since the local truncation error for the Green’s matrix is $O(K^{-2})$. □

Now we can use Lemma 8.5 to prove Lemma 5.4.

Proof of Lemma 5.4 To prove Lemma 5.4, it suffices to prove the a priori estimate (5.8). let $\gamma = -(\mathbf{b} \cdot \mathbf{q}_0)/(\mathbf{q}_1 \cdot \mathbf{q}_0)$. By Lemma 4.2, for \mathbf{q} satisfies (2.9), we have

$$R^{-1}\mathbf{q}_1 = c_0\mathbf{q}_0 + O(KR^{-m}),$$

which implies that $\|R^{-1}\mathbf{q}_1\|_\infty \leq C$ and $|R^{-1}\mathbf{q}_1 \cdot \mathbf{q}_0| \geq CK$. Hence $\|\gamma\mathbf{q}_1\|_\infty \leq C\|\mathbf{b}\|_\infty$. Therefore, by Lemma 8.5, we have

$$\|\widetilde{\mathbf{q}}\|_* \leq C(\ln K)^2\|\mathbf{b} + \gamma\mathbf{q}_1\|_\infty \leq C(\ln K)^2\|\mathbf{b}\|_\infty.$$

□

Appendix B: Energy expansion

In this section, we give the energy expansion of $\mathcal{E}(U + \phi)$ and prove Lemma 3.9.

Proof of Lemma 3.9 By (1.4), we get

$$\begin{aligned} \mathcal{E}(u) &= \underbrace{\sum_{j=1}^K \mathcal{E}(w_{Q_j})}_{J_1} + \underbrace{\frac{1}{2} \sum_{i \neq j} \int_{\mathbb{R}^N} (\nabla w_{Q_i} \nabla w_{Q_j} + V(x) w_{Q_i} w_{Q_j}) dx}_{J_2} \\ &\quad + \underbrace{\frac{1}{p+1} \int_{\mathbb{R}^N} \left\{ \sum_{j=1}^K w_{Q_j}^{p+1} - \left(\sum_{j=1}^K w_{Q_j} \right)^{p+1} \right\} dx}_{J_3}. \end{aligned}$$

Claim 1: By (1.11), there are positive constants I_0 and a_0 such that

$$J_1 = KI_0 + a_0(1 + o(1)) \sum_{j=1}^K |Q_j|^{-m}. \tag{9.1}$$

Indeed, by the definition of the energy functional, (1.11) and Taylor’s expansion,

$$J_1 = KI_0 + \frac{1}{2} \sum_{j=1}^K \int_{\mathbb{R}^N} (V(x) - 1) w_{Q_j}^2 dx = KI_0 + a_0 \sum_{j=1}^K (|Q_j|^{-m} + O(R^{-m-\sigma})),$$

where

$$I_0 = \left(\frac{1}{2} - \frac{1}{p+1} \right) \int_{\mathbb{R}^N} w^{p+1} dx, \text{ and } a_0 = \frac{a}{2} \int_{\mathbb{R}^N} w^2 dx.$$

Claim 2: By (1.11) we have

$$J_2 = \frac{1}{2} \sum_{i \neq j} \int_{\mathbb{R}^N} w_{Q_j}^p w_{Q_i} dx + O(KR^{-2m}). \tag{9.2}$$

Indeed, the term J_2 can be divided into two parts:

$$J_2 = \underbrace{\frac{1}{2} \sum_{i \neq j} \int_{\mathbb{R}^N} w_{Q_j}^p w_{Q_i} dx}_{J_{21}} + \underbrace{\frac{1}{2} \sum_{i \neq j} \int_{\mathbb{R}^N} (V(x) - 1) w_{Q_j} w_{Q_i} dx}_{J_{22}}.$$

For J_{22} , by (1.11), we have

$$|J_{22}| \leq CR^{-m} \sum_{i \neq j} e^{-|Q_j - Q_i|} |Q_j - Q_i|^{-(N-3)/2} \leq CKR^{-m} e^{-d} d^{-(N-3)/2} \leq CKR^{-2m}.$$

Claim 3: Let $\mathbf{Q} \in \Lambda_K$, we have

$$J_3 = - \sum_{i \neq j} \int_{\mathbb{R}^N} w_{Q_j}^p w_{Q_i} dx + O(K e^{-\min\{2, \frac{p+1}{2}\}d} d^{-\frac{N-3}{2}}). \tag{9.3}$$

Indeed, write

$$J_3 = \frac{1}{p+1} \int_{\mathbb{R}^N} \left\{ \sum_{j=1}^K w_{Q_j}^{p+1} - \left(\sum_{j=1}^K w_{Q_j} \right)^{p+1} \right\} dx = - \int_{\mathbb{R}^N} E_3 dx.$$

For $x \in \Omega_{K+1}^\ell$, where $\ell \in \mathbb{N}$ chosen later, we have

$$|E_3(x)| \leq \frac{1}{p+1} \left\{ \sum_{j=1}^K w_{Q_j}^{p+1} + \left(\sum_{j=1}^K w_{Q_j} \right)^{p+1} \right\} \leq \frac{1}{p+1} \left(\sum_{j=1}^K w_{Q_j}^{p+1} + K^p \sum_{j=1}^K w_{Q_j}^{p+1} \right).$$

By choosing $\ell \geq \frac{4(p+2m+3)}{pm}$ (but independent of K), we get

$$\int_{\Omega_{K+1}^\ell} |E_3| dx \leq CK^p \sum_{j=1}^K \int_{\Omega_{K+1}^\ell} w_{Q_j}^{p+1} dx \leq CK^{p+1} w^p(\ell\rho/2) \leq CK^{-2m-2}.$$

For $x \in \Omega_j^\ell, j = 1, 2, \dots, K$, by a similar argument in the proof of Lemma 4.4, we have

$$\left| E_3 - w_{Q_j}^p \sum_{i \neq j} w_{Q_i} \right| \leq C \ell^{(N-1)(p-1)} w_{Q_j}^{p-1} \left(\sum_{i \neq j} w_{Q_i} \right)^2.$$

Applying Lemma 3.8, we get

$$\begin{aligned} \left| J_3 + \sum_{i \neq j} \int_{\mathbb{R}^N} w_{Q_j}^p w_{Q_i} \right| &\leq C \sum_{j=1}^K \int_{\Omega_j^\ell} w_{Q_j}^{p-1} \left(\sum_{i \neq j} w_{Q_i} \right)^2 dx + CK^{-2m-2} \\ &\leq CK e^{-\min\{2, \frac{p+1}{2}\}d} d^{-\frac{N-3}{2}} \leq CK^{-\min\{2, \frac{p+1}{2}\}m+1} (\ln K)^{1/2}, \end{aligned}$$

which implies Claim 3.

Combining our Claim 1, Claim 2 and Claim 3, the desired result follows from Lemma 3.8. □

At the last, by (1.4), Lemma 4.4 and Proposition 4.1,

$$\begin{aligned} \mathcal{E}(U + \phi) &= \mathcal{E}(U) + \frac{1}{2} \int_{\mathbb{R}^N} \{ |\nabla \phi|^2 + V(x)\phi^2 \} dx + \int_{\mathbb{R}^N} (\nabla U \nabla \phi + V(x)U\phi) dx \\ &\quad - \frac{1}{p+1} \int_{\mathbb{R}^N} \{ (U + \phi)_+^{p+1} - U^{p+1} \} dx \\ &= \mathcal{E}(U) + \frac{1}{2} \int_{\mathbb{R}^N} \{ (U + \phi)_+^p - U^p + E \} \phi dx \\ &\quad - \frac{1}{p+1} \int_{\mathbb{R}^N} \{ (U + \phi)_+^{p+1} - U^{p+1} - (p+1)U^p \phi \} dx \\ &= \mathcal{E}(U) + O(K \|\phi\|_{**}^2) + O(K \|\phi\|_{**} \|E\|_{**}) \end{aligned}$$

$$\begin{aligned}
&= KI_0 + (a_0 + o(1)) \sum_{j=1}^K |Q_j|^{-m} - \frac{1}{2} \sum_{i \neq j} (\gamma_0 + o(1)) w(|Q_i - Q_j|) \\
&\quad + O\left(K^{-\min\{2, \frac{p+1}{2}\}m+1} (\ln K)^{1/2}\right).
\end{aligned}$$

References

- Ambrosetti, A., Badiale, M., Cingolani, S.: Semiclassical states of nonlinear Schrödinger equations. *Arch. Ration. Mech. Anal.* **140**(3), 285–300 (1997)
- Ambrosetti, A., Colorado, E., Ruiz, D.: Multi-spike solitons to linearly coupled systems of nonlinear Schrödinger equations. *Calc. Var. Partial Differ. Equ.* **30**(1), 85–112 (2007)
- Ambrosetti, A., Malchiodi, A., Ni, W.-M.: Singularly perturbed elliptic equations with symmetry: existence of solutions concentrating on spheres. I. *Commun. Math. Phys.* **235**(3), 427–466 (2003)
- Ao, W., Wei, J.: Infinitely many positive solutions for nonlinear equations with non-symmetric potential. *Calc. Var. Partial Differ. Equ.* (preprint)
- Ao, W., Musso, M., Pacard, F., Wei, J.: Solutions without any symmetry for semilinear elliptic problems (preprint)
- Bahri, A., Li, Y.Y.: On a min-max procedure for the existence of a positive solution for certain scalar field equations in \mathbb{R}^N . *Rev. Mat. Iberoamericana* **6**(1–2), 1–15 (1990)
- Bahri, A., Lions, P.L.: On the existence of a positive solution of semilinear elliptic equations in unbounded domains. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **14**(3), 365–413 (1997)
- Berestycki, H., Wei, J.: On least energy solutions to a semilinear elliptic equation in a strip. *Discrete Contin. Dyn. Syst.* **28**(3), 1083–1099 (2010)
- Cao, D.: Positive solution and bifurcation from the essential spectrum of a semilinear elliptic equation on \mathbb{R}^n . *Nonlinear Anal.* **15**(11), 1045–1052 (1990)
- Cao, D., Noussair, E.S., Yan, S.: Existence and uniqueness results on single-peaked solutions of a semilinear problem. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **15**(1), 73–111 (1998)
- Cao, D., Noussair, E.S., Yan, S.: Solutions with multiple peaks for nonlinear elliptic equations. *Proc. R. Soc. Edinb. Sect. A* **129**(2), 235–264 (1999)
- Berestycki, H., Lions, P.-L.: Nonlinear scalar field equations. I. Existence of a ground state. *Arch. Ration. Mech. Anal.* **82**(4), 313–345 (1983)
- Cerami, G., Devillanova, G., Solimini, S.: Infinitely many bound states for some nonlinear scalar field equations. *Calc. Var. Partial Differ. Equ.* **23**(2), 139–168 (2005)
- Cerami, G., Passaseo, D., Solimini, S.: Infinitely many positive solutions to some scalar field equations with nonsymmetric coefficients. *Commun. Pure Appl. Math.* **66**(3), 372–413 (2013)
- Cerami, G., Molle, R.: On some Schrödinger equations with non regular potential at infinity. *Discrete Contin. Dyn. Syst.* **28**(2), 827–844 (2010)
- Coti Zelati, V., Rabinowitz, P.: Homoclinic type solutions for a semilinear elliptic PDE on \mathbb{R}^n . *Commun. Pure Appl. Math.* **45**(10), 1217–1269 (1992)
- del Pino, M., Felmer, P.: Local mountain passes for semilinear elliptic problems in unbounded domains. *Calc. Var. Partial Differ. Equ.* **4**(2), 121–137 (1996)
- del Pino, M., Felmer, P.: Semi-classical states for nonlinear Schrödinger equations. *J. Funct. Anal.* **149**(1), 245–265 (1997)
- del Pino, M., Felmer, P.: Multi-peak bound states of nonlinear Schrödinger equations. *Ann. Inst. H. Poincaré, Anal. Non Linéaire* **15**(2), 1271–1299 (1998)
- del Pino, M., Felmer, P.: Semi-classical states of nonlinear Schrödinger equations: a variational reduction method. *Math. Ann.* **324**(1), 1–32 (2002)
- del Pino, M., Kowalczyk, M., Wei, J.: Concentration on curves for nonlinear Schrödinger equations. *Commun. Pure Appl. Math.* **70**(1), 113–146 (2007)
- del Pino, M., Kowalczyk, M., Wei, J.: On de Giorgi conjecture in dimension $n \geq 9$. *Ann. Math.* **174**(3), 1485–1569 (2011)
- Devillanova, G., Solimini, S.: Min-Max solutions to some scalar field equations. *Adv. Nonlinear Stud.* **12**(1), 173–186 (2012)
- Ding, W., Ni, W.-M.: On the existence of positive entire solutions of a semilinear elliptic equation. *Arch. Ration. Mech. Anal.* **91**(4), 283–308 (1986)
- Floer, A., Weinstein, A.: Nonspreading wave packets for the cubic Schrödinger equation with a bounded potential. *J. Funct. Anal.* **69**(3), 397–408 (1986)

26. Gidas, B., Ni, W.-M., Nirenberg, L.: Symmetry of positive solutions of nonlinear elliptic equations in \mathbb{R}^n . In: *Mathematical Analysis and Applications. Part A. Advances in Mathematical Supplementary Studies*, vol. 7A, pp. 369–402. Academic Press, London (1981)
27. Kang, X., Wei, J.: On interacting spikes of semi-classical states of nonlinear Schrödinger equations. *Adv. Differ. Equ.* **5**(7–9), 899–928 (2000)
28. Kra, I., Simanca, S.R.: On circulant matrices. *Notices AMS* **59**(3), 368–377 (2012)
29. Kwong, M.K.: Uniqueness of positive solutions of $-\Delta u + u = u^p$ in \mathbb{R}^N . *Arch. Ration. Mech. Anal.* **105**(3), 243–266 (1989)
30. Lin, F.H., Ni, W.-M., Wei, J.: On the number of interior peak solutions for a singularly perturbed Neumann problem. *Commun. Pure Appl. Math.* **60**(2), 252–281 (2007)
31. Lions, P.L.: The concentration-compactness principle in the calculus of variations. The locally compact case. I. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **1**(2), 109–145 (1984)
32. Lions, P.L.: The concentration-compactness principle in the calculus of variations. The locally compact case. II. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **1**(4), 223–283 (1984)
33. Mahmoudi, F., Malchiodi, A., Montenegro, M.: Solutions to the nonlinear Schrödinger equation carrying momentum along a curve. *Commun. Pure Appl. Math.* **62**(9), 1155–1264 (2009)
34. Malchiodi, A.: Some new entire solutions of semilinear elliptic equations on \mathbb{R}^n . *Adv. Math.* **221**(6), 1843–1909 (2009)
35. Musso, M., Pacard, F., Wei, J.: Finite-energy sign-changing solutions with dihedral symmetry for the stationary nonlinear Schrödinger equation. *J. Eur. Math. Soc.* **14**(6), 1923–1953 (2012)
36. Ni, W.-M., Takagi, I.: Locating the peaks of least-energy solutions to a semilinear Neumann problem. *Duke Math. J.* **70**(2), 247–281 (1993)
37. Oh, Y.J.: On positive multi-lump bound states of nonlinear Schrödinger equations under multiple well potential. *Commun. Math. Phys.* **133**(2), 223–253 (1990)
38. Pacard, F., Ritore, M.: From constant mean curvature hypersurfaces to the gradient theory of phase transitions. *J. Differ. Geom.* **64**, 359–423 (2003)
39. Rabinowitz, P.H.: On a class of nonlinear Schrödinger equations. *Z. Angew. Math. Phys.* **43**(2), 207–291 (1992)
40. Strauss, W.A.: Existence of solitary waves in higher dimensions. *Commun. Math. Phys.* **55**(2), 149–162 (1977)
41. Wang, L., Wei, J., Yang, J.: On Ambrosetti–Malchiodi–Ni conjecture for general hypersurfaces. *Commun. Partial Differ. Equ.* **36**(12), 2117–2161 (2011)
42. Wang, X.: On concentration of positive bound states of nonlinear Schrödinger equations. *Commun. Math. Phys.* **153**(2), 229–243 (1993)
43. Wei, J., Yan, S.: Infinitely many positive solutions for the nonlinear Schrödinger equations in \mathbb{R}^N . *Calc. Var. Partial Differ. Equ.* **37**(3–4), 423–439 (2010)