

# Tabla de contenido

CAPITULO 1 .....	1
INTRODUCCIÓN.....	1
1.1 Estructura de la Tesis.....	1
1.2 Motivación.....	1
1.3 Trabajos Anteriores.....	3
1.4 Hipótesis de trabajo .....	6
1.5 Objetivos.....	6
1.5.1 Objetivo general .....	6
1.5.2 Objetivos específicos.....	6
CAPITULO 2 .....	7
GEOLOGÍA SECTOR PUCHULDIZA.....	7
2.1 Generalidades .....	7
2.2 Estratigrafía .....	7
2.2.1 Depósitos y Rocas Estratificadas.....	8
2.2.2 Volcanes y Secuencias Volcánicas.....	12
2.2.3 Rocas Intrusivas.....	12
2.3 Geología estructural.....	13
2.4 Hidrogeoquímica .....	14
2.4.1 Metodología.....	14
2.4.2 Geoquímica de aguas termales .....	15
2.4.5 Geotermometría.....	17

CAPITULO 3 .....	19
GEOCHEMISTRY OF METALS AND METALLOIDS IN SILICEOUS SINTER DEPOSITS: IMPLICATIONS FOR ELEMENTAL PARTITIONING INTO SILICA PHASES .....	19
Abstract.....	19
3.1 Introduction .....	20
3.2 Geological background and samples .....	23
3.3 Analytical Methods .....	24
3.4 Results .....	27
3.4.1 Sinter mineralogy and micro-morphology .....	27
3.4.2 Sinter geochemistry and hydrothermal fluid composition .....	30
3.5 Discussion.....	31
3.5.1 Silica precipitation and diagenetic maturation effects on trace element uptake..	31
3.5.2 Trace metal and metalloid enrichment of silica phases in sinter.....	35
3.6 Summary and concluding remarks .....	39
3.7 Acknowledgements.....	41
3.8 References.....	56
CAPITULO 4 .....	65
CONCLUSIONES.....	65
BIBLIOGRAFÍA .....	67
ANEXOS .....	73
Sitios de Muestreo .....	73
Resultados LA-ICP-MS.....	76

# Índice de figuras

Figura 1. Mapa geológico del sector de Puchuldiza. Adaptado de Ortiz et al., 2008 y Cortés et al., 2014. ....	8
Figura 2. Campo geotérmico de Puchuldiza. Se presenta la distribución de las fuentes termales muestreadas junto con las temperaturas medidas. ....	14
Figura 3. Diagrama Piper con datos analíticos medidos. Se representan las aguas de Puchuldiza como aguas maduras de tipo clorurado sódicas. ....	16
Figura 4. Diagrama de cationes (K, Na y Mg), basado en Giggenbach, 1988. Según el gráfico, la temperatura del sistema hidrotermal fluctuaría entre 240 a 270 °C. ....	18
Figura 5. Location map of the Puchuldiza geothermal field in the Altiplano of northern Chile. Other geothermal areas are shown (purple circles), as well as active volcanoes (black triangles). Figure modified from Nicolau et al. (2014) and Aravena et al. (2016). ....	46
Figura 6. (A) Location of the 13 sinter sampling sites, distributed throughout the Puchuldiza field. (B and D) sample sites of active sinter related to bubbling pools. The pool in B has ~1 m diameter in sampling site 7 and in D the pool has ~0.3 m, with discharge channels and temperatures of 84°C and 54°C, respectively. (C and E) Paleo-sinter samples showing bands and red nodules (sampling site 9) and multiple color bands (site 13, E). (F and G): active sinter samples showing high porosity (site 6, F) and nodules of a sulfide precipitate (site 11, G). ...	47
Figura 7. Representative X-Ray diffractogram traces of sinter samples from the Puchuldiza. (A) Stack of 5 diffractograms of opal A- bearing samples: the broadband is centered at 23.5-24 °2θ, with FWHM values between 7- and 9.5°2θ. (B) Stack of 4 diffractograms of paracrystalline opal A/CT – CT-bearing samples: the broadband is centered at ~21.5°2θ, and FWHM values vary between 1.6 and 6.59°2θ. (C) Stack of 3 diffractograms of opal CT/C-C/quartz: the well-defined peaks of quartz are representative of high crystallinity degree. Main accessory minerals phases are labeled and correspond to anorthite (an), hematite (hem), realgar (rlg), halite (hl), detrital quartz (qz), cinnabar (cin), native sulphur (s), trypidite (trd) and cristobalite (crd). ....	48
Figura 8. Morphology and micro-textures of amorphous silica phases in sinters from Puchuldiza. (A) opal A micro-spheres (1-4 μm) forming a honey-comb arrangement. (B) Micro-botroidal arrangement of opal a spheres of diameter ~4-5 μm (C) Agglomeration and packs of aligned opal a micro-spheres (2-6 μm), showing a fracture surface. (D) Opal a micro-spheres aligned in tubular rows as filaments. (E) agglomeration of opal-CT micro-spheres ~6-15 μm diameter,	

showing lepispheres with roughness surfaces. (F) opal-CT lepispheres and micro-particles encrusted between silica platelets on larger micro-spheres..... 49

Figura 9. Morphology and micro-textures and microbial components of the Puchuldiza sinters. (A) microorganisms between microspheres of opal A show well-defined forms and display silica platelets on their surfaces. (B) Tubular microorganisms with micro-spheres of opal A encrusted on the surface. (C) Micro-spheres show rough surfaces due to large amounts of silica platelets. An amorphous silica band can be observed at the bottom. (D) Micro-spheres and silica platelets form conglomerates of different sizes..... 50

Figura 10. Morphology and micro-textures of the more crystalline silica phases in sinters from Puchuldiza. (A) Agglomeration of opal C micro-spheres with rough surfaces. (B) Agglomerated micro-spheres of opal-CT/C with quartz pseudo-microcrystals on the surface. (C) Typical morphology of opal C micro-spheres in sinter. (D) quartz microcrystals with prismatic faces. (E) microcrystals of quartz on the surface of a micro-sphere. (F) well-defined bipyramidal microcrystals of quartz..... 51

Figura 11. Box and whisker chart showing the concentration of metals (Au, Ag, Cu) and metalloids (As, Sb, B) in sinters from Puchuldiza measured by LA-ICP-MS and plotted against the degree of crystallinity of silica. Measured concentrations in opal A, -opal CT and opal C/quartz are shown in red, blue and green boxes, respectively. As a comparison, elemental concentrations in sinters reported in the literature are shown on the right-hand side (black lines, no corresponding crystallinity data available). References: (1): Steamboat Springs and Opal Mound, US (Lynne et al., 2007); (2): Deseado Massif, Argentina (Guido et al., 2002); (3):Champagne Pool, New Zealand (Jones et al., 2001), (4):Hoshino area, Japan (Belhadi et al., 2002); (5): El Tatio, Chile (Landrum et al., 2009); (6):Waiotapu, New Zealand (Pope et al., 2005); (7):Drummond Basin, Australia (Uysal et al., 2011)..... 52

Figura 12. Box and whisker chart showing the full width half maximum (FWHM, or “degree of crystallinity”) values measured on XRD traces of opal A. The data shown includes samples from Puchuldiza (this work) and from previously published studies, spanning a wide range of altitudes (m a.s.l.). The high FWHM is a key characteristic of sinter deposits from high-altitude setting (e.g., chilean Altiplano). References: El Tatio, Chile (Nicolau et al., 2014), Opal Mound, US (Lynne et al., 2005), Steamboat Spring, USA (Lynne et al., 2007), Te Kopia (Lynne and Campbell, 2004; Rodgers et al., 2004), Waiotapu, New Zealand (Lynne and Campbell, 2004; Rodgers et al., 2004), Orakei Korako, New Zealand (Lynne and Campbell, 2004) and Sinter Island, New Zealand (Lynne et al., 2007)..... 53

Figura 13. Representative time vs. intensity signals for Si, Au, and Ag obtained during LA-ICP-MS analyses in sample M2.6r. (a) Profile shows homogeneous distribution of Si, Au and Ag, (b) Ablation of Au and Ag-bearing inclusions hosted in the silica matrix show as spikes in the transient signal (red rectangle). Intensity is in counts per second (cps) and time in seconds (s).  
 ..... 54

Figura 14. Conceptual model of metal incorporation into silica phases during sinter formation. (a): At the nano-scale, the initial stage of sinter formation starts with over saturation of geothermal water with respect to silica as they discharge and cool at the surface, resulting in silica precipitation; (b): Amorphous opal A nanospheres agglomerate and form micro-spheres. During this stage, gold and silver are incorporated into silica spheres either as cationic species and/or metal nanoparticles or colloids (c), while arsenic and boron are incorporated into accessory minerals Fe-oxyhydroxides (D, brown). (e) As diagenesis progress, metals are enriched while metalloids are depleted from the sinter as a result of structural changes underwent by the silica host (i.e., maturation or increase in crystallinity), among other kinetically controlled processes..... 55

## Índice de tablas

Tabla 1. Mineralogical features of silica sinters from Puchuldiza. The FWHM values and the corresponding structural classification are provided, along with a general macroscopic description and location information. Accessory mineralogy is described for each sample.... 41

Tabla 2. Selected trace elements analyses of different silica phases determined by LA-ICP-MS. Concentration are in ppm unless specified otherwise..... 44

Tabla 3. Geochemistry of thermal water. The temperatures are in °C and concentrations in ppm, unless specified otherwise. .... 45