



Tissue factor activity in women with preeclampsia or SGA: a potential explanation for the excessive thrombin generation in these syndromes

Offer Erez, Roberto Romero, Edi Vaisbuch, Nandor Gabor Than, Juan Pedro Kusanovic, Shali Mazaki-Tovi, Francesca Gotsch, Pooja Mittal, Zhong Dong, Tinnakorn Chaiworapongsa, Chong Jai Kim, Chia-Ling Nhan-Chang, Sun Kwon Kim, Lami Yeo, Moshe Mazor & Sonia S. Hassan

To cite this article: Offer Erez, Roberto Romero, Edi Vaisbuch, Nandor Gabor Than, Juan Pedro Kusanovic, Shali Mazaki-Tovi, Francesca Gotsch, Pooja Mittal, Zhong Dong, Tinnakorn Chaiworapongsa, Chong Jai Kim, Chia-Ling Nhan-Chang, Sun Kwon Kim, Lami Yeo, Moshe Mazor & Sonia S. Hassan (2018) Tissue factor activity in women with preeclampsia or SGA: a potential explanation for the excessive thrombin generation in these syndromes, *The Journal of Maternal-Fetal & Neonatal Medicine*, 31:12, 1568-1577, DOI: [10.1080/14767058.2017.1320543](https://doi.org/10.1080/14767058.2017.1320543)

To link to this article: <https://doi.org/10.1080/14767058.2017.1320543>



Published online: 19 May 2017.



Submit your article to this journal [↗](#)



Article views: 115



View related articles [↗](#)



View Crossmark data [↗](#)

Tissue factor activity in women with preeclampsia or SGA: a potential explanation for the excessive thrombin generation in these syndromes

Offer Erez^{a,b}, Roberto Romero^{a,c,d,e}, Edi Vaisbuch^{a,b}, Nandor Gabor Than^{a,b,f,g,h}, Juan Pedro Kusanovic^{a,i,j}, Shali Mazaki-Tovi^{a,b}, Francesca Gotsch^{a,k}, Pooja Mittal^{a,b}, Zhong Dong^{a,b}, Tinnakorn Chaiworapongsa^{a,b}, Chong Jai Kim^{a,l}, Chia-Ling Nhan-Chang^{a,b,m}, Sun Kwon Kim^{a,b}, Lami Yeo^{a,b}, Moshe Mazorⁿ and Sonia S. Hassan^{a,b}

^aPerinatology Research Branch, NICHD/NIH/DHHS, Bethesda, MD, and Detroit, MI, USA; ^bDepartment of Obstetrics and Gynecology, Wayne State University School of Medicine, Detroit, MI, USA; ^cDepartment of Obstetrics and Gynecology, University of Michigan, Ann Arbor, MI, USA; ^dDepartment of Epidemiology and Biostatistics, Michigan State University, East Lansing, MI, USA; ^eCenter for Molecular Medicine and Genetics, Wayne State University, Detroit, MI, USA; ^fMaternity Private Department, Kutvolgyi Clinical Block, Semmelweis University, Budapest, Hungary; ^gSystems Biology of Reproduction Lendulet Research Group, Institute of Enzymology, Research Centre for Natural Sciences, Hungarian Academy of Sciences, Budapest, Hungary; ^hFirst Department of Pathology and Experimental Cancer Research, Semmelweis University, Budapest, Hungary; ⁱDepartment of Obstetrics and Gynecology, Center for Research and Innovation in Maternal-Fetal Medicine (CIMAF), S tero del R o Hospital, Santiago, Chile; ^jDivision of Obstetrics and Gynecology, Faculty of Medicine, Pontificia Universidad Cat lica de Chile, Santiago, Chile; ^kDepartment of Obstetrics and Gynecology, Azienda, Ospedaliera Universitaria Integrata, Verona, Italy; ^lDepartment of Pathology, University of Ulsan College of Medicine, Seoul, Republic of Korea; ^mDepartment of Obstetrics and Gynecology, Columbia University, New York, NY, USA; ⁿDepartment of Obstetrics and Gynecology, Ben-Gurion University, Beer-Sheva, Israel

ABSTRACT

Objective: The aim of this study was to determine whether the activity of tissue factor (TF) and tissue factor pathway inhibitor (TFPI) in the plasma of women with preeclampsia (PE) and small for gestational age (SGA) neonate differ from that of normal pregnant women and whether they are related to specific placental lesions.

Methods: This cross-sectional study included the following groups: (1) normal pregnancy ($n = 68$); (2) PE ($n = 128$); and (3) SGA ($n = 56$). Maternal plasma TF and TFPI activity was determined with chromogenic assays.

Results: (1) The median maternal plasma TF activity, but not TFPI activity, differed among the study groups ($p < .0001$ and $p = .4$, respectively); (2) patients with PE had a higher median maternal plasma TF activity than women with normal pregnancies ($p < .0001$) and mothers with SGA fetuses ($p = .002$); (3) among patients with PE, those with distal villous hypoplasia had a higher median maternal TF activity than those without these placental lesions ($p = .018$); and (4) following adjustment for confounding variables, maternal plasma TF and TFPI activity were not associated with an SGA neonate.

Conclusions: Plasma TF activity is higher in women with PE than in those with SGA or normal pregnancies. We propose that these changes may be responsible, at least in part, for the increased *in-vivo* thrombin generation observed in this obstetrical syndrome.

ARTICLE HISTORY

Received 2 February 2017
Revised 13 April 2017
Accepted 14 April 2017

KEYWORDS

Early-onset preeclampsia; extrinsic pathway of coagulation; maternal underperfusion; obstetrical syndromes; thrombin generation

Introduction

The extrinsic pathway of the coagulation cascade is modulated by two major proteins; it is activated by tissue factor (TF) [1]. Upon its exposure on the endothelium or activated monocytes, tissue factor activates factor VII (FVII), that in turn transforms factor X to its activated form; the latter leads to the generation of thrombin from pro-thrombin [1]. The major inhibitor of TF is tissue factor pathway inhibitor (TFPI) that exerts its action on TF/FVIIa and Factor X through its

three Kunitz domains [2,3]. TF has a role in systemic inflammation [4–8], placental implantation [9,10], and angiogenesis [11–16]. During normal pregnancy, TF is abundant in the uterine decidua, fetal membranes, and amniotic fluid [9–11], resulting in an efficient hemostatic mechanism activated during implantation and after delivery [17]. TF also has a soluble form – blood-borne TF – that can be detected in the maternal plasma. The plasma concentration of TF during normal pregnancy is comparable to the non-pregnant state

[17,18] and increases during labor [19,20]. There is a positive correlation between TF concentration and activity in women with a normal pregnancy [21], and TF activity is higher than the non-pregnant state [22].

Further support for the pro-coagulant state reported in preeclampsia and preterm prelabor rupture of the membranes (PROM) [23,24] is the higher median maternal plasma TF concentration observed in these patients in comparison to women with normal pregnancies [25–28], as well as the low plasma protein Z concentrations observed in women with preeclampsia [29] and those with idiopathic intrauterine bleeding and preterm labor [30]. In contrast, in spite of their increased thrombin generation, patients who delivered a small-for-gestational-age (SGA) neonate had a lower median maternal plasma TF concentration than that of women with a normal pregnancy [19]. Patients with both preeclampsia [16,17,25–29] and SGA neonates [16] have increased thrombin generation, as evident by elevated maternal plasma thrombin anti-thrombin (TAT) complexes concentrations [24,31–33]. However, patients with these obstetrical syndromes differ in their maternal TF concentrations. Therefore, we conducted a study that was aimed to test (1) whether TF and TFPI activity can explain the increased thrombin generation in these syndromes and (2) whether the TF and TFPI activity are related to specific placental lesions in each study group.

Material and methods

Study groups and inclusion criteria

This cross-sectional study included patients with singleton gestations in the following groups: (1) women with normal pregnancies ($n=68$); (2) patients with preeclampsia ($n=128$); and (3) women who delivered SGA neonates without preeclampsia ($n=56$). Patients with fetuses who had congenital and/or chromosomal anomalies were excluded.

Samples and data were retrieved from the Bank of Biological Materials and the clinical databases of the Perinatology Research Branch, Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD), Wayne State University, and the Detroit Medical Center. Many of these samples have been employed to study the biology of inflammation, hemostasis, angiogenesis regulation, and growth factor concentrations in non-pregnant women, women who had normal pregnancies, and those with pregnancy complications.

All women provided written informed consent prior to the collection of maternal blood. The Institutional

Review Boards of Wayne State University and NICHD approved the collection and utilization of samples for research purposes.

Clinical definitions

Women with normal pregnancies met the following criteria: (1) no medical, obstetrical, or surgical complications at the time of the study; (2) gestational age ranging from 20 to 41 weeks; and (3) delivery of a term infant, appropriate for gestational age, without complications. Preeclampsia was defined as the presence of hypertension (systolic blood pressure ≥ 140 mmHg and/or diastolic blood pressure ≥ 90 mmHg on at least two occasions, 4 h to 1 week apart) and proteinuria (≥ 300 milligrams in a 24 h urine collection or one dipstick measurement $\geq 2+$) [34]. Preeclampsia diagnosed < 34 weeks of gestation was defined as early onset. A small for gestational age neonate was defined as birth-weight $< 10^{\text{th}}$ percentile [35]. Placental histologic findings were classified according to a diagnostic schema proposed by Redline et al. [36].

Sample collection and immunoassays

All blood samples were collected into a vacutainer with 0.109 M trisodium citrate anticoagulant solution (BD Biosciences, San Jose, CA). The samples were centrifuged at 1300g for 10 min at 4°C and stored at -70°C until assay.

Human tissue factor immunoassay

TF concentrations were determined by sensitive and specific immunoassays obtained from American Diagnostica (Greenwich, CT), which recognize TF-apo, TF, and TF-FVII complexes. The assays were conducted according to the manufacturer's recommendations. The coefficient of variation (CV) calculated in our laboratory was 5.3%, and the sensitivity was 10 pg/ml.

Human tissue factor activity assay

TF activity was determined by a chromogenic assay obtained from American Diagnostica (Greenwich, CT). The assays were conducted according to the manufacturer's recommendations. The calculated intra-assay CV was 1.83%, while the inter-assay CV was 3.58%, and the sensitivity of this assay was 0.596 pM.

Human tissue factor pathway inhibitor immunoassay

The concentrations of TFPI in maternal plasma were determined by sensitive and specific immunoassays

obtained from American Diagnostica (Greenwich, CT). The TFPI ELISA employs, as the capture antibody, a murine anti-TFPI monoclonal antibody directed against the Kunitz-1 domain of the TFPI molecule; therefore, it detects both TFPI-1 and TFPI-2, while measuring the total TFPI plasma concentrations. The assay was conducted according to the manufacturer's recommendations. The CV calculated in our laboratory was 6.6%, and the sensitivity was ~ 10 ng/ml.

Human tissue factor pathway inhibitor activity assay

TFPI activity was determined by a chromogenic assay obtained from American Diagnostica (Greenwich, CT). The assays were conducted according to the manufacturer's recommendations. The calculated intra-assay CV was 5.51%, while the inter-assay CV was 8.74% and the sensitivity was 0.017 unit/ml.

Statistical analysis

Maternal plasma concentrations and activity of TF and TFPI were not normally distributed. Thus, the Kruskal–Wallis test with *post-hoc* analysis by the Mann–Whitney *U* test was used for the comparison of continuous variables. Comparisons of proportions were performed by the Chi-square and Fisher's exact tests. The Spearman's rho test was used to detect a correlation between the concentrations and activity of TF, TFPI, and TFPI/TF ratio to the gestational age at sample collection. The ratio of TFPI/TF concentration was previously calculated [26] and included in the logistic regression models. A *p* value $< .05$ was considered

statistically significant. Analysis was performed with SPSS, version 12 (SPSS Inc., Chicago, IL).

Results

Demographic and clinical characteristics

In comparison to women with normal pregnancies, patients with preeclampsia and those who delivered an SGA neonate had a higher median gestational age at sample collection, lower median gestational age at delivery, and lower median birthweight. Women with preeclampsia had lower gestational age at delivery than women who delivered an SGA neonate (Table 1).

Among women with a normal pregnancy (1) gestational age at sample collection positively correlated with maternal plasma TFPI activity ($r = 0.3$, $p = .01$); (2) the correlation between maternal plasma TFPI concentration and activity did not reach statistical significance ($r = 0.3$, $p = .06$); and (3) maternal plasma TF activity was negatively correlated with the TFPI/TF ratio ($r = -0.3$, $p = .006$).

Changes in the median plasma activity of tissue factor and tissue factor pathway inhibitor in the different study groups

The median maternal plasma TF activity, but not TFPI activity, differed among the study groups (Kruskal–Wallis, $p < .0001$, $p = .4$, respectively). Patients with preeclampsia had higher median maternal plasma TF activity than women with normal pregnancies (16.1 pM, range 2.9–91.2 vs. 9.9 pM, range 0.7–37.6,

Table 1. Demographic and clinical characteristics of the study population.

	Normal pregnancy (<i>n</i> = 68)	Preeclampsia (<i>n</i> = 128)	SGA (<i>n</i> = 56)
Maternal age (years)	24 (21–27)	25 (20–31)	25 (20–30)
Gravidity ^a			
1	14 (21.2)	43 (33.9)	11 (20.4)
2–5	42 (63.6)	70 (55.1)	36 (66.7)
≥ 6	10 (15.2)	14 (11.0)	7 (13.0)
Parity ^b			
1	37 (55.2)	93 (73.2)	34 (61.8)
2–5	29 (43.3)	30 (23.6)	20 (36.4)
≥ 6	1 (1.5)	4 (3.1)	1 (1.8)
Ethnic origin ^c			
African-American	50 (76.9)	106 (84.1)	46 (86.8)
Caucasian	11 (16.9)	13 (10.3)	4 (7.5)
Hispanic	2 (3.1)	5 (4.0)	1 (1.9)
Asian	2 (3.1)	1 (0.8)	1 (1.9)
Other	0	1 (0.8)	1 (1.9)
Gestational age at blood collection (weeks)	31.8 (27.7–34.8)	34.3* (30.2–37.6)	37.2* (31.7–38.4)
Gestational age at delivery (weeks)	39.6 (38.5–40.5)	34.6*‡ (31.0–37.9)	37.6* (34.4–39.0)
Neonatal birthweight (grams)	3335 (3014–3695)	1920* (1215–2702)	2110* (1490–2510)

Data are presented as median (minimum, maximum) or numbers (%). SGA: small for gestational age.

^aNormal pregnancy (*n* = 66); preeclampsia (*n* = 127); SGA (*n* = 54).

^bNormal pregnancy (*n* = 67); preeclampsia (*n* = 127); SGA (*n* = 55).

^cNormal pregnancy (*n* = 65); preeclampsia (*n* = 126); SGA (*n* = 53).

* $p < 0.05$ in comparison to normal pregnancy.

‡ $p < 0.05$ in comparison to SGA.

respectively; $p < .0001$), and patients with SGA fetuses (12.9 pM, range 2.5–74.4, $p = .002$). The latter group had higher median maternal plasma TF activity than women with normal pregnancies ($p = .0001$) (Figure 1).

Early-onset preeclampsia was associated with higher median maternal plasma TF activity than term preeclampsia (18.7 pM, range 8.4–65.9 vs. 15.3 pM, range 2.9–91.2, respectively; $p = .016$) (Figure 2).

To determine the association between maternal plasma TF and TFPI activity and preeclampsia or SGA, we constructed three multivariate logistic regression models and demonstrated that (1) maternal plasma TF activity and gestational age at sample collection, but not TFPI activity, were independently associated with the development of preeclampsia (Table 2); (2) the independent association between TF activity and the development of preeclampsia remained significant even after the introduction of the TFPI/TF ratio into the model (Table 3); and (3) by contrast, only gestational age at sample collection, but not TF or TFPI

activity, was associated with the delivery of an SGA neonate (Table 4).

Placental histological findings in patients with preeclampsia and SGA and their association with the changes in tissue factor and tissue factor pathway inhibitor concentrations and their ratio (TFPI/TF)

Placental histology was available from 88.3% (113/128) of patients in the preeclampsia group and 78.6% (44/56) of patients from the SGA group, and the specific findings are presented in Table 5. Increased syncytial knots were more frequent in the placentae of patients with preeclampsia than in those of patients with SGA neonates [54% (61/113) vs. 27.3% (12/44), respectively; $p = .004$]. Among patients with preeclampsia, those with distal villous hypoplasia had a higher median maternal TF activity than those without




Figure 1. Maternal plasma tissue factor activity in women with normal pregnancies, preeclampsia or SGA neonates.

this lesion (20.1 pM, range 8.4–91.2 vs. 16.1 pM, range 2.9–59.3, respectively; $p = .018$).

Discussion

Principal findings of the study

(1) Patients with preeclampsia or women who delivered an SGA neonate had higher median maternal plasma TF activity than those with normal pregnancies; (2) by contrast, the maternal plasma TFPI activity did not differ significantly among the study groups; (3) patients who developed early-onset preeclampsia had higher median maternal plasma TF activity than those who developed this syndrome at term; (4) maternal plasma TF activity was independently associated with the development of preeclampsia even after adjustment for gestational age at sample collection and with the maternal TFPI/TF ratio; and (5) there was no association between the delivery of an SGA neonate with maternal plasma TF and TFPI activity after adjustment for gestational age at sample collection.

Changes in maternal plasma tissue factor during normal and complicated pregnancies

During normal pregnancy, TF is abundant in the uterine decidua [37,38], fetal membranes, and amniotic

fluid [9,10,20,21], resulting in an efficient hemostatic mechanism activated during implantation and after delivery [17]. The maternal plasma concentration of TF during normal pregnancy is similar to the non-pregnant state and increase during labor [11,15].

Table 2. Multiple logistic regression analysis of the association of maternal plasma tissue factor pathway inhibitor and tissue factor concentrations — activity as well as the interactions between each analyte concentration and activity and preeclampsia.

Factor	OR (95% CI)
Gestational age at sample collection (weeks)	1.09 (1.03–1.14)
TF activity (pM)	1.07 (1.04–1.11)
TFPI activity (unit/ml)	1.6 (0.8–3.2)

OR: odds ratio; CI: confidence interval; TF: tissue factor; TFPI: tissue factor pathway inhibitor.

Table 3. Multiple logistic regression analysis of the association of maternal plasma tissue factor pathway inhibitor/tissue factor ratio and preeclampsia.

Factor	OR (95% CI)
Gestational age at sample collection (weeks)	1.13 (1.06–1.22)
TF activity (pM)	1.08 (1.02–1.13)
TFPI/TF ratio	0.97 (0.96–0.98)
TFPI activity (unit/ml)	2.8 (0.94–8.35)

TFPI/TF: Tissue factor pathway inhibitor/Tissue factor; OR: odds ratio; CI: confidence interval.




Figure 2. Maternal plasma tissue factor activity in women with early-onset or term preeclampsia.

Changes in the maternal plasma TF concentration have been reported in several pregnancy complications. Indeed, in comparison to normal pregnancy, the median maternal plasma TF concentration increases in patients with preeclampsia [39] or those with preterm PROM [25], decreases in those who deliver an SGA neonate [39], and does not change significantly in patients with preterm labor [19] or in women with a fetal demise [21]. Along with the increased maternal plasma TAT III complexes concentration reported in these obstetrical syndromes [23,25–27,29,40], the changes in the maternal plasma TF concentration were regarded as additional evidence for the presence of a procoagulant state in preeclampsia and preterm PROM.

The association between maternal plasma tissue factor concentration and procoagulant activity

It is unclear whether the maternal plasma TF concentration reflects a state of activation of the coagulation cascade, systemic maternal inflammation, or both. Indeed, the pro-coagulant activity of plasma immunoreactive TF (blood-borne TF) of non-pregnant patients is a controversial topic. While Butenas et al. [41] demonstrated that blood-borne TF has very little or no pro-coagulant activity, and that only the

administration of exogenous, active TF generates a whole blood and plasma clot after the inhibition of the contact factor (factor XIIa) [41], others [42–46] have proposed that blood-borne TF does not initiate the coagulation cascade, but rather propagates clot formation by attaching to activated platelets, further enhancing this process.

The assay used in this study measures TF activity through the generation of factor Xa (FXa) in a given sample. Thus, increased TF activity is reflected in elevated FXa formation that leads to increased thrombin generation. Therefore, the positive correlation between maternal plasma TF concentration and activity in normal pregnant women, previously reported by our group [19] suggests that during gestation circulating TF has pro-coagulant properties that may be part of the hypercoagulable state of normal pregnancy.

Preeclampsia, SGA and tissue factor activity

The finding that maternal plasma TF activity was independently associated with the development of preeclampsia, but not with the delivery of an SGA neonate, is novel.

What are the possible mechanisms leading to increased tissue factor activity among patients with preeclampsia?

The following are possible factors contributing to the higher TF plasma concentration among patients with preeclampsia: (1) the exaggerated systemic maternal inflammation characterizing patients with preeclampsia [47–55]. Indeed, patients with this syndrome have increased monocyte activation when compared to those with a normal pregnancy [49,55–64], as well as a higher monocyte metabolic activity and oxidative

Table 4. Multiple logistic regression analysis of the association of maternal plasma tissue factor pathway inhibitor and tissue factor concentrations, activity as well as the interaction between each analyte concentration and activity and SGA.

Factor	OR (95% CI)
Gestational age at sample collection (weeks)	1.12 (1.05–1.2)
TFPI activity (unit/ml)	0.93 (0.38–2.25)
TF activity (Pm)	0.99 (0.96–1.2)

OR: odds ratio; CI: confidence interval; TF: tissue factor; TFPI: tissue factor pathway inhibitor.

Table 5. A comparison of placental histologic lesions between patients with preeclampsia and patients who delivered an SGA neonate.

Placental histologic findings	Preeclampsia (n = 113)	SGA (n = 44)	p value
Findings consistent with amniotic fluid infection			
Chorioamnionitis, maternal response	7 (6.2)	4 (9)	.5
Funisitis, fetal response	2 (1.8)	4 (9)	.05
Findings consistent with maternal underperfusion			
Remote villous infarcts	21 (18.6)	5 (11.4)	.3
Recent villous infarcts	6 (5.3)	0	.2
Increased syncytial knots	61 (54)	12 (27.3)	.004
Villous agglutination	14 (12.4)	7 (15.9)	.6
Increased intervillous fibrin	26 (23.0)	7 (15.7)	.4
Distal villous hypoplasia	28 (24.8)	6 (13.6)	.19
Persistent muscularization of basal plate arteries	4 (3.5)	2 (4.5)	.6
Mural hypertrophy of decidual arterioles	8 (7.1)	4 (9.1)	.7
Acute atherosclerosis of basal plate arteries/decidual arterioles	16 (14.2)	2 (4.5)	.1
Findings consistent with fetal vascular thrombo-occlusive disease			
Villous changes	9 (6.1)	4 (9.1)	.8
Chronic villitis with obliterative fetal vasculopathy	9 (8)	5 (11.4)	.5

Data are presented as numbers (%).

burst than those delivering an SGA neonate [65]. Peripheral blood monocytes from the uterine vein of patients with preeclampsia showed a higher degree of activation in comparison to those obtained from the cubital vein [60], and this has been proposed to be the consequence of maternal monocyte activation during their passage through the placental bed of patients with preeclampsia [60]. (2) A second mechanism to consider is the production of microparticles containing TF and their release into the maternal circulation. Activated monocytes are a possible source of such microparticles that express TF on their membranes [5,41,66–72] and shed microparticles containing TF [41,46,73–76] into the plasma during systemic inflammation. The placenta may be an additional source for microparticles. Indeed, patients with preeclampsia have increased syncytiotrophoblast and endothelial macrovesicle production [77–82]. Additionally, a higher apoptosis rate was observed in cultured cytotrophoblasts of patients with preeclampsia and fetal growth restriction, and the primary placental tissues of these patients had 1.8- and 1.9-fold increases in the apoptosis rate compared to the controls [83]. Placentas of patients with preeclampsia have higher degrees of ischemia [84] and apoptosis [82] associated with microparticles generation. Placental microparticles that would carry the placental isoform of TF, which is different than recombinant TF and has a higher affinity for FVIIa [85], and syncytiotrophoblast microvesicles released from the placentas of patients with preeclampsia, exhibit increased TF activity than that of women who had normal pregnancies; a recent study suggested that macrovesicles released from the placentas of patients with preeclampsia exhibit increased TF activity, and when they were added to platelet-poor plasma, they led to increased thrombin generation [86]. (3) Patients with preeclampsia had a significant reduction in their microparticle-induced thrombin generation after treatment with anti-FVII antibodies [87]. Thus, the authors [87] concluded that a higher proportion of thrombin generation is derived from the extrinsic pathway of coagulation in patients with preeclampsia. This is in accord with our own findings, since TF is the activator of the extrinsic pathway of coagulation, and the increased maternal plasma activity of this coagulation factor among patients with preeclampsia reported herein may contribute to the observation made by VanWijk et al. [87]. Moreover, maternal plasma TF activity is increased in patients who developed early-onset preeclampsia, suggesting

an association between TF activity and the severity of this syndrome.

Tissue factor and maternal thrombin generation in women who delivered an SGA neonate

The finding that women who delivered an SGA neonate have higher median maternal plasma TF activity than women who had normal pregnancies is novel. TF activity, however, was not independently associated with the delivery of an SGA neonate. The association between the higher TAT III complexes concentration reported in these patients [24] and the activation of TF in the maternal plasma is not clear. Our group reported that although women who delivered SGA neonates have faster thrombin generation kinetics, they do not generate significantly more thrombin than women who had normal pregnancies [88]. In addition, the median maternal plasma TF concentration was lower in the SGA group than in the preeclampsia as well as the normal pregnancy groups, while TFPI concentration and activity did not differ from that of women with a normal pregnancy [39]. Collectively, these findings suggest that patients with an SGA neonate have higher TF activity compared to women who had normal pregnancies, and this may contribute to the faster thrombin generation kinetics in these patients. However, the low maternal plasma TF concentration and the normal TFPI concentration result in a higher TFPI/TF ratio that may lead to less activation of the extrinsic pathway of coagulation.

In conclusion, maternal plasma TF activity, although high in patients who developed preeclampsia and those who delivered an SGA neonate, is independently associated only with the development of preeclampsia, and is higher in patients with the early-onset form than in those who developed preeclampsia later during gestation. This finding supports the procoagulant activity of blood-borne TF in complicated pregnancies, especially among patients with preeclampsia, and further emphasizes the role of the extrinsic pathway of coagulation in this obstetrical syndrome.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This research was supported, in part, by the Perinatology Research Branch, Program for Perinatal Research and Obstetrics, Division of Intramural Research, Eunice Kennedy Shriver National Institute of Child Health and Human

Development, National Institutes of Health, Department of Health and Human Services (NICHD/NIH/DHHS); and, in part, with Federal funds from NICHD/NIH/DHHS under Contract No. HHSN275201300006C.

References

- [1] White CG II, Marder VJ, Schulman S, et al., editors. Overview of basic coagulation and fibrinolysis. In: *Thrombosis and hemostasis*. 6th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2013. p. 103–110.
- [2] Broze GJ Jr, Girard TJ, Novotny WF. Regulation of coagulation by a multivalent Kunitz-type inhibitor. *Biochemistry*. 1990;29:7539–7546.
- [3] Broze GJ Jr, Warren LA, Novotny WF, et al. The lipoprotein-associated coagulation inhibitor that inhibits the factor VII-tissue factor complex also inhibits factor Xa: insight into its possible mechanism of action. *Blood*. 1988;71:335–343.
- [4] Osterud B. Tissue factor expression by monocytes: regulation and pathophysiological roles. *Blood Coagul Fibrinolysis*. 1998;9:S9–S14.
- [5] Osterud B, Bjorklid E. Sources of tissue factor. *Semin Thromb Hemost*. 2006;32:11–23.
- [6] Schouten M, Wiersinga WJ, Levi M, et al. Inflammation, endothelium, and coagulation in sepsis. *J Leukoc Biol*. 2008;83:536–545.
- [7] Gando S, Kameue T, Nanzaki S, et al. Participation of tissue factor and thrombin in posttraumatic systemic inflammatory syndrome. *Crit Care Med*. 1997;25:1820–1826.
- [8] Green J, Doughty L, Kaplan SS, et al. The tissue factor and plasminogen activator inhibitor type-1 response in pediatric sepsis-induced multiple organ failure. *Thromb Haemost*. 2002;87:218–223.
- [9] Lockwood CJ, Krikun G, Rahman M, et al. The role of decidualization in regulating endometrial hemostasis during the menstrual cycle, gestation, and in pathological states. *Semin Thromb Hemost*. 2007;33:111–117.
- [10] Dusse LM, Carvalho MG, Cooper AJ, et al. Tissue factor and tissue factor pathway inhibitor: a potential role in pregnancy and obstetric vascular complications? *Clin Chim Acta*. 2006;372:43–46.
- [11] Ruf W, Mueller BM. Tissue factor in cancer angiogenesis and metastasis. *Curr Opin Hematol*. 1996;3:379–384.
- [12] Wojtukiewicz MZ, Sierko E, Klement P, et al. The hemostatic system and angiogenesis in malignancy. *Neoplasia*. 2001;3:371–384.
- [13] Nash GF, Walsh DC, Kakkar AK. The role of the coagulation system in tumour angiogenesis. *Lancet Oncol*. 2001;2:608–613.
- [14] Brodsky SV. Coagulation, fibrinolysis and angiogenesis: new insights from knockout mice. *Exp Nephrol*. 2002;10:299–306.
- [15] Schatz F, Krikun G, Caze R, et al. Progestin-regulated expression of tissue factor in decidual cells: implications in endometrial hemostasis, menstruation and angiogenesis. *Steroids*. 2003;68:849–860.
- [16] Belting M, Ahamed J, Ruf W. Signaling of the tissue factor coagulation pathway in angiogenesis and cancer. *Arterioscler Thromb Vascular Biol*. 2005;25:1545–1550.
- [17] Hahn L. On fibrinolysis and coagulation during parturition and menstruation. *Acta Obstet Gynecol Scand Suppl*. 1974;28:7–40.
- [18] Holmes VA, Wallace JM. Haemostasi in normal pregnancy: a balancing act? *Biochem SocTrans*. 2005;33:428–432.
- [19] Erez O, Romero R, Vaisbuch E, et al. High tissue factor activity and low tissue factor pathway inhibitor concentrations in patients with preterm labor. *J Matern Fetal Neonatal Med*. 2010;23:23–33.
- [20] Uszynski M, Zekanowska E, Uszynski W, Kuczynski J. Tissue factor (TF) and tissue factor pathway inhibitor (TFPI) in amniotic fluid and blood plasma: implications for the mechanism of amniotic fluid embolism. *Eur J Obstet Gynecol Reprod Biol*. 2001;95:163–166.
- [21] Erez O, Gotsch F, Mazaki-Tovi S, et al. Evidence of maternal platelet activation, excessive thrombin generation, and high amniotic fluid tissue factor immunoreactivity and functional activity in patients with fetal death. *J Matern Fetal Neonatal Med*. 2009;22:672–687.
- [22] Rousseau A, Favier R, Van Dreden P. Elevated circulating soluble thrombomodulin activity, tissue factor activity and circulating procoagulant phospholipids: new and useful markers for pre-eclampsia? *Eur J Obstet Gynecol Reprod Biol*. 2009;146:46–49.
- [23] Chaiworapongsa T, Espinoza J, Yoshimatsu J, et al. Activation of coagulation system in preterm labor and preterm premature rupture of membranes. *J Matern Fetal Neonatal Med*. 2002;11:368–373.
- [24] Chaiworapongsa T, Yoshimatsu J, Espinoza J, et al. Evidence of in vivo generation of thrombin in patients with small-for-gestational-age fetuses and pre-eclampsia. *J Matern Fetal Neonatal Med*. 2002;11:362–367.
- [25] Erez O, Espinoza J, Chaiworapongsa T, et al. A link between a hemostatic disorder and preterm PROM: a role for tissue factor and tissue factor pathway inhibitor. *J Matern Fetal Neonatal Med*. 2008;21:732–744.
- [26] Erez O, Romero R, Hoppensteadt D, et al. Tissue factor and its natural inhibitor in pre-eclampsia and SGA. *J Matern Fetal Neonatal Med*. 2008;21:855–869.
- [27] Bellart J, Gilabert R, Angles A, et al. Tissue factor levels and high ratio of fibrinopeptide A:D-dimer as a measure of endothelial procoagulant disorder in pre-eclampsia. *Br J Obstet Gynaecol*. 1999;106:594–597.
- [28] Schjetlein R, Abdelnoor M, Haugen G, et al. Hemostatic variables as independent predictors for fetal growth retardation in preeclampsia. *Acta Obstet Gynecol Scand*. 1999;78:191–197.
- [29] Erez O, Hoppensteadt D, Romero R, et al. Preeclampsia is associated with low concentrations of protein Z. *J Matern Fetal Neonatal Med*. 2007;20:661–667.
- [30] Kusanovic JP, Espinoza J, Romero R, et al. Plasma protein Z concentrations in pregnant women with idiopathic intrauterine bleeding and in women with spontaneous preterm labor. *J Matern Fetal Neonatal Med*. 2007;20:453–463.
- [31] Cadroy Y, Grandjean H, Pichon J, et al. Evaluation of six markers of haemostatic system in normal

- pregnancy and pregnancy complicated by hypertension or pre-eclampsia. *Br J Obstet Gynaecol.* 1993;100:416–420.
- [32] de Boer K, ten Cate JW, Sturk A, et al. Enhanced thrombin generation in normal and hypertensive pregnancy. *Am J Obstet Gynecol.* 1989;160:95–100.
- [33] Higgins JR, Walshe JJ, Darling MR, et al. Hemostasis in the uteroplacental and peripheral circulations in normotensive and pre-eclamptic pregnancies. *Am J Obstet Gynecol.* 1998;179:520–526.
- [34] ACOG Practice Bulletin. Diagnosis and management of preeclampsia and eclampsia. Number 33, January 2002. *Obstet Gynecol.* 2002;99:159–167.
- [35] Alexander GR, Himes JH, Kaufman RB, et al. A United States national reference for fetal growth. *Obstet Gynecol.* 1996;87:163–168.
- [36] Redline RW, Heller D, Keating S, Kingdom J. Placental diagnostic criteria and clinical correlation – a workshop report. *Placenta.* 2005;26:S114–S117.
- [37] Lockwood CJ, Krikun G, Schatz F. The decidua regulates hemostasis in human endometrium. *Semin Reprod Endocrinol.* 1999;17:45–51.
- [38] Lockwood CJ, Krikun G, Caze R, et al. Decidual cell-expressed tissue factor in human pregnancy and its involvement in hemostasis and preeclampsia-related angiogenesis. *Ann NY Acad Sci* 2008;1127:67–72.
- [39] Erez O, Hoppensteadt D, Than NG, et al. Tissue factor and its natural inhibitor in preeclampsia and SGA. *Reprod Sci.* 2007;14:182A.
- [40] Elovitz MA, Baron J, Phillippe M. The role of thrombin in preterm parturition. *Am J Obstet Gynecol.* 2001;185:1059–1063.
- [41] Butenas S, Bouchard BA, Brummel-Ziedins KE, et al. Tissue factor activity in whole blood. *Blood.* 2005;105:2764–2770.
- [42] Balasubramanian V, Grabowski E, Bini A, Nemerson Y. Platelets, circulating tissue factor, and fibrin colocalize in ex vivo thrombi: real-time fluorescence images of thrombus formation and propagation under defined flow conditions. *Blood.* 2002;100:2787–2792.
- [43] Balasubramanian V, Vele O, Nemerson Y. Local shear conditions and platelet aggregates regulate the incorporation and activity of circulating tissue factor in ex-vivo thrombi. *Thromb Haemost.* 2002;88:822–826.
- [44] Chou J, Mackman N, Merrill-Skoloff G, et al. Hematopoietic cell-derived microparticle tissue factor contributes to fibrin formation during thrombus propagation. *Blood.* 2004;104:3190–3197.
- [45] Kobayashi M, Wada H, Wakita Y, et al. Decreased plasma tissue factor pathway inhibitor levels in patients with thrombotic thrombocytopenic purpura. *Thromb Haemost.* 1995;73:10–14.
- [46] Osterud B. Cellular interactions in tissue factor expression by blood monocytes. *Blood Coagul Fibrinolysis.* 1995;6(Suppl 1):S20–S25.
- [47] Gervasi MT, Chaiworapongsa T, Naccasha N, et al. Phenotypic and metabolic characteristics of maternal monocytes and granulocytes in preterm labor with intact membranes. *Am J Obstet Gynecol.* 2001;185:1124–1129.
- [48] Carr DB, McDonald GB, Brateng D, et al. The relationship between hemodynamics and inflammatory activation in women at risk for preeclampsia. *Obstet Gynecol.* 2001;98:1109–1116.
- [49] Redman CW, Sargent IL. The pathogenesis of preeclampsia. *Gynecol Obstet Fertil.* 2001;29:518–522.
- [50] Chaiworapongsa T, Gervasi MT, Refuerzo J, et al. Maternal lymphocyte subpopulations (CD45RA+ and CD45RO+) in preeclampsia. *Am J Obstet Gynecol.* 2002;187:889–893.
- [51] Redman CW, Sargent IL. Pre-eclampsia, the placenta and the maternal systemic inflammatory response – a review. *Placenta.* 2003;24:S21–S27.
- [52] Sargent IL, Germain SJ, Sacks GP, et al. Trophoblast deportation and the maternal inflammatory response in pre-eclampsia. *J Reprod Immunol.* 2003;59:153–160.
- [53] Borzychowski AM, Sargent IL, Redman CW. Inflammation and pre-eclampsia. *Semin Fetal Neonatal Med.* 2006;11:309–316.
- [54] Saito S, Sakai M. Th1/Th2 balance in preeclampsia. *J Reprod Immunol.* 2003;59:161–173.
- [55] Redman CW, Sacks GP, Sargent IL. Preeclampsia: an excessive maternal inflammatory response to pregnancy. *Am J Obstet Gynecol.* 1999;180(2 Pt 1):499–506.
- [56] Bailey K, Herrod HG, Younger R, Shaver D. Functional aspects of T-lymphocyte subsets in pregnancy. *Obstet Gynecol.* 1985;66:211–215.
- [57] Haeger M, Unander M, Norder-Hansson B, et al. Complement, neutrophil, and macrophage activation in women with severe preeclampsia and the syndrome of hemolysis, elevated liver enzymes, and low platelet count. *Obstet Gynecol.* 1992;79:19–26.
- [58] Daniel Y, Kupferminc MJ, Baram A, et al. Plasma interleukin-12 is elevated in patients with preeclampsia. *Am J Reprod Immunol.* 1998;39:376–380.
- [59] Gervasi MT, Chaiworapongsa T, Pacora P, et al. Phenotypic and metabolic characteristics of monocytes and granulocytes in preeclampsia. *Am J Obstet Gynecol.* 2001;185:792–797.
- [60] Mellembakken JR, Aukrust P, Olafsen MK, et al. Activation of leukocytes during the uteroplacental passage in preeclampsia. *Hypertension.* 2002;39:155–160.
- [61] Sakai M, Tsuda H, Tanebe K, et al. Interleukin-12 secretion by peripheral blood mononuclear cells is decreased in normal pregnant subjects and increased in preeclamptic patients. *Am J Reprod Immunol.* 2002;47:91–97.
- [62] Holthe MR, Lyberg T, Staff AC, Berge LN. Leukocyte-platelet interaction in pregnancies complicated with preeclampsia. *Platelets.* 2005;16:91–97.
- [63] Schiessl B. Inflammatory response in preeclampsia. *Mol Aspects Med.* 2007;28:210–219.
- [64] Saito S, Shiozaki A, Nakashima A, et al. The role of the immune system in preeclampsia. *Mol Aspects Med.* 2007;28:192–209.
- [65] Ogge G, Romero R, Chaiworapongsa T, et al. Leukocytes of pregnant women with small-for-gestational age neonates have a different phenotypic and metabolic activity from those of women with

- preeclampsia. *J Matern Fetal Neonatal Med.* 2010;23:476–487.
- [66] Satta N, Toti F, Feugeas O, et al. Monocyte vesiculation is a possible mechanism for dissemination of membrane-associated procoagulant activities and adhesion molecules after stimulation by lipopolysaccharide. *J Immunol.* 1994;153:3245–3255.
- [67] Osterud B. The role of platelets in decrypting monocyte tissue factor. *Semin Hematol.* 2001;38(4 Suppl 12):2–5.
- [68] Sabatier F, Roux V, Anfosso F, et al. Interaction of endothelial microparticles with monocytic cells in vitro induces tissue factor-dependent procoagulant activity. *Blood.* 2002;99:3962–3970.
- [69] Shet AS, Aras O, Gupta K, et al. Sickle blood contains tissue factor-positive microparticles derived from endothelial cells and monocytes. *Blood.* 2003;102:2678–2683.
- [70] Eilertsen KE, Osterud B. The role of blood cells and their microparticles in blood coagulation. *Biochem Soc Trans.* 2005;33(Pt 2):418–422.
- [71] Baroni M, Pizzirani C, Pinotti M, et al. Stimulation of P2 (P2X7) receptors in human dendritic cells induces the release of tissue factor-bearing microparticles. *FASEB J.* 2007;21:1926–1933.
- [72] Poitevin S, Cochery-Nouvellon E, Dupont A, Nguyen P. Monocyte IL-10 produced in response to lipopolysaccharide modulates thrombin generation by inhibiting tissue factor expression and release of active tissue factor-bound microparticles. *Thromb Haemost.* 2007;97:598–607.
- [73] Bach RR, Moldow CF. Mechanism of tissue factor activation on HL-60 cells. *Blood.* 1997;89:3270–3276.
- [74] Butenas S, Mann KG. Blood coagulation. *Biochemistry (Mosc).* 2002;67:3–12.
- [75] Egorina EM, Sovershaev MA, Bjorkoy G, et al. Intracellular and surface distribution of monocyte tissue factor: application to intersubject variability. *Arterioscler Thromb Vasc Biol.* 2005;25:1493–1498.
- [76] Rivers RP, Hathaway WE, Weston WL. The endotoxin-induced coagulant activity of human monocytes. *Br J Haematol.* 1975;30:311–316.
- [77] Chen Y, Huang Y, Jiang R, Teng Y. Syncytiotrophoblast-derived microparticle shedding in early-onset and late-onset severe pre-eclampsia. *Int J Gynaecol Obstet.* 2012;119:234–238.
- [78] Guller S, Ma YY, Fu HH, et al. The placental syncytium and the pathophysiology of preeclampsia and intrauterine growth restriction: a novel assay to assess syncytial protein expression. *Ann NY Acad Sci.* 2008;1127:129–133.
- [79] Marques FK, Campos FM, Sousa LP, et al. Association of microparticles and preeclampsia. *Mol Biol Rep.* 2013;40:4553–4559.
- [80] Orozco AF, Jorgez CJ, Ramos-Perez WD, et al. Placental release of distinct DNA-associated microparticles into maternal circulation: reflective of gestation time and preeclampsia. *Placenta.* 2009;30:891–897.
- [81] Petrozella L, Mahendroo M, Timmons B, et al. Endothelial microparticles and the antiangiogenic state in preeclampsia and the postpartum period. *Am J Obstet Gynecol.* 2012;207:140–146.
- [82] Redman CW, Sargent IL. Microparticles and immunomodulation in pregnancy and pre-eclampsia. *J Reprod Immunol.* 2007;76:61–67.
- [83] Langbein M, Strick R, Strissel PL, et al. Impaired cytotrophoblast cell-cell fusion is associated with reduced Syncytin and increased apoptosis in patients with placental dysfunction. *Mol Reprod Dev.* 2008;75:175–183.
- [84] Reddy A, Zhong XY, Rusterholz C, et al. The effect of labour and placental separation on the shedding of syncytiotrophoblast microparticles, cell-free DNA and mRNA in normal pregnancy and pre-eclampsia. *Placenta.* 2008;29:942–949.
- [85] Butenas S, Krudysz-Amblo J, Mann KG. Posttranslational modifications and activity of natural and recombinant tissue factor. *Thromb Res.* 2010;125(Suppl 1):S26–S28.
- [86] Gardiner C, Tannetta DS, Simms CA, et al. Syncytiotrophoblast microvesicles released from pre-eclampsia placentae exhibit increased tissue factor activity. *PLoS One.* 2011;6:e26313.
- [87] VanWijk MJ, Boer K, Berckmans RJ, et al. Enhanced coagulation activation in preeclampsia: the role of APC resistance, microparticles and other plasma constituents. *Thromb Haemost.* 2002;88:415–420.
- [88] Erez O, Romero R, Vaisbuch E, Kusanovic JP, Mazaki-Tovi S, Gotsch F, et al. The pattern and magnitude of in-vivo thrombin generation is different in women with preeclampsia and in those with SGA fetuses without preeclampsia. *Reprod Sci.* 2009;16:170A.