

RESEARCH

Open Access



# Pectinase secreted by psychrotolerant fungi: identification, molecular characterization and heterologous expression of a cold-active polygalacturonase from *Tetracladium* sp.

Mario Carrasco<sup>1</sup>, Juan Manuel Rozas<sup>1</sup>, Jennifer Alcaíno<sup>2</sup>, Víctor Cifuentes<sup>2</sup> and Marcelo Baeza<sup>2\*</sup> 

## Abstract

**Background:** Pectinolytic enzymes, which are used in several industries, especially in the clarification process during wine and fruit juice production, represent approximately 10% of the global enzyme market. To prevent the proliferation of undesired microorganisms, to retain labile and volatile flavor compounds, and to save energy, the current trend is to perform this process at low temperatures. However, the commercially available pectinases are highly active at temperatures approximately 50 °C and poorly active at temperatures below 35 °C, which is the reason why there is a constant search for cold-active pectinases. In preliminary studies, pectinolytic activity was detected in cold-adapted yeasts and yeast-like microorganisms isolated from Antarctica. The aim of the present work was to characterize pectinases secreted by these microorganisms and to express the best candidate in *Pichia pastoris*.

**Results:** Degradation of pectin by extracellular protein extracellular extracts obtained from 12 yeast cultures were assayed in plates at 4 °C to 37 °C and pH from 5.4 to 7.0, obtaining positive results in samples obtained from *Dioszegia* sp., *Phenoliferia glacialis* and *Tetracladium* sp. An enzyme was purified from *Tetracladium* sp., analyzed by peptide mass fingerprinting and compared to genome and transcriptome data from the same microorganism. Thus, the encoding gene was identified corresponding to a polygalacturonase-encoding gene. The enzyme was expressed in *Pichia pastoris*, and the recombinant polygalacturonase displayed higher activity at 15 °C than a mesophilic counterpart.

**Conclusions:** Extracellular pectinase activity was found in three yeast and yeast-like microorganisms from which the highest activity was displayed by *Tetracladium* sp., and the enzyme was identified as a polygalacturonase. The recombinant polygalacturonase produced in *P. pastoris* showed high activity at 15 °C, representing an attractive candidate to be applied in clarification processes in the production of fermented beverages and fruit juices.

## Background

Currently, there is a high demand for enzymes used in several industrial applications for food, detergent, paper, textile and synthesis of organic compounds because they are highly efficient and environmentally friendly [1–3]. Additionally, these enzymes constitute a well-established global market projected to reach US\$6.3 billion in 2021

[1, 4]. The current trend is to use cold-adapted or cold-active enzymes to decrease the temperature of the industrial processes, allowing energy savings and diminishing their carbon footprint and to manufacture products with better performance at ambient or lower temperatures [5–7]. Approximately 10% of the enzyme market is represented by pectinolytic enzymes [8, 9], which are used in the wine, food, paper and textile industries [9, 10]. The reduction of cloudiness and bitterness of fruit juices and grape must in the juice and wine industries is performed at low temperatures ( $\leq 15$  °C) to prevent the proliferation of undesired microorganisms, to retain the labile

\*Correspondence: mbaeza@uchile.cl

<sup>2</sup> Laboratorio de Genética, Departamento de Ciencias Ecológicas, Facultad de Ciencias, Universidad de Chile, Las Palmeras 3425, Casilla 653, Santiago, Chile

Full list of author information is available at the end of the article



and volatile flavor compounds, and to save energy [7, 9]. These requirements have led to the search for pectinases with high performance at lower temperatures but also at low pH, as the pH of fruit juices and grape must be in the range from 2.5 to 3.5 [11]. Currently, the industrially available pectinases are obtained from mesophilic filamentous fungi, mainly from *Aspergillus* species; however, they work poorly at temperatures  $\leq 35$  °C [12]. Thus, cold-active enzymes have higher enzymatic activities at lower temperatures than their mesophilic counterparts [13].

Microorganisms that thrive in cold environments have evolved several adaptations to live under this condition, including the synthesis of cold-active enzymes [14–17]. In particular, cold-adapted fungi secrete cold-active enzymes that hydrolyze the complex compounds available in the environment to use as nutrients [18–21]. The production of pectinases has been reported in cold-adapted bacteria and fungi [11, 22–36]; however, in most of these studies no purification or biochemical characterization of the enzyme were performed. Pectinolytic enzymes, or pectinases, are classified according to their mode of action and to their substrate: polygalacturonases, which are subclassified as endo-polygalacturonases (E.C. 3.2.1.15) and exo-polygalacturonases (E.C. 3.2.1.67); lyases, which are classified into pectate lyases (E.C. 4.2.2.9 and EC. 4.2.2.2) or pectin lyases (E.C. 4.2.2.10); and pectin methylesterases (E.C. 3.1.1.11). It is recommended the use of a combination of different kind of pectinases that degrade different parts of the polymer, to achieve a maximal degradation of pectin in various raw materials [37]. Bacteria produce alkaline pectinases, most frequently polygalacturonases and pectate lyases [38]. In fungi, the production of acidic pectinases has been described, mainly exo- and endo- polygalacturonases [38]. The most frequently used pectinases in industry are the polygalacturonases, which belong to glycosyl hydrolase family 28 (GH28). Although there have been attempts to isolate cold-active pectinases from different sources, to the best of our knowledge, there are no commercially available cold-active pectinases.

In previous work, pectinolytic activity was detected in 12 yeasts and yeast-like microorganisms isolated from soils of King George Island in the sub-Antarctic region [22]. In this work, these microorganisms were further studied to characterize the pectinases secreted by them. Secretion of pectinase was confirmed in three of them, and among these, *Tetracladium* sp. showed the highest pectinase activity. The enzyme from *Tetracladium* sp. was purified and analyzed by peptide mass fingerprinting. The peptide sequences were compared to genome and transcriptome data, and the gene, which encodes a polygalacturonase, was identified. The gene

was expressed in *Pichia pastoris*, and the recombinant polygalacturonase displayed higher activity at 15 °C than a mesophilic counterpart.

## Methods

### Strains, plasmid and growth conditions

Yeast strains used in this work and their relevant characteristics are listed in Table 1, which were isolated, identified and characterized by our group from soil samples of King George Island at the sub-Antarctic [22]. The yeasts were routinely cultivated in yeast-malt medium (YM): 3 g L<sup>-1</sup> yeast extract, 3 g L<sup>-1</sup> malt extract and 5 g L<sup>-1</sup> peptone, supplemented with 10 g L<sup>-1</sup> of glucose or 10 g L<sup>-1</sup> of pectin (pectin of citrus peel, Sigma-Aldrich Corporation, MO, USA). *E. coli* strains were cultivated in lysogeny broth medium (LB): 10 g L<sup>-1</sup> tryptone, 5 g L<sup>-1</sup> yeast extract and 5 g L<sup>-1</sup> NaCl, supplemented with 1 g L<sup>-1</sup> glucose; ampicillin was added at 100 mg ml<sup>-1</sup> when necessary. Synthetic defined medium (SD: 20 g L<sup>-1</sup> glucose and 6.7 g L<sup>-1</sup> yeast nitrogen base) was used for selection of recombinant *P. pastoris* strains. Buffered complex medium (BCM: 10 g L<sup>-1</sup> yeast extract, 20 g L<sup>-1</sup> peptone, 13.4 g L<sup>-1</sup> yeast nitrogen base, 0.004 g L<sup>-1</sup> biotin, 10 g L<sup>-1</sup> glycerol and 100 mM potassium phosphate buffer pH 6.0), supplemented with 10 g L<sup>-1</sup> glycerol or 0.5% methanol when necessary, was used for gene expression in *P. pastoris*. For semisolid-media, 15 g L<sup>-1</sup> of agar was used.

### Molecular and biochemical methods

Standard molecular and biochemical procedures, such as plasmid DNA purification, digestion with restriction enzymes, cloning procedures, PCR assays, SDS-PAGE, protein quantification and electrotransformation, were performed according to standard protocols [39]. Protein quantification was made using the BCA Kit Assay (Thermo Scientific, IL, USA). Plasmid and genomic DNAs were purified using the GeneJet Plasmid Miniprep Kit (Thermo Scientific, Waltham, MA, USA) and the Wizard Genomic DNA Purification kit (Promega, WI, USA) according to manufacturer's instructions.

### Extraction and purification of extracellular proteins

Yeast cultures (100–300 mL) at the initial stationary phase of growth were centrifuged at 7000×g for 10 min at 4 °C and filtered through sterile 0.45-μm pore size polyvinylidene fluoride membrane filters (Millipore, Billerica, MA, USA). Ammonium sulfate was added to the cell-free supernatants to reach 80% of saturation to obtain total proteins or in the case of protein fractioning, to reach increasing saturation from 20 to 80%. Samples were incubated at 4 °C for 2 h and centrifuged at 10,000×g for 15 min at 4 °C. The protein pellets were suspended in 2–4 mL of 20 mM potassium phosphate and

**Table 1 Microorganism, plasmids and oligonucleotides used in this work**

Microorganism/plasmid/oligonucleotide	Description	References
Yeast/yeast-like		
<i>Tetracladium</i> sp.	30 °C <sup>a</sup>	[22]
<i>Dioszegia</i> sp.	15 °C <sup>a</sup>	[22]
<i>Rhodotorula glacialis</i> ( <i>Phenoliferia glacialis</i> ) (T8Rg)	22 °C <sup>a</sup>	[22]
<i>Leucosporium</i> sp. T17Cd1	15 °C <sup>a</sup>	[22]
<i>Leucosporidiella fragaria</i> ( <i>Leucosporidium fragarium</i> )	22 °C <sup>a</sup>	[22]
<i>Metschnikowia</i> sp.	10 °C <sup>a</sup>	[22]
<i>Mrakia psychrophila</i>	10 °C <sup>a</sup>	[22]
<i>Mrakia robertii</i>	15 °C <sup>a</sup>	[22]
<i>Rhodotorula glacialis</i> ( <i>Phenoliferia glacialis</i> ) (T11Rs)	22 °C <sup>a</sup>	[22]
<i>Rhodotorula</i> ( <i>Cystobasidium</i> ) <i>laryngis</i>	30 °C <sup>a</sup>	[22]
<i>Sporidiobolus salmonicolor</i>	22 °C <sup>a</sup>	[22]
<i>Wickerhamomyces anomalus</i>	30 °C <sup>a</sup>	[22]
PichiaPink strain 4	<i>P. pastoris</i> strain: <i>ade2, prb1, pep4</i>	Invitrogen
PichiaPink strain 2	<i>P. pastoris</i> strain: <i>ade2, pep4</i>	Invitrogen
Bacteria		
<i>Escherichia coli</i> Top10	F-mcrA Δ(mrr-hsdRMS-mcrBC) Φ80lacZΔM15 ΔlacX74 recA1 araD139 Δ(araI)7697 galU galK rpsL (StrR) endA1 nupG	Thermo Fisher
Plasmids		
pPinkα-HC	<i>P. pastoris</i> integrating vector for high-copy expression of a secreted protein	Invitrogen
Oligonucleotides		
Pectfw	5'-GCACCTACAGTCTCATCA TTG-3'	This work
Pectrev	5'-GCAGGAAGCAGGGGA TGGGAA-3'	This work

<sup>a</sup> Optimal temperature for growth. The current taxonomic classification is given in parentheses

150 mM NaCl pH 7.0, and samples were desalted using a HiTrap Desalting column (GE, Schenectady, Nueva York, USA) in AKTA Prime purification equipment (GE). For ion exchange protein purification, the protein sample was loaded onto a DEAE-Sephadex column equilibrated with 50 mM Tris-HCl pH 7.0, proteins were eluted using a NaCl gradient from 0 to 200 mM as the mobile phase at 0.5 mL min<sup>-1</sup>, and 1 mL fractions were collected. For

gel filtration purification, 0.2 mL of concentrated protein extracellular extracts were loaded onto a Superdex 200 Increase 10/300 GL column (GE Healthcare, IL, USA) equilibrated with 20 mM sodium phosphate buffer, using the same buffer as the mobile phase at flow rate of 0.5 mL min<sup>-1</sup>. In both cases, proteins of the fractions were monitored at an absorbance of 280 nm. The relative molecular weight (rMW) of proteins were calculated from SDS-PAGE by comparison its relative mobilities to that of the proteins standard.

#### Determination of pectinase activity

For semiquantitative determinations, 100 μL of the protein extracellular extracts was deposited into wells cut into agar plates containing 1% w/v pectin with pH of 5.4, 6.2 or 7.0, adjusted with phosphate-citrate buffer. The plates were incubated at 4 °C, 10 °C, 15 °C, 22 °C, 30 °C and 37 °C for 1 to 5 days, and the appearance of a clear halo around the well indicated pectinase activity. For quantitative determinations, the release of reducing sugars from pectin was quantified using the 3-amino-5-nitrosalicylic acid (DNS) method [40]. Briefly, 50 μL of the protein sample was mixed with 50 μL of 10 mg mL<sup>-1</sup> pectin, incubated at 30 °C for 1 h and then 100 μL of DNS solution was added. After incubation at 100 °C for 10 min, the samples were incubated on ice for 5 min, and the absorbance at 540 nm was measured. For comparative purposes, commercial polygalacturonase was used in activity assays.

#### Peptide mass fingerprinting

Protein extracellular extracts were separated by SDS-PAGE and stained with Coomassie Blue G-250. The protein band of interest was cut from the gel and analyzed using the protein analysis service of Alphasys (Palo Alto, CA, USA). Briefly, the protein sample was reduced and alkylated with carbamidomethylation and subsequently digested with trypsin. The resulting peptides were spotted onto an anchorchip target for analysis on a Bruker Autoflex Speed MALDI-TOF/TOF instrument. The obtained data were analyzed by Mascot, and results having a score greater than 54 ( $P < 0.05$ ) were considered statistically significant.

#### Next-generation Sequencing (NGS)

Cultures of *Tetracladium* sp. were centrifuged at 7000×g at 4 °C for 10 min. The cell pellets were used for DNA and RNA purification with the Wizard Genomic DNA Purification Kit (Promega, WI, USA) and RiboMinus Yeast Kit (Thermo Fisher, MA, USA), respectively. The quality and quantity of the samples were determined by absorbance at 260 and 280 nm, and those having a 260/280 ratio of 1.7 to 1.9 and a 260/230 ratio > 2 were used for whole

genome sequencing by NGS at Macrogen Inc. (Seoul, South Korea) using the Hiseq 2000 platform.

#### Assemblies, ORFs and gene prediction, annotation and expression level analysis

Assemblies were done with ALLPATHS-LG [41]. RNA-seq reads were mapped to the assembled genome with TopHat software [42] and STAR aligner [43]. To capture all junctions, the RNA-seq reads were assembled according to Grabherr et al. [44]. The gene model prediction was made with Augustus [45, 46] and Pasa [47]. Functional annotation was performed using the standard BLAST and InterPro databases. CAZY annotation was made using dbCAN [48] and HMMER3 [49]. ClustalW analysis and gene expression analysis were conducted using Geneious version 10.0.9 [50] and the included plugins.

#### Alignment, modeling and bioinformatics analysis

Amino acid sequence alignments were conducted using MEGA7 software [51]. The polygalacturonase sequences chosen for the amino acid sequence comparison had a minimum of 50% similarity and 50% coverage, and corresponded to: *Venturia nashicola* (BAG72101), *Colletotrichum fiorinae* (EXF76863), *Colletotrichum lupini* (ABL01533), *Colletotrichum simmondsii* (KXH46697), *Achaetomium* sp. (AGR51994), *Colletotrichum higginsianum* (XP\_018155590), *Venturia pyrina* (BAG72133), *Pestalotiopsis fici* (XP0\_07836731), *Talaromyces cellulosus* (GAM33350), *Cadophora* sp. (PVH80831), *Pseudomassariellva xata* (ORY56346).

*Phialocephalus balpina* (CZR53206), *Penicillium freii* (KUM62405), *Lepidopterella palustris* (OCK84102), *Thielavia arenaria* (AIZ95162), *Fusarium avenaceum* (KIL90067), *Neonectria ditissima* (KPM736\_08), *Fusarium venenatum* (CEI70336), *Penicillium griseoroseum* (MF06810), *Colletotrichum gloeosporioides* (ELA24368), *Verticillium alfalfa* (XP0\_03002875), *Verticillium dahlia* (XP0\_09653008), *Diplodia corticola* (XP\_020130427), *Pezoloma ericae* (PMD23778), *Penicillium camemberti* (CRL23357), *Fusarium langsethiae* (KPA37956), *Nectria haematococca* (XP\_003040641), *Meliniomyces bicolor* (XP\_024742782), *Ustilaginomycotina* sp. (PWN47943), *Talaromyces arneffeii* (KFX46954), *Bipolaris zeicola* (XP\_007711986), *Penicillium subrubescens* (OKO96599), *Penicillium brasilianum* (OOQ88122). The polygalacturonase model was constructed using the Swiss-model platform [52], using PDB: 2iq7.1 as the template, which has 91% coverage and 63% identity with the polygalacturonase from *Tetracladium* sp. For validation of the model structure the programs Verify 3D and ERRAT were used through the AVES v5.0 server (<http://servicesn.mbi.ucla.edu/SAVES/>), and the values were 96.43% and 89.13%, respectively.

#### Heterologous expression of pectinase

The coding sequence of *TPG1* identified in this work (see “Results and discussion” section) was analyzed bioinformatically and modified to generate a sequence of 1066 nt that lacks the first 57 nucleotides, which encode the signal peptide. *Mly* I and *Kpn* I restriction sites were added at the 5' and 3' end, respectively, and the codon usage was optimized to the one from *P. pastoris*. The modified gene was synthesized by Genescript company (New York, USA), and the cloning and expression of pectinase was performed using the PichiaPink™ Expression System (Invitrogen, CA, USA) according to manufacturers' instructions. Briefly, synthetic CDS was ligated to a pPink $\alpha$ -HC vector and transformed into *E. coli* Top10. The obtained transformants were selected on LB-ampicillin plates, and the presence of recombinant plasmid was confirmed by colony PCR using the primer pair Pectfw/Pectrev (Table 1). Plasmid DNA was purified from selected clones, digested with *Afl*III and transformed into *P. pastoris* PichiaPink strain 2. Several transformants developed on SD plates were selected, and genomic DNA was extracted and checked by PCR using the primer pair Pectfw/Pectrev. Amplicon-positive clones were grown overnight in BCM supplemented with glycerol at 30 °C and then centrifuged at 1500 g for 5 min. Then, the yeast pellets were suspended in 10 ml of BMMY medium and incubated overnight at 30 °C, and methanol was added to reach 4% v/v final concentration. Ten 100  $\mu$ L culture aliquots were collected at different times from 0 to 24 h and centrifuged at 1.500 g for 5 min, and cellular pellets were suspended in 50 mM sodium phosphate pH 7.4, 1 mM PMSF, 1 mM EDTA and 5% glycerol, and vortexed for 3 min. Samples were directly analyzed by SDS-PAGE.

## Results and discussion



### Screening, selection and identification of secreted pectinases

The concentration of protein extracellular extracts obtained from twelve yeasts and yeast-like species cultured in liquid YM medium supplemented with glucose ranged from 5 to 12  $\mu$ g mL<sup>-1</sup> (Fig. 1). The obtained protein extracellular extracts were used to test pectin degradation in plates at 4 °C to 37 °C and pH from 5.4 to 7.0, and positive results were observed in three of them (Fig. 1). Pectin degradation was detected from 4 °C to 37 °C and pH from 5.4 and 6.2 in samples from *Dioszegia* sp. and *Phenoliferia glacialis* D7 and from 4 °C to 37 °C and pH from 5.4 and 7.0 in samples from *Tetracladium* sp., which are in the range of described pectinases from other microorganisms [10, 53]. Pectinolytic activity has been previously described in *Rhodotorula mucilaginosa*, *Cystofilobasidium capitatum* [32], *Candida sake*, *Debaromyces vanrijae* (now *Schwanniomyces vanrijae*),

*Saccharomyces cerevisiae*, *Candida* sp., *Debaryomyces* sp., *Kluyveromyces marxianus*, *Pichia* sp., *Saccharomyces* sp., *Zygosaccharomyces* sp., [54], *Cryptococcus cylindricus* (now *Piskurozyma cylindrica*), *Mrakia frigida*, *Cystofilobasidium capitatum*, *Cystofilobasidium macerans* and *Rhodotorula mucilaginosa* [55] isolates. In this work, secreted pectinolytic activity is described for the first time in *Dioszegia* sp., *Phenoliferia glacialis* and *Tetracladium* sp. isolates. The highest pectin degradation was observed for protein extracellular extracts from *Tetracladium* sp. at 37 °C and pH 6.2. Therefore, *Tetracladium* sp. was selected for further enzyme purification, identification and heterologous expression. To evaluate if pectin induces the production of pectinase in *Tetracladium* sp., the yeast was grown in YM medium supplemented with glucose or pectin. As shown in Fig. 2, the growth curves were similar in both media, reaching growth rates of 1.2 h<sup>-1</sup> and 1.1 h<sup>-1</sup> and maximum biomass of 13 g L<sup>-1</sup> and 12 g L<sup>-1</sup> in medium supplemented with glucose and pectin, respectively. However, extracellular pectinase activity at the initial stationary phase of growth was almost seven-fold higher in the medium supplemented with pectin than with glucose, indicative of an induction of the production of pectinolytic enzymes in presence of pectin. Similarly, an induction effect of pectin in the production of endo-PG in *Aspergillus niger* T0005007-2 was observed [56].

To purify the pectinase enzyme, *Tetracladium* sp. was cultivated in YM medium supplemented with pectin,

and culture supernatant proteins were fractionated using increasing saturation of ammonium sulfate (20 to 80%). Pectinase activity was detected in fractions corresponding to 40, 60 and 80% ammonium sulfate having 7, 33 and




60% total activity, respectively. The protein sample corresponding to 80% ammonium sulfate precipitation was separated by ionic exchange chromatography (Fig. 3), and two main peaks (fractions 14 and 42) were obtained; pectinase activity was detected in fractions surrounding fraction 14. According to the SDS-PAGE analysis of these fractions, the pectinase activity was associated with a protein band of rMW 38,000 (arrow in Fig. 3), which was purified, concentrated and analyzed by peptide mass fingerprinting (PMFP). The MW of described microbial polygalacturonases is widely variable ranging from 6500 to 320,000 according to the BRENDA enzyme database [57]. From the analysis of the results and comparison using the Mascot database search engine, it was found that peptides ARAACTFSGATG, IKNSDNGVRIKTIEG, DIVYKDITLVNIA and KVTMNNVAGDTKGGHNTDAF matched with 91, 87, 77 and 79% identity with polygalacturonases described from *Fusarium avenaceum* (KIL93978.1), *Sclerotinia sclerotiorum* (CAA74019.1), *Ascochyta rabiei* (KZM23605.1) and *A. niger* CBS 513.88 (XP\_001397067.1), respectively.

#### Identification of the putative pectinase-encoding gene

The putative CDS were predicted in the genomic scaffolds of *Tetracladium* sp. and its transcriptomes, when grown in YM medium supplemented with glucose or pectin were determined. The ORFs predicted in the transcriptomes were mapped to genomic scaffolds, and those having 100% identity and a correct exon–intron

structure were considered for comparison to the NCBI database and annotation. Thus, 7 putative polygalacturonase genes (lengths from 1325 to 4827 bp) and 14 putative pectate lyase genes (lengths from 993 to 2712 bp) were found. The expression at the transcript level of these putative genes was determined in both culture conditions (Table 2). The majority of the putative pectate lyase-encoding genes did not show significant differences in the expression level between both culture conditions, except for scaffold\_2.g849.t1 and scaffold\_3.g812.t1, which had a 28- and 112-fold expression increase respectively, in cultures supplemented with pectin in comparison to glucose. In contrast, all the putative polygalacturonase-encoding genes increased their expression in medium supplemented with pectin compared to glucose, with scaffold\_10.g413.t1 (12,228-fold) and scaffold\_6.g590.t1 (12,360-fold) having the highest increases. Similar to these results, a higher transcript level of gene *pgg1* that encodes a pectinase in *Penicillium griseoroseum* was observed in the presence of pectin with in comparison to without yeast extract [58]. As mentioned above concerning the analysis by PMFP of the purified pectinolytic enzyme, it was found four peptides that matched with polygalacturonases from other fungi. These peptides were compared to the translated sequences of all of the putative CDS for polygalacturonases and pectate lyases identified in *Tetracladium* sp., found that all peptides mapped only with translated sequence from a CDS named scaffold\_10.g413.t1. Is important to mention that




**Table 2 Expression, at the transcript level, of putative pectate lyase- and polygalacturonase-encoding genes from *Tetracladium* sp**


Putative	CDS	Expression (FPKM)		P/G <sup>a</sup>
		YM-G	YM-P	
PL	scaffold_4.g450.t1	3.6	0.8	-4
PL	scaffold_5.g260.t1	22.9	6.5	-4
PL	scaffold_4.g243.t1	2.2	0.8	-3
PL	scaffold_2.g1041.t1	2.4	0.9	-3
PL	scaffold_0.g2020.t1	5.6	3.3	-2
PL	scaffold_1.g759.t1	6.5	4.2	-2
PL	scaffold_1.g52.t1	29.9	22.0	-1
PL	scaffold_0.g2289.t1	1.1	0.8	-1
PL	scaffold_11.g858.t1	26.0	45.2	2
PL	scaffold_7.g463.t1	7.0	15.0	2
PL	scaffold_3.g58.t1	18.0	60.9	3
PL	scaffold_0.g2293.t1	5.5	19.7	4
PG	scaffold_3.g273.t1	1.6	11.8	7
PG	scaffold_3.g424.t1	19.1	243.4	13
PL	scaffold_2.g849.t1	2.6	74.9	28
PG	scaffold_3.g812.t1	0.6	66.3	112
PL	scaffold_3.g812.t1	0.6	66.3	112
PG	scaffold_0.g606.t1	3.9	1012.9	258
PG	scaffold_3.g1104.t1	1.0	291.1	288
PG	scaffold_10.g413.t1	0.1	1762.5	12,228
PG	scaffold_6.g590.t1	0.0	566.6	12,360

Expression in medium supplemented with glucose (YM-G) or pectin (YM-P). PL pectate lyase, PG polygalacturonase. <sup>a</sup>Expression fold changes between cultures in YM-P (P) and YM-G (G) media

along with our purification methodologies the focus was in finding the enzyme having the highest activity; therefore the other CDSs may encode for putative pectinolytic enzymes with a minor activity that were not considered in this work. Therefore, the enzyme purified from *Tetracladium* sp. is a polygalacturonase and the encoding gene was named *TPG1*, which is composed of 5 exons and encodes a protein of 368 residues (Additional file 1: Fig. S1). Figure 4 shows the alignment of the translated sequence from *TPG1* and pectinases described in other fungi. Globally, there are several conserved residues (Fig. 4a), including those in the active and substrate binding sites (Fig. 4b) [59]. A three-dimensional model of the *Tetracladium* sp. TPG1 was constructed (Fig. 5) using the crystal structure of the endopolygalacturonase from the phytopathogenic fungus *Colletotrichum lupini* (CluPG1) as the template [60]. The overall predicted structure of TPG1 corresponded to a right-handed beta-helical structure that forms a large cleft (Fig. 5b), which is a common feature in other pectinases, such as endo-polygalacturonases from *Fusarium moniliforme* [61] and from *A. niger* [62, 63], and polygalacturonases from *Ewinia carotovora* [64], *Aspergillus aculeatus* [65], and *C. lupini* [60]. The putative catalytic residues in TPG1 corresponded to QDD and GHG and the substrate binding residues to RIK and NTD (Fig. 5). Furthermore, there were conserved cysteines (Fig. 5c), which are important for disulfide bridge formation, especially in fungal polygalacturonases [59].



**Fig. 4** Alignment of fungal polygalacturonases primary structures. Global (a) and detailed (b) comparison of fungal polygalacturonases. The conserved regions corresponding to the active site and to the substrate binding site are enclosed in continuous and discontinuous squares, respectively. The conserved residues in each region are indicated by arrows



**Fig. 5** 3D structure model of TPG1. In **a** and **c**, the N-terminus of the protein is on the right and in **b**, it is in front. Conserved motifs are shown in color and/or in balls-and-sticks: catalytic residues QDD are in green and GHG in blue; substrate binding residues RIK are in orange and NTD in yellow. Conserved cysteine residues are shown in red balls-and-sticks in **c**. AT, amino terminus; CT, carboxyl terminus

#### Heterologous expression and characterization of TPG1


The synthetic sequence of *TPG1* was inserted in the pPink $\alpha$ -HC vector and transformed in *P. pastoris* strain PichiaPink. Clones were analyzed by colony PCR using the primers pair Pectfw and Pectrev, the plasmid DNA was purified from one of the clones showing amplicons of the expected length and PCR-amplified with the same primers pair, obtaining an amplicon with a size expected for synthetic *TPG1* (Fig. 6a). The selected clone was cultured in BMGY medium at 30 °C for 24 h, then the production of pectinase was induced by changing the medium to BMMY and incubation at the same temperature for 6 to 72 h. Extracellular proteins were obtained from cultures and analyzed by SDS-PAGE observing an intense protein band of approximately 38,000 rMW (Fig. 6b). The recombinant TPG1 and a commercial polygalacturonase were used in pectinase

activity assays, using purified pectin and wine must as substrates; the hydrolysis of pectin was evaluated as the liberation of glucuronic acid at 15 °C and pH 3.0 in 1 h. The amount of glucuronic acid released by the recombinant TPG1 was twofold higher than the amount released by the commercial polygalacturonase using either substrate (Fig. 6c).

#### Conclusion

The polygalacturonase TPG1 was identified in *Tetradcladium* sp., and its expression, determined at the transcriptional and enzyme activity levels, is induced by pectin. The TPG1-encoding gene was successfully expressed in *P. pastoris*, and a recombinant polygalacturonase highly active at 15 °C was obtained.





**Fig. 6** Characterization of the recombinant TPG1. **a** Analysis by PCR of a recombinant clone (lane 2) in which an amplicon with a size expected for synthetic *TPG1* is observed (arrow); Lane 1, DNA ladder. **b** SDS-PAGE. The arrow indicates the protein band corresponding to the recombinant TPG1; M, molecular marker. **c** Pectinase activity of TPG1 and a commercial polygalacturonase, evaluated at pH 3.0 and 15 °C using pectin or must as substrate. Data shown are the average values of three independent enzymatic assays

Furthermore, the feasibility to cultivate *P. pastoris* in high-cell density fermentations facilitates the efficient production of pectinase at amount and purity required to be applied in industrial processes such as clarification processes of fermented beverage and fruit juice production. Therefore, the recombinant pectinase described in this work is economically attractive by both its efficient production and its activity at low temperatures allowing energy-saving in the processes.

## Additional file

[Additional file 1: Fig. S1.](#) TPG1 sequence and gene structure.

### Authors' contributions

MC carried out the experiments. MC, JMR and MB contributed to the design of the experiment and discussion of the results. MC, JA, VC and MB drafted the manuscript. All authors read and approved the final manuscript.

### Author details

<sup>1</sup>Innocold S.A, Santiago, Chile. <sup>2</sup>Laboratorio de Genética, Departamento de Ciencias Ecológicas, Facultad de Ciencias, Universidad de Chile, Las Palmeras 3425, Casilla 653, Santiago, Chile.

### Acknowledgements

Not applicable.

### Competing interests

The authors declare a conflict of interest. Innovacold S.A. applied for a patent based on the results presented in this article.

### Availability of data and materials

The datasets supporting this work are included in the manuscript and additional file.

### Consent for publication

Not applicable.

### Ethics approval and consent to participate

Not applicable.

### Funding

The research was supported by the Fundación para la Innovación Agraria (Grant NoFIA PYT - 2014 - 0030).

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 8 September 2018 Accepted: 22 February 2019

Published online: 07 March 2019

## References

- Polaina J, MacCabe AP. Industrial enzymes. Structure, function and applications. Dordrecht: Springer; 2007.
- Schäfer T, Borchert TW, Nielsen VS, Skagerlind P, Gibson K, Wenger K, Hatzack F, Nilsson LD, Salmon S, Pedersen S, Heldt-Hansen HP, Poulsen PB, Lund H, Oxenbøll KM, Wu GF, Pedersen HH, Xu H. Industrial enzymes. Adv Biochem Eng Biotechnol. 2007;105:59–131.

3. Singh R, Mittal A, Kumar M, Kumar P. Amylases: a note on current applications. *Int Res J Biol Sci*. 2016;5:27–32.
4. Shahani Dewan S. Global markets for enzymes in industrial applications (report overview). *BIO030J. BCC Research*. 2017; 1–5.
5. Sarmiento F, Peralta R, Blamey JM. Cold and hot extremozymes: industrial relevance and current trends. *Front Bioeng Biotechnol*. 2015;3:148.
6. Białkowska A, Turkiewicz M. Miscellaneous cold-active yeast enzymes of industrial importance. In: Buzzini P, Margesin R, editors. *Cold-adapted yeasts: biodiversity, adaptation strategies and biotechnological significance*. Berlin, Heidelberg: Springer; 2014. p. 377–95.
7. Pulicherla KK, Ghosh M, Kumar PS, Rao KRSS. Psychrozymes—the next generation industrial enzymes. *J Marine Sci Res Dev*. 2011;1:1.
8. Semenova MV, Sinityn OA, Morozova VV, Fedorova EA, Gusakov AV, Okunev ON, Sokolova LM, Koshelev AV, Bubnova TV, Vinetskii YP, Sinityn AP. Use of a preparation from fungal pectin lyase in the food industry. *Appl Biochem Microbiol*. 2006;42:598–602.
9. Kashyap DR, Vohra PK, Chopra S, Tewari R. Applications of pectinases in the commercial sector: a review. *Bioresour Technol*. 2001;77:215–27.
10. Garg G, Singh A, Kaur A, Singh R, Kaur J, Mahajan R. Microbial pectinases: an ecofriendly tool of nature for industries. *3 Biotech*. 2016;6:47.
11. Adapa V, Ramya LN, Pulicherla KK, Rao KR. Cold active pectinases: advancing the food industry to the next generation. *Appl Biochem Biotechnol*. 2014;172:2324–37.
12. Gummadi SN, Kumar DS. Microbial pectic transeliminases. *Biotechnol Lett*. 2005;27:451–8.
13. D'Amico S, Claverie P, Collins T, Georgette D, Gratia E, Hoyoux A, Meuwis MA, Feller G, Gerday C. Molecular basis of cold adaptation. *Philos Trans R Soc Lond B Biol Sci*. 2002;357:917–25.
14. D'Amico S, Collins T, Marx JC, Feller G, Gerday C. Psychrophilic microorganisms: challenges for life. *EMBO Rep*. 2006;7:385–9.
15. Barria C, Malecki M, Arraiano CM. Bacterial adaptation to cold. *Microbiology*. 2013;159:2437–43.
16. Margesin R, Miteva V. Diversity and ecology of psychrophilic microorganisms. *Res Microbiol*. 2011;162:346–61.
17. Cavicchioli R. Cold-adapted archaea. *Nat Rev Microbiol*. 2006;4:331–43.
18. Baeza M, Alcaíno J, Cifuentes V, Turchetti B, Buzzini P. Cold-active enzymes from cold-adapted yeasts. In: Sibirny AA, editor. *Biotechnology of yeasts and filamentous fungi*. Cham: Springer International Publishing; 2017. p. 297–324.
19. Alcaíno J, Cifuentes V, Baeza M. Physiological adaptations of yeasts living in cold environments and their potential applications. *World J Microbiol Biotechnol*. 2015;31:1467–73.
20. Buzzini P, Margesin R. Cold-adapted yeasts: a lesson from the cold and a challenge for the XXI century. In: Buzzini P, Margesin R, editors. *Cold-adapted yeasts*. Berlin, Heidelberg: Springer Berlin Heidelberg; 2014. p. 3–22.
21. Buzzini P, Margesin R. Cold-adapted yeasts. *Biodiversity, adaptation strategies and biotechnological significance*. Berlin: Springer; 2014.
22. Carrasco M, Rozas JM, Barahona S, Alcaíno J, Cifuentes V, Baeza M. Diversity and extracellular enzymatic activities of yeasts isolated from King George Island, the sub-Antarctic region. *BMC Microbiol*. 2012;12:251.
23. Birgisson H, Delgado O, García Arroyo L, Hatti-Kaul R, Mattiasson B. Cold-adapted yeasts as producers of cold-active polygalacturonases. *Extremophiles*. 2003;7:185–93.
24. Yadav AN, Sachan SG, Verma P, Kaushik R, Saxena AK. Cold active hydrolytic enzymes production by psychrotrophic *Bacilli* isolated from three sub-glacial lakes of NW Indian Himalayas. *J Basic Microbiol*. 2016;56:294–307.
25. Ramya LN, Pulicherla KK. Molecular insights into cold active polygalacturonase enzyme for its potential application in food processing. *J Food Sci Technol*. 2015;52:5484–96.
26. Merín MG, Morata de Ambrosini VI. Highly cold-active pectinases under wine-like conditions from non-*Saccharomyces* yeasts for enzymatic production during winemaking. *Lett Appl Microbiol*. 2015;60:467–74.
27. Venkatachalam S, Gowdaman V, Prabakaran SR. Culturable and culture-independent bacterial diversity and the prevalence of cold-adapted enzymes from the Himalayan mountain ranges of India and Nepal. *Microb Ecol*. 2015;69:472–91.
28. Apolinar-Valiente R, Williams P, Romero-Cascales I, Gómez-Plaza E, López-Roca JM, Ros-García JM, Doco T. Polysaccharide composition of Monastrell red wines from four different Spanish terroirs: effect of wine-making techniques. *J Agric Food Chem*. 2013;61:2538–47.
29. Cabeza MS, Baca FL, Puentes EM, Loto F, Baigorri MD, Morata VI. Selection of psychrotolerant microorganisms producing cold-active pectinases for biotechnological processes at low temperature. *Food Technol Biotechnol*. 2011;49:187.
30. Merín MG, Mendoza LM, Fariás ME, Morata de Ambrosini VI. Isolation and selection of yeasts from wine grape ecosystem secreting cold-active pectinolytic activity. *Int J Food Microbiol*. 2011;147:144–8.
31. Pathan AA, Bhadra B, Begum Z, Shivaji S. Diversity of yeasts from puddles in the vicinity of midre lovenbreen glacier, arctic and bioprospecting for enzymes and fatty acids. *Curr Microbiol*. 2010;60:307–14.
32. Sahay S, Hamid B, Singh P, Ranjan K, Chauhan D, Rana RS, Chaurse VK. Evaluation of pectinolytic activities for oenological uses from psychrotrophic yeasts. *Lett Appl Microbiol*. 2013;57:115–21.
33. Naga Padma P, Anuradha K, Reddy G. Pectinolytic yeast isolates for cold-active polygalacturonase production. *Innov Food Sci Emerg Technol*. 2011;12:178–81.
34. Nakagawa T, Nagaoka T, Miyaji T, Tomizuka N. A cold-active pectin lyase from the psychrophilic and basidiomycetous yeast *Cystoflobasidium capitatum* strain PPY-1. *Biotechnol Appl Biochem*. 2005;42:193–6.
35. Nakagawa T, Nagaoka T, Taniguchi S, Miyaji T, Tomizuka N. Isolation and characterization of psychrophilic yeasts producing cold-adapted pectinolytic enzymes. *Lett Appl Microbiol*. 2004;38:383–7.
36. Nakagawa T, Yamada K, Miyaji T, Tomizuka N. Cold-active pectinolytic activity of psychrophilic-basidiomycetous yeast *Cystoflobasidium capitatum* strain PPY-1. *J Biosci Bioeng*. 2002;94:175–7.
37. Lara-Márquez A, Zavala-Páramo MG, López-Romero E, Camacho HC. Biotechnological potential of pectinolytic complexes of fungi. *Biotechnol Lett*. 2011;33:859–68.
38. Pedrolli DB, Monteiro AC, Gomes E, Carmona EC. Pectin and pectinases: production, characterization and industrial application of microbial pectinolytic enzymes. *Open Biotechnol J*. 2009;3:9–18.
39. Sambrook J, Russell D. *Molecular cloning: a laboratory manual*. 3rd ed. New York: Cold Spring Harbor Lab Press; 2001.
40. Miller GL. Use of DNS reagent for the measurement of reducing sugar. *Anal Chem*. 1959;31:426–8.
41. Butler J, MacCallum I, Kleber M, Shlyakhter IA, Belmonte MK, Lander ES, Nusbaum C, Jaffe DB. ALLPATHS: de novo assembly of whole-genome shotgun microreads. *Genome Res*. 2008;18:810–20.
42. Trapnell C, Pachter L, Salzberg SL. TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics*. 2009;25:1105–11.
43. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*. 2013;29:15–21.
44. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L, Raychowdhury R, Zeng Q, Chen Z, Muceli E, Hacohen N, Gnirke A, Rhind N, di Palma F, Birren BW, Nusbaum C, Lindblad-Toh K, Friedman N, Regev A. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat Biotechnol*. 2011;29:644–52.
45. Stanke M, Morgenstern B. AUGUSTUS: a web server for gene prediction in eukaryotes that allows user-defined constraints. *Nucleic Acids Res*. 2005;33:W465–7.
46. Stanke M, Tzvetkova A. AUGUSTUS at EGASP: using EST, protein and genomic alignments for improved gene prediction in the human genome. *Genome*. 2006;7:511.
47. Haas BJ, Zeng Q, Pearson MD, Cuomo CA, Wortman JR. Approaches to Fungal Genome Annotation. *Mycology*. 2011;2:118–41.
48. Yin Y, Mao X, Yang J, Chen X, Mao F, Xu Y. dbCAN: a web resource for automated carbohydrate-active enzyme annotation. *Nucleic Acids Res*. 2012;40:W445–51.
49. Mistry J, Finn RD, Eddy SR, Bateman A, Punta M. Challenges in homology search: HMMER3 and convergent evolution of coiled-coil regions. *Nucleic Acids Res*. 2013;41:e121.
50. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics*. 2012;28:1647–9.
51. Kumar S, Stecher G, Tamura K. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol*. 2016;33:1870–4.

52. Biasini M, Bienert S, Waterhouse A, Arnold K, Studer G, Schmidt T, Kiefer F, Gallo Cassarino T, Bertoni M, Bordoli L, Schwede T. SWISS-MODEL: modelling protein tertiary and quaternary structure using evolutionary information. *Nucleic Acids Res.* 2014;42:252–8.
53. Abdulrachman D, Thongkred P, Kocharin K, Nakpathom M, Somboon B, Narumol N, Champreda V, Eurwilaichitr L, Suwanto A, Nimchua T, Chantasingh D. Heterologous expression of *Aspergillus aculeatus* endopolygalacturonase in *Pichia pastoris* by high cell density fermentation and its application in textile scouring. *BMC Biotechnol.* 2017;17:15.
54. Arevalo-Villena M, Briones-Perez A, Corbo MR, Sinigaglia M, Bevilacqua A. Biotechnological application of yeasts in food science: starter cultures, probiotics and enzyme production. *J Appl Microbiol.* 2017;123:1360–72.
55. Burhan H, Ravinder SR, Deepak C, Poonam S, Fayaz AM, Sanjay S, Ishfaq A. Psychrophilic yeasts and their biotechnological applications—a review. *Afr J Biotechnol.* 2014;13:2188–97.
56. Dartora AB, Bertolin TE, Bilibio D, Silveira MM, Costa JA. Evaluation of filamentous fungi and inducers for the production of endo-polygalacturonase by solid state fermentation. *Z Naturforsch C.* 2002;57:666–70.
57. Jeske L, Placzek S, Schomburg I, Chang A, Schomburg D. BRENDA in 2019: a European ELIXIR core data resource. *Nucleic Acids Res.* 2019;47:D542–9.
58. Ribon AO, Queiroz MV, Coelho JL, de Araújo EF. Differential expression of polygalacturonase-encoding genes from *Penicillium griseoroseum* in different carbon sources. *J Ind Microbiol Biotechnol.* 2002;29:145–8.
59. Niture S. Comparative biochemical and structural characterizations of fungal polygalacturonases. *Biologia.* 2008;63:1–19.
60. Bonivento D, Pontiggia D, Matteo AD, Fernandez-Recio J, Salvi G, Tsernoglou D, Cervone F, Lorenzo GD, Federici L. Crystal structure of the endopolygalacturonase from the phytopathogenic fungus *Colletotrichum lupini* and its interaction with polygalacturonase-inhibiting proteins. *Proteins.* 2008;70:294–9.
61. Federici L, Caprari C, Mattei B, Savino C, Di Matteo A, De Lorenzo G, Cervone F, Tsernoglou D. Structural requirements of endopolygalacturonase for the interaction with PGIP (polygalacturonase-inhibiting protein). *Proc Natl Acad Sci U S A.* 2001;98:13425–30.
62. van Pouderooyen G, Snijder HJ, Benen JAE, Dijkstra BW. Structural insights into the processivity of endopolygalacturonase I from *Aspergillus niger*. *FEBS Lett.* 2003;554:462–6.
63. van Santen Y, Benen JAE, Schröter K-H, Kalk KH, Armand S, Visser J, Dijkstra BW. 1.68-Å Crystal structure of endopolygalacturonase II from and identification of active site residues by site-directed mutagenesis. *J Biol Chem.* 1999;274:30474–80.
64. Pickersgill R, Smith D, Worboys K, Jenkins J. Crystal structure of polygalacturonase from *Erwinia carotovora* ssp. *carotovora*. *J Biol Chem.* 1998;273:24660–4.
65. Cho SW, Lee S, Shin W. The X-ray structure of *Aspergillus aculeatus* polygalacturonase and a modeled structure of the polygalacturonase-octagalacturonate complex. *J Mol Biol.* 2001;311:863–78.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

