

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	The MW's satellite population . . . . .	3
1.1.1	Dwarf galaxies . . . . .	4
1.1.2	The classical dwarf spheroidal galaxies . . . . .	5
1.1.3	The ultra-faint dwarf galaxies . . . . .	7
1.1.4	Dark Matter content in dwarf spheroidal galaxies . . . . .	8
1.1.5	Globular clusters . . . . .	11
1.1.6	The classification of satellite objects . . . . .	14
1.1.7	Tidal effects on satellite objects . . . . .	16
1.2	The data . . . . .	18
1.3	Aims . . . . .	24
<b>2</b>	<b>Scaling relations of Milky Way's satellites</b>	<b>25</b>
2.1	Parameter distributions . . . . .	26
2.1.1	Effect of low numbers of member stars on measured parameters. . . . .	27
2.2	$n - R_e$ correlation . . . . .	30
2.3	Discussion . . . . .	32
2.3.1	Origin of the $n - R_e$ relation . . . . .	32
2.3.2	Surface brightness versus absolute magnitude . . . . .	33
2.3.3	Sérsic index versus Absolute magnitude . . . . .	33
2.3.4	Linear fits to the $\mu_0 - M_V$ and $n - M_V$ relations . . . . .	34
2.3.5	Two separate GC populations . . . . .	36
2.3.6	DM in GCs . . . . .	41
2.3.7	Possible Origin of the HSB and LSB GC groups . . . . .	41
<b>3</b>	<b>Morphology of Milky Way's outer halo satellites</b>	<b>45</b>
3.1	Determination of satellite's morphology . . . . .	46
3.1.1	Matched-filter formulation . . . . .	46
3.1.2	Estimating the member star distributions . . . . .	48
3.1.3	Estimating the background distribution . . . . .	49
3.1.4	Counting member stars . . . . .	49
3.2	Results . . . . .	52
3.2.1	Morphology of classical dSph galaxies . . . . .	54
3.2.2	Morphology of UFD galaxies . . . . .	54
3.2.3	Morphology of outer halo GCs . . . . .	55
3.2.4	Morphology of not classified objects . . . . .	56

3.3 Discussion . . . . .	56
<b>Summary</b>	<b>117</b>
<b>Bibliography</b>	<b>119</b>
<b>Appendix</b>	<b>130</b>
A.1 The MCMC algorithm and the Bayes' theorem . . . . .	131