

CONTENT

CHAPTER 1 – Introduction	1
1.1. Overview and Motivation.....	1
1.2. Geological Setting of Stratabound Cu-(Ag) Deposits.....	5
1.3. Research Objectives and Hypothesis	7
1.4. Methodology	7
1.5. Previous Studies	9
1.6. Publications Resulting from this Thesis.....	10
CHAPTER 2 – Assessing the role of bitumen in the formation of stratabound Cu-(Ag) deposits: Insights from the Lorena deposit, Las Luces district, northern Chile	12
2.1. Introduction	13
2.2. Geological Background.....	15
2.2.1. Upper Jurassic and Lower Cretaceous stratabound Cu-(Ag) deposits.....	15
2.2.2. The Lorena stratabound Cu-(Ag) deposit.....	16
2.3. Samples and Methods	17
2.3.1. SEM, FE-SEM and EMPA methods.....	17
2.3.2. XRF elemental mapping.....	18
2.3.3. Sulfur X-ray absorption near edge structure (XANES) spectroscopy	18
2.4. Results	19
2.4.1. Variations with depth.....	19
2.4.2. Hydrothermal alteration and mineralization	20
2.4.3. Pyrobitumen inclusions in pyrite and chalcocite	21
2.4.4. Elemental composition and pyrobitumen textures	21
2.4.5. Pyrobitumen XRF maps.....	22
2.4.6. Sulfur XANES data	22
2.5. Discussion	23
2.5.1. Paragenetic sequence and ore-forming processes	23
2.5.2. Evidence for hydrocarbon-hydrothermal fluid emulsions	25
2.5.3. Key metals in bitumen.....	26
2.5.4. Pyrobitumen as source of organic sulfur.....	28
2.6. Genetic model and concluding remarks	30
2.7. Acknowledgements	31
2.8. References	31
2.9. Figures	40

CHAPTER 3 – Trace element geochemistry of pyrite from bitumen-bearing stratabound Cu-(Ag) deposits, northern Chile	58
3.1. Introduction	59
3.2. Geological Background.....	60
3.3. Samples and methods	62
3.4. Results	63
3.4.1. Micro-textural observations	63
3.4.2. Chemical composition of pyrite	64
3.5. Discussion	64
3.5.1. Trace element incorporation in pyrite from stratabound Cu-(Ag) deposits	64
3.5.2. Comparisons with IOA and IOCG deposits within the Coastal Cordillera.....	67
3.6. Concluding remarks	68
3.7. Acknowledgements	69
3.8. References	69
3.9. Figures.....	79
CHAPTER 4 – Sulfur isotope and trace element signature of pyrite from bitumen-bearing stratabound Cu-(Ag) deposits, northern Chile: Implications for fluid sources and mineralization processes ...88	
4.1. Introduction	89
4.2. Geological Background.....	91
4.2.1. Las Luces district	91
4.2.2. Copiapó district	91
4.2.3. El Soldado district	92
4.3. Samples and methods	93
4.3.1. LA-ICP-MS	93
4.3.2. In-situ sulfur isotope analysis - Secondary-ion mass spectrometry (SIMS)	94
4.4. Results	95
4.4.1. Chemical composition of pyrite	95
4.4.2. Sulfur isotope composition of pyrite	95
4.4.3. LA-ICP-MS and SIMS transects.....	96
4.5. Discussion	96
4.5.1. Pyrite textures and geochemical variations.....	96
4.5.2. Source(s) of sulfur and fractionation processes	97
4.5.3. Implications for ore mineralization processes.....	99
4.6. Final remarks.....	101

4.7. Acknowledgements	101
4.8. References	101
4.9. Figures	113
4.10. Tables	121
CHAPTER 5 – Conclusions	125
BIBLIOGRAPHY	128
ANNEXES	134
ANNEXES A: Supplementary Material of Chapter 2	134
A.1. Sulfur K-edge spectra for sulfur standards (Bolin et al., 2016)	135
A.2. Representative electron microprobe analyses (EMPA) data (wt.%) of bitumen in Lorena deposit carried out at the Peter Hooper GeoAnalytical Laboratory of Washington State University (WSU), USA and LAMARX Laboratory of the Universidad Nacional de Córdoba, Argentina. b.d.l: below detection limit. Detection limits (wt.%) are shown below each element analyzed	136
A.3. Concentration boxplot for selected minor elements in pyrobitumen from Lorena. Electron microprobe analyses (EMPA) including all data are plotted in weight percent (wt.%) on a vertical logarithmic scale. In the boxplot, mean and median concentrations are shown. A circle outlier is a data point above $1.5 \times (75\text{th percentile} - 25\text{th percentile})$ while a square outlier is above $3.0 \times (75\text{th percentile} - 25\text{th percentile})$	139
A.4. Boxplot showing the concentration of Fe, Cu, S, Cl, Ag and Au in pyrobitumen plotted as a function of sample depth (LO-62, LO-7-1 and LO-15: intermediate-shallow samples, LO-24: deep sample). All EMPA data are plotted in weight percent (wt.%) on a vertical logarithmic scale. In each boxplot, mean concentrations are marked. A circle outlier is above $1.5 \times (\text{quartile } 3 - \text{quartile } 1)$ while a square outlier is above $3.0 \times (75\text{th percentile} - 25\text{th percentile})$	140
ANNEXES B: Supplementary Material of Chapter 3	141
B.1. Electron probe microanalysis (EPMA) data of pyrite from stratabound Cu-(Ag) deposits, Chile. Concentrations are reported in weight percent (wt.%). Detection limits (in wt.%) are shown below each element. LO: Lorena deposit; MTCUL: La Culebra deposit; MTC: Manto Cobrizo deposit. b.d.l: below detection limit.	142
B.2. Laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) analyses of Ag, Au, Cu, Bi, Hg, Tl, Pb, Zn, Se, Te, As, Sb, Ge, Cd, In, Sn, Co, Ni, Mn, Cr, V and Mo of pyrite from stratabound Cu-(Ag) deposits, Chile. Concentrations are reported in parts per million (ppm). Detection limits (in ppm) are shown below each element. LO: Lorena deposit; MTCUL: La Culebra deposit; MTC: Manto Cobrizo deposit. b.d.l: below detection limit.	147
ANNEXES C: Supplementary Material of Chapter 4	150
C.1. BSE images showing pyrite grains from Lorena deposit analyzed by SIMS. In-situ sulfur isotope values are shown.	151

C.2. BSE images showing pyrite grains from La Culebra deposit analyzed by SIMS. In-situ sulfur isotope values are shown.....152

C.3. BSE images showing pyrite grains from Lorena deposit analyzed by SIMS. In-situ sulfur isotope values are shown.....153