

Contents

Contents	vii
List of Tables	ix
List of Figures	xi
1 Introduction	1
1.1 Molecular clouds	1
1.1.1 Formation	2
1.1.2 Classification and Composition	2
1.1.3 Structure	3
1.2 Star formation	4
1.2.1 Low-mass star formation	4
1.2.2 High-mass star formation	5
1.3 Identification and characterization of High-mass clumps	7
1.3.1 Basic parameters of radiation	7
1.3.2 Dust thermal emission	10
1.3.3 Molecular line emission	12
2 Aims and observations	17
2.1 Thesis aims	17
2.1.1 Source selection	17
2.2 The data	18
2.2.1 Single dish observations	18
2.2.2 ALMA observations	20
3 Large scale (~ 0.3 pc) properties of the gas	22
3.1 Results	22
3.1.1 Line profiles	27
3.1.2 Velocity fields	29
3.2 Discussion	35
3.2.1 Virial Masses	35
3.2.2 Evolutionary stages	36
3.2.3 Collapse	36
3.2.4 Radial velocity dispersion profiles	37
4 G305: Small scale (~ 0.02 pc) structure with ALMA	41

4.1	Results	41
4.1.1	Continuum emission	41
4.1.2	Molecular line emission	43
4.2	Discussion	55
4.2.1	Dust column density N-PDF	55
4.2.2	Physical parameters of cores	57
4.2.3	Fragmentation and the Core Mass Function	59
4.2.4	Constraints on models of high-mass star formation	61
5	Conclusions	63
	Bibliography	65

List of Tables

1.1	Classification of Interstellar Cloud Types from Snow and McCall (2006) . . .	3
1.2	Average properties of the different types of molecular clouds: Giant Molecular Clouds (GMC), typical clouds, clumps and low-mass and high-mass cores (Blitz 1993; Garay and Lizano 1999; Bergin and Tafalla 2007; Beuther et al. 2007).	4
2.1	Coordinates of the four massive and dense cold clumps (MDCCs) observed with the APEX telescope.	18
2.2	Summary of the observed molecular transitions with the APEX telescope. . .	20
2.3	Summary of the molecular transitions observed with the ALMA interferometer.	21
3.1	Molecular lines detected towards each of the observed clumps.	22
3.3	Observed parameters of the HCN(3-2), HCO ⁺ (3-2), HNC(3-2), CH ₃ OH(5-4) and CH ₃ OH(9-8) emission for G305.136+0.068, G333.125-0.562, G18.606-0.076 and G34.458+0.121 (if they follow a Gaussian-like shape). The first column lists the molecule, columns 2-5 list, respectively, the antenna temperature, line center velocity, velocity dispersion and velocity integrated antenna temperature at the peak emission; columns 6-9 list, respectively, the antenna temperature, line center velocity from, sigma and velocity integrated antenna temperature of the source averaged emission, and column 10 the FWHM angular size.	29
3.4	Fitted parameters of the N ₂ H ⁺ (3-2) emission for G305.136+0.068, G18.606-0.076 and G34.458+0.121. The fit was done using the CLASS program hfs for hyperfine structure. Columns 2-4 list, respectively, the line center velocity, velocity dispersion and opacity at the peak emission; columns 5-7 list, respectively, the line center velocity, velocity dispersion and opacity of the source averaged emission, and column 8 the FWHM angular size.	34
3.5	Derived physical parameters of MDCC. Columns from left to right: source name, average radius, average line-width, virial mass, dust mass and virial parameter.	35
3.6	Results of the HILL5 model fitting for G305.136+0.068 and G333.125-0.562.	37
3.7	Results of power-law fitting ($\Delta v \propto r^\alpha$) from the radial velocity dispersion profiles from Figure 3.13. The values in column 6 are the average from the columns 2,3,4 and 5.	38
4.1	Observed parameters of the dust continuum cores1.	44
4.2	Observed parameters of the N ₂ H ⁺ (1-0) emission from cores.1	54

4.3	Column densities of cores with a single velocity component in the N_2H^+ $J = 1-0$ transition.	55
4.4	Derived parameters of cores.	58

List of Figures

1.1	Relation between density and molecular and atomic abundance in molecular clouds. The proposed classification of Interstellar Cloud Types from Snow and McCall (2006) is illustrated.	3
1.2	The four stages of star formation from Shu et al. (1987). (a) Cores form within molecular clouds as magnetic and turbulent support is lost, (b) A protostar with a surrounding disk forms at the center of the core collapsing from inside-out. (c) Stellar wind breaks out along the rotational axis of the disk. (d) The infall finishes revealing a newly formed star with a circumstellar disk.	5
1.3	Schematic illustration of the discussed evolutionary sequence for high-mass star formation from Battersby et al. (2010).	6
2.1	Spitzer-IRAC three-color images ($3.6 \mu m$ = blue, $4.5 \mu m$ = green, and $8.0 \mu m$ = red) of the four MDCC showing different star formation activity, increasing from the top left to the bottom right. Overlaid are contours of the dust emission (ATLASGAL $870 \mu m$; levels are drawn at 30%, 40%, 50%, 60%, 70%, 80%, and 90% of the peak value). From top left to bottom right: G18.606-0.076, G305.136+0.068, G333.125-0.562, G34.458+0.25.	19
3.1	Spectral maps of $\text{HCO}^+(3-2)$, $\text{HNC}(3-2)$, $\text{HCN}(3-2)$, $\text{N}_2\text{H}^+(3-2)$, $\text{CH}_3\text{OH}(5-4)$ and $\text{CH}_3\text{OH}(9-8)$ molecular line emission toward G305.136+0.068. The grid spacing is $20''$. The velocity scale ranges from -60 to -10 km/s. The antenna temperature scale ranges from -0.3 to 4.5 K for all molecules, except for the CH_3OH transitions where it ranges from -0.3 to 1.5 K.	23
3.2	Spectral maps of $\text{HCO}^+(3-2)$, $\text{HNC}(3-2)$, $\text{HCN}(3-2)$, $\text{N}_2\text{H}^+(3-2)$, $\text{CH}_3\text{OH}(5-4)$ and $\text{CH}_3\text{OH}(9-8)$ molecular line emission toward G333.125-0.562. The grid spacing is $20''$. The velocity scale ranges from -85 to -35 km/s. The antenna temperature scale ranges from -0.3 to 4 K for all molecules, except for the CH_3OH transitions where it ranges from -0.3 to 1.3 K.	24
3.3	Spectral maps of $\text{HCO}^+(3-2)$, $\text{HNC}(3-2)$, $\text{HCN}(3-2)$, $\text{N}_2\text{H}^+(3-2)$ and $\text{CH}_3\text{OH}(5-4)$ molecular line emission toward G18.606-0.076. The grid spacing is $20''$. The velocity scale ranges from 20 to 70 km/s. The antenna temperature scale ranges from -0.3 to 3 K for all molecules, except for $\text{CH}_3\text{OH}(5-4)$ where it ranges from -0.3 to 1 K.	25

3.4	Spectral maps of $\text{HCO}^+(3-2)$, $\text{HNC}(3-2)$, $\text{HCN}(3-2)$, $\text{N}_2\text{H}^+(3-2)$, $\text{CH}_3\text{OH}(5-4)$ and $\text{CH}_3\text{OH}(9-8)$ molecular line emission toward G34.458+0.121. The grid spacing is $20''$. The velocity scale ranges from 35 to 85 km/s. The antenna temperature scale ranges from -0.3 to 3 K for all molecules, except for the CH_3OH transitions where it ranges from -0.3 to 1 K.	26
3.5	Spectra of $\text{CH}_3\text{OH}(5-4)$ (orange), $\text{HCN}(3-2)$ (red), $\text{HCO}^+(3-2)$ (blue), $\text{HNC}(3-2)$ (green) and $\text{N}_2\text{H}^+(3-2)$ (purple) emission observed at the peak position of G305.136+0.068, G333.125-0.562, G18.606-0.076 and G34.458+0.121.	27
3.6	Integrated spectra of $\text{CH}_3\text{OH}(5-4)$ (orange), $\text{HCN}(3-2)$ (red), $\text{HCO}^+(3-2)$ (blue), $\text{HNC}(3-2)$ (green) and $\text{N}_2\text{H}^+(3-2)$ (purple) emission toward G305.136+0.068, G333.125-0.562, G18.606-0.076 and G34.458+0.121.	28
3.7	G305.136+0.068 moment maps of the $\text{HCN}(3-2)$, $\text{HCO}^+(3-2)$, $\text{HNC}(3-2)$ and $\text{N}_2\text{H}^+(3-2)$ emission (from top to bottom). The bottom left panel shows the ATLASGAL continuum emission. The panels from left to right: integrated intensity (zeroth moment, units of K kms^{-1}), intensity weighted velocity field (first moment, units of kms^{-1}), intensity weighted velocity dispersion (second moment, units of kms^{-1}). Overlaid with white contours are the contours of the integrated intensity (levels are 20, 40, 60 and 80% of the peak).	30
3.8	G333.125-0.562 integrated intensity moment maps (zeroth moment, units of K kms^{-1}) of the $\text{HCN}(3-2)$, $\text{HCO}^+(3-2)$, $\text{HNC}(3-2)$ and $\text{N}_2\text{H}^+(3-2)$ emission (from top to bottom). The bottom left panel shows the ATLASGAL continuum emission.	31
3.9	G18.606-0.076 integrated intensity moment maps (zeroth moment, units of K kms^{-1}) of the $\text{HCN}(3-2)$, $\text{HCO}^+(3-2)$, $\text{HNC}(3-2)$ and $\text{N}_2\text{H}^+(3-2)$ emission (from top to bottom). The bottom left panel shows the ATLASGAL continuum emission. Also shown are the intensity weighted velocity field (first moment, units of kms^{-1}) and the intensity weighted velocity dispersion (second moment, units of kms^{-1}) for $\text{HNC}(3-2)$ and $\text{N}_2\text{H}^+(3-2)$. Overlaid with white contours are the contours of the integrated intensity (levels are 20, 40, 60 and 80% of the peak).	32
3.10	G34.458+0.121 integrated intensity moment maps (zeroth moment, units of K kms^{-1}) of the $\text{HCN}(3-2)$, $\text{HCO}^+(3-2)$, $\text{HNC}(3-2)$ and $\text{N}_2\text{H}^+(3-2)$ emission (from top to bottom). The bottom left panel shows the ATLASGAL continuum emission. Also shown are the intensity weighted velocity field (first moment, units of kms^{-1}) and the intensity weighted velocity dispersion (second moment, units of kms^{-1}) for $\text{HCO}^+(3-2)$, $\text{HNC}(3-2)$ and $\text{N}_2\text{H}^+(3-2)$. Overlaid with white contours are the contours of the integrated intensity (levels are 20, 40, 60 and 80% of the peak).	33
3.11	$\text{HCN}(3-2)$ (solid line) and $\text{HNC}(3-2)$ (dotted line) peak position spectra toward G305.136+0.068. The red line shows the best result of the fitting of the HILL5 model.	36
3.12	$\text{HNC}(3-2)$ (solid line) and $\text{CH}_3\text{OH}(5-4)$ (dotted line) peak position spectra toward G333.125-0.562. The red line shows the best result of the fitting of the HILL5 model.	37

3.13	Radial velocity dispersion profiles from G305.136+0.068, G333.125-0.562, G18.606-0.076 and G34.458+0.121 using the HCO ⁺ (blue), HNC (magenta) and N ₂ H ⁺ (green) data. The dashed lines are linear fittings to the points. The results are listed in Table 3.7.	39
3.14	Radial velocity dispersion profiles for the 15 sources from the SuperMALT and G305.136+0.068, G333.125-0.562, G18.606-0.076 and G34.458+0.121, grouped following their evolutionary classification: quiescent (left panel), protostellar (center panel) and H II region (right panel).	40
4.1	Images of the millimetre and infrared emission toward G305.137+0.069 at ~2'' angular resolution. Left panel: 3 mm image from ALMA. Right panel: 4.5 μm image from <i>Spitzer Space Telescope</i> . In both images the white contour levels correspond to single dish dust continuum emission (ATLASGAL 870 μm, ~19'' angular resolution, levels are drawn at 20, 50 and 90% of the peak emission). The stars show the position of three Spitzer sources reported by Garay et al. (2015). The scale bar marks a size of 0.1 pc on the bottom right corner of the left panel.	42
4.2	ALMA 3mm dust continuum compact structures (shown as orange ellipses) extracted using the GaussClump algorithm (12 cores). The parameters used in the search are <i>threshold</i> = 3.5σ, <i>step</i> = 1σ and <i>size</i> = 1 beam.	44
4.3	Images of the dust continuum emission and velocity integrated line emission from G305.136+0.068. Upper left: 3mm continuum. Upper right: HCO ⁺ . Lower left: N ₂ H ⁺ . Lower right: CS. In each panel overlaid in white contours is the 3 mm dust continuum emission at 20%, 50% and 80% of the peak. The color scale is in units of mJy beam ⁻¹ in the continuum image and in mJy beam ⁻¹ km s ⁻¹ for the line images. The peak of each line emission is shown by the black crosses. The angular resolution is shown in the lower left corner of the continuum emission image.	45
4.4	Spectra of the line emission observed at the peak position of the three brighter mm sources (see Table 4.1 for more details). From left to right: mm-8, mm-9 and mm-10. From top to bottom: HCO ⁺ , CS and N ₂ H ⁺ . The dashed vertical line shows the clump velocity of -36.5 km/s (Garay et al. 2015). The brackets show the velocity intervals used for the integration of the moment maps. . .	46
4.5	30''×30'' spectral map of the HCO ⁺ emission towards the central region of G305.137+0.069. The grid spacing is 2'' (roughly the beam size).	47
4.6	Moment maps of the HCO ⁺ , CS and N ₂ H ⁺ emission (from top to bottom). From left to right: integrated intensity (zeroth moment, units of Jy beam ⁻¹ km s ⁻¹), intensity weighted velocity field (first moment, units of km s ⁻¹), intensity weighted velocity dispersion (second moment, units of km s ⁻¹). Overlaid in white are contour levels of the integrated intensity (20, 40, 60 and 80% of the peak). The beam is shown in the lower left corner.	49
4.7	Position velocity diagrams of HCO ⁺ , N ₂ H ⁺ and CS. The cuts were made along two perpendicular directions shown in the upper panels. The south-east to north-west cut is shown on the left column, and the north-east to south-west cut in the right column. The positions of the mm-sources crossed by the cuts are shown with arrows. The center of the cuts, from which the offset is measured, are shown by blue crosses on the top panels.	50

4.8	CS maps of intermediate (IV) and high velocity(HV) gas emission overlaid on the ALMA dust continuum (gray scale). The range of flow velocity integration for the IV map (upper panel) is from -5 to -9 km s $^{-1}$ for the blueshifted gas (blue contours) and from 5 to 9 km s $^{-1}$ for the redshifted gas (red contours) and for the HV map (bottom panel) is from -14 to -9 km s $^{-1}$ for the blueshifted gas and from 9 to 14 km s $^{-1}$ for the redshifted gas.	52
4.9	Spectra of the integrated N $_2$ H $^+$ (1-0) emission from the cores. The red line shows the result of an hyperfine fit using a single velocity component (labeled S) or two velocity components (labeled D). The fitted parameters are given in Table 4.2.	53
4.10	Column density PDF Gaussian kernel density estimation for G305.136+0.068 (blue line). Blue shaded regions show the uncertainties obtained through the bootstrap method (Efron 1982). The top panel shows the observed (blue line and shaded region) and inferred (gray line and shaded region) PDFs. The inferred PDF is a combination of three distributions: a Gaussian for the noise, a log-normal for the turbulent gas, and a power-law for the self-gravitating gas. The bottom panel shows the PDF with the noise subtracted, and the inferred contributions of the log-normal (orange) and power-law (green) distributions. The vertical red line shows the point at which the distributions deviate from log-normal, $N(\text{H}_2)=(8.4 \pm 0.23)\times 10^{22}$ cm $^{-2}$. The thickness of the lines represents 1σ errors. The 1σ noise level is equivalent to $N(\text{H}_2)=1.4\times 10^{22}$ cm $^{-2}$	56
4.11	Dust mass versus virial mass for the seven cores with derived line widths from hyperfine fitting of N $_2$ H $^+$ J(1-0) emission (see Table 4.4. The solid line represents $\alpha_{\text{vir}} = 1$).	59
4.12	Core mass function (CMF) for the cores within G305.137+0.069. The blue line shows the cumulative distribution function of the cores, the orange line shows the initial mass function distribution (IMF) from Kroupa (2001). The green line and green shaded area show the maximum likelihood estimation of the power law index ($\gamma = 0.02$) and its 90% confidence interval ($\gamma \in [-0.60, 0.64]$), respectively.	60