

Table of Content

| | | |
|----------|---|-----------|
| 1 | Introduction | 1 |
| 1.1 | Supernovae | 1 |
| 1.1.1 | Light curves of SNe | 2 |
| 1.1.2 | SNe LC modelling | 3 |
| 1.1.3 | Type II SNe | 5 |
| 1.2 | Hydrodynamical models | 6 |
| 1.2.1 | Moriya wind acceleration models | 7 |
| 1.3 | Surveys | 9 |
| 2 | Data | 11 |
| 2.1 | ZTF | 11 |
| 2.2 | ATLAS | 11 |
| 3 | Methods | 15 |
| 3.1 | Data selection | 15 |
| 3.2 | Light curve cleaning | 16 |
| 3.3 | Physical parameters | 18 |
| 3.3.1 | Synthetic light curves | 18 |
| 3.3.2 | Interpolation | 19 |
| 3.3.3 | MCMC | 20 |
| 4 | Results | 26 |

| | | |
|-------------------------------|--|-----------|
| 4.1 | Posterior distribution | 26 |
| 4.2 | Sample distribution | 34 |
| 4.3 | Inferred redshift vs host redshift | 35 |
| 5 | Analysis | 41 |
| 5.1 | Validating our results | 41 |
| 5.2 | Our method as a distance indicator | 44 |
| 5.3 | Limitations of our method | 45 |
| 5.4 | Implementation on LSST | 46 |
| 6 | Conclusions | 48 |
| Bibliography | | 50 |
| ANNEX Host association | | 60 |