

# LEAF ALKALOIDS OF *SOPHORA MACROCARPA*

ROSA NEGRETE and NADINE BACKHOUSE

*Department of Pharmaceutical Sciences, Faculty of Basic and Pharmaceutical Sciences,  
University of Chile, Santiago, Chile*

and

BRUCE K. CASSELS

*Department of Chemistry, Faculty of Science, University of Santiago, Santiago, Chile*

The seeds of *Sophora macrocarpa* Sm. (Fabaceae) are a promising source of edible oil (1), and contain matrine, *N*-methylecytisine and baptifoline (2). Glc analyses of the leaves of this species suggested that matrine is always, by far, the major alkaloid, accompanied by several unidentified minor bases (3). Fractionation of the crude leaf alkaloids has now afforded matrine, matrine *N*-oxide, sophoranol, *N*-methylecytisine and cytisine.

## EXPERIMENTAL<sup>1</sup>

**PLANT MATERIAL.**—*Sophora macrocarpa* leaves were harvested in November, 1978, after flowering, at the eastern foot of La Dormida Pass, about 40 km north of Santiago, Chile. Voucher specimens are deposited at the Natural History Museum in Santiago.

**EXTRACTION AND ISOLATION OF ALKALOIDS.**<sup>2</sup>—Air-dried leaves of *Sophora macrocarpa* (9.0 kg), when worked up by standard procedures, yielded 130 g of crude alkaloids. Matrine, *N*-methylecytisine, cytisine, matrine *N*-oxide and sophoranol were identified by mp,  $[\alpha]_D$ , uv, ir, ms, <sup>1</sup>H and <sup>13</sup>C nmr. The first three alkaloids were further characterized by the comparison with reference samples and mp of derivatives. Matrine *N*-oxide was compared with a synthetic sample prepared from matrine and H<sub>2</sub>O<sub>2</sub>.

## ACKNOWLEDGMENTS

We are grateful to Professors L.E. Navas for the identification of the plant material and J. Bartulin (ms and <sup>13</sup>C nmr), G. Eckhardt (ms), E. Breitmaier (<sup>13</sup>C nmr), M. Shamma and F. Horn (<sup>1</sup>H nmr), and J. Medina (ir) for spectra. We acknowledge the technical assistance of N.E. Creuz, H. Fuentes, H. Maldonado, P.J. Manriquez, V. Soto and S. Tortello. This work was supported by SDCACI (University of Chile), DICYT (University of Santiago), and the Organization of American States.

Received 20 November 1981

## LITERATURE CITED

1. A. de Mayo and B. K. Cassels, *Contribuciones (UTE)*, (5), 33 (1971).
2. M. Silva, M. Medina and P. G. Sammes, *Phytochemistry*, 7, 661 (1968).
3. A. Riquelme, Thesis, University of Chile, 1979.

<sup>1</sup>Melting points were determined on a Reichert Kofler hot plate and are uncorrected. Uv spectra were recorded on a Zeiss DMR-21 spectrophotometer, and ir spectra were determined on a Leitz III G instrument. <sup>1</sup>H and <sup>13</sup>C nmr spectra were recorded with Varian HA-100 and CFT-20 spectrometers, respectively. Electron impact ms were obtained at 70 eV with a Varian Mat CH-7 instrument. Optical rotations were measured on Perkin-Elmer 141 and 241 polarimeters.

<sup>2</sup>Full details of the isolation and identification of the compounds are available on request to R.N.