

Mitochondrial DNA Polymorphisms in Chilean Aboriginal Populations: Implications for the Peopling of the Southern Cone of the Continent

MAURICIO L. MORAGA,¹ PAOLA ROCCO,¹ JUAN F. MIQUEL,²
FLAVIO NERVI,² ELENA LLOP,¹ RANAJIT CHAKRABORTY,³
FRANCISCO ROTHHAMMER,¹ AND PILAR CARVALLO^{1*}

¹*Programa de Genética Humana, Instituto de Ciencias Biomédicas, Facultad de Medicina, Universidad de Chile, Santiago 7, Chile*

²*Departamento de Gastroenterología, Facultad de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile*

³*Human Genetic Center, University of Texas School of Public Health, Houston, Texas 77225*

KEY WORDS Amerindian mtDNA haplogroups; sequence analysis; peopling of the southern cone

ABSTRACT The mitochondrial DNAs (mtDNAs) from individuals belonging to three Chilean tribes, the Mapuche, the Pehuenche, and the Yaghan, were studied both by RFLP analysis and D-loop (control region) sequencing. RFLP analysis showed that 3 individuals (1.3%) belonged to haplogroup A, 19 (8%) to haplogroup B, 102 (43%) to haplogroup C, and 113 (47.7%) to haplogroup D. Among the 73 individuals analyzed by D-loop sequencing, we observed 37 different haplotypes defined by 52 polymorphic sites. Joint analysis of data obtained by RFLP and sequencing methods demonstrated that, regardless of the method of analysis, the mtDNA haplotypes of these three contemporary South American aborigine groups clustered into four main haplogroups, in a way similar to those previously described for other Amerindians. These results further revealed the absence of haplogroup A in both the Mapuche and Yaghan as well as the absence of haplogroup B in the Yaghan. These results suggest that the people of Tierra del Fuego are related to tribes from south-central South America. *Am J Phys Anthropol* 113:19–29, 2000. © 2000 Wiley-Liss, Inc.

The analysis of mitochondrial DNA variation has been extensively used in the recent past, for genetic characterization of contemporary aboriginal populations of the Americas. The primary goals of this analysis have been to determine the origins, relationships, and migrational patterns of New World populations. In this respect, the study of the frequency distribution of the four founding Amerindian haplogroups, suggested by Schurr et al. (1990) and defined by Torroni et al. (1992), has proven useful.

The majority of these studies have focused on aboriginal populations from North

and Central America. Of those studies conducted with South American groups, almost all have involved populations from the northern region of the continent (Torroni et al., 1993; Easton et al., 1996; Santos et al., 1996). Most of the data generated from these studies were obtained through RFLP

Grant sponsor: FONDECYT; Grant number: 198-1111; Grant sponsor: Ministero degli Affari Esteri d'Italia; Grant number: 1091-G224/ICU/CILE; Grant sponsor: NIH; Grant number: GM 41399.

*Correspondence to: Pilar Carvalho, Departamento de Biología Celular y Molecular, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Casilla 114-D, Santiago, Chile. E-mail: pcarvall@genes.bio.puc.cl

Received 23 November 1998; accepted 28 March 2000.

mapping, rather than by D-loop sequencing, although some D-loop sequence data from these populations are available for analysis. Similarly, the mtDNAs taken from southern South American groups were primarily characterized by RFLP analysis (Merriwether et al., 1995; Lalueza et al., 1997). Thus, there remains limited D-loop sequence information from groups inhabiting the southern part of South America.

Studies of South Amerindians using RFLP and D-loop sequence analysis have revealed a concordance between the distribution of haplogroups A–D, and haplogroups (lineage clusters) I–IV as defined by Horai et al. (1993), based on the nucleotide polymorphisms occurring in the D-loop of each haplogroup (Torroni et al., 1993; Easton et al., 1996; Santos et al., 1996). However, Santos et al. (1996) found that 7% of individuals belonging to tribes from the Brazilian Amazon Region comprise unusual haplogroups because of their lack of correspondence between RFLP and sequencing analysis. There is also extensive information referring to RFLP analysis in these populations. In addition, Bailliet et al. (1994) described a subdivision of the classical Amerindian haplogroups A, C, and D on the basis of the presence or absence of the *HaeIII* np 16517 site. This division has led to the creation of haplotypes A1/A2, C1/C2, and D1/D2. However, the *HaeIII* np 16517 site has been shown to be highly polymorphic in African (Chen et al., 1995), European (Torroni et al., 1996), and Asian (Ballinger et al., 1992) populations, indicating that it may not be useful for distinguishing within Native American haplogroups A, C, and D. Similarly, Forster et al. (1996) also postulated subdivisions of founding haplotypes, using D-loop sequence data. In this case, they divided haplotypes A1 and A2 by a C→T transition at np 16111 and defined D1 and D2 haplotypes by the presence or absence of the 16271 T→C transition, with this transition being present only in North American Indian mtDNAs. Hence, there would appear to be specific nucleotide changes in the D-loop, which distinguish between North-Central and South American populations.

To expand our knowledge about the extent of mtDNA variation in southern South America and to test the hypothesis of a single wave of migration to the southern part of South America, we performed RFLP and sequence analysis on three Chilean aboriginal populations (the Pehuenche, the Mapuche, and the Yaghan).

MATERIALS AND METHODS

Population samples

The samples include three different Chilean aboriginal populations: 105 Pehuenche from the community of Trapa Trapa in the Province of Bio Bio, 8th region (37° 43' S; 71° 16' W), 111 Mapuche from the Huapi Island, Province of Valdivia, 10th region (40° 15' S; 72° 25' W), and the 21 last surviving Yaghan from Puerto Williams, Navarino Island, Province of Antartica, 12th region (55° S; 67° 40' W). The three groups analyzed correspond to Amerinds showing a percentage of admixture between 2–13% (Llop et al., 1993, 1994). The geographic locations of the subjects are shown in Figure 1.

DNA extraction and PCR amplification

Total DNA was prepared from peripheral blood lymphocytes (Gustincich et al., 1991). DNA amplification of the four polymorphic mtDNA regions, defining the four Amerindian haplogroups, was carried out by using the following specific primers:

Haplogroup A:

L604 GTAGCTTACCTCCTCAAAGCAA

H708 AGGGTGAACCTCACTGGAACG

Haplogroup B:

L8214 CACAGTTTCATGCCCATCGT

H8294 ATGCTAAGTTAGCTTTACAGTGG

Haplogroup C:

L13232 CGCCCTTACACAAAATGACATCAA

H13344 GGAGCACATAAATAGTATGGC

Haplogroup D:

L5120 CCTAACTACTACCGAATTCCTA

H5255 ATTCTTCGATAATGGCCCATTTG

PCR was performed in a final volume of 50 µl, containing 300 ng genomic DNA, 1 U Taq DNA polymerase (Promega), 25 pmoles of each primer, 200 nM of each deoxynucleotide, and the appropriate buffer. Samples



Fig. 1. Map of Chile, showing the geographic location of the three Chilean tribes included in the present study.

were processed under the following PCR conditions: 1 cycle at 95°C for 5 min, followed by 30 cycles at 95°C for 1 min, 55°C for 1 min, and 72°C for 1 min, and finally one cycle at 72°C for 5 min. Haplotypes A, C, and D were analyzed by restriction digestion, using *Hae*III for haplotype A, *Hinc*II for haplotype C, and *Alu*I for haplotype D.

The resulting restriction fragments and the PCR product including region V, defining haplotype B, were analyzed by electrophoresis on 3% NuSieve-Agarose (2:1) (FMC BioProducts) gels. PCR amplification of the D-loop region was carried out with two specific primers, spanning a region of 1,021 bp: L15997, CACCATTAGCACCCA-AAGCT; and H408, CTGTTAAAAGTGCAT-ACCGCCA.

Direct sequencing of PCR products

DNA sequencing was carried out by using the PCR dsDNA Cycle Sequencing System (Life Technologies), with the prim-

ers L29, GGTCTATCACCTATTAACCAC; and H16401, CACCATCCTCCGTGAAATCA, labeled at the 5' end with γ -³²P-ATP. DNA sequence analysis was performed by electrophoresis on high-resolution 6% polyacrylamide gels, followed by autoradiography.

Data analysis

Sequence alignment was performed by using the Clustal W computer program (Thompson et al., 1994). Substitutions and deletions were inferred from direct sequence comparisons. The phylogenetic trees were obtained through two different methods: neighbor joining (Saitou and Nei, 1987), using the Clustal W program, and parsimony analysis (Swofford and Olson, 1990). Maximum parsimony trees were generated through random addition of sequences using the tree bisection and reconnection (TBR) algorithm, with the PAUP (Swofford, 1993) computer program.

RESULTS

RFLP analysis

All individuals analyzed showed one of the characteristic haplogroups, A, B, C, or D. As shown in Table 1, haplogroups C and D were the most frequent in the three populations, appearing at very similar frequencies in all of them. Haplogroup A was present in the Pehuenche at a very low frequency (2.8%), while both haplogroups A and B were absent in the Yaghan group. We tested the gain of the *Hae*III site at position 16517 only in individuals belonging to haplogroup B, since this is a specific marker to differentiate Amerindians from non-Amerindian B individuals (Torrioni et al., 1992, 1993). The gain of the *Hae*III site was present in all individuals exhibiting haplogroup B, including 8 Mapuche and 11 Pehuenches.

A 3 × 4 contingency table analysis suggests that in terms of haplogroup frequencies, these three populations cannot be statistically distinguished (chi-square = 6.78, exact *P*-value = 0.32 with 10,000 permutations). However, in terms of haplotype diversity (Nei, 1978) defined by these four haplogroups, the Yaghan have the lowest diversity (52.4%), and the Pehuenche the

TABLE 1. MTDNA haplogroup distribution in three Chilean aboriginal populations

Populations	Haplogroups (%)				
	N	A	B	C	D
Mapuche	111	0.0	7.2	44.1	48.7
Pehuenche (Trapa-Trapa)	105	2.8	10.5	41.0	45.7
Yaghan	21	0.0	0.0	47.6	52.4
Total	237	1.3	8.0	43.0	47.7

highest (61.7%), with the Mapuche being intermediate (56.8%), the differences being statistically insignificant ($P = 0.11$).

Sequence analysis

The two hypervariable regions of the D-loop were analyzed by direct sequencing of the DNA from 73 individuals from the three native populations, which were representative of the four haplogroups described previously. As shown in Table 2, 37 different haplotypes were observed, defined by 52 polymorphic sites. It is important to note that two haplotypes shared by individuals from different populations are shown separately. This gives a total of 40 lines in Table 2, corresponding to only 37 haplotypes. In haplotype C, lineages PE3C, YA4C, and HU4C are the same, as well as YA2C and HU1C.

Only nucleotide changes at np 73 and np 263 are shared by all haplogroups. One of these, transition A→G at position 73, has also been described in all individuals from an Argentinean Mapuche group (Ginther et al., 1993), as well as in half of the individuals from a Huétar sample of Costa Rica (Santos et al., 1994). Two types of adenine deletions, an A at np 248 and an AA at 286–291, were found at hypervariable region II (HVS-II). These A deletions have been found in all haplogroup C individuals from the three Chilean aboriginal populations, as well as the Argentinean Mapuche sample (Ginther et al., 1993). Since these A deletions have not been described for other populations (Handt et al., 1998), they may be characteristic nucleotide changes at least for South American aboriginal populations.

In these three Chilean groups, as with haplogroup diversity, we found the highest sequence diversity in the Pehuenche sample, showing 38 out of 52 total polymorphic sites at the D-loop, amounting to 93.2% se-

quence diversity. The Yaghan, likewise, showed the smallest sequence diversity of 91.4%, although this was not significantly smaller than the others.

Disregarding the phylogenetic relationships of the observed sequences, at the sequence level, the Yaghan appear to be almost 3–5 times as distant from the Pehuenche and the Mapuche, (the standard distance of Nei (1978) being 0.31 between the Pehuenche and the Mapuche, 1.55 between the Pehuenche and the Yaghan, and 1.09 between the Mapuche and the Yaghan). The contingency table analysis of sequence-lineage frequency differences suggests that the genetic distances among these three populations are statistically significant (chi-square = 15.12, $P < 10^{-5}$, with 10,000 permutations).

Haplogroup D was the most diverse at the sequence level, revealing 30 polymorphic sites, in contrast to haplogroups A with 13 changes, B with 14 changes, and C with 16 changes. We also found that, from all individuals classified in the four haplogroups A–D (Torroni et al., 1992), only haplogroups A–C corresponded perfectly to 3 of the 4 clusters described by Horai et al. (1993). The T nucleotide at position 16290 and A 16319 define haplogroup A/cluster III. Haplogroup B/cluster I is defined by C 16189 and C 16217. Haplogroup C/cluster IV is defined by a C 16298 and T 16327.

Regarding haplogroup D/cluster II, C nucleotides at position 16325 and 16362 have been described by Horai et al. (1993) as characteristics of cluster II, since they were shared by 72 native Americans, including 45 Chilean aborigines. From our study, one Mapuche individual lacked the 16362 T→C transition, and the other 13 individuals belonging to the three Chilean populations did not show the characteristic 16325 T→C transition. Hence, our results do not corrob-

TABLE 2. Polymorphic mtDNA D-loop sites in a sample of Chilean aborigines¹

																										Anderson													
																										Individuals	Lineages												
	T	T	C	G	G	C	T	A	T	G	T	C	C	A	A	T	C	C	A	T	T	G	T	A	T	G	A	A	A	A	C	A	A	A					
16086																																						3	PE1A
16092				T	A																																2	PE6B	
16111							C																														1	PE5B	
16129							C																														1	PE4B	
16153							C																														1	PE1B	
16187							C																														4	HU1B	
16189							C																														3	HU3B	
16207							C																														1	HU2B	
16209							C																														1	PE2B	
16213							C																														1	PE3B	
16217							C																														1	PE2C	
16218							C																														1	YA3C	
16223							C																														1	PE1C	
16234							C																													6	PE3C		
16241							C																														1	YA4C	
16249							C																														7	HU4C	
16270							C																														2	YA2C	
16286							C																														3	HU1C	
16290							C																														1	HU2C	
16291							C																														1	HU3C	
16298							C																														2	YA1C	
16301							C																														2	YA4D	
16304							C																														2	YA3D	
16311							C																														1	HU10D	
16319							C																														2	HU9D	
16325							C																														1	HU1D	
16327							C																														1	PE3D	
16342							C																														1	PE4D	
16343							C																														2	PE5D	
16362							C																														1	PE2D	
53							C																														1	PE1D	
55							C																														1	HU7D	
56							C																														1	HU4D	
64							C																														1	HU2D	
73							C																														3	HU5D	
77							C																														1	HU6D	
94							C																														2	HU8D	
129							C																														1	HU3D	
143							C																														4	YA2D	
146							C																														1	YA1D	
150							C																																
152							C																																
153							C																																
195							C																																
207							C																																
215							C																																
234							C																																
248							C																																
258							C																																
263							C																																
286-291							C																																
							C																																
							C																																

¹The number of individuals for each lineage is shown. In the right column each lineage is named by its corresponding population (PE, Pehuenche; HU, Mapuche (Huapi Island); YA, Yaghan) the sample number, and the RFLP haplogroup. Individuals highlighted in bold share the same sequence and belong to different groups (PE3C, YA4C, HU4C, and YA2C, HU1C). The sequence published by Anderson et al. (1981) was used as a reference.

orate the classification of haplogroup D/cluster II on the basis of these mutations, but are in agreement with the results of Ginther et al. (1993), who also observed the absence of the 16325 T→C mutation in haplogroup D mtDNAs from the Argentinean Mapuche group.

It is interesting to note that one of these nucleotide changes, C 16325, is present in all individuals of haplogroup C/cluster IV from our study, in the Argentinean Mapuche, and in Amazonian individuals (Santos et al., 1996). Also, in our study, C in position 16362 is present in individuals from cluster III/haplogroup A. Therefore it seems that there are no characteristic and exclusive nucleotide changes in either hypervariable region defining haplogroup D, at least for these aboriginal populations from South America. However, we found a C→T change at nucleotide 16187 which is present in the majority of South Amerindians of haplogroup D, which is absent from haplogroups A–C. This change is also present in 6/10 Argentinean Mapuche (Ginther et al., 1993), and in 14/18 native Americans (Horai et al., 1993). From this analysis, we suggest that this change is the most characteristic one for South Amerindian populations.

Phylogenetic analysis

The D-loop DNA sequences, spanning hypervariable regions I and II from the 37 Chilean lineages, were aligned and submitted to phylogenetic analysis through neighbor-joining (NJ) (Fig. 2) and parsimony methods (data not shown). In the NJ tree, we can see that all lineages fall into four differentiated branches, including haplogroups A–D. Nevertheless, it is noteworthy that three D lineages, HU8D, HU9D, and YA4D, cluster close to the C branch, while HU10D and YA3D cluster close to branch A. All the other D lineages fall into one clade.

The majority rule consensus (data not shown) shows that two branches B and C cluster independently in 100% of maximum parsimony trees. Seven out of 19 lineages corresponding to haplogroup D cluster in one branch, with 74 % consensus from a total of 500 maximum parsimony trees. The remaining lineages, also corresponding to haplogroup D, appeared unresolved in this

analysis. D-loop sequences from the 3 Pehuenche individuals leading to haplogroup A correspond to one lineage, clustering obviously in one branch of 100% consensus.

DISCUSSION

In the present paper we report on the results of a mtDNA RFLP and sequence polymorphism analysis of three Chilean aboriginal populations. The RFLP analysis showed that all individuals belong to one of the previously defined Amerindian haplogroups, A, B, C, or D. In terms of haplogroup frequencies, the three Chilean aboriginal groups studied here cannot be genetically distinguished. However, as shown in Table 3, combining the present data with other published haplogroup frequencies of South American aborigines, it is noteworthy that the frequencies of haplogroups A and B decrease from north to south in meridian South America, whereas haplogroups C and D tend to increase. This trend is most likely the result of founder effect, which occurred during the initial peopling of the southern cone of the continent. As Paleoindians moved south, new territory was probably colonized by small bands that were carriers of a subsample of genes from the ancestral populations they left behind. Haplogroups A and B probably got lost as a consequence of this process. An alternative hypothesis, e.g., gene flow from migrant populations carrying A and B into more ancient groups possessing C and D, would imply that more than one migration event occurred during the peopling of the southern cone of South America. This hypothesis is not supported by recently published gene geography maps of South America (Rothhammer and Silva, 1992; Rothhammer et al., 1997).

Turning now to the sequence data, we note that the three Chilean aboriginal groups, can be genetically distinguished at the D-loop sequence level, making the Yaghan most distant (as well as least diverse) from the Pehuenche and the Mapuche.

Further, the nucleotide changes described by Forster et al. (1996), including one base substitution at nucleotide 16111 leading to subhaplogroups A1/A2, and another substitution at 16271 defining subhaplogroups D1/D2, have been tested in the present




Fig. 2. Tree obtained by the neighbor-joining method for 37 Chilean haplogroups. Individuals highlighted in bold share the same sequence and belong to different groups (PE3C, YA4C, HU4C, and YA2C, HU1C). An African sequence was used as an outgroup in order to root the tree. The alignment was performed over 549 bp, including hypervariable regions I and II. Letters on the side indicate respective RFLP haplogroups.

study (Table 2). All individuals from haplogroup A did have the C→T transition at 16111, and therefore are subhaplotype A2 of

Forster et al. (1996). Indeed, all haplotype D individuals lacked the T→C transition and, therefore, are haplotype D1.

TABLE 3. Frequency distribution of founding haplogroups in aboriginal populations of Chile and Argentina¹

Populations	Haplogroups						References
	N	A	B	C	D	Others	
Aymara	172	0.06	0.68	0.12	0.14		Merriwether et al., 1995
Atacameño	50	0.12	0.72	0.10	0.06		Merriwether et al., 1995
Whichi	72	0.06	0.63	0.03	0.26	0.03	Bravi et al., 1995
Chorote	20	0.15	0.40	0.30	0.15		Bravi et al., 1995
Mapuche 1	58	0.05	0.31	0.21	0.30	0.14	Ginther et al., 1993
Mapuche 2	111		0.07	0.44	0.49		This study
Pehuenche 1	100	0.02	0.09	0.37	0.52		Merriwether et al., 1995
Pehuenche 2	105	0.03	0.10	0.41	0.46		This study
Huilliche	80	0.04	0.28	0.19	0.49		Merriwether et al., 1995
Tehuelche	29		0.21	0.24	0.55		Bravi (pers. Comm.)
Fuegian	45			0.42	0.56	0.02	Lalueza et al., 1997
Yaghan	21			0.48	0.52		This study

¹ Populations are listed from top to bottom in order of distribution along Chile and Argentina, from northernmost to southernmost.

Comparison of the two hypervariable D-loop regions reveals a high variability of haplogroup D, in comparison to the other three haplogroups, and especially with haplogroup C, including a similar number of individuals, as shown in Table 2. This variability is expressed by 19 different lineages defined by 30 polymorphic sites. We were unable to find any nucleotide change shared by 100% of D individuals, as described previously by Horai et al. (1993) in a similar sample. Nevertheless, our results indicate a C→T 16187 change which is present in the majority of D individuals, and is absent in haplogroups A–C (Table 2), concordant with lineages found by Ginther et al. (1993) in an Argentinean Mapuche sample, as well as Chilean lineages described by Horai et al. (1993). However, this change seems to be exclusive to Chilean and Argentinean southern aborigines, since it is not present in other populations from South America.

As shown in Figure 3, it is noteworthy that 26/26 individuals from this study belonging to haplogroup C, two lineages from Chilean aborigines described by Horai et al. (1993), and 7/8 Argentinean aborigine individuals (Ginther et al. 1993), are grouped in one cluster. This extreme low diversity within haplogroup C for southern Chilean and Argentinean groups could also be explained by a founder effect.

Comparing our findings with other Amerindian mtDNA sequences from South America, we found that the retrieved sequences of the three Chilean groups cluster into the four basic mtDNA haplogroups, together with the Amazonian tribes and the Argen-

tinean Mapuche (Fig. 3). These findings are noteworthy in relation to unconventional speculations concerning the origin of the aborigines of Tierra del Fuego. More than half a century ago, it was proposed that people of Tierra del Fuego had direct ancestral links to Australoid populations (Frenguelli, 1963; Imbelloni, 1938). This hypothesis was based on the unusual morphological characteristics, particularly the cranial robustness (supposedly a sign of antiquity), of the aborigines of Patagonia and Tierra del Fuego, that is to say the Ona (Selk'nam), Yaghan (Yamana), and Alacaluf (Kaweskar).

In the interim, a consensus among anthropologists was reached in the sense that Amerinds were derived from Mongoloids who, moving from Asia, crossed the Bering Land Bridge as suggested by Hrdlicka (Frenguelli, 1963). There are, however, still anthropologists with different views. Recently, on the basis of the statistical manipulation of craniometric means, Neves and Pucciarelli (1991) suggested the existence of a pre-Mongoloid migration to South America, coming close to reviving the old ideas of Imbelloni (1938) in a more current context.

In contrast, our results indicate that the Yaghan do not exhibit divergent D-Loop sequences, clustering together with the corresponding haplogroup sequences from other South Amerindian populations (Fig. 3), and presenting similar haplogroup frequencies compared to the Mapuche and Pehuenche. In other words, our data do not support the hypothesis that the Yaghan descend from a different Paleoindian migration lacking haplogroups A and B, as suggested by Lalueza




Fig. 3. Twenty-seven distinct Chilean haplogroups, as well as 105 previously published Amerindian lineages from 9 different South American populations, were analyzed using the neighbor-joining method with a 254-bp fragment, including only the hypervariable region I. An African sequence was used as an outgroup in order to root the tree. Letters on the side indicate respective RFLP haplogroups. PE, Pehuenche; HU, Ma-

puche (Huapi Island); YA, Yaghan; HOCH, Chilean aborigine; HOCO, Colombian aborigine; HOBR, Brazilian aborigine (Horai et al., 1993); WBR, Brazilian aborigine (Handt et al., 1998); GMA, Argentinean Mapuche (Ginther et al., 1993); SBR, Brazilian aborigine (Santos et al., 1996); TARG, Argentinean aborigine; TBR, Brazilian aborigine (Torroni et al., 1993); EYA, Yanomama (Easton et al., 1996).

eza et al. (1997). Our results are rather in agreement with the view that the Yaghan are related to tribes from south-central Chile and Argentina such as the Huilliche, Pehuenche, and Tehuelche, as previously suggested by Rothhammer et al. (1986) and Llop (1996). Furthermore, we note that the haplogroup distribution of the Yaghan is similar to the distribution obtained by Lalueza Fox (1996) for extinct groups from Tierra del Fuego-Patagonia. All samples from these geographic areas lack haplogroups A and B and are therefore most probably descendants of one common ancestral Paleoindian population. In this context, it is interesting to note that Bennett and Bird (1964) reported a complete archeological sequence linking the first Paleoindian migrants to contemporary Ona (Shelknam) populations.

LITERATURE CITED

- Anderson S, Bankier AT, Barrell BG, de Bruijn MHL, Coulson AR, Drouin J, Eperon IC, Nierlich DP, Roe BA, Sanger F, Schreier PH, Smith AJH, Staden R, Young IG. 1981. Sequence and organization of the human mitochondrial genome. *Nature* 290:457-465.
- Bailliet G, Rothhammer F, Carnese F, Bravi C, Bianchi NO. 1994. Founder mitochondrial haplogroups in Amerindian populations. *Am J Hum Genet* 55:27-33.
- Ballinger SW, Schurr TG, Torroni A, Gan YY, Hodge JA, Hassan K, Chen KH, Wallace DC. 1992. Southern Asian mitochondrial DNA analysis reveals genetic continuity of ancient Mongoloid migrations. *Genetics* 130:139-152.
- Bernnett WC, Bird JB. 1964. *Andean culture history*. New York: Natural History Press. Doubleday & Co., Inc.
- Chen Y-S, Torroni A, Excoffier L, Santachiara-Benerecetti AS, Wallace DC. 1995. Analysis of mtDNA variation in African populations reveals the most ancient of all human continent-specific haplogroups. *Am J Hum Genet* 57:133-149.
- Easton R, Merriwether, DA, Crews D, Ferrell R. 1996. mtDNA variation in the Yanomami: evidence for additional New World founding lineages. *Am J Hum Genet* 59:213-225.
- Forster P, Harding R, Torroni A, Bandelt H-J. 1996. Origin and evolution of Native American mtDNA variation: a reappraisal. *Am J Hum Genet* 59:935-945.
- Frenguelli J. 1963. The present status of the theories concerning primitive man in Argentina. In: Steward JH, editor. *Handbook of South American Indians*, Vol 6. New York: Cooper Square Publisher, Inc. p 11-17.
- Ginther C, Corach D, Penacino GA, Rey JA, Carnese FR, Hutz MH, Anderson A, Just J, Salzano FM, King MC. 1993. Genetic variation among the Mapuche Indians from the Patagonian region of Argentina: mitochondrial DNA sequence variation and allele frequencies of several nuclear genes. *EXS* 67:211-219.
- Gustincich S, Manfoletti G, Del Sal G, Schneider C, Carninci P. 1991. A fast method for high quality genomic DNA extraction from whole blood. *Biotechniques* 11:298-302.
- Handt O, Meyer S, von Haeseler A. 1998. Compilation of human mtDNA control region sequences. *Nucleic Acids Res* 26:126-129.
- Horai S, Kondo R, Nakagawa-Hattori Y, Hayashi S, Sonoda S, Tajima K. 1993. Peopling of the Americas, founded by four major lineages of mitochondrial DNA. *Mol Biol Evol* 10:23-47.
- Imbelloni G. 1938. Tabla clasificatoria de los Indios: regiones biologicas y grupos raciales en America. *Physis* 12:229-249.
- Lalueza C, Perez-Perez A, Prats E, Cornudella L, Turbon D. 1997. Lack of founding Amerindian mitochondrial DNA lineages in extinct aborigines from Tierra del Fuego-Patagonia. *Hum Mol Genet* 6:41-46.
- Lalueza Fox C. 1996. Mitochondrial DNA haplogroups in four tribes from Tierra del Fuego-Patagonia: inferences about the peopling of the Americas. *Hum Biol* 68:853-871.
- Llop E. 1996. Genetic composition of Chilean aboriginal populations: HLA and other genetic marker variation. *Am J Phys Anthropol* 101:325-332.
- Llop E, Harb Z, Acuña M, Moreno R, Barton S, Aspillaga E, Rothhammer F. 1993. Composición genética de la población chilena: los Pehuenches de Trapa-Trapa. *Rev Med Chil* 121:494-498.
- Llop E, Harb Z, Acuña M, Moreno R, Aspillaga E, Van de Maele M, Rothhammer F. 1994. Composición genética de la población chilena: los Yamanas de Ukika. *Rev Med Chil* 122:979-985.
- Merriwether DA, Rothhammer F, Ferrell RE. 1995. Distribution of the four founding lineage haplotypes in Native Americans suggests a single wave of migration for the New World. *Am J Phys Anthropol* 98:411-430.
- Nei M. 1978. Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics* 89:583-590.
- Neves W, Pucciarelli H. 1991. Morphological affinities of the first Amerindian: an exploratory analysis based on early South American human remains. *J Hum Evol* 21:261-273.
- Rothhammer F, Silva C, Cocilovo J, Quevedo S. 1986. Una hipótesis sobre el poblamiento de Chile basada en el análisis multivariado de medidas craneométricas. *Rev Chung* 16-17:115-118.
- Rothhammer F, Silva C. 1992. Gene geography of South America: testing models of population displacement based on archeological evidence. *Am J Phys Anthropol* 89:441-446.
- Rothhammer F, Silva C, Callegari-Jacques SM, Llop E, Salzano FM. 1997. Gradients of HLA diversity in South American Indians. *Ann Hum Biol* 24:197-208.
- Saitou N, Nei M. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406-425.
- Santos M, Ward RH, Barrantes R. 1994. mtDNA variation in the Chibcha Amerindian Huetar from Costa Rica. *Hum Biol* 66:963-977.
- Santos SE, Ribeiro-Dos-Santos AK, Meyer D, Zago MA. 1996. Multiple founder haplotypes of mitochondrial DNA in Amerindians revealed by RFLP and sequencing. *Ann Hum Genet* 60:305-319.
- Schurr TG, Ballinger SW, Gan YY, Hodge JA, Merriwether DA, Lawrence DN, Knowler WC. 1990. Amerindian mitochondrial DNAs have rare Asian mutations at high frequencies suggesting they derived from four primary maternal lineages. *Am J Hum Genet* 46:613-623.

- Swofford DL. 1993. Phylogenetic analysis using parsimony (PAUP), version 3.1. Champaign: Illinois Natural History Survey.
- Swofford DL, Olsen GJ. 1990. Phylogeny reconstruction. In: Hillis DM, Moritz C, editors. *Molecular systematics*. Sunderland, MA: Sinauer. p 411–500.
- Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice. *Nucleic Acids Res* 22:4673–4680.
- Torrioni A, Schurr TG, Yang CC, Szathmary EJE, Williams RC, Schanfield MS, Troup GA, Knowler WC, Lawrence DN, Weiss KM, Wallace DC. 1992. Native American mitochondrial DNA analysis indicated that the Amerind and the Nadene populations were founded by two independent migrations. *Genetics* 130:153–162.
- Torrioni A, Schurr TG, Cabell MF, Brown MD, Neel JV, Larsen M, Smith DG, Vullo CM, Wallace DC. 1993. Asian affinities and continental radiation of the four founding Native American mtDNAs. *Am J Hum Genet* 53:563–590.
- Torrioni A, Huoponen K, Francalacci P, Petrozzi M, Morelli L, Scozzari R, Obinu D. 1996. Classification of European mtDNAs from an analysis of three European populations. *Genetics* 144:1835–1850.