

The enigma of crustal zircons in upper-mantle rocks: Clues from the Tumut ophiolite, southeast Australia

Elena A. Belousova¹, José María González Jiménez^{1,2}, Ian Graham³, William L. Griffin¹, Suzanne Y. O'Reilly¹, Norman Pearson¹, Laure Martin⁴, Stephen Craven¹, and Cristina Talavera⁵

¹ARC Centre of Excellence for Core to Crust Fluid Systems (CCFS) and GEMOC, Macquarie University, Sydney, NSW 2109, Australia

²Department of Geology and Andean Geothermal Center of Excellence, Universidad de Chile, Plaza Ercilla # 803, 8370450 Santiago de Chile, Chile

³School of Biological, Earth and Environmental Sciences, University of New South Wales, Sydney, NSW 2052, Australia

⁴Centre for Microscopy Characterisation and Analysis, The University of Western Australia, Crawley, WA 6009, Australia

⁵Department of Imaging and Applied Physics, Curtin University, Perth, WA 6102, Australia

ABSTRACT

We suggest a new explanation for the presence of crustally derived zircons in the upper-mantle rocks of ophiolitic complexes, as an alternative to subduction-related models. Integrated isotopic (U-Pb, Hf, and O isotopes) and trace-element data for zircons from the Tumut ophiolitic complex (southeast Australia) indicate that these grains are related to granitic magmatism and were introduced into the mantle rocks after their emplacement into the crust. These observations emphasize that a clear understanding of the origin of individual zircon populations and their relationship to the host rock is essential to interpretations of the tectonic history of upper-mantle rocks and the dynamics of crust-mantle interactions.

INTRODUCTION

Several studies have reported U-Pb ages on zircons recovered from upper-mantle rocks in ophiolitic complexes (e.g., Grieco et al., 2001; Saveleva et al., 2007; Stern et al., 2010; Badanina et al., 2013; Yamamoto et al., 2013; González-Jiménez et al., 2013). Did these zircons crystallize in the mantle from metasomatic fluids (e.g., Grieco et al., 2001; Zaccarini et al., 2004; Zheng et al., 2006), or are they xenocrysts of crustal material recycled during subduction (e.g., Yamamoto et al., 2013; Robinson et al., 2014)? Crucially, what is their significance in the framework of ophiolite evolution?

We have studied the geochemical and isotopic signatures of zircons recovered from mantle and crustal rocks of the Coolac serpentinite belt, part of the Tumut ophiolite complex in the Lachlan fold belt of southeastern Australia (Fig. 1). The integrated analysis of U-Pb, Lu-Hf, and O isotopes has provided critical information about the origin of zircons in both mantle and crustal rocks. The zircon data are complemented by field observations and in situ Re-Os isotopic information on laurite grains in the chromitites of the mantle section of this ophiolite.

This case study emphasizes that deciphering the origin of zircons requires integration of data sets beyond zircon U-Pb geochronology alone. It reveals that crustally derived zircons can be introduced into mantle rocks after their obduction via microscopic melt networks, and sends a cautionary message about the interpretation of zircon ages from ultramafic rocks in general.

GEOLOGICAL SETTING AND SAMPLE BACKGROUND

The Coolac serpentinite belt, Wambidgee serpentinite belt, Tumut Ponds serpentinite belt, and smaller bodies form the ~250-km-long Tumut ophiolitic complex in the Lachlan fold belt of southeastern Australia (Fig. 1; Graham et al., 1996a, 1996b; Spaggiari et al., 2003). The Coolac serpentinite belt is ≥63 km long, up to 3.5 km wide, and ≥5 km thick. It was metamorphosed mostly to greenschist facies and consists of variably serpentinized porphyroclastic meta-harzburgites with lenses of dunite that host chromitite bodies. The ultramafic rocks are cut by dikes of gabbro and plagiogranite, and dike-like bodies, veinlets, and pod-like masses of rodingite. On its eastern contact, it is either faulted against or intruded by the Silurian S-type Young Granodiorite; to the west, it is faulted against




Figure 1. Location of Tumut region in Lachlan fold belt (southeastern Australia; after Graham et al., 1996b). Solid black strip separating Tumut Trough and Young Granodiorite zones is Coolac serpentinite belt. ML—Mount Lightning sample site.

metavolcanic and metasedimentary units of the Silurian Tumut trough (Stuart-Smith, 1990; Graham et al., 1996b).

The age of formation of the ultramafic rocks from the Tumut region is unclear. Previous suggestions range from Cambrian-Ordovician (Stuart-Smith, 1990) to Devonian (Graham et al., 1996b). Graham et al. (1996b) suggested that a Devonian U-Pb age (ca. 400 Ma), obtained on zircons from plagiogranite dikes, indicated that the ophiolitic rocks and granitic magmas were generated during the same major tectonothermal event in the Lachlan fold belt (430–390 Ma; Chappell, 1994). However, this age is considerably younger than those of the Young Granodiorite (zircon U-Pb age of 428 ± 1.9 Ma; Lyons and Percival, 2002) and the North Mooney Complex (horn-

blende K-Ar age 425 ± 6 Ma; Stuart-Smith, 1990), leaving the relationships of the ultramafic rocks to the rest of the Lachlan fold belt ambiguous.

We separated zircons from samples of two (high-Al, high-Cr) massive chromitites as well as from samples of leucogabbro, plagiogranite, and rodingite. We also sampled sediments in gullies draining outcrops of weakly serpentinized massive porphyroclastic harzburgite. Zircons from the Young Granodiorite sampled at the contact were analyzed to refine the timing of granitic magmatism and its relationship to the ultramafic-mafic rocks.

Several grains of laurite (RuS_2 ; up to 10–20 μm) were identified in polished thin sections of massive (high-Cr) chromitite from the Quilter's open-cut mine in the Coolac serpentinite belt. Their predominantly euhedral shapes and occurrence within fresh chromite grains (Fig. DR1 in the GSA Data Repository¹) suggest a primary magmatic origin. Laurite grains were identified by their X-ray spectra, and their Os isotope compositions were analyzed in situ.

Detailed information on sample locations, sample processing, and analytical techniques are provided in Appendix DR1 and Tables DR1–DR5 in the Data Repository.

RESULTS

Analytical data are given in Tables DR2–DR5. Table 1 provides a summary of the information collected on zircons.

Zircon U-Pb Age Dating

The most important observation from the U-Pb data is that the main zircon populations from the Coolac belt rocks (430.5 ± 4.5 Ma for plagiogranite, 429.4 ± 4.7 Ma for leucogabbro; Figs. DR2 and DR3) are indistinguishable from the age defined for the Young Granodiorite during this study (428 ± 3 Ma; Fig. DR4) and previously (428 ± 1.9 Ma; Lyons and Percival, 2002). Detrital zircons from the stream samples are also dominated by Early Silurian age populations (432.9 ± 3.1 and 426.6 ± 6.3 Ma; Figs. DR5–DR7), indistinguishable from the ages defined for the Coolac and Young Granodiorite samples. The inherited populations in both Coolac and Young Granodiorite samples are also similar: ages range from Ordovician to Paleoproterozoic and Archean with clusters around 442–470 Ma, 1.0–1.3 Ga, 1.7–1.8 Ga, and 2.2–2.4 Ga (Table 1; Figs. 2 and 3; Fig. DR8). Moreover, the youngest (ca. 370–385 Ma) populations of zircons from the Coolac samples have ages identical to the rims on zircons from the Young Granodiorite.

Sixteen (16) U-Pb analyses on 12 zircon grains recovered from the chromitite samples also show ages ranging from Ordovician (442 ± 5 Ma, 1σ) to Paleoproterozoic (2289 ± 25 Ma), with a small cluster ($n =$

3) around 595 Ma (Figs. 2 and 3; Fig. DR9); grain morphology and the scatter in age support their xenocrystic origin. Zircons from the rodingite show ages from Devonian (376 ± 9 Ma) to Paleoproterozoic (1718 ± 22 Ma) with a population at 390 ± 5 Ma (Figs. 2 and 3; Fig. DR10); this younger age is indistinguishable from the age of the younger overprint observed in zircons from the other Coolac samples and the Young Granodiorite (Table 1).

Zircon O- and Hf-Isotope Data

Most of the Coolac belt zircons show $\delta^{18}\text{O}$ ranging from 6.5‰ to almost 10‰ (Fig. 2; Table 1), clearly outside the typical mantle range ($5.3\text{‰} \pm 0.3\text{‰}$; Valley, 2003) and much higher than the range defined by zircons from plagiogranites of other ophiolitic complexes (3.9‰–5.6‰; Grimes et al., 2013). By and large, the Coolac zircons are dominated by crustal O isotope compositions, typical of zircons from granitoids of the Lachlan fold belt (Kemp et al., 2009). They overlap the main magmatic population from the Young Granodiorite (7.7‰–10‰; Fig. 2). A more restricted range of $\delta^{18}\text{O}$ (7.5‰–8.4‰; Fig. 2B) is observed in the younger population of zircons from the rodingite sample; this coincides with values in the rims of the granodiorite zircons (8.0‰ and 8.2‰).

The crustal O-isotope signatures of the studied zircons are consistent with their non-radiogenic Hf isotope compositions; most grains show negative ϵ_{Hf} (Fig. 3; Fig. DR11; Table 1), mainly from 0 to –16 for the Coolac samples and from –2 to –9.5 for the Young Granodiorite. The mean Hf crustal model ages (T_{DM}^{C}) of the main magmatic populations range from 1.6 to 1.7 Ga and from 1.5 to 1.8 Ga for the Coolac rocks and granodiorite, respectively. However, in grains from the plagiogranite sample, crustal $\delta^{18}\text{O}$ values contrast with a distinctive mantle Hf isotope signature (ϵ_{Hf} ranging from +9 to +13, and mean T_{DM}^{C} of 0.76 Ga).

Several (mainly inherited) grains from the Coolac samples plot within or close to the mantle field in Figure 2. However, a few zircons from the main Early Silurian population, including one from the plagiogranite, and two detrital zircons, have both juvenile ϵ_{Hf} and mantle $\delta^{18}\text{O}$ values. These two detrital zircons (grains 24 and 41; Tables DR2–DR4) have ϵ_{Hf} of +15.5 and +14.8 and are dated at 439 ± 10 Ma and 442 ± 12 Ma, respectively.

Zircon Trace-Element Composition

Zircons from all the Coolac belt samples have trace-element compositions typical of continental-crust zircons (Belousova et al., 2002); none plot within the U/Yb–Y field that is unique to oceanic-crust zircons (Fig. DR12; Grimes et al., 2007). Both magmatic and inherited populations show nega-

TABLE 1. SUMMARY OF THE U-Pb AGE AND Hf- AND O-ISOTOPE INFORMATION FOR ZIRCONS FROM THE COOLAC SERPENTINITE BELT, SOUTHEAST AUSTRALIA

Sample	Rock type	Main population/ inferred crystallization age (Ma) (no. of analyses)	Younger overprint (Ma)	Inheritance (Ma)	Hf model ages* T_{DM}^{C} & T_{DM}^{D} (Ga)	Hf isotope signature*	O isotope signature*
M104-M13	Chromitite (high-Cr)	–	–	442–2289	T_{RD}^{**} 1.18–1.64	Mainly crustal	Mainly crustal
ML101A	Chromitite (high-Al)	–	–	470–1953	–	Mainly crustal	Crustal
RH106	Rodingite	390 ± 5 (?) (1)	376–390 (?)	424(?)–1718	1.24 & 1.59 (?)	Crustal	Crustal
#238	Leucogabbro	429.4 ± 4.7 (15)	361–416	447–2408 (3412?)	1.26 & 1.63	Mainly crustal	Mainly crustal
#20	Plagiogranite	430.5 ± 4.5 (11) / 439 ± 10 (1)	375–420	458 \pm 6	0.66 & 0.76	Strongly juvenile	Mantle to crustal
TC12-ML01	Stream sample	432.9 ± 3.1 (15)	385 \pm 6	456–1877	1.30 & 1.68	Mainly crustal	Mainly crustal
TC12-BG01	Stream sample	426.6 ± 6.3 (4)	388 \pm 5	471–2257	1.27 & 1.66	Crustal	Crustal
Young Granodiorite							
YG202	Granodiorite	428 ± 3 (11)	367–385	450–3053	1.25 & 1.62	Crustal	Crustal

Note: T_{DM}^{D} —depleted mantle model age; T_{DM}^{C} —mean Hf crustal model age.
*For main magmatic population (excluding inherited grains) or youngest populations.
** T_{RD}^{D} (Re-depletion model age); Re-Os model age from laurite.

¹GSA Data Repository item 2015049, Appendix DR1, Tables DR1–DR5, and Figures DR1–DR17, is available online at www.geosociety.org/pubs/ft2015.htm, or on request from editing@geosociety.org or Documents Secretary, GSA, P.O. Box 9140, Boulder, CO 80301, USA.




Figure 2. $\delta^{18}\text{O}$ versus U-Pb age for zircons. $\delta^{18}\text{O}$ values are quoted in per mil (‰) with respect to Vienna Standard Mean Ocean Water (VSMOW). **A:** From Coolac serpentinite belt and Young Granodiorite. **B:** Populations younger than 1200 Ma. **C:** Plagiogranite sample (main population). Dark- and light-band deviations on the mantle value (5.3‰; Valley, 2003) show 1σ and 2σ . Typical uncertainty of individual $\delta^{18}\text{O}$ analyses is $<0.3\%$ (2 standard deviations).




Figure 3. ϵ_{Hf} versus U-Pb age in zircons from Coolac serpentinite belt and Young Granodiorite, southeastern Australia (sample symbols as in Fig. 2). DM—depleted mantle; CHUR—chondritic uniform reservoir.

tive Eu anomalies, suggesting crystallization with plagioclase. Many grains show different degrees of light rare earth element (LREE) enrichment, typical of altered or metamict zircons. The range in trace-element compositions defined by the zircons of the Coolac samples overlaps that of zircons from the Young Granodiorite (Figs. DR13 and DR14; Table DR4).

In contrast, the trace-element signatures and REE distribution patterns of zircons from the plagiogranite are clearly distinct from those of the rest of the zircons of the Coolac belt and the Young Granodiorite: they have lower Hf contents (<1 wt%, common in zircons from mafic rocks), and lower Th/U. Apart from a single grain, they lack Eu anomalies (Fig. DR14A), suggesting that they crystallized before plagioclase. The only feature in common with the rest of the samples is variable LREE enrichment. The grain (15) that appears to be least affected by alteration (no LREE enrichment) is the only one with a mantle O-isotope composition ($\delta^{18}\text{O} = 5.61\%$) and that also yielded a slightly older U-Pb age (439 ± 10 Ma) than the other zircons in the main population from this plagiogranite. Furthermore, $\delta^{18}\text{O}$ in the plagiogranite sample increases with decreasing U-Pb age (Fig. 2C), suggesting that zircons from this sample have experienced progressive alteration by ^{18}O -enriched supracrustal fluids soon after their crystallization. Thus, the lone zircon “survivor”, with no LREE enrichment and a mantle O-isotope composition, suggests that the plagiogranite is at least 439 Ma old. It is important that the two detrital grains (grains 24 and 41) with mantle isotopic signatures also have “undisturbed” REE patterns (Fig. DR14A).

Laurite Re-Os Data

The Os-isotope data for 10 laurite grains from the chromitites (Table DR5) show a range of unradiogenic $^{187}\text{Os}/^{188}\text{Os}$ (0.1146–0.1220) and low $^{187}\text{Re}/^{188}\text{Os}$ ratios. Their T_{RD} (Re-depletion) model ages (relative to Enstatite Chondrite Reservoir; Walker et al., 2002) vary between 0.84 Ga and 1.89 Ga, with a main peak at around 1.18 Ga and a shoulder at ca. 1.64 Ga (Fig. DR15). The older T_{RD} model ages overlap the range of zircon Hf crustal model ages (ca 1.62–1.68 Ga), except for the plagiogranite sample (mean T_{DM}^{c} of 0.77 Ga). The youngest “event” ($T_{\text{RD}} = 0.84$ Ga) recorded by the Re-Os system is very close to the age of the protolith reflected in the Lu-Hf systematics of the plagiogranite zircons.

DISCUSSION

Origin of Crustal Zircons in Upper-Mantle Rocks of the Coolac Belt

The combined isotopic and trace-element data indicate that most zircons are of crustal origin. Previous studies of crustal zircons in upper-mantle rocks (including chromitites) have interpreted such grains as remnants of crustal rocks subducted into the mantle or as material picked up by ascending melts and/or fluids released during dehydration of subducting slabs containing crustal sedimentary rocks (e.g., Stern et al., 2010; Yamamoto et al., 2013; Robinson et al., 2014; Zhou et al., 2014). However, in this case the striking resemblance to zircon populations from the adjacent Young Granodiorite suggests that the grains recovered from the Coolac belt rocks are related to the granitic magmatism. Indeed, the isotopic and trace-element signatures recorded in the Coolac zircons are consistent with the timing and the nature of granitic magmatism within the larger Lachlan fold belt (420–390 Ma; Chappell, 1994). This similarity includes not only the grains of the main magmatic populations, but the inherited populations and the younger overprint (Table 1); this could hardly be a coincidence.

The presence of similar populations of zircons in ultramafic rocks, chromitites, and granitic rocks suggests that late melts from the granite batholith penetrated the ultramafic rocks. Granodioritic and aplitic dikes from the Young Granodiorite intrude the Coolac belt peridotites (Stuart-Smith, 1990; Graham et al., 1996b), and zircon could have been more widely introduced via intergranular networks percolating from the larger fluid channels (Fig. DR16). These fluids precipitated zircons of the latest generations (e.g., 375–390 Ma) and transported grains of the main magmatic and inherited populations. Such intergranular networks could have healed later, leaving little visible record under conventional microscopy.

These crustally derived fluids were most likely responsible for the alteration of primary magmatic zircons in the plagiogranite, where enrich-

ment in LREE and increase in $\delta^{18}\text{O}$ correlates with a progressive resetting of the U-Pb isotope system. The fact that the Hf-isotope system in these zircons retained the original mantle signature emphasizes the robustness of this system.

Implication for the Local and Regional Geology

The Re-Os data from the laurites indicate that the mantle section in the Coolac serpentinite belt is at least Mesoproterozoic in age, with a main tectonomagmatic event recorded at ca 1.2 Ga. The coincidence of Os model ages on laurite and Hf crustal model ages on zircon at 1.64–1.68 Ga implies that Proterozoic lithosphere (or at least ribbons of it) extends as far as the Tumut region of the Lachlan fold belt and supports existing models (Betts et al., 2002, and references therein) allowing for incorporation of an older continental ribbon into the Paleozoic crust during back-arc inversions.

Our observations suggest that the zircon U-Pb age information reported here and in previous studies (e.g., Graham et al., 1996b) cannot be used directly to constrain the timing of magmatic events that have produced the upper-mantle rocks of the Coolac belt. The U-Pb age of the zircon “survivor” from the plagiogranite suggests that the ultramafic rocks were part of an oceanic lithosphere ≥ 439 m.y. ago. This is supported by the ages (439 ± 10 Ma and 442 ± 12 Ma) of two detrital zircons with mantle isotopic compositions, recovered from the Mount Lightning stream sample TC12-ML01 (Fig. 1; Table DR1). Even if we consider a subduction-related model to explain the presence of crustal zircons in upper-mantle rocks, then the youngest xenocrystic grain would provide us with a maximum age estimate for subduction. The youngest zircon from the chromitites that has a supracrustal O-isotope composition ($\delta^{18}\text{O} = 7.9 \pm 0.4$, 2σ) was dated at 442 ± 10 Ma, which is consistent with this timing.

Broader Implications

We have presented a new explanation for the presence of crustal zircons in the upper-mantle sections of ophiolitic complexes, as an alternative to the existing subduction-related models. In this case, the granitoid-related melts and/or fluids, injected into already-emplaced mafic-ultramafic rocks, apparently carried existing zircons (including inherited ones) and possibly crystallized new grains. This study emphasizes that integrated information on zircons (not U-Pb data alone) is critical to the interpretation of events recorded by upper-mantle rocks, including their emplacement and subsequent tectonic history. A clear understanding of the origin of xenocrystic zircons and their relationship with the host rock increases the probability of correct conclusions about the history of upper-mantle rocks (e.g., the upper-mantle sequences of ophiolitic complexes) and interpretations of the dynamics of crust-mantle interaction.

ACKNOWLEDGMENTS

Funding for this work was provided by Australian Research Grants (Belousova, Griffin, and O'Reilly) and Macquarie University, Australia. We are grateful for positive and constructive comments made by four anonymous reviewers that led to clarification and improvement of the manuscript. This is contribution 519 from the ARC Centre of Excellence for Core to Crust Fluid Systems (www.ccfsmq.edu.au) and contribution 967 from the GEMOC ARC National Key Centre (www.gemoc.mq.edu.au).

REFERENCES CITED

- Badanina, I.Y., Malitch, K.N., and Belousova, E.A., 2013, U-Pb and Hf isotope characteristics of zircon from chromitites at Finero [abstract]: *Mineralogical Magazine*, v. 77, p. 639.
- Belousova, E.A., Griffin, W.L., O'Reilly, S.Y., and Fisher, N.I., 2002, Igneous zircon: Trace element composition as an indicator of source rock type: *Contributions to Mineralogy and Petrology*, v. 143, p. 602–622, doi:10.1007/s00410-002-0364-7.
- Betts, P.G., Giles, D., Lister, G.S., and Frick, L.R., 2002, Evolution of the Australian lithosphere: *Australian Journal of Earth Sciences*, v. 49, p. 661–695, doi:10.1046/j.1440-0952.2002.00948.x.
- Chappell, B.W., 1994, Lachlan and New England: Fold belts of contrasting magmatic and tectonic development: *Royal Society of New South Wales Journal and Proceedings*, v. 127, p. 47–59.

- González-Jiménez, J.M., Locmelis, M., Belousova, E., Griffin, W.L., Gervilla, F., Kerestedjian, T.N., O'Reilly, S.Y., Pearson, N.J., and Sergeeva, I., 2013, Genesis and tectonic implications of podiform chromitites in the metamorphosed ultramafic massif of Dobromiritsi (Bulgaria): *Gondwana Research*, doi:10.1016/j.gr.2013.09.020 (in press).
- Graham, I.T., Franklin, B.J., and Marshall, B., 1996a, Chemistry and mineralogy of podiform chromitite deposits, southern NSW, Australia: A guide to their origin and evolution: *Mineralogy and Petrology*, v. 57, p. 129–150, doi:10.1007/BF01162355.
- Graham, I.T., Franklin, B.J., Marshall, B., Leitch, E.C., and Fanning, M., 1996b, Tectonic significance of 400 Ma zircon ages for ophiolitic rocks from the Lachlan fold belt, eastern Australia: *Geology*, v. 24, p. 1111–1114, doi:10.1130/0091-7613(1996)024<1111:TSMZMA>2.3.CO;2.
- Grieco, G., Ferrario, A., Von Quadt, A., Koepfel, V., and Mathez, E.A., 2001, The zircon-bearing chromitites of the phlogopite peridotite of Finero (Ivrea zone, Southern Alps): Evidence and geochronology of a metasomatized mantle slab: *Journal of Petrology*, v. 42, p. 89–101, doi:10.1093/petrology/42.1.89.
- Grimes, C.B., John, B.E., Kelemen, P.B., Mazdab, F.K., Wooden, J.L., Cheadle, M.J., Hanghoj, K., and Schwartz, J.J., 2007, Trace element chemistry of zircons from oceanic crust: A method for distinguishing detrital zircon provenance: *Geology*, v. 35, p. 643–646, doi:10.1130/G23603A.1.
- Grimes, C.B., Ushikubo, T., Kozdon, R., and Valley, J.W., 2013, Perspectives on the origin of plagiogranite in ophiolites from oxygen isotopes in zircon: *Lithos*, v. 179, p. 48–66, doi:10.1016/j.lithos.2013.07.026.
- Kemp, A.I.S., Hawkesworth, C.J., Collins, W.J., Gray, C.M., Blevin, P.L., and EIMF, 2009, Isotopic evidence for rapid continental growth in an extensional accretionary orogen: The Tasmanides, eastern Australia: *Earth and Planetary Science Letters*, v. 284, p. 455–466, doi:10.1016/j.epsl.2009.05.011.
- Lyons, P., and Percival, I.G., 2002, Middle to Late Ordovician age for the Jindalee Group of the Lachlan Fold Belt, New South Wales: Conodont evidence and some tectonic problems: *Australian Journal of Earth Sciences*, v. 49, p. 801–808, doi:10.1046/j.1440-0952.2002.00951.
- Robinson, P.T., Trumbull, R.B., Schmitt, A., Yang, J.-S., Li, J.-W., Zhou, M.-F., Erzinger, J., Dare, S., and Xiong, F., 2014, The origin and significance of crustal minerals in ophiolitic chromitites and peridotites: *Gondwana Research*, doi:10.1016/j.gr.2014.06.003 (in press).
- Savelieva, G.N., Suslov, P.V., and Larionov, A.N., 2007, Vendian tectono-magmatic events in mantle ophiolitic complexes of the Polar Urals: U-Pb dating of zircon from chromitite: *Geotectonics*, v. 41, p. 105–113, doi:10.1134/S0016852107020021.
- Spaggiari, C.V., Gray, D.R., and Foster, D.A., 2003, Tethyan- and Cordilleran-type ophiolites of eastern Australia: Implications for the evolution of the Tasmanides: *Ophiolites in Earth History*, v. 218, p. 517–539.
- Stern, R.J., Ali, K.A., Liegeois, J.P., Johnson, P.R., Kozdrojs, W., and Kattan, F.H., 2010, Distribution and significance of pre-Neoproterozoic zircons in juvenile Neoproterozoic igneous rocks of the Arabian–Nubian shield: *American Journal of Science*, v. 310, p. 791–811, doi:10.2475/09.2010.02.
- Stuart-Smith, P.G., 1990, The emplacement and fault history of the Coolac Serpentinite, Lachlan Fold Belt, southeastern Australia: *Journal of Structural Geology*, v. 12, p. 621–638, doi:10.1016/0191-8141(90)90078-D.
- Valley, J.W., 2003, Oxygen isotopes in zircon, in Hanchar, J.M., and Hoskin, P.W.O., eds., *Zircon: Reviews in Mineralogy and Geochemistry*, v. 53, p. 343–385.
- Walker, R.J., Horan, M.F., Morgan, J.W., Becker, H., Grossman, J.N., and Rubin, A.E., 2002, Comparative ^{187}Re – ^{187}Os systematics of chondrites: Implications regarding early solar system processes: *Geochimica et Cosmochimica Acta*, v. 66, p. 4187–4201, doi:10.1016/S0016-7037(02)01003-7.
- Yamamoto, S., et al., 2013, Recycled crustal zircons from podiform chromitites in the Luobusa ophiolite, southern Tibet: *The Island Arc*, v. 22, p. 89–103, doi:10.1111/iar.12011.
- Zaccarini, F., Pushkarev, E.V., Fershtater, G.B., and Garuti, G., 2004, Composition and mineralogy of PGE-rich chromitites in the Nurali lherzolite–gabbro complex: *Canadian Mineralogist*, v. 42, p. 545–562, doi:10.2113/gscanmin.42.2.545.
- Zheng, J.P., Griffin, W.L., O'Reilly, S.Y., Yang, J.S., and Zhang, R.Y., 2006, A refractory mantle protolith in younger continental crust, east-central China: Age and composition of zircon in the Sulu ultrahigh-pressure peridotite: *Geology*, v. 34, p. 705–708, doi:10.1130/G22569.1.
- Zhou, M.-F., Robinson, P.T., Ben-Xun, S., Gao, J.-F., Li, J.-W., Yang, J.-S., and Malpas, J., 2014, Compositions of chromite, associated minerals, and parental magmas of podiform chromite deposits: The role of slab contamination of asthenospheric melts in suprasubduction zone environments: *Gondwana Research*, v. 26, p. 262–283, doi:10.1016/j.gr.2013.12.011.

Manuscript received 31 August 2014

Revised manuscript received 17 November 2014

Manuscript accepted 21 November 2014

Printed in USA

Geology

The enigma of crustal zircons in upper-mantle rocks: Clues from the Tumut ophiolite, southeast Australia

Elena A. Belousova, José María González Jiménez, Ian Graham, William L. Griffin, Suzanne Y. O'Reilly, Norman Pearson, Laure Martin, Stephen Craven and Cristina Talavera

Geology 2015;43;119-122
doi: 10.1130/G36231.1

Email alerting services click www.gsapubs.org/cgi/alerts to receive free e-mail alerts when new articles cite this article

Subscribe click www.gsapubs.org/subscriptions/ to subscribe to *Geology*

Permission request click <http://www.geosociety.org/pubs/copyrt.htm#gsa> to contact GSA

Copyright not claimed on content prepared wholly by U.S. government employees within scope of their employment. Individual scientists are hereby granted permission, without fees or further requests to GSA, to use a single figure, a single table, and/or a brief paragraph of text in subsequent works and to make unlimited copies of items in GSA's journals for noncommercial use in classrooms to further education and science. This file may not be posted to any Web site, but authors may post the abstracts only of their articles on their own or their organization's Web site providing the posting includes a reference to the article's full citation. GSA provides this and other forums for the presentation of diverse opinions and positions by scientists worldwide, regardless of their race, citizenship, gender, religion, or political viewpoint. Opinions presented in this publication do not reflect official positions of the Society.

Notes