



## Mathematics of Operations Research

Publication details, including instructions for authors and subscription information:  
<http://pubsonline.informs.org>

### Cut-Generating Functions and S-Free Sets

Michele Conforti, Gérard Cornuéjols, Aris Daniilidis, Claude Lemaréchal, Jérôme Malick

To cite this article:

Michele Conforti, Gérard Cornuéjols, Aris Daniilidis, Claude Lemaréchal, Jérôme Malick (2015) Cut-Generating Functions and S-Free Sets. *Mathematics of Operations Research* 40(2):276-391. <http://dx.doi.org/10.1287/moor.2014.0670>

Full terms and conditions of use: <http://pubsonline.informs.org/page/terms-and-conditions>

This article may be used only for the purposes of research, teaching, and/or private study. Commercial use or systematic downloading (by robots or other automatic processes) is prohibited without explicit Publisher approval, unless otherwise noted. For more information, contact [permissions@informs.org](mailto:permissions@informs.org).

The Publisher does not warrant or guarantee the article's accuracy, completeness, merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications, or inclusion of an advertisement in this article, neither constitutes nor implies a guarantee, endorsement, or support of claims made of that product, publication, or service.

Copyright © 2015, INFORMS

Please scroll down for article—it is on subsequent pages



INFORMS is the largest professional society in the world for professionals in the fields of operations research, management science, and analytics.

For more information on INFORMS, its publications, membership, or meetings visit <http://www.informs.org>

# Cut-Generating Functions and $S$ -Free Sets

Michele Conforti

Department of Mathematics, University of Padova, 35121 Padova, Italy, [conforti@math.unipd.it](mailto:conforti@math.unipd.it)

G rard Cornu jols

Tepper School of Business, Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, [gc0v@andrew.cmu.edu](mailto:gc0v@andrew.cmu.edu)

Aris Daniilidis

Department of Mathematical Engineering, University of Chile, Santiago, Chile, [arisd@dim.uchile.cl](mailto:arisd@dim.uchile.cl)

Claude Lemar chal

INRIA-Bipop, Montbonnot 38334 Saint Ismier, France, [claude.lemarechal@inria.fr](mailto:claude.lemarechal@inria.fr)

J r me Malick

CNRS, Laboratoire J. Kuntzmann, 38041 Grenoble, France, [jerome.malick@inria.fr](mailto:jerome.malick@inria.fr)

We consider the separation problem for sets  $X$  that are pre-images of a given set  $S$  by a linear mapping. Classical examples occur in integer programming, as well as in other optimization problems such as complementarity. One would like to generate valid inequalities that cut off some point not lying in  $X$ , without reference to the linear mapping. To this aim, we introduce a concept: cut-generating functions (CGF) and we develop a formal theory for them, largely based on convex analysis. They are intimately related to  $S$ -free sets and we study this relation, disclosing several definitions for minimal CGF's and maximal  $S$ -free sets. Our work unifies and puts into perspective a number of existing works on  $S$ -free sets; in particular, we show how CGF's recover the celebrated Gomory cuts.

*Keywords:* integer programming; convex analysis; separation; generalized gauges;  $S$ -free sets

*MSC2000 subject classification:* Primary: 52A41, 90C11; secondary: 52A40

*OR/MS subject classification:* Primary: combinatorics, convexity; secondary: cutting plane

*History:* Received February 28, 2013; revised February 2, 2014. Published online in *Articles in Advance* August 20, 2014.

## 1. Introduction. In this paper, we consider sets of the form

$$X = X(R, S) := \{x \in \mathbb{R}_+^n : Rx \in S\}, \quad \text{where} \tag{1a}$$

$$\begin{cases} R = [r_1 \dots r_n] \text{ is a real } q \times n \text{ matrix,} \\ S \subset \mathbb{R}^q \text{ is a nonempty closed set with } 0 \notin S. \end{cases} \tag{1b}$$

In other words, our set  $X$  is the intersection of a closed convex cone with a pre-image by a linear mapping. This model goes back to Johnson [18], where  $S$  was a finite set: constraints  $Rx = b$  were considered for several right-hand sides  $b$ . Here, we rather consider a general (possibly infinite) set  $S$  and a varying constraint matrix  $R$ . The closed convex hull of  $X$  does not contain 0 (see Lemma 2.1) and we are then interested in *separating* 0 from  $X$ : we want to generate *cuts*, i.e., inequalities that are valid for  $X$ , which we write as

$$c^\top x \geq 1, \quad \text{for all } x \in X. \tag{2}$$

### 1.1. Motivating examples. Our first motivation comes from (mixed) integer linear programming.

**EXAMPLE 1.1 (AN INTEGER LINEAR PROGRAM).** Let us first consider a pure integer program, which consists in optimizing a linear function over the set defined by the constraints

$$Dz = d \in \mathbb{R}^m, \quad z \in \mathbb{Z}_+^p. \tag{3}$$

Set  $n := p - m$ , assume the matrix  $D$  to have full row-rank  $m$  and select  $m$  independent columns (a *basis*). The corresponding decomposition  $z = (x, y)$  into nonbasic and basic variables amounts to writing the above feasible set as the intersection of  $\mathbb{Z}^n \times \mathbb{Z}^m$  with the polyhedron

$$P := \{(x, y) \in \mathbb{R}_+^n \times \mathbb{R}_+^m : Ax + y = b\} \tag{4}$$

for suitable  $m \times n$  matrix  $A$  and  $m$ -vector  $b$ .

Downloaded from informs.org by [200.89.68.74] on 15 July 2015, at 13:40. For personal use only, all rights reserved.

Relaxing the nonnegativity constraint on the basic variables  $y$ , we obtain the classical *corner polyhedron* (Gomory [14]), namely, the convex hull of

$$\{(x, y) \in \mathbb{Z}_+^n \times \mathbb{Z}^m: Ax + y = b\}.$$

This model has the form (1) if we set

$$q = n + m, \quad R = \begin{bmatrix} I \\ -A \end{bmatrix}, \quad S = \mathbb{Z}^n \times (\mathbb{Z}^m - \{b\}), \quad (5)$$

where  $\mathbb{Z}^m - \{b\}$  denotes the translation of  $\mathbb{Z}^m$  by the vector  $-b$ . Assuming  $b \notin \mathbb{Z}^m$ , the above  $S$  is a closed set not containing the origin.

For  $m = 1$ , (4) has a single constraint

$$\sum_{j=1}^n a_j x_j + y = b, \quad y \in \mathbb{Z}, \quad x \in \mathbb{Z}_+^n;$$

the celebrated Gomory cut (Gomory [13]) is

$$\sum_{j:f_j \leq f_0} \frac{f_j}{f_0} x_j + \sum_{j:f_j > f_0} \frac{1-f_j}{1-f_0} x_j \geq 1, \quad (6)$$

where  $f_j = a_j - \lfloor a_j \rfloor$  and  $f_0 = b - \lfloor b \rfloor$ . Inequality (6) is valid for the corner polyhedron and cuts off the basic solution  $(x = 0, y = b)$ . In the  $x$ -space  $\mathbb{R}^n$ , this inequality is a cut as defined in (2). We will demonstrate in Example 2.8 how to recover such a cut from our formalism.

Except for the translation by the basic solution  $(0, b)$ ,  $S$  is quasi instance independent. This is actually a crucial feature; it determines the approach developed in this paper, namely, cut-generating functions to be developed below.

EXAMPLE 1.2 (A MIXED-INTEGER LINEAR PROGRAM). In our integer program (3), let us now relax not only nonnegativity of the basic variables but also integrality of the nonbasic variables: the corner polyhedron is further relaxed to the convex hull of

$$\{(x, y) \in \mathbb{R}_+^n \times \mathbb{Z}^m: Ax + y = b\}.$$

We are still in the context of (1) with

$$q = m, \quad R = -A, \quad S = \mathbb{Z}^m - b;$$

this is the model considered in Andersen et al. [1] for  $m = 2$ , and in Borozan and Cornuéjols [8] for general  $m$ . Other relevant references are Basu et al. [5, 6], Dey and Wolsey [12], Gomory and Johnson [15], and Johnson [18].

This type of relaxation can be used when (3) becomes a mixed-integer linear program

$$Dz = d \in \mathbb{R}^m, \quad z \geq 0, \quad z_j \in \mathbb{Z}, \quad j \in J,$$

where  $J$  is some subset of  $\{1, \dots, p\}$ . Extract a basis as before and choose a subset of basic variables indexed in  $J$ ; call  $m' \leq m$  the number of rows in this restriction and  $b' \in \mathbb{R}^{m'}$  the resulting restriction of  $b$  (in other words, ignore a number  $m - m'$  of linear constraints). Relax nonnegativity of the  $m'$  remaining basic variables, as well as integrality of the nonbasic variables indexed in  $J$ . This results in (1), with

$$q = m', \quad R = -A, \quad S = \mathbb{Z}^{m'} - b'.$$

Any cut for this set  $X$  is *a fortiori* a valid inequality for the original mixed-integer linear program.

When  $m' = 1$ , a classical example of such inequalities is

$$\sum_{j:a_j > 0} \frac{a_j}{f_0} x_j - \sum_{j:a_j < 0} \frac{a_j}{1-f_0} x_j \geq 1. \quad (7)$$

Actually, Gomory's mixed-integer cuts (Gomory [13]) combine (6) for the integer nonbasic variables with the above formula for the continuous ones.

Model (1) occurs in other areas than integer programming and we give another example.

EXAMPLE 1.3 (COMPLEMENTARITY PROBLEM). Still using  $P$  of (4), let

$$E \subset \{1, 2, \dots, m\} \times \{1, 2, \dots, m\} \quad \text{and} \quad C := \{y \in \mathbb{R}_+^m : y^i y^j = 0, (i, j) \in E\}$$

(in this paper,  $\subset$  stands for inclusion and  $\subsetneq$  for strict inclusion).

The set of interest is then  $P \cap (\mathbb{R}^n \times C)$ . It can be modeled by (1), where

$$q = m, \quad R = -A, \quad S = C - b.$$

Cuts have been used for complementarity problems of this type, for example, in Júdice et al. [19].

We will retain from these examples the dissymmetry between  $S$  (a very particular and highly structured set) and  $R$  (an arbitrary matrix). Keeping this in mind, we will consider that  $(q, S)$  is given and fixed, whereas  $(n, R)$  is instance-dependent data: our cutting problem can be viewed as *parametrized* by  $(n, R)$ . This point of view is natural for the last two examples; but some preprocessing (to be seen in Example 2.8) is needed to apply it to Example 1.1: by (5),  $S$  does depend on the data through its dimension  $q$ , which depends on  $n$ .

**1.2. Introducing cut-generating functions.** To generate cuts in the present situation, it would be convenient to have a mapping, taking instances of (1) as input, and producing cuts as output. What we need for this is a function

$$\mathbb{R}^q \ni r \mapsto \rho(r) \in \mathbb{R},$$

which, applied to the columns  $r_j$  of a  $q \times n$  matrix  $R$  (an arbitrary matrix, with an arbitrary number of columns) will produce the  $n$  coefficients  $c_j := \rho(r_j)$  of a cut (2). We stress the fact that  $\rho$  must assign a number  $\rho(r)$  to *any*  $r \in \mathbb{R}^q$ : the function  $\rho$  is defined on the whole space.

Thus, we require from our  $\rho$  to satisfy

$$x \in X \implies \sum_{j=1}^n \rho(r_j) x_j \geq 1, \quad (8)$$

for every instance  $X$  of (1). Such a  $\rho$  can then justifiably be called a *cut-generating function* (CGF). The notation  $\rho$  refers to *representation*, which will appear in Definition 2.6. One of the most well-known cut-generating functions in integer programming is the so-called Gomory function (Gomory [13]), which we presented in Examples 1.1 and 1.2. The corresponding cuts can be generated quickly, so they are a powerful tool in computations; indeed, they drastically speed up integer-programming solvers (Bixby and Rothberg [7]).

So far, a CGF is a rather abstract object, as it lies in the (vast!) set of functions from  $\mathbb{R}^q$  to  $\mathbb{R}$ ; but the following observation allows a drastic reduction of this set.

REMARK 1.4 (DOMINATING CUTS). Consider in (2) a vector  $c'$  with  $c'_j \leq c_j$  for  $j = 1, \dots, n$ ; then  $c'^T x \leq c^T x$  whenever  $x \geq 0$ . If  $c'$  is a cut, it is tighter than  $c$  in the sense that it cuts a bigger portion of  $\mathbb{R}_+^n$ . We can impose some “minimal” character to a CGF, in order to reach some “tightness” of the resulting cuts.

With this additional requirement, the decisive Theorem 2.3 will show that a CGF can be imposed to be *convex positively homogeneous* (and defined on the whole of  $\mathbb{R}^q$ ; positive homogeneity means  $\rho(tr) = t\rho(r)$  for all  $r \in \mathbb{R}^q$  and  $t > 0$ ). This is a fairly narrow class of functions indeed, which is fundamental in convex analysis. Such functions are in correspondence with closed convex sets and in our context, this correspondence is based on the mapping  $\rho \mapsto V$  defined by

$$V = V(\rho) := \{r \in \mathbb{R}^q : \rho(r) \leq 1\}, \quad (9)$$

which turns out to be a cornerstone: via Theorem 2.5, (9) establishes a correspondence between the CGF's and the so-called *S-free sets*. As a result, cut-generating functions can alternatively be studied from a geometric point of view, involving sets  $V$  instead of functions  $\rho$ . This situation, common in convex analysis, is often very fruitful. With regard to Remark 1.4, observe that  $V(\rho)$  increases when  $\rho$  decreases: small  $\rho$ 's give large  $V$ 's. However the converse is not true because the mapping in (9) is many-to-one and therefore has no inverse. A first concern will therefore be to specify appropriate correspondences between (cut-generating) functions and (*S-free*) sets.

**1.3. Scope of the paper.** The aim of the paper is to present a formal theory of minimal cut-generating functions and maximal  $S$ -free sets, valid independently of the particular  $S$ . Such a theory would gather and synthesize a number of papers dealing with the above problem for various special forms for  $S$ : Lovász [21], Andersen et al. [1], Borozan and Cornuéjols [8], Dey and Wolsey [12], Basu et al. [5, 6], and references therein. For this, we use basic tools from convex analysis and geometry. Readers not familiar with this field may use Hiriart-Urruty and Lemaréchal [17, Chapter C] for an elementary introduction, although Hiriart-Urruty and Lemaréchal [16], Rockafellar [23] are more complete.

The paper is organized as follows:

- Section 2 states more accurately the concepts of CGF's and  $S$ -free sets.
- Section 3 studies the mapping (9). We show that the pre-images of a given  $V$  (the representations of  $V$ ) have a unique largest element  $\gamma_V$  and a unique smallest element  $\mu_V$ ; in view of Remark 1.4, the latter then appears as the relevant inverse of  $\rho \mapsto V(\rho)$ .
- In §4, we study the correspondence  $V \leftrightarrow \mu_V$ . We show that different concepts of minimality come into play for  $\rho$  in Remark 1.4. Geometrically they correspond to different concepts of maximality for  $V$ .
- We also show in §5 that these minimality concepts coincide in a number of cases.
- Finally we have a conclusion section, with some suggestions for future research.

The ideas in §§2 and 3 extend, in a natural way, the earlier works mentioned above. However, §§4 and 5 contain new results.

**2. Cut-generating functions: Definitions and first results.** We begin with making sure that our framework is consistent. We will use  $\text{conv}(X)$  [respectively,  $\overline{\text{conv}}(X)$ ] to denote the convex hull [respectively, closed convex hull] of a set  $X$ .

LEMMA 2.1. *With  $X$  given as in (1),  $0 \notin \overline{\text{conv}}(X)$ .*

PROOF. Assume  $X \neq \emptyset$ , otherwise we have nothing to prove. Since 0 does not lie in the closed set  $S$ , there is  $\varepsilon > 0$  such that  $s \in S$  implies  $\|s\|_1 \geq \varepsilon$ ; and by continuity of the mapping  $x \mapsto Rx$ , there is  $\eta > 0$  such that  $\|x\|_1 \geq \eta$  for all  $x \in X \subset \mathbb{R}_+^n$ . This means

$$\|x\|_1 = \sum_{j=1}^n |x_j| = \sum_{j=1}^n x_j \geq \eta, \quad \text{for all } x \in X.$$

In other words, the hyperplane  $\sum_j x_j \geq \eta$  separates 0 from  $X$ , hence from  $\overline{\text{conv}}(X)$ .  $\square$

Remember that we are interested in functions  $\rho$  satisfying (8) for any  $(n, R)$  in (1). There are too many such functions; we now proceed to specify exactly which ones are relevant.

**2.1. Sublinear cut-generating functions suffice.** The following lemma, inspired from Claim 1 in the proof of Basu et al. [5, Lemma 23], is instrumental for our purpose.

LEMMA 2.2. *Let  $\rho$  be a CGF. For all sets of  $K$  vectors  $r_k \in \mathbb{R}^q$  and nonnegative coefficients  $\alpha_k$ , the following relation holds:*

$$\sum_{k=1}^K \alpha_k r_k = 0 \implies \sum_{k=1}^K \alpha_k \rho(r_k) \geq 0.$$

PROOF. Call  $e \in \mathbb{R}^q$  the vector of all ones and  $\alpha \in \mathbb{R}^K$  the vector of  $\alpha_k$ 's; take  $t \geq 0$  and define the vectors in  $\mathbb{R}^{K+q}$

$$x := \begin{bmatrix} 0 \\ e \end{bmatrix}, \quad d := \begin{bmatrix} \alpha \\ 0 \end{bmatrix}, \quad \text{so that } x + td = \begin{bmatrix} t\alpha \\ e \end{bmatrix} \in \mathbb{R}_+^{K+q}.$$

Then pick  $s \in S$ ; make an instance of (1) with  $n = K + q$  and  $R := [r_1 \dots r_K \mid D(s)]$ , where the  $q \times q$  matrix  $D(s)$  is the diagonal matrix whose diagonal is the vector  $s$ . Observing that

$$R(x + td) = t \sum_k \alpha_k r_k + D(s)e = s,$$

$x + td$  is feasible in the resulting instance of (1a): (8) becomes

$$t \sum_{k=1}^K \alpha_k \rho(r_k) \geq 1 - z,$$

where  $z$  is a fixed number gathering the result of applying  $\rho$  to the columns of  $D(s)$ . Letting  $t \rightarrow +\infty$  proves the claim.  $\square$

Now we introduce some notation. The *domain* and *epigraph* of a function  $\rho: \mathbb{R}^q \rightarrow \mathbb{R} \cup \{+\infty\}$  are

$$\text{dom } \rho := \{r \in \mathbb{R}^q: \rho(r) < +\infty\} \quad \text{and} \quad \text{epi } \rho := \{(r, z) \in \mathbb{R}^{q+1}: z \geq \rho(r)\}.$$

If  $\text{dom } \rho$  is the whole of  $\mathbb{R}^q$  (i.e.,  $\rho(r)$  is a finite real number for all  $r \in \mathbb{R}^q$ ), we say that  $\rho$  is *finite valued*; a convex finite-valued function is continuous on  $\mathbb{R}^q$ . A function is said to be *sublinear* if it is convex and positively homogeneous; or equivalently if its epigraph is a convex cone. The *conical hull* cone ( $\text{epi } \rho$ ) of  $\text{epi } \rho$  is the set of nonnegative combinations of points  $(r, z) \in \text{epi } \rho$ :

$$r = \sum_{k=1}^K \alpha_k r_k, \quad z = \sum_{k=1}^K \alpha_k z_k, \quad \text{with } z_k \geq \rho(r_k), \quad \alpha_k \geq 0, \quad k = 1, \dots, K,$$

where  $K$  is an arbitrary integer. This conical hull is itself the epigraph of a sublinear function  $\bar{\rho}$ , called the *sublinear hull* of  $\rho$ . Its value at  $r$  is the smallest possible of the above  $z$ 's:

$$\bar{\rho}(r) := \inf \left\{ \sum_{k=1}^K \alpha_k \rho(r_k) : \sum_{k=1}^K \alpha_k r_k = r, \alpha_k \geq 0 \right\}. \tag{10}$$

Of course  $\bar{\rho} \leq \rho$ ; in the spirit of Remark 1.4, our next result establishes that a CGF can be improved by taking its sublinear hull.

**THEOREM 2.3.** *If  $\rho$  is a CGF, then  $\bar{\rho}$  of (10) is nowhere  $-\infty$  and is again a CGF.*

**PROOF.** Express every  $r \in \mathbb{R}^q$  as a nonnegative combination:  $\sum_k \alpha_k r_k - r = 0$ , hence (Lemma 2.2)  $\sum_{k=1}^K \alpha_k \rho(r_k) + \rho(-r) \geq 0$  and  $\bar{\rho}(r) \geq -\rho(-r) > -\infty$ .

Then take an instance  $R = [r_j]_{j=1}^n$  of (1b). If it produces  $X = \emptyset$  in (1a), there is nothing to prove. Otherwise fix  $\bar{x} \in X$ .

Any positive decomposition  $r_j = \sum_k \alpha_{j,k} r_{j,k}$  of each column of  $R$  satisfies

$$\bar{s} := R\bar{x} = \sum_{j=1}^n \bar{x}_j r_j = \sum_{j=1}^n \bar{x}_j \sum_{k=1}^K \alpha_{j,k} r_{j,k} = R_+ x_+,$$

where  $x_+ \in \mathbb{R}^{nK}$  denotes the vector with coordinates  $\alpha_{j,k} \bar{x}_j \geq 0$  and  $R_+$  the matrix whose  $nK$  columns are  $r_{j,k}$ . Then  $R_+$  is a possible instance of (1b) and  $R_+ x_+ = \bar{s} \in S$ , so the CGF  $\rho$  separates  $x_+$  from 0:

$$1 \leq \sum_{j,k} \rho(r_{j,k}) (\alpha_{j,k} \bar{x}_j) = \sum_{j=1}^n \left( \sum_{k=1}^K \alpha_{j,k} \rho(r_{j,k}) \right) \bar{x}_j. \tag{11}$$

Apply the definition of an infimum: for each  $\varepsilon > 0$  we can choose our decompositions  $(r_{j,k}, \alpha_{j,k})$  so that

$$\sum_{k=1}^K \alpha_{j,k} \rho(r_{j,k}) \leq \bar{\rho}(r_j) + \varepsilon, \quad \text{for } j = 1, \dots, n$$

which yields with (11)

$$1 \leq \sum_{j=1}^n (\bar{\rho}(r_j) + \varepsilon) \bar{x}_j = \sum_{j=1}^n \bar{\rho}(r_j) \bar{x}_j + \varepsilon \sum_{j=1}^n \bar{x}_j.$$

Since  $\varepsilon$  is arbitrarily small—and  $\bar{x}$  is fixed—we see that  $\bar{\rho}$  does satisfy (8).  $\square$

In view of Remark 1.4, Theorem 2.3 allows us to restrict our attention to CGF's that are sublinear; and their domain is the whole space by definition. We are now in a position to explain the use of the operation (9) in our context.

**2.2. Cut-generating functions as representations.** From now on, a CGF  $\rho$  will always be understood as a (finite-valued) sublinear function. By continuity and because  $\rho(0) = 0$ ,  $V(\rho)$  in (9) is a closed convex neighborhood of 0 in  $\mathbb{R}^q$ . Besides, its interior and boundary are, respectively,

$$\text{int}(V(\rho)) = \{r \in V: \rho(r) < 1\}, \quad \text{bd}(V(\rho)) = \{r \in V: \rho(r) = 1\}. \tag{12}$$



This follows from the Slater property  $\rho(0) = 0$  (see, e.g., Hiriart-Urruty and Lemaréchal [17, Proposition D.1.3.3]); it can also be checked directly:

- by continuity,  $\rho(\bar{r}) < 1$  implies  $\rho(r) \leq 1$  for  $r$  close to  $\bar{r}$ ; and
- by positive homogeneity,  $\rho(\bar{r}) = 1$  implies  $\rho(r) = 1 + \varepsilon$  for  $r = (1 + \varepsilon)\bar{r}$ .

The relevant neighborhoods for our purpose are the following:

**DEFINITION 2.4 (S-FREE SET).** Given a closed set  $S \subset \mathbb{R}^q$  not containing the origin, a closed convex neighborhood  $V$  of  $0 \in \mathbb{R}^q$  is called  $S$ -free if its interior contains no point in  $S$ :  $\text{int}(V) \cap S = \emptyset$ .

Let us make clear the importance of this definition.

**THEOREM 2.5.** Let the sublinear function  $\rho: \mathbb{R}^q \rightarrow \mathbb{R}$  and the closed convex neighborhood  $V$  (of  $0 \in \mathbb{R}^q$ ) satisfy (9). Then  $\rho$  is a CGF for (1) if and only if  $V$  is  $S$ -free.

**PROOF.** Let  $V$  be  $S$ -free; in view of (12),  $\rho(r) \geq 1$  for all  $r \in S$ . In particular, take a  $q \times n$  matrix  $R$ ,  $x \in X$  of (1a) and set  $r := Rx \in S$ . Then, using sublinearity,

$$1 \leq \rho(Rx) = \rho\left(\sum_{j=1}^n x_j r_j\right) \leq \sum_{j=1}^n x_j \rho(r_j);$$

$\rho$  is a CGF.

Conversely, suppose  $V$  is not  $S$ -free: again from (12), there is some  $r_1 \in S$  such that  $\rho(r_1) < 1$ . Take in (1b) the instance  $(n, R) = (1, [r_1])$ . Then  $1 \in X$  ( $r_1 \in S$ ), so  $c_1 := \rho(r_1) < 1$  cannot be a cut.  $\square$

This allows a new definition of CGF's, much more handy than the original one.

**DEFINITION 2.6 (CGF AS REPRESENTATION).** Let  $V \subset \mathbb{R}^q$  be a closed convex neighborhood of the origin. A representation of  $V$  is a finite-valued sublinear function  $\rho$  such that

$$V = \{r \in \mathbb{R}^q: \rho(r) \leq 1\}. \tag{13}$$

We will say that  $\rho$  represents  $V$ .

A sublinear cut-generating function for (1) is a representation of an  $S$ -free set.

A finite-valued sublinear function  $\rho$  represents a unique  $V = V(\rho)$ , well-defined by (13). One easily checks monotonicity of the mapping  $V(\cdot)$ :

$$\rho \leq \rho' \implies V(\rho) \supset V(\rho'). \tag{14}$$

Conversely, one may ask whether a given closed convex neighborhood of the origin  $V$  always has a representation. In fact, (13) fixes via (12) the value  $\rho(r) = 1$  on the boundary of  $V$ ; whether this set of prescribed values can be extended to make a sublinear function on the whole of  $\mathbb{R}^q$  is not obvious. This will be the subject of §3, where we will see that this is indeed possible; there may even be infinitely many extensions, and we are interested in the small ones. Now we illustrate the material introduced so far with some examples.

**2.3. Examples.** We start with a simple one-dimensional example supporting the claim that the mapping  $\rho \rightarrow V$  of (13) is many-to-one—or equivalently that a given  $V$  may have several representations.

**EXAMPLE 2.7.** With  $q = 1$ , consider  $V = ]-\infty, 1]$ . In  $\mathbb{R}^1$ , the positively homogeneous functions have the form

$$\rho(r) = \begin{cases} \alpha r & \text{for } r \geq 0, \\ \beta r & \text{for } r \leq 0, \end{cases}$$

they are convex when  $\alpha \geq \beta$ .

Taking  $r = 1 \in V$  in (13) imposes  $\alpha \leq 1$ , and taking  $r = 1 + \varepsilon \notin V$  ( $\varepsilon > 0$ ) imposes  $\alpha > 1/(1 + \varepsilon)$ . Altogether  $\alpha = 1$ . On the other hand, letting  $r \rightarrow -\infty$ , the property  $\beta r \leq 1$  imposes  $\beta \geq 0$ .

Conversely, we easily see that, for any  $\beta \in [0, 1]$ , the function

$$\rho(r) = \begin{cases} r & \text{for } r \geq 0, \\ \beta r & \text{for } r \leq 0, \end{cases}$$

is sublinear and satisfies (13). Thus, the representations of  $V$  are exactly the functions of the form  $\rho(r) = \max\{r, \beta r\}$ , for  $\beta \in [0, 1]$ .

This example suggests—and Lemma 3.2 will confirm—that nonuniqueness appears when  $V$  is unbounded.




FIGURE 1. Two  $S$ -free sets for  $q = 2$ .

Example 1.2 is quite suitable for illustration, Figure 1 visualizes it for  $q = m = 2$ . The dots are the set  $S = \mathbb{Z}^2 - \{b\}$ . The stripe  $V$  of the left part, called a *split set*, is used in the framework of disjunctive cuts. Other neighborhoods can be considered, for example, triangles (right part of the picture) as in Andersen et al. [1].

With  $q = 1$ , no need for a picture and the calculations in Example 1.2 can be worked out. In this case,  $X \subset \mathbb{R}_+^n$  is defined by  $a^\top x \in \mathbb{Z} - b$ , i.e., (1) with  $r = -a$  and  $S = \mathbb{Z} - \{b\}$ . The only possible  $S$ -free neighborhoods of the origin are the segments  $r \in [-r_-, r_+]$  with

$$-f_0 = \lfloor b \rfloor - b \leq -r_- < 0 < r_+ \leq \lceil b \rceil - b = 1 - f_0.$$

For a representation  $\rho$  of this segment, the equations  $\rho(r_+) = 1$  and  $\rho(-r_-) = 1$  fix in a unique way

$$\rho(r) = \begin{cases} \frac{r}{r_+} & \text{if } r \geq 0, \\ -\frac{r}{r_-} & \text{if } r \leq 0. \end{cases}$$

Choose the extreme values for  $r_+$  and  $r_-$  to obtain

$$c_j = \rho(-a_j) = \begin{cases} \frac{a_j}{f_0} & \text{if } a_j \leq 0, \\ \frac{-a_j}{1 - f_0} & \text{if } a_j \geq 0, \end{cases}$$

which is just (7).

Finally, let us show how Gomory cuts (6) can be obtained as CGF's.

EXAMPLE 2.8. Still in Example 1.2, take  $q = m = 1$ ; we want to separate the set defined by

$$\sum_{j=1}^n a_j x_j + y = b, \quad y \in \mathbb{Z}, \quad x \in \mathbb{Z}_+^n$$

from the origin (remember that  $b \notin \mathbb{Z}$ ). This set has the form (1) with

$$q = n + 1, \quad R = \begin{bmatrix} I \\ -a^\top \end{bmatrix}, \quad S = \mathbb{Z}^n \times (\mathbb{Z} - \{b\}).$$

Introduce the vector  $\pi \in \mathbb{R}^{n+1}$  defined by

$$\pi_{n+1} := 1 \quad \text{and} \quad \text{for } j = 1, \dots, n: \pi_j := \begin{cases} \lfloor a_j \rfloor & \text{if } f_j \leq f_0, \\ \lceil a_j \rceil & \text{if } f_j > f_0, \end{cases}$$

and its scalar product  $\pi^\top r = \sum_{j=1}^n \pi_j x_j + y$  with  $r = (x, y) \in \mathbb{R}^{n+1}$ . Then define

$$V := \{r: \lfloor b \rfloor - b \leq \pi^\top r \leq \lceil b \rceil - b\}. \tag{15}$$

The assumption  $b \notin \mathbb{Z}$  implies that  $(0, 0) \in \text{int}(V)$ ; therefore  $V$  is a closed convex neighborhood of the origin. Furthermore,  $V$  is  $S$ -free: in fact,  $b + \pi^\top r$  is an integer for every  $r = (x, y) \in S$  and therefore it cannot be strictly between the two consecutive integers  $\lfloor b \rfloor$  and  $\lceil b \rceil$ . We claim that any representation of  $V$  produces Gomory cuts.

Downloaded from informs.org by [200.89.68.74] on 15 July 2015, at 13:40. For personal use only, all rights reserved.



Call  $e_j$  the  $j$ th unit vector of  $\mathbb{R}^n$ , so that the  $n$  columns of  $R$  are

$$r_j = \begin{pmatrix} e_j \\ -a_j \end{pmatrix}.$$

Direct calculations give

$$\pi^\top r_j = \begin{cases} \lfloor a_j \rfloor - a_j = -f_j & \text{if } f_j \leq f_0, \\ \lceil a_j \rceil - a_j = 1 - f_j & \text{if } f_j > f_0. \end{cases}$$

For each  $j = 1, \dots, n$ , consider three cases.

(i) If  $\pi^\top r_j > 0$  (which implies  $f_j > f_0$ ), there is  $t > 0$  such that  $t\pi^\top r_j = \lceil b \rceil - b > 0$ , namely,

$$t = \frac{\lceil b \rceil - b}{\pi^\top r_j} = \frac{\lceil b \rceil - b}{\lceil a_j \rceil - a_j} = \frac{1 - f_0}{1 - f_j}.$$

(ii) If  $\pi^\top r_j < 0$  (which implies  $0 < f_j \leq f_0$ ), there exists likewise  $t > 0$  such that  $t\pi^\top r_j = \lfloor b \rfloor - b < 0$ , therefore

$$t = \frac{f_0}{f_j}.$$

(iii) If  $\pi^\top r_j = 0$  (which implies  $a_j \in \mathbb{Z}$ ),  $tr_j \in V$  for any  $t > 0$ .

In (i) and (ii), the computed value of  $t$  puts  $tr_j$  on the boundary of  $V$ . Let  $\rho$  represent  $V$ ; then by (12) and positive homogeneity,  $\rho(r_j) = (1/t)\rho(tr_j) = 1/t$  in cases (i) and (ii) and  $\rho(r_j) = 0$  in case (iii). Altogether,

$$\rho(r_j) = \begin{cases} \frac{f_j}{f_0} & \text{if } f_j \leq f_0, \\ \frac{1 - f_j}{1 - f_0} & \text{if } f_j > f_0, \end{cases}$$

for  $j = 1, \dots, n$ ; we recognize Gomory's formula (6).

As mentioned after Definition 2.6, the  $n$  values  $\rho(r_j)$  can be extended to make a sublinear function on the whole of  $\mathbb{R}^{n+1}$ . This will be confirmed in the next section but can be accepted here, thanks to the simple form (15) of  $V$ : a stripe orthogonal to  $\pi$ . Indeed, the above calculations are designed so as to construct  $\rho(r) = 1$  for each  $r$  such that  $\pi^\top r = \lceil b \rceil - b > 0$  as in (i) (respectively,  $\pi^\top r = \lfloor b \rfloor - b < 0$  as in (ii)). Then  $\rho(r)$  is given by positive homogeneity for any  $r$  such that  $\pi^\top r \neq 0$ ; and  $\rho \equiv 0$  on  $\pi^\perp$ .

**3. Largest and smallest representations.** In this section, we study the representation operation introduced in Definition 2.6. The main result is that our closed convex neighborhood  $V$  has a largest and a smallest representation. This result was already given in Caprari and Zaffaroni [9], Basu et al. [4], and Zaffaroni [24], with weaker assumptions in the latter work (which came to our knowledge only after the present paper was completed). Here we emphasize the geometric counterpart of the result, we put the proof of Basu et al. [4] in perspective, and we take advantage of our stricter assumptions to develop finer results that will be useful in the sequel.

**3.1. Some elementary convex analysis.** First recall some basic theory (see, e.g., Hiriart-Urruty and Lemaréchal [17, Chapter C]), which will be central in our development. In what follows,  $V$  will always be a closed convex neighborhood of  $0 \in \mathbb{R}^q$ .

A common object in convex analysis is the *gauge*

$$\mathbb{R}^q \ni r \mapsto \gamma_V(r) := \inf \{ \lambda > 0 : r \in \lambda V \}, \tag{16}$$

a (nonnegative) finite-valued sublinear function. Applying for example Hiriart-Urruty and Lemaréchal [17, Theorem C.1.2.5] with the notation  $(x, C, r)$  replaced by  $(r, V, 1)$ , we obtain the relation

$$V = \{ r \in \mathbb{R}^q : \gamma_V(r) \leq 1 \}.$$

Thus  $\gamma_V$  represents  $V$ ; this first confirms that Definition 2.6 is consistent.

Another fundamental object is the *support function* of an arbitrary set  $G \subset \mathbb{R}^q$ , defined by

$$\mathbb{R}^q \ni r \mapsto \sigma_G(r) := \sup_{d \in G} d^\top r. \tag{17}$$

This function is easily seen to be sublinear, to grow when  $G$  grows, and to remain unchanged if  $G$  is replaced by its closed convex hull:  $\sigma_G = \sigma_{\text{conv}(G)}$ . Besides, it is finite valued if (and only if)  $G$  is bounded.

Conversely, every (finite-valued) sublinear function  $\sigma$  is the support function of a (bounded) closed convex set, unambiguously defined by

$$G = G_\sigma := \{d \in \mathbb{R}^q : d^\top r \leq \sigma(r) \text{ for all } r \in \mathbb{R}^q\} \quad (18)$$

(note:  $G_\sigma$  is closed and convex because it is an intersection of half spaces; actually,  $G_\sigma$  is just the *subdifferential* of  $\sigma$  at 0). We then say that  $\sigma$  supports  $G_\sigma$ . The correspondence  $\sigma \leftrightarrow G$  defines a one-to-one mapping between finite-valued sublinear functions and bounded closed convex sets (the mapping  $\sigma \mapsto G$  of (18) extends to sublinear functions in  $\mathbb{R} \cup \{+\infty\}$  but such an extension is not needed here).

**REMARK 3.1 (PRIMAL-DUAL NOTATION).** Equation (17) involves two variables,  $d$  and  $r$ , both written as column vectors; nevertheless, they lie in two mutually dual spaces. In this paper, we keep going back and forth between these two spaces; even though they are the same  $\mathbb{R}^q$ , we make a point to distinguish between the two. The notation  $r, V, \dots$  [respectively,  $d, G, \dots$ ] will generally be used for primal elements [respectively, for dual ones]. Most of the time, we will deal with support functions  $\sigma_G(r)$  of dual sets; but we will also consider the support function  $\sigma_V(d)$  of our primal neighborhood  $V$ .

Being finite-valued sublinear, the gauge of  $V$  supports a compact convex set, obtained by replacing  $\sigma$  by  $\gamma_V$  in (18). Since  $\gamma_V \geq 0$ , we guess from positive homogeneity that this set is just the *polar* of  $V$ :

$$\{d \in \mathbb{R}^q : d^\top r \leq \gamma_V(r) \text{ for all } r \in \mathbb{R}^q\} = \{d \in \mathbb{R}^q : d^\top r \leq 1 \text{ for all } r \in V\} =: V^\circ. \quad (19)$$

Write (19) as  $V^\circ = \{d \in \mathbb{R}^q : \sigma_V(d) \leq 1\}$  to see that  $\sigma_V$  represents  $V^\circ$ ; thus, the support function of  $V$  is the gauge of  $V^\circ$ , so that the polar of  $V^\circ$  is  $V$  itself:  $(V^\circ)^\circ = V$ . These various properties are rather classical; see, for example, Hiriart-Urruty and Lemaréchal [17, Proposition C.3.2.4, Corollary C.3.2.5], with  $(d, C, s)$  replaced by  $(r, V, d)$ .

Now remember Example 2.7:  $V$  may have several representations. Any such representation  $\rho$  supports a set  $G_\rho$  and we will see that the polar of  $G_\rho$  is again  $V$  itself;  $G_\rho$  is a pre-image of  $V$  for the polarity mapping. We thus obtain a new concept: a *prepolar* of  $V$  is a set  $G$  such that

$$G^\circ := \{r \in \mathbb{R}^q : \sigma_G(r) \leq 1\} = V,$$

or equivalently  $\sigma_G$  represents  $V$ .

The property  $(V^\circ)^\circ = V$  means that the standard polar  $V^\circ$  is itself a prepolar—which is somewhat confusing; and it turns out to be the largest one (Corollary 3.3); or equivalently, its support function  $\sigma_{V^\circ} = \gamma_V$  turns out to be the largest representation of  $V$ . The main result of this section states that  $V$  has also a smallest prepolar, or equivalently a smallest representation (Proposition 3.6); keeping Remark 1.4 in mind, this is exactly what we want. This result is actually Theorem 1 in Basu et al. [4]; here we use elementary convex analysis and we insist more on the geometric aspect.

**3.2. Largest representation.** Introduce the recession cone  $V_\infty$  of  $V$ . Using the property  $0 \in V$ , it can be defined as

$$V_\infty = \{r \in \mathbb{R}^q : tr \in V \text{ for all } t > 0\} = \bigcap_{\lambda > 0} \lambda V,$$

and the second relation shows that  $V_\infty$  is closed; taking in particular  $\lambda = 1$  shows that

$$V_\infty \subset V. \quad (20)$$

One then easily sees from (16) that  $\gamma_V(r) = 0$  if  $r \in V_\infty$ . Yet, for any other representation  $\rho$  of  $V$ , (13) just imposes  $\rho(r) \leq 0$  at this  $r$  and we may a priori have  $\rho(r) < 0$ : the possible representations of  $V$  may differ on  $V_\infty$ ; see Example 2.7 again. We make this more precise.

**LEMMA 3.2 (REPRESENTATIONS AND RECESSON CONE).** *For all representations  $\rho$  of the closed convex neighborhood  $V$ ,*

$$\rho(r) \leq 0 \iff r \in V_\infty \quad \text{and} \quad \rho(r) < 0 \implies r \in \text{int}(V_\infty).$$

*Besides, all representations coincide on the complement of  $\text{int}(V_\infty)$  in  $\mathbb{R}^q$ .*




FIGURE 2. All representations coincide except in  $\text{int}(V_\infty)$ .

PROOF. By positive homogeneity,  $\rho(r) \leq 0$  implies  $\rho(tr) \leq 0 < 1$  (hence  $tr \in V$ ) for all  $t > 0$ ; this implies  $r \in V_\infty$ . Conversely,  $\rho(r) > 0$  implies  $\rho(tr) > 1$  for  $t$  large enough: using  $0 \in V$  again,  $r$  cannot lie in  $V_\infty$ .

To prove the second implication, invoke continuity of  $\rho$ : if  $\rho(r) < 0$ ,  $\rho$  is still negative in a neighborhood of  $r$ , this neighborhood is contained in  $V_\infty$ .

Besides, take a half line emanating from 0 but not contained in  $V_\infty$ ; it certainly meets the boundary of  $V$ , at a point  $\bar{r}$ , which is unique (see, e.g., Hiriart-Urruty and Lemaréchal [17, Remark A.2.1.7]). By (12), every representation  $\rho$  satisfies  $\rho(\bar{r}) = 1$ ; and by positive homogeneity, the value of this representation is determined all along the half line. In other words, all possible representations of  $V$  coincide on the complement  $W$  of  $V_\infty$ ; and by continuity, they coincide also on the closure of  $W$ , which is the complement of  $\text{int}(V_\infty)$ .  $\square$

Figure 2 illustrates the difference between the recession cone (where the gauge is “maximal”) and the rest of the space (where it is *the* representation). Altogether, the gauge appears as the largest representation:

COROLLARY 3.3 (MAXIMALITY OF THE GAUGE). *All representations  $\rho$  of  $V$  satisfy  $\rho \leq \gamma_V$ , with equality on the complement of  $\text{int}(V_\infty)$ .*

*Geometrically, all prepolars  $G$  are contained in the polar of  $V$ :*

$$G^\circ = V \implies G \subset V^\circ.$$

*In particular,  $V$  has a unique representation  $\rho = \gamma_V$  (and a unique prepolar  $V^\circ$ ) whenever  $\text{int}(V_\infty) = \emptyset$ .*

PROOF. Just apply Lemma 3.2, observing from (16) that the gauge is nonnegative.

Geometrically, the inequality between support functions becomes an inclusion: the set  $G$  supported by  $\rho$  is included in the set  $V^\circ$  supported by  $\gamma_V$  (see, e.g., Hiriart-Urruty and Lemaréchal [17, Theorem C.3.3.1]).  $\square$

The next subsection will use the support function  $\sigma_V$ . It is positive on  $\mathbb{R}^q \setminus \{0\}$ , and even more: for some  $\varepsilon > 0$ ,  $V$  contains the ball  $B(\varepsilon)$  centered at 0 of radius  $\varepsilon$ , hence

$$\varepsilon \|d\| = \sigma_{B(\varepsilon)}(d) \leq \sigma_V(d), \quad \text{for all } d \in \mathbb{R}^q. \quad (21)$$

Then  $V^\circ$  is bounded since the relation  $\sigma_V(d) \leq 1$  implies  $\|d\| \leq 1/\varepsilon$ .

**3.3. Smallest representation.** The previous subsection dealt with polarity in the usual sense, viewing the gauge as a special representation. However, we are rather interested in *small* representations. Geometrically, we are interested in small prepolars, and the following definitions are indeed relevant:

$$\begin{cases} \tilde{V}^\circ := \{d \in V^\circ : d^\top r = \sigma_V(d) = 1 \text{ for some } r \in V\}, \\ \hat{V}^\circ := \{d \in V^\circ : \sigma_V(d) = 1\}. \end{cases} \quad (22)$$

From (12),  $\hat{V}^\circ \neq \emptyset$  if  $V$  has a boundary, i.e., if  $V \neq \mathbb{R}^q$ . Obviously,  $\tilde{V}^\circ \subset \hat{V}^\circ$ . Besides, (21) implies that the two sets are bounded. There is a slight difference between the two, suggested by Figure 2 and specified on Figure 3, where the dashed line represents them both. We see that  $d_1$  lies in  $\tilde{V}^\circ$  but not in  $\hat{V}^\circ$ ; and  $d_2$  lies in both. On this example,  $\hat{V}^\circ$  is closed but Figure 5 will show that it need not be so. Although quite similar, we introduce the two sets for technical reasons, when proving that they have the same closed convex hull—which is our required smallest prepolar.

LEMMA 3.4. *The sets in (22) satisfy  $\tilde{V}^\circ \subset \hat{V}^\circ \subset \text{cl}(\tilde{V}^\circ)$ . It follows that  $\hat{V}^\circ$  and  $\tilde{V}^\circ$  have the same closed convex hull. In particular,  $\tilde{V}^\circ \neq \emptyset$  whenever  $\hat{V}^\circ \neq \emptyset$ .*




FIGURE 3. Activity in  $V^\circ$ .

PROOF. The first inclusion is clear. To prove the second inclusion, recall two properties:

- the domain  $\text{dom } \partial\sigma_V$  of a subdifferential is dense in the domain  $\text{dom } \sigma_V$  of the function itself: see, e.g., Hiriart-Urruty and Lemaréchal [17, Theorem E.1.4.2];
- the subdifferential  $\partial\sigma_V(d)$  is the face of  $V$  exposed by  $d$ : see, e.g., Hiriart-Urruty and Lemaréchal [17, Proposition C.3.1.4].

Thus,  $d \notin \tilde{V}^\circ$  implies  $\partial\sigma_V(d) = \emptyset$ ; in other words,  $\tilde{V}^\circ \supset \text{dom } \partial\sigma_V$ . Taking closures,

$$\text{cl } \tilde{V}^\circ \supset \text{cl}(\text{dom } \partial\sigma_V) \supset \text{dom } \sigma_V;$$

the required inclusion follows, since the last set obviously contains  $\hat{V}^\circ$ .

It follows from the second inclusion that

$$\overline{\text{conv}}(\hat{V}^\circ) \subset \overline{\text{conv}}(\text{cl}(\tilde{V}^\circ)).$$

On the other hand, the first inclusion implies that  $\overline{\text{conv}}(\hat{V}^\circ)$  (a closed set) contains the closure of  $\tilde{V}^\circ$ :  $\text{cl}(\tilde{V}^\circ) \subset \overline{\text{conv}}(\hat{V}^\circ)$ . This inclusion remains valid by taking the closed convex hulls:

$$\overline{\text{conv}}(\text{cl}(\tilde{V}^\circ)) \subset \overline{\text{conv}}(\hat{V}^\circ);$$

the two sets coincide. The last statement is clear since the closure of the empty set is the empty set.  $\square$

To help understand this construction, consider the polyhedral case, say  $V = \text{conv}\{p_i\}_i + \text{cone}\{r_i\}_i$ . Then the linear program defining  $\sigma_V(d)$

- has no finite solution if some  $d^\top r_i$  is positive, i.e., if  $d \notin (V_\infty)^\circ$ ,
- is solved at some extreme point  $p_i$  otherwise.

In this situation, the two sets in (22) coincide and are closed; they are a union of hyperplanes of equation  $d^\top p_i = 1$  (facets of  $V^\circ$ ), for  $p_i$  describing the extreme points of  $V$ . Besides, the polar  $V^\circ$  is defined by

$$d^\top p_i \leq 1 \quad \text{and} \quad d^\top r_i \leq 0.$$

EXAMPLE 3.5. For later use, we detail the calculation on a simple instance. Take for  $V$  the polyhedron of Figure 4, defined by the three inequalities

$$\phi \leq 1, \quad \psi \leq 1, \quad \psi \leq 2 + \phi;$$




FIGURE 4. Constructing  $\tilde{V}^\circ$  or  $\hat{V}^\circ$ .

Downloaded from informs.org by [200.89.68.74] on 15 July 2015, at 13:40. For personal use only, all rights reserved.

here  $(\phi, \psi)$  denotes a primal point in  $\mathbb{R}^2$  (we take row vectors for typographical convenience). The two extreme points  $p_1 = (1, 1)$  and  $p_2 = (-1, 1)$  of  $V$  define the two segments (facets of  $V^\circ$ )  $[A, B]$  and  $[B, C]$ .

As for  $V^\circ$ , it has first the two constraints  $d^\top p_i \leq 1$  (yielding the above two segments). Besides, the two extreme rays  $r_1 = (0, -1)$  and  $r_2 = (-1, -1)$  of  $V_\infty$  make two more constraints  $d^\top r_i \leq 0$ , so that  $V^\circ$  is the convex hull of  $A, B, C$ , and  $0$ . If  $V$  had a fourth constraint, say  $\psi \geq -1$ , then  $0$  would be moved down to  $D = (0, -1)$ —and enter  $\tilde{V}^\circ$  and  $\hat{V}^\circ$ .

The closed convex hull thus revealed deserves a notation, as well as its support function: we set

$$V^\bullet := \overline{\text{conv}}(\tilde{V}^\circ) = \overline{\text{conv}}(\hat{V}^\circ) \quad \text{and} \quad \mu_V := \sigma_{V^\bullet} = \sigma_{\tilde{V}^\circ} = \sigma_{\hat{V}^\circ}. \quad (23)$$

For example in Figure 4,  $V^\bullet$  is the triangle  $\text{conv}\{A, B, C\}$ . In fact, the next result shows that  $\mu_V$  is the small representation we are looking for. From now on, we assume  $V \neq \mathbb{R}^q$ , otherwise  $V^\bullet = \emptyset$ ,  $\mu_V \equiv -\infty$ ; a degenerate situation, which lacks interest anyway.

**PROPOSITION 3.6 (SMALLEST REPRESENTATION).** *Any  $\rho$  representing  $V \neq \mathbb{R}^q$  satisfies  $\rho \geq \mu_V$ . Geometrically,  $V^\bullet$  is the smallest closed convex set whose support function represents  $V$ .*

**PROOF.** Our assumption implies that neither  $\hat{V}^\circ$  nor  $\tilde{V}^\circ$  is empty (recall Lemma 3.4). Then take an arbitrary  $d$  in  $\tilde{V}^\circ$ . We have to show that  $d^\top r \leq \rho(r)$  for all  $r \in \mathbb{R}^q$ ; this inequality will be transmitted to the supremum over  $d$ , which is  $\mu_V(r)$ .

*Case 1.* First let  $r$  be such that  $\rho(r) > 0$ . Then  $\bar{r} := r/\rho(r)$  lies in  $V$ , so that  $d^\top \bar{r} \leq \sigma_V(d) = 1$ . In other words,  $d^\top \bar{r} = (d^\top r)/\rho(r) \leq 1$ , which is the required inequality.

*Case 2.* Now let  $r$  be such that  $\rho(r) \leq 0$ , so that  $r \in V_\infty$  by Lemma 3.2. Since  $d \in \tilde{V}^\circ$ , we can take  $r_d \in V$  such that  $d^\top r_d = 1$ . Being exposed,  $r_d$  lies on the boundary of  $V$ : by (12),  $\rho(r_d) = 1$ .

By definition of the recession cone,  $r_d + tr \in V$  for all  $t > 0$  and, by continuity of  $\rho$ ,  $\rho(r_d + tr) > 0$  for  $t$  small enough. Apply Case 1:

$$d^\top r_d + td^\top r = d^\top (r_d + tr) \leq \rho(r_d + tr) \leq \rho(r_d) + t\rho(r),$$

where we have used sublinearity. This proves the required inequality since the first term is  $1 + td^\top r$  and the last one is  $1 + t\rho(r)$ .

The geometric counterpart is proved just as in Corollary 3.3.  $\square$

Thus,  $V$  does have a smallest representation, which is the support function of  $V^\bullet$ . Piecing together our results, we can now fully describe the polarity operation.

**3.4. The set of prepolars.** First of all, it is interesting to link the two extreme representations/prepolars introduced so far, and to confirm the intuition suggested by Figure 4:

**PROPOSITION 3.7.** *Appending 0 to  $V^\bullet$  gives the standard polar:*

$$\gamma_V = \max\{\mu_V, 0\}, \quad \text{i.e., } V^\circ = \overline{\text{conv}}(V^\bullet \cup \{0\}) = [0, 1]V^\bullet.$$

**PROOF.** For  $r \in V_\infty$ ,  $\gamma_V(r) = 0$ , while  $\mu_V(r) \leq 0$  (Proposition 3.6). For  $r \notin V_\infty$ , Lemma 3.2 gives  $\gamma_V(r) = \mu_V(r) > 0$  because  $\gamma_V$  and  $\mu_V$  are two particular representations.

Altogether, the first equality holds. Its geometric counterpart is Hiriart-Urruty and Lemaréchal [17, Theorem C.3.3.2]; and because  $V^\bullet$  is convex compact, its closed convex hull with  $0$  is the sets of  $\alpha d + (1 - \alpha)0$  for  $\alpha \in [0, 1]$ .  $\square$

In summary, the set of representations—or of prepolars—is fully described as follows.

**THEOREM 3.8.** *The representations of  $V$  (a closed convex neighborhood of the origin) are the finite-valued sublinear functions  $\rho$  satisfying*

$$\sigma_{V^\bullet} = \mu_V \leq \rho \leq \gamma_V = \sigma_{V^\circ} = \max\{0, \mu_V\}. \quad (24)$$

*Geometrically, the prepolars of  $V$ , i.e., the sets  $G$  whose support function represents  $V$ , are the sets sandwiched between the two extreme prepolars of  $V$ :*

$$G^\circ = V \iff V^\bullet \subset \overline{\text{conv}}(G) \subset V^\circ = \overline{\text{conv}}(V^\bullet \cup \{0\}) = [0, 1]V^\bullet.$$




FIGURE 5. Trouble appears if the neighborhood has no asymptote.

PROOF. In view of Corollary 3.3 and Propositions 3.6, 3.7, we just have to prove that a  $\rho$  satisfying (24) does represent  $V$ . Indeed, if  $r \in V$  then  $\rho(r) \leq \gamma_V(r) \leq 1$ ; if  $r \notin V$ , then  $1 < \mu_V(r) \leq \rho(r)$ . The geometric counterpart is again standard calculus with support functions.  $\square$

We end this section with a deeper study of prepolars, which will be useful in the sequel. The next result introduces the polar cone  $(V_\infty)^\circ$ . When  $G$  is a cone, positive homogeneity can be used to replace the right-hand side “1” in (19) by any positive number, or even by “0”: in particular,

$$V_\infty^\circ := (V_\infty)^\circ = \{r \in \mathbb{R}^q : \sigma_{V_\infty}(r) \leq 0\}. \tag{25}$$

The notation  $V_\infty^\circ$  is used for simplicity, although it is somewhat informal;  $(V_\infty)^\circ$  and  $(V^\circ)_\infty$  differ, the latter is  $\{0\}$  since  $V^\circ$  is bounded.

LEMMA 3.9 (ADDITIONAL PROPERTIES OF PREPOLARS). *With the notation (22), (23), (25),*

- (i)  $V_\infty^\circ$  is the closure of  $\text{dom } \sigma_V$ , and
- (ii)  $\mathbb{R}_+ \hat{V}^\circ = \mathbb{R}_+ V^\bullet = \mathbb{R}_+ V^\circ = \text{dom } \sigma_V$ .

PROOF. First of all, let  $d \notin V_\infty^\circ$ : there is  $r \in V_\infty$  ( $\mathbb{R}_+ r \in V$ ) and  $d^\top r > 0$ ; then  $d^\top(tr) \rightarrow +\infty$  for  $t \rightarrow +\infty$  and  $\sigma_V(d)$  cannot be finite, i.e.,  $d \notin \text{dom } \sigma_V$ . Thus,  $\text{dom } \sigma_V \subset V_\infty^\circ$ ; hence  $\text{cl}(\text{dom } \sigma_V) \subset V_\infty^\circ$  because  $V_\infty^\circ$  is closed.

To prove the converse inclusion, take  $r \notin (\text{dom } \sigma_V)^\circ$ : there is  $d$  such that  $\sigma_V(d) < +\infty$  and  $d^\top r > 0$ . Then  $d^\top(tr) \rightarrow +\infty$  when  $t \rightarrow +\infty$ ; if  $r$  were in  $V_\infty$ , then  $tr$  would lie in  $V$  and  $\sigma_V(d)$  would be  $+\infty$ , a contradiction. Thus we have proved  $V_\infty^\circ \subset (\text{dom } \sigma_V)^\circ$ . Taking polars and knowing that  $\text{dom } \sigma_V$  is a cone,  $V_\infty^\circ \supset (\text{dom } \sigma_V)^\circ = \text{cl}(\text{dom } \sigma_V)$  (see Hiriart-Urruty and Lemaréchal [17, Proposition A.4.2.6]). This proves (i).

To prove (ii), observe first that  $\hat{V}^\circ \subset V^\bullet \subset V^\circ \subset \text{dom } \sigma_V$ ; and because  $\text{dom } \sigma_V$  is a cone,

$$\mathbb{R}_+ \hat{V}^\circ \subset \mathbb{R}_+ V^\bullet \subset \mathbb{R}_+ V^\circ \subset \text{dom } \sigma_V. \tag{26}$$

On the other hand, take  $0 \neq d \in \text{dom } \sigma_V$ , so that  $\sigma_V(d) > 0$  by (21) and  $(1/\sigma_V(d))d \in \hat{V}^\circ$ :  $d \in \mathbb{R}_+ \hat{V}^\circ$ . Since 0 also lies in  $\mathbb{R}_+ \hat{V}^\circ$ , we do have  $\text{dom } \sigma_V \subset \mathbb{R}_+ \hat{V}^\circ$ ; (26) is actually a chain of equalities. To complete the proof, observe from Proposition 3.7 that  $\mathbb{R}_+ V^\circ = \mathbb{R}_+ V^\bullet$ .  $\square$

Beware that really pathological prepolars can exist, Figure 5 illustrates a well-known situation. Its left part displays the parabolic neighborhood  $V = P \subset \mathbb{R}^2$  defined by the constraint  $\psi \leq 1 - \frac{1}{2}\phi^2$ . A direction  $d = (u, v)$  with  $v > 0$  exposes the point  $r(d)$ . When  $v \downarrow 0$ , the component of  $r(d)$  along  $d$  (namely,  $\phi$ ) goes to  $+\infty$ , which does bring trouble. Computing  $r(d)$  is an exercise resulting in

$$\sigma_P(d) = \sigma_P(u, v) = \begin{cases} 0 & \text{if } d = 0, \\ v + \frac{u^2}{2v} & \text{if } v > 0, \\ +\infty & \text{if } v \leq 0; \end{cases} \tag{27}$$

two phenomena are then revealed.

— First,  $\hat{V}^\circ$  is defined by the equation

$$v + \frac{u^2}{2v} = 1, \quad \text{i.e., } 2(v^2 - v) + u^2 = 0.$$

This is an ellipse passing through the origin (right part of Figure 5); yet 0 cannot lie in  $\hat{V}^\circ$ , since  $\sigma_P(0) = 0 \neq 1$ . Thus,  $\hat{P}^\circ$  is not closed and, more importantly,  $0 \in P^\bullet$ .

— The second phenomenon is a violent discontinuity of  $\sigma_P$  at 0. In fact, fix  $\alpha > 0$  and let  $d_k = (\alpha/k, 1/k^2)$ ; then  $d_k \rightarrow 0$ , while  $\sigma_P(d_k) \rightarrow \alpha^2/2$ , an arbitrary positive number.

Downloaded from informs.org by [200.89.68.74] on 15 July 2015, at 13:40. For personal use only, all rights reserved.



Both phenomena are due to (local) unboundedness of  $\sigma_p$  on its domain, which is thus not closed; if  $(u_k, v_k) \in \text{dom } \sigma_p$  tends to any  $(u, 0)$  with  $u \neq 0$ , then  $\sigma_p(u_k, v_k) \rightarrow +\infty$ . Ruling out such a behaviour brings additional useful properties.

COROLLARY 3.10 (SAFE PREPOLARS). *If  $0 \notin V^\bullet$ , then*

$$\mathbb{R}_+ \hat{V}^\circ = \mathbb{R}_+ V^\bullet = \mathbb{R}_+ V^\circ = \text{dom } \sigma_V = V_\infty^\circ \quad (28)$$

and  $\text{int } V_\infty^\circ \neq \emptyset$  (the polar  $V_\infty^\circ$  is a so-called pointed cone).

PROOF. When  $0 \notin V^\bullet$ ,  $\mathbb{R}_+ V^\bullet$  is closed (Hiriart-Urruty and Lemaréchal [17, Proposition A.1.4.7]). Then apply Lemma 3.9: by (ii)  $\text{dom } \sigma_V$  is closed and (28) follows from (i).

Now we separate 0 from  $V^\bullet$ : there is some  $r$  such that  $\sigma_{V^\bullet}(r) < 0$ . By continuity of the finite-valued convex function  $\sigma_{V^\bullet}$ , this inequality is still valid in a neighborhood of  $r$ :  $\sigma_{V^\bullet} \leq 0$  over some nonzero ball  $B$  around  $r$ . By Lemma 3.9(ii),

$$\sigma_{V_\infty^\circ}(d) = \sigma_{\mathbb{R}_+ V^\bullet}(d) = \sup_{t \geq 0} \sup_{d \in V^\bullet} t d^\top r = \sup_{t \geq 0} t \sigma_{V^\bullet}(d),$$

so that  $\sigma_{V_\infty^\circ}$  enjoys the same property: by (25),  $B$  is contained in  $(V_\infty^\circ)^\circ$ . Hiriart-Urruty and Lemaréchal [17, Proposition A.4.2.6] finishes the proof.  $\square$

Property (28) means closedness of  $\text{dom } \sigma_V$  and is rather instrumental. We mention another simple assumption implying it.

PROPOSITION 3.11. *If  $V = U + V_\infty$ , where  $U$  is bounded, then  $\text{dom } \sigma_V = V_\infty^\circ$ .*

PROOF. The support function of a sum is easily seen to be the sum of support functions:  $\sigma_V = \sigma_U + \sigma_{V_\infty}$ . Every  $d \in V_\infty^\circ$  then satisfies  $\sigma_V(d) = \sigma_U(d)$ , a finite number when  $U$  is bounded.  $\square$

Let us put this section in perspective. The traditional gauge theory defines via (16), (19) the polarity correspondence  $V \leftrightarrow V^\circ$  for compact convex neighborhoods of the origin. We generalize it to unbounded neighborhoods, whose standard gauge is replaced via Definition 2.6 by their family of representations. Each representation  $\rho$ , which may assume negative values, gives rise to  $\partial\rho(0)$ —which we call a prepolar of  $V$ . Theorem 3.8 establishes the existence of a largest element (the usual polar  $V^\circ$ ) and of a smallest element ( $V^\bullet$ ) in the family of (closed convex) prepolars of  $V$ . Gauge theory is further generalized in Zaffaroni [24], in which 0 may lie on the boundary of  $V$ . Our stricter framework allows a finer analysis of the smallest prepolar; in particular, the property  $0 \notin V^\bullet$  helps avoid nasty phenomena.

**4. Minimal CGF's, maximal  $S$ -free sets.** Remembering Remark 1.4, our goal in this section is to study the concept of minimality for CGF's. Geometrically, we study the concept of maximality for  $S$ -free sets. In fact, the two concepts are in correspondence via (14); but a difficulty arises because the reverse inclusion does not hold in (14). As a result, several definitions of minimality and maximality are needed.

**4.1. Minimality, maximality.** In our quest for small CGF's, the following definition is natural.

DEFINITION 4.1 (MINIMALITY). A CGF  $\rho$  is called minimal if the only possible CGF  $\rho' \leq \rho$  is  $\rho$  itself.

Knowing that a CGF  $\rho$  represents  $V(\rho)$  and that  $\mu_{V(\rho)} \leq \rho$  represents the same set, a minimal CGF is certainly a smallest representation:

$$\rho \text{ is a minimal CGF} \implies \rho = \mu_{V(\rho)} = \sigma_{V(\rho)^\bullet}. \quad (29)$$

In addition,  $V(\rho)$  must of course be a special  $S$ -free set when  $\rho$  is minimal. Take for example  $S = \{1\} \subset \mathbb{R}$ ,  $V = [-1, +1]$ ; then  $\rho(r) := |r|$  is the smallest (because unique) representation of  $V$  but is not minimal:  $\rho'(r) := \max\{0, r\}$  is also a CGF, representing  $V' = ]-\infty, +1]$ .

From (14), a smaller  $\rho$  describes a larger  $V$ ; so Definition 4.1 has its geometrical counterpart:

DEFINITION 4.2 (MAXIMALITY). An  $S$ -free set  $V$  is called maximal if the only possible  $S$ -free set  $V' \supset V$  is  $V$  itself.

The two objects are indeed related.

PROPOSITION 4.3. *If  $V$  is a maximal  $S$ -free set, then its smallest representation  $\mu_V$  is a minimal CGF.*

PROOF. Take a CGF  $\rho'$ , representing the  $S$ -free set  $V' = V(\rho')$ . If  $\rho' \leq \mu_V$ , then  $V' \supset V$ ; and if  $V$  is maximal,  $V' = V$ . Then  $\rho' \geq \mu_V = \mu_{V'}$  by Proposition 3.6.  $\square$






FIGURE 6. A linear CGF is always maximal.

Besides, these objects do exist.

**THEOREM 4.4.** *Every  $S$ -free set is contained in a maximal  $S$ -free set. It follows that there exists a maximal  $S$ -free set and a minimal CGF.*

**PROOF.** Let  $V$  be an  $S$ -free set. In the partially ordered family  $(\mathcal{F}, \subset)$  of all  $S$ -free sets containing  $V$ , let  $\{W_i\}_{i \in I}$  be a totally ordered subfamily (a chain) and define  $W := \bigcup_{i \in I} W_i$ . Clearly,  $W$  is a neighborhood of the origin; its convexity is easily established, let us show that its closure is  $S$ -free.

Remember from Hiriart-Urruty and Lemaréchal [17, Theorem C.3.3.2(iii)] that the support function of a union is the (closure of the) supremum of the support functions:

$$\sigma_{\text{int}(W)} = \sigma_W = \text{cl}\left(\sup_{i \in I} \sigma_{W_i}\right) = \text{cl}\left(\sup_{i \in I} \sigma_{\text{int}(W_i)}\right) = \sigma_{\bigcup_i \text{int}(W_i)}.$$

Having the same support function, the two open convex sets  $\text{int}(W)$  and  $\bigcup_i \text{int}(W_i)$  coincide:  $r \in \text{int}(W)$  means  $r \in \text{int}(W_i)$  for some  $i$ ; because  $W_i$  is  $S$ -free,  $r \notin S$  and our claim is proved. Thus, the chain  $\{W_i\}$  has an upper bound in  $\mathcal{F}$ ; in view of Zorn’s lemma,  $\mathcal{F}$  has a maximal element.

Now (1b) implies that a ball centered at 0 with a small enough radius is  $S$ -free; and there exists a maximal  $S$ -free set containing it. Proposition 4.3 finishes the proof.  $\square$

The maximal  $S$ -free sets can be explicitly described for some special  $S$ ’s:  $\mathbb{Z}^q$  (Lovász [21]), the intersection of  $\mathbb{Z}^q$  with an affine subspace (Basu et al. [5]), with a rational polyhedron (Basu et al. [6]), or with an arbitrary closed convex set (Morán and Dey [22], Averkov [2]). Unfortunately, the “duality” between minimal CGF’s and maximal  $S$ -free sets is deceiving, as the two definitions do not match: the set represented by a minimal CGF need not be maximal. In fact, when  $\rho$  is linear, the property introduced in Definition 4.1 holds vacuously: no sublinear function can properly lie below a linear function. Thus, a linear CGF  $\rho$  is always minimal; yet, a linear  $\rho$  represents a neighborhood  $V(\rho)$  (a half space), which is  $S$ -free but has no reason to be maximal. See Figure 6 with  $n = 1$ , the set  $V = ]-\infty, 1]$  (represented by  $\rho(x) = x$ ) is  $\{2\}$ -free but is obviously not maximal.

A more elaborate example reveals the profound reason underlying the trouble: for an  $S$ -free set  $W$  containing  $V$ ,  $\mu_W$  need not be comparable to  $\mu_V$ .

**EXAMPLE 4.5.** In Example 3.5, take for  $S$  the union of the three lines with respective equations

$$\phi = 1, \quad \psi = 1, \quad \psi = 2 + \phi,$$

so that  $V$  is clearly maximal  $S$ -free.

Now shrink  $V$  to  $V_t$  (left part of Figure 7) by moving its right vertical boundary to  $\phi \leq 1 - t$ . Then  $A$  is moved to  $A_t = (1/(1 - t), 0)$ ; there is no inclusion between the new  $V_t^\bullet = \text{conv}\{A_t, B, C\}$  and the original  $V^\bullet = \text{conv}\{A, B, C\}$ ; this is the key to our example.

Let us show that  $\mu_{V_t}$  is minimal, even though  $V_t$  is not maximal. Take for this a CGF  $\rho \leq \mu_{V_t}$ , which represents an  $S$ -free set  $W$ ; by (14),  $W \supset V_t$ . We therefore have

$$\sigma_{W^\bullet} = \mu_W \leq \rho \leq \mu_{V_t} = \sigma_{V_t^\bullet}, \quad \text{i.e., } W^\bullet \subset V_t^\bullet$$

and we proceed to show that equality does hold, i.e., the three extreme points of  $V_t^\bullet$  do lie in  $W^\bullet$ .




FIGURE 7. The mapping  $V \mapsto V^\bullet$  is not monotonic.

— If  $A_i \notin W^\bullet$ , the right part of Figure 7 shows that  $W^\bullet$  is included in the open upper half space. Knowing that

$$W = (W^\bullet)^\circ = \{r: d^\top r \leq 1 \text{ for all } d \in W^\bullet\},$$

this implies that  $W_\infty$  has a vector of the form  $r_A = (\varepsilon, -1)$  ( $\varepsilon > 0$ );  $W$  cannot be  $S$ -free.

— If  $C \notin W^\bullet$ , there is  $r_C \in \mathbb{R}^2$  such that  $C^\top r_C > \sigma_{W^\bullet}(r_C) = \mu_W(r_C)$  (we denote also by  $C$  the 2-vector representing  $C$ ). For example  $r_C = (-2, 0) \in \text{bd}(V)$  (see the right part of Figure 7), so that

$$C^\top r_C = 1 > \sigma_{W^\bullet}(-2, 0) = \mu_W(-2, 0).$$

By continuity,  $\mu_W(-2 - \varepsilon, 0) \leq 1$  for  $\varepsilon > 0$  small enough. Since  $\mu_W$  represents  $W$ , this implies that  $(-2 - \varepsilon, 0) \in W$ ;  $W$  (which contains  $V_i$ ) is not  $S$ -free.

— By the same token, we prove that  $B \in W^\bullet$  (the separator  $r_B = (0, 1) \in \text{bd}(V)$  does the job).

We have therefore proved that  $W^\bullet = V_i^\bullet$ , i.e.,  $\mu_W = \mu_{V_i}$ , i.e.,  $\mu_{V_i}$  is minimal.

The next section makes a first step toward a theory relating small CGF's and large  $S$ -free sets.

**4.2. Strong minimality, asymptotic maximality.** First, let us give a name to those minimal CGF's corresponding to maximal  $S$ -free sets.

**DEFINITION 4.6 (STRONGLY MINIMAL CGF).** A CGF  $\rho$  is called *strongly minimal* if it is the smallest representation of a maximal  $S$ -free set.

The strongly minimal CGF's can be characterized without any reference to the geometric space.

**PROPOSITION 4.7.** A CGF  $\rho$  is strongly minimal if and only if, for every CGF  $\rho'$ ,

$$\rho' \leq \max\{0, \rho\} [= \gamma_{V(\rho)} = \sigma_{V(\rho)^\circ}] \implies \rho' \geq \rho. \quad (30)$$

**PROOF.** Take first a maximal  $V$ . Every CGF  $\rho' \leq \gamma_V$  represents an  $S$ -free set  $V'$ , which contains  $V$ —see (13)—so that  $V' = V$  by maximality, i.e.,  $\rho'$  represents  $V$  as well; hence  $\rho' \geq \mu_V$  by Proposition 3.6. Thus,  $\rho (= \mu_V)$  satisfies (30).

Now let  $\rho$  satisfy (30), we have to show that  $V := V(\rho)$  is maximal. Taking in particular  $\rho' = \mu_V$  in (30) shows that  $\rho$  must equal  $\mu_V$ . Let  $V' \supset V$  be  $S$ -free; we have  $(V')^\circ \subset V^\circ$ , i.e.,

$$\gamma_{V'} = \sigma_{(V')^\circ} \leq \sigma_{V^\circ} = \gamma_V = \max\{0, \rho\}.$$

Now  $\rho' := \gamma_{V'}$  is a CGF, so  $\rho' \geq \rho = \mu_V$  by (30); by Theorem 3.8,  $\rho'$  represents not only  $V'$  but also  $V$ , i.e.,  $V' = V$ :  $V$  is maximal.  $\square$

In §3 we have systematically developed the geometric counterpart of representations; this exercise can be continued here. In fact, the concept of minimality involves two properties from a sublinear function:

- it must be the *smallest* representation of some neighborhood  $V$ —remember (29),
- this neighborhood must enjoy some maximality property.

In view of the first property, a CGF can be imposed to be not only sublinear but also to support a set that is a *smallest* prepolar. Then Definition 4.1 has a geometric counterpart: minimality of  $\rho = \mu_V = \sigma_{V^\circ}$  means

$$\begin{array}{lll} G' \subset V^\bullet & \text{and} & (G')^\circ \text{ is } S\text{-free} \implies G' = V^\bullet, \text{ i.e., } (G')^\circ = V. \\ [\rho' = \sigma_{G'} \leq \rho] & & [\rho' \text{ is a CGF}] \quad [\rho' = \rho] \end{array}$$

Likewise for Definition 4.6: strong minimality of  $\rho = \gamma_V = \sigma_{V^\circ}$  means

$$\begin{array}{lll} G' \subset V^\circ & \text{and} & (G')^\circ \text{ is } S\text{-free} \implies G' \supset V^\bullet, \text{ i.e., } (G')^\circ \subset V. \\ [\rho' = \sigma_{G'} \leq \gamma_V] & & [\rho' \text{ is a CGF}] \quad [\rho' \geq \rho] \end{array}$$

These observations allow some more insight into the  $(\cdot)^\bullet$  operation.

**PROPOSITION 4.8.** Let  $\rho = \mu_V = \sigma_{V^\circ}$  be a minimal CGF. If an  $S$ -free set  $W$  satisfies  $W^\bullet \subset V^\bullet$ , then  $W = V$ .

**PROOF.** The smallest representation  $\rho' := \mu_W = \sigma_{W^\circ}$  of the  $S$ -free set  $W$  is a CGF; and from monotonicity of the support operation,  $\rho' \leq \rho$ . Then minimality of  $\rho$  implies  $\rho' = \rho$ , i.e.,  $W^\bullet = V^\bullet$ , an equality transmitted to the polars:  $W = (W^\bullet)^\circ = (V^\bullet)^\circ = V$ .  $\square$

This result confirms that nonequivalence between minimal CGF’s and maximal  $S$ -free sets comes from nonmonotonicity of the mapping  $V \mapsto V^\bullet$ —or of  $V \mapsto \mu_V$ . To construct Example 4.5, we do need a  $W \supset V$  such that  $W^\bullet \not\subset V^\bullet$ .

Then comes a natural question: how maximal are the  $S$ -free sets represented by minimal CGF’s? For this, we introduce one more concept.

DEFINITION 4.9. An  $S$ -free set  $V$  is called *asymptotically maximal* if every  $S$ -free set  $V' \supset V$  satisfies  $V'_\infty = V_\infty$ .

It allows a partial answer to the question.

THEOREM 4.10 (MINIMAL  $\Rightarrow$  ASYMPTOTICALLY MAXIMAL). *The  $S$ -free set represented by a minimal CGF is asymptotically maximal.*

PROOF. Let  $\mu_V$  be a minimal CGF and take an  $S$ -free set  $V' \supset V$ . Introduce the set  $G := V^\bullet \cap (V'_\infty)^\circ$ . Inclusions translate to inequalities between support functions:

$$\sigma_G \leq \sigma_{V^\bullet} = \mu_V, \tag{31}$$

and we proceed to prove that this is actually an equality. Let us compute the set  $W := G^\circ$  represented by  $\sigma_G$ . The support function of an intersection is obtained via an inf-convolution (formula (3.3.1) in Hiriart-Urruty and Lemaréchal [17, Chapter C] for example):  $\sigma_G(\cdot)$  is the closure of the function

$$r \mapsto \inf \{ \sigma_{V^\bullet}(r_1) + \sigma_{(V'_\infty)^\circ}(r_2) : r_1 + r_2 = r \}.$$

In this formula,  $\sigma_{V^\bullet} = \mu_V$  and the support function of the closed convex cone  $(V'_\infty)^\circ$  is the indicator of its polar  $V'_\infty$ : the above function is

$$r \mapsto \inf \{ \mu_V(r_1) : r_1 + r_2 = r, r_2 \in V'_\infty \}.$$

Now use (12): because  $\sigma_G$  represents  $W$ , to say that  $r \in \text{int}(W)$  is to say that the above infimum is strictly smaller than 1, i.e., that there are  $r_1, r_2$  such that

$$r_1 + r_2 = r, r_2 \in V'_\infty, \mu_V(r_1) < 1, \quad \text{i.e., } r_1 + r_2 = r, r_2 \in V'_\infty, r_1 \in \text{int } V.$$

In a word,

$$\text{int}(W) = V'_\infty + \text{int}(V) \supset \text{int}(V) \ni 0,$$

where we have used the property  $0 \in V'_\infty$ . Remembering the inclusion  $V \subset V'$  and the definition of a recession cone, we also have

$$\text{int}(W) = V'_\infty + \text{int}(V) \subset V'_\infty + \text{int}(V') \subset V'_\infty + V' \subset V'.$$

Altogether,  $0 \in \text{int}(W) \subset \text{int}(V')$ . As a result,  $W (= G^\circ)$  is an  $S$ -free closed convex neighborhood of the origin: its representation  $\sigma_G$  is a CGF and minimality of  $\mu_V = \sigma_{V^\bullet}$  implies with (31) that  $\sigma_G = \sigma_{V^\bullet}$ .

By closed convexity of both sets  $V^\bullet$  and  $G = V^\bullet \cap (V'_\infty)^\circ$ , this just means  $G = V^\bullet$ , i.e.,  $(V'_\infty)^\circ \supset V^\bullet$ . By polarity,  $V'_\infty \subset (V^\bullet)^\circ = V$  (invoke Theorem 3.8). The cone  $V'_\infty$ , contained in the neighborhood  $V$ , is also contained in its recession cone:  $V'_\infty \subset V_\infty$ . Since the converse inclusion is clear from  $V' \supset V$ , we have proved  $V'_\infty = V_\infty$ :  $V$  is asymptotically maximal.  $\square$

**5. Favourable cases.** Despite Example 4.5, a number of papers have established the equivalence between maximal  $S$ -free sets and minimal CGF’s, for various forms of  $S$ . This equivalence is indeed known to hold in a number of situations:

- (a) when  $S$  is a finite set of points in  $\mathbb{Z}^q - b$ ; see Johnson [18] and more recently Dey and Wolsey [12];
- (b) when  $S$  is the intersection of  $\mathbb{Z}^n$  with an affine space; this was considered in Borozan and Cornuéjols [8] and Basu et al. [5];
- (c) when  $S = P \cap (\mathbb{Z}^q - b)$  for some rational polyhedron  $P$ ; and this was considered in Dey and Wolsey [12] and Basu et al. [6].

Accordingly, we investigate in this section the question: when does minimality imply strong minimality? So we consider an  $S$ -free set  $V$ , whose smallest representation  $\mu_V = \sigma_{V^\bullet}$  is minimal, hence  $V$  is asymptotically maximal (Theorem 4.10); we want to exhibit conditions under which  $V$  is maximal. We denote by  $L = (-V_\infty) \cap V_\infty$  the *lineality space* of  $V$  (the largest subspace contained in the closed convex cone  $V_\infty$ ) and our result is the following.




FIGURE 8. Constructing in  $S$  an unbounded sequence “tending to”  $V$ .

**THEOREM 5.1.** Suppose  $0 \in \bar{S} := \overline{\text{conv}}(S)$ . A minimal  $\mu_V$  is strongly minimal whenever one of the following two properties (i) and (ii) holds:

- (i)  $V_\infty \cap \bar{S}_\infty = \{0\}$  (in particular  $S$  bounded),
- (ii)  $\left\{ \begin{array}{l} \text{(ii)}_1 \bar{S} = U + \bar{S}_\infty \text{ with } U \text{ bounded, and} \\ \text{(ii)}_2 V_\infty \cap \bar{S}_\infty = L \cap \bar{S}_\infty. \end{array} \right.$

This theorem generalizes the above-mentioned results: (i) is a weakening of (a) and (ii) weakens (b) or (c). Note that (ii)<sub>2</sub> generalizes (i) (to an unbounded  $V_\infty \cap \bar{S}$ ); the price to pay is assumption (ii)<sub>1</sub>, whose role is to exclude an asymptotic behaviour of  $\bar{S}$  similar to that of  $P$  in Figure 5 (see Proposition 3.11).

However, the interesting point does not lie in the above assumptions (a)–(ii). Recalling that the whole issue lies in unboundedness of  $V$ , our proof of Theorem 5.1 uses Theorem 4.10 as follows. Starting from an  $S$ -free set  $V$ , which is asymptotically maximal but not maximal, we construct a sequence of neighborhoods  $V^k$  satisfying  $V_\infty \supsetneq V_\infty$ . Then  $V^k$  is not  $S$ -free: there is some  $r^k \in S \cap \text{int}(V^k)$ ; see Figure 8.

Besides, our construction is organized in such a way that  $V^k$  “tends to”  $V$  and, by nonmaximality of  $V$ ,  $r^k$  is unbounded but “tends to”  $V$ . More precisely,

the cluster points of the normalized sequence  $\{r^k\}$  lie in  $\bar{S}_\infty \cap V_\infty$ .

Decomposing  $r^k = l^k + u^k$  along  $L$  and  $L^\perp$ , we also prove that  $u^k$  is unbounded but “tends to”  $\bar{S} \cap L^\perp$ , more precisely

the cluster points of the normalized sequence  $\{u^k\}$  lie in  $\bar{S}_\infty \cap L^\perp$ .

We believe that these are key properties of nonmaximal  $S$ -free sets. Having established them, the whole business is to find appropriate assumptions under which existence of our unbounded sequences is impossible; (a)–(ii) are such *ad hoc* assumptions.

Obtaining  $r^k$  and  $u^k$  is a fairly complicated operation, which we divide into a series of lemmas. For a reason that will appear in (39), we may assume  $0 \notin V^\bullet$ . Then we enlarge  $V$  to  $V^k$  by chopping off a bit of  $V^\bullet$  as follows. Take an extreme ray  $\mathbb{R}_+ d_V$  of  $V_\infty^\circ$ . By (28), its intersection with  $V^\bullet$  is a nonempty segment  $[d_V, t_V d_V]$ , with  $1 \leq t_V < +\infty$ . Given a positive integer  $k$ , we introduce the open neighborhood of  $[d_V, t_V d_V]$ :

$$N^k := [d_V, t_V d_V] + B(0, 1/k) = \bigcup_{1 \leq t \leq t_V} B(td_V, 1/k), \tag{32}$$

where  $B(d, \delta)$  is the open ball of center  $d$  and radius  $\delta$ . We remove  $N^k$  from  $V^\bullet$ , thus obtaining a set  $C$ , closed hence compact; its convex hull

$$G^k := \text{conv } C, \quad \text{with } C := V^\bullet \setminus N^k = \{d \in V^\bullet: \|d - td_V\| \geq 1/k \text{ for all } t \in [1, t_V]\} \tag{33}$$

is convex compact. Figure 9 illustrates our construction.

Note for future use that the distance from every  $d \in [d_V, t_V d_V]$  to  $C$  does not exceed  $1/k$ ; and the same holds for  $G^k \supset C$ . Formally,

$$\forall \bar{d} \in [d_V, t_V d_V], \exists d_k \in G^k \text{ such that } \|d_k - \bar{d}\| \leq 1/k. \tag{34}$$




FIGURE 9. Chopping off  $V^\bullet$  near an extreme ray.

REMARK 5.2. The above construction would become substantially simpler, and  $N^k$  would reduce to the open ball  $B(d_V, 1/k)$ , if  $V^\bullet \cap \mathbb{R}_+ d_V$  reduced to a singleton, i.e., if  $t_V = 1$ ; but this property need not hold when  $\sigma_V$  is not continuous.

To make a counterexample, start from the parabola of Figure 5. We already know that  $\sigma_p(d_k)$  can tend to any nonnegative value when  $d_k \rightarrow 0$ . However  $0 \in P^\bullet$ ; alternatively, the domain of  $\sigma_p$  is not closed. In fact, we need a discontinuous sublinear function that is locally bounded on its domain—and this requires three variables. Thus, we first bound  $\sigma_p$  by defining

$$f(d) := 1 + \begin{cases} \sigma_p(d) & \text{if } \sigma_p(d) \leq 1, \\ +\infty & \text{otherwise,} \end{cases}$$

(the “1+” appearing above is just aimed at getting 0 in the interior of  $V$ ). Although no longer positively homogeneous,  $f$  is still convex, its domain is the compact convex set  $P^\bullet$ , on which  $1 \leq f \leq 2$ ; when  $d_k \in P^\bullet$  tends to 0,  $f(d_k)$  can tend to any value in  $[1, 2]$ . To complete the construction, we take the so-called *perspective* of  $f$ :

$$\mathbb{R}^2 \times \mathbb{R} \ni (d, w) \mapsto \sigma(d, w) := \begin{cases} wf(d/w) & \text{if } w > 0, \\ 0 & \text{if } (d, w) = (0, 0), \\ +\infty & \text{otherwise,} \end{cases}$$

whose positive homogeneity is clear. Actually,  $\sigma$  is known to be convex and to support a closed convex set  $V$ ; see Hiriart-Urruty and Lemaréchal [17, §B.2.2] (in particular Remark 2.2.3), where our  $(d, w)$  is called  $(x, u)$ . Besides, the property  $f \geq 1$  implies that  $V$  is a neighborhood of the origin; remember (21).

Now take  $(d, w) \in \hat{V}^\circ \subset \text{dom } \sigma$ , so that  $d' := (d/w) \in \text{dom } f$  and  $w > 0$ . Then use positive homogeneity:

$$1 = \sigma(d, w) \implies \frac{1}{w} = \sigma(d', 1) = f(d') \in [1, 2] \implies w \geq \frac{1}{2}.$$

Thus,  $\hat{V}^\circ$  is separated from the origin (by the hyperplane  $w \geq \frac{1}{2}$ ) and this property is transmitted to its closed convex hull  $V^\bullet$ . On the other hand,  $\sigma$  inherits the discontinuities of  $f$ . In fact, choose  $\alpha \in [1, 2]$  and construct a sequence  $\{d_k\}$  in  $\text{dom } f$  tending to 0, such that  $f(d_k) \rightarrow \alpha$ . Since  $\sigma(d_k, 1) = f(d_k) > 0$ , positive homogeneity gives

$$\sigma\left(\frac{d_k}{f(d_k)}, \frac{1}{f(d_k)}\right) = 1, \quad \text{hence } \left(\frac{d_k}{f(d_k)}, \frac{1}{f(d_k)}\right) \in \hat{V}^\circ.$$

Pass to the limit,

$$\left(\frac{d_k}{f(d_k)}, \frac{1}{f(d_k)}\right) \rightarrow \left(0, \frac{1}{\alpha}\right) \in \text{cl } \hat{V}^\circ \subset V^\bullet.$$

Since  $\alpha$  was arbitrary in  $[1, 2]$ , the intersection of  $V^\bullet$  with the ray  $\{0\} \times \mathbb{R}_+$  contains the whole segment  $\{0\} \times [\frac{1}{2}, 1]$ .

Viewing  $G^k$  of (33) as a prepolars, we set

$$V^k := (G^k)^\circ.$$

Of course,  $V^\bullet \supset G^{k+1} \supset G^k$  and  $V \subset V^{k+1} \subset V^k$ . The closed convex neighborhood  $V^k$  enjoys all of the properties listed in §3, in particular those coming from  $0 \notin G^k$ .

LEMMA 5.3 (ENLARGING  $V_\infty$ ). Assume  $0 \notin V^\bullet$ ; let  $\mathbb{R}_+ d_V$  be an extreme ray of  $V_\infty^\circ$  and assume that  $\mathbb{R}_+ d_V \subsetneq V_\infty^\circ$  ( $\mathbb{R}_+ d_V$  is properly contained in  $V_\infty^\circ$ ). Given an integer  $k > 0$ , construct  $N^k, G^k, V^k$  as above. Then  $G^k \neq \emptyset$  for  $k$  large enough (say  $k \geq k_0$ ) and

- (i)  $V_\infty \subsetneq V_\infty^k$  for  $k \geq k_0$ ,
- (ii)  $\bigcap_{k \geq k_0} V^k = V$ .

PROOF. If  $G^k$  were empty for all  $k$ , we would have  $V^\bullet \subset N^k$  for all  $k$ , hence  $V^\bullet$  would reduce to  $[d_V, t_V d_V]$ . In view of (28), this would imply  $\mathbb{R}_+ d_V = V_\infty^\circ$ , which our assumption rules out.

Every  $d \in G^k$  is a convex combination  $\sum_i \alpha_i d_i$  with each  $d_i$  in  $V^\bullet \setminus N^k \subset V_\infty^\circ$ . None of these  $d_i$ 's can lie in  $[d_V, t_V d_V] \subset N^k$ , and none of their convex combinations either because of extremality of  $\mathbb{R}_+ d_V$ . We conclude that

$$G^k \cap [d_V, t_V d_V] = \emptyset. \tag{35}$$

Now, we see from Theorem 3.8 that

$$\mathbb{R}_+(V^k)^\bullet \subset \mathbb{R}_+ G^k \subset \mathbb{R}_+(V^k)^\circ;$$

but from Proposition 3.7, this is actually a chain of equalities:

$$\mathbb{R}_+(V^k)^\bullet = \mathbb{R}_+ G^k. \tag{36}$$

Besides,  $(V^k)^\bullet \subset G^k \subset V^\bullet$ , hence  $0 \notin (V^k)^\bullet$  and we can apply (28) to  $V^k$ . Then we write

$$\begin{aligned} (V_\infty^k)^\circ &= \mathbb{R}_+(V^k)^\bullet && [(28)] \\ &= \mathbb{R}_+ G^k && [(36)] \\ &\subsetneq \mathbb{R}_+ V^\bullet && [\text{consequence of (35)}] \\ &= V_\infty^\circ && [(28) \text{ again}] \end{aligned}$$

Thus,  $(V_\infty^k)^\circ \subsetneq V_\infty^\circ$ , which implies (i) since polarity is an involution between closed convex cones.

To prove (ii), take  $\bar{r}$  in  $\bigcap_k V^k$ ; we have to prove that  $\bar{r} \in V$  (the other inclusion being obvious). If  $\bar{r} \notin V$  there is a separating hyperplane  $\bar{d}$ :  $\sigma_V(\bar{d}) < \bar{d}^\top \bar{r}$ . Normalizing  $\bar{d}$  via (28), we have altogether

$$\bar{r} \in \bigcap_k V^k, \quad \bar{d} \in \hat{V}^\circ, \quad \bar{d}^\top \bar{r} > 1; \tag{37}$$

but  $\sigma_{G^k}$  represents  $V^k$ , so (37) gives

$$\sigma_{G^k}(\bar{r}) \leq 1 < \bar{d}^\top \bar{r}, \quad \text{hence } \bar{d} \notin G^k.$$

Then  $\bar{d} \in V^\bullet \cap N^k$  for all  $k$  (large enough), i.e.,  $\bar{d} \in [d_V, t_V d_V]$ . Introduce  $d_k \in G^k$  from (34):

$$\|d_k - \bar{d}\| \leq \frac{1}{k} \quad \text{and} \quad d_k^\top \bar{r} \leq \sigma_{G^k}(\bar{r}) \leq 1.$$

Passing to the limit,  $\bar{d}^\top \bar{r} \leq 1$ ; a contradiction to (37). Therefore  $\bar{r} \in V$ .  $\square$

Now we assume the existence of an  $S$ -free set  $W$  containing  $V$ ; it satisfies in particular

$$W^\bullet \subset W^\circ \subset V^\circ = [0, 1]V^\bullet. \tag{38}$$

If  $W^\bullet \subset V^\bullet$ , this  $W$  is of no use to disprove maximality of  $V$  (Proposition 4.8). We are therefore in the situation

$$W^\bullet \not\subset V^\bullet, \quad \text{which implies from (38): } 0 \notin V^\bullet. \tag{39}$$

Thus,  $W^\bullet$  contains some points out of  $V^\bullet$ . The key argument for our analysis is that one of these points lies on an extreme ray of  $V_\infty^\circ$ —which will be the  $d_V$  of Lemma 5.3, crucial to construct the unbounded sequence  $\{r^k\}$  of Figure 8.

LEMMA 5.4 (CONSTRUCTING AN APPROPRIATE EXTREME RAY). Let  $W \supset V$  satisfy (39). There is an extreme ray  $\mathbb{R}_+ d_V$  of  $V_\infty^\circ$  such that the set  $N^k$  defined by (32) satisfies  $W^\circ \cap N^k = \emptyset$  for  $k$  large enough.




FIGURE 10. The extreme ray  $\mathbb{R}_+ b_{j_0}$  contains some point in  $V^* \setminus W^*$ .

PROOF. From (39), we are in the framework of Corollary 3.10; Figure 10 is helpful to follow the proof. If  $\hat{W}^\circ \subset V^*$  then  $W^* = \overline{\text{conv}}(\hat{W}^\circ) \subset V^*$ , contradiction. So there is  $e \in \hat{W}^\circ$  (hence  $\sigma_W(e) = 1$ ) which does not lie in  $V^*$ ; because  $V \subset W$ , i.e.,  $\sigma_V \leq \sigma_W$ , this  $e$  satisfies  $\sigma_V(e) < 1$  (otherwise  $\sigma_V(e) = 1$ , hence  $e \in \hat{V}^\circ \subset V^*$ ).

Then construct  $d_e := (1/\sigma_V(e))e \in \hat{V}^\circ$  (remember (21):  $\sigma_V(e) > 0$ ). For every  $e' \in [0, e]$ , the segment  $[e', d_e]$  contains  $e$ . Being a convex set,  $V^*$  cannot contain such an  $e'$  (otherwise it would contain  $e$  as well). As a result, the compact convex sets  $V^*$  and  $[0, e]$  can be separated: there is  $l \in \mathbb{R}^q$  (appropriately scaled) such that

$$\max\{0, e^\top l\} < 1 < \min_{d \in V^*} d^\top l. \tag{40}$$

Observe that

$$1 > e^\top l = \sigma_V(e) d_e^\top l > 0. \tag{41}$$

Now introduce the closed convex set

$$B := \{b \in V_\infty^\circ : b^\top l = 1\}.$$

Clearly,  $\mathbb{R}_+ B \subset V_\infty^\circ$ . Conversely, apply (28): every nonzero  $d \in V_\infty^\circ$  can be scaled to some  $td \in V^*$ . By (40),  $td^\top l > 1$ , then  $d$  can be scaled again to  $td/(td^\top l)$ , which lies in  $B$ . We have shown

$$\mathbb{R}_+ B = V_\infty^\circ. \tag{42}$$

By (28), every  $b \in B$  can be obtained by scaling some  $d \in \hat{V}^\circ$ :  $b = td$ ; and  $t = 1/(d^\top l) \in ]0, 1[$  by (40). This means that

$$B \subset ]0, 1[\hat{V}^\circ \subset V^\circ; \tag{43}$$

$B$  is therefore bounded (and closed because  $V_\infty^\circ$  is closed), hence compact.

Using (41), scale  $e$  to  $\bar{b} := (1/(e^\top l))e \in B$  and express  $\bar{b} = \sum_j \alpha_j b_j$  as a convex combination of extreme points  $b_j$  of  $B$  (Minkowski's Theorem). Then

$$\sigma_W(\bar{b}) = \frac{1}{e^\top l} \sigma_W(e) = \frac{1}{e^\top l} > 1.$$

By convexity of  $\sigma_W$ , there is some  $j_0$  such that  $\sigma_W(b_{j_0}) > 1$  (we may have  $\sigma_W(b_{j_0}) = +\infty$ ). Altogether, we have exhibited

$$b_{j_0} \text{ extreme in } B \text{ and satisfying } 1 < \sigma_W(b_{j_0}).$$

Extremality of  $b_{j_0}$  in  $B$  implies extremality of the ray  $\mathbb{R}_+ b_{j_0}$  in  $\mathbb{R}_+ B$ , i.e., in  $V_\infty^\circ$  because of (42). The intersection of  $W^\circ$  with this extreme ray is some  $[0, d_w]$  ( $d_w$  may be 0) which, by definition of a polar, does not contain  $b_{j_0}$ . Since  $b_{j_0}^\top l = 1$  (because  $b_{j_0} \in B$ ),  $d^\top l < 1$  for all  $d \in [0, d_w]$ . Then, (40) shows that  $[0, d_w]$  and  $[d_v, t_v d_v]$  are separated.

As a result, the two compact sets  $W^\circ$  and  $[d_v, t_v d_v]$  are disjoint. If there were  $d^k \in W^\circ \cap N^k$  for all  $k$ , then the bounded sequence  $\{d^k\}$  would have some cluster point  $d^*$ ; but  $W^\circ$  is closed:  $d^*$  would lie in  $W^\circ \cap [d_v, t_v d_v]$ , contradiction.  $\square$

The set  $B$  constructed in the above proof is a so-called basis of the pointed cone  $V_\infty^\circ$ . The case  $\sigma_W(b_{j_0}) = +\infty$ ,  $d_w = 0$  corresponds to a  $W$  as in Figure 5; it occurs in Figure 10. This latter picture is still helpful to follow the next proof. Recall that  $L$  is the lineality space of  $V$ .



PROPOSITION 5.5. Assume  $0 \in \bar{S} = \overline{\text{conv}}(S)$ . If a minimal CGF  $\rho$  represents the  $S$ -free set  $V = V(\rho)$ , which is not maximal, then  $V^k$  exists as described by Lemma 5.3. There is  $r^k \in V^k \cap S$ , decomposed as  $r^k = l^k + u^k$  with  $l^k \in L$  and  $u^k \in L^\perp$ , such that

$$\text{for some } K \subset \mathbb{N}, \quad \lim_{k \in K} \|r^k\| = +\infty \quad \text{and} \quad \lim_{k \in K} \|u^k\| = +\infty.$$

PROOF. If all of the  $S$ -free sets  $W$  containing  $V$  satisfy  $W^\bullet \subset V^\bullet$ , then  $V$  is maximal (Proposition 4.8). Thus, there is an  $S$ -free set  $W \supset V$  satisfying (39) and we can construct  $d_V$  as in Lemma 5.4.

If  $\mathbb{R}_+ d_V = V_\infty^\circ$ , then  $\hat{V}^\circ = V^\bullet = \{d_V\}$  and  $V^\circ = [0, d_V]$  (Proposition 3.7): the  $S$ -free set  $V$ , represented by  $\sigma_{V^\circ}$ , is the half space  $\{r: d_V^\top r \leq 1\}$ , which separates 0 from  $\bar{S}$ ; this is ruled out by assumption.

Otherwise,  $\mathbb{R}_+ d_V \subsetneq V_\infty^\circ$ : we can apply Lemma 5.3 and construct the sequence of neighborhoods  $V^k$ . By minimality of  $\mu_V$ ,  $V^k$  cannot be  $S$ -free (Lemma 5.3(i) and Theorem 4.10): there exists  $r^k$  lying

— in  $\text{int } V^k$ , hence from (12)

$$1 > \sigma_{G^k}(r^k), \tag{44}$$

— and in  $S$ , hence  $r^k \notin \text{int } W$ :  $\sigma_{W^\bullet}(r^k) \geq 1$ ; since  $W^\bullet$  is compact,

$$\exists e_k \in W^\bullet \quad \text{such that } e_k^\top r^k \geq 1. \tag{45}$$

Now we claim that there is  $\delta > 0$  such that

$$t_k e_k \in V^\bullet \cap N^k, \quad \text{for some } t_k \geq 1 + \delta \text{ and all } k \text{ large enough.} \tag{46}$$

Using (28), scale  $e_k$  (nonzero from its definition) to  $t_k e_k \in V^\bullet$ ; and note from (38) that  $t_k \geq 1$ . Then (45) implies that  $t_k e_k \notin G^k$ : otherwise

$$1 \leq e_k^\top r^k \leq t_k e_k^\top r^k \leq \sigma_{G^k}(r^k)$$

by definition of a support function; this contradicts (44). It follows that  $t_k e_k \in V^\bullet \cap N^k$ , which is far from  $W^\bullet$  (Lemma 5.4); (46) is proved.

Now we can conclude. First, let  $\bar{d} \in [d_V, t_V d_V]$  be a cluster point of the bounded sequence  $\{t_k e_k\}$ . Next, use (46), (45), (44) to write for all  $d \in G^k$

$$1 + \delta \leq t_k \leq t_k e_k^\top r^k = (t_k e_k - d)^\top r^k + d^\top r^k < (t_k e_k - d)^\top r^k + 1.$$

This holds in particular for  $d = d_k$  stated in (34):

$$\delta < (t_k e_k - d_k)^\top r^k. \tag{47}$$

Then we obtain with the Cauchy-Schwarz inequality

$$\delta < \|t_k e_k - \bar{d} + \bar{d} - d_k\| \|r^k\| \leq \left( \|t_k e_k - \bar{d}\| + \frac{1}{k} \right) \|r^k\|.$$

Furthermore, decompose  $r^k = l^k + u^k$  in (47) and observe that both  $e_k^\top l^k$  and  $d_k^\top l^k$  are 0 ( $l^k \in L$  and  $e^k$  and  $d^k$  lie in  $V_\infty^\circ \subset L^\perp$ ). So (47) gives also

$$\delta < (t_k e_k - d_k)^\top u^k \leq \left( \|t_k e_k - \bar{d}\| + \frac{1}{k} \right) \|u^k\|.$$

Both statements are proved since there is  $K \subset \mathbb{N}$  such that  $\lim_{k \in K} \|t_k e_k - \bar{d}\| = 0$ .  $\square$

As suggested in the beginning of this section, proving Theorem 5.1 is now easy. An  $S$ -free set represented by a minimal CGF will be automatically maximal under any assumption contradicting the existence of our unbounded sequences.

PROOF OF THEOREM 5.1. Construct the sequences  $\{r^k\}$  and  $\{u^k\}$  of Proposition 5.5.

Case (i). Extract a cluster point  $\hat{r}$  of the normalized subsequence  $\{r^k\}_{k \in K}$ : for some  $K' \subset K$ ,

$$\lim_{k \in K'} \frac{r^k}{\|r^k\|} = \hat{r}.$$

Then take an arbitrary  $M > 0$ . We know that  $M/\|r^k\| \leq 1$  if  $k$  is large enough in  $K'$  so, because both  $0$  and  $r^k$  lie in  $V^k \cap \bar{S}$ ,

$$\frac{M}{\|r^k\|} r^k \in V^k \cap \bar{S}, \quad \text{for large enough } k \in K'.$$

By closedness, this implies  $M\hat{r} \in \bar{S}$ , hence  $\hat{r} \in \bar{S}_\infty$  because  $M$  is arbitrary. The same argument using Lemma 5.3(ii) gives  $\hat{r} \in V_\infty$ .

Let us sum up. If  $V$  is not maximal, then  $V_\infty \cap \bar{S}_\infty$  contains a vector  $\hat{r}$  of norm 1; this contradicts (i).

Case (ii). Write  $u^k = r^k - l^k \in V^k - L = V^k + L \subset V^k + V_\infty \subset V^k$ . Then proceed as in Case (i): extract a cluster point  $\hat{u}$  of  $\{u^k/\|u^k\|\}_K$  and argue that  $(M/\|u^k\|)u^k \in V^k \cap L^\perp$  to exhibit

$$\hat{u} \in V_\infty \cap L^\perp \quad \text{and} \quad \|\hat{u}\| = 1. \tag{48}$$

Besides,  $u^k$  is the projection onto  $L^\perp$  (a linear operator) of  $r^k \in S \subset U + \bar{S}_\infty$ ; hence

$$u^k \in \text{Proj}_{L^\perp} U + \text{Proj}_{L^\perp} \bar{S}_\infty.$$

By (ii)<sub>1</sub>,  $\text{Proj}_{L^\perp} U$  is a bounded set, so our cluster direction  $\hat{u}$  lies in  $\text{Proj}_{L^\perp} \bar{S}_\infty$ :

$$\hat{u} = \hat{s} - \hat{l}, \quad \text{for some } \hat{s} \in \bar{S}_\infty \text{ and } \hat{l} \in L.$$

Use (48):

$$\bar{S}_\infty \ni \hat{s} = \hat{u} + \hat{l} \in V_\infty + L = V_\infty;$$

then use (ii)<sub>2</sub>:

$$\hat{s} \in V_\infty \cap \bar{S}_\infty = L \cap \bar{S}_\infty.$$

As a result,  $\hat{u} = \hat{s} - \hat{l}$  lies in  $L$ ; use (48) again:  $\hat{u} \in L \cap L^\perp$  cannot have norm 1.

Thus, in this case also,  $V$  has to be maximal.  $\square$

Let us insist once more: the core of our proof is Proposition 5.5. Then (i) and (ii) appear as *ad hoc* assumptions to contradict the existence of the stated unbounded sequences; other similar assumptions might be designed.

**6. Conclusion and perspectives.** In this paper, we have laid down some basic theory toward studying the cutting paradigm for sets of the form (1). We have introduced for this the concept of cut-generating functions, which allowed us to put in perspective an abundant literature devoted to  $S$ -free sets. We have revealed the discrepancy between minimality and maximal  $S$ -freeness; and we have recovered existing theorems (Johnson [18], Borozan and Cornuéjols [8], Basu et al. [5], Dey and Wolsey [12], Basu et al. [6]), dealing with mere minimality, exhibiting the intrinsic arguments allowing their proofs. Our theory necessitated a generalization of the polarity correspondence to certain unbounded sets; we have conducted it via a systematic exploitation of the correspondence between sublinear functions and closed convex sets.

A number of questions arise from this theoretical work. Some are suggested by §3:

*Question 1.* Given a convex compact set  $G$ , can we detect whether it is the minimal prepolar of  $V := G^\circ$ ? And if not, can we compute  $(G^\circ)^\circ$ ?

*Question 2.* Knowing that our generalization of polarity goes along with that of Zaffaroni [24], linking the two works should certainly be instructive. For example, we define the prepolar by (23), which looks quite different from the set  $Q$  in Zaffaroni [24, Proposition 5.1]. Yet the two sets have to coincide, at least when  $0 \in \text{int } V$ ; can this be clarified? And can we explain what happens when  $0$  becomes a boundary point of  $V$ ? Also, does this other definition help answer question 1?

These are limited to pure convex analysis; concerning the CGF theory itself, some other questions have a concrete interest:

*Question 3.* Is it possible to characterize exactly the  $S$ -free sets represented by minimal CGF's? A converse form of Theorem 4.10 should be desirable.

*Question 4.* One might want to consider more general models. For example, it should not be too difficult to replace the “ground set”  $\mathbb{R}_+^n$  of (1a) by some other closed convex cone; say the cone of positive semi-definite matrices, which would open the way toward cutting SDP relaxations. Kılınç-Karzan [20] opens interesting perspectives for this. Another generalization would be inspired by the approach of Gomory [14] of Example 1.1: there,  $X$  has the form

$$\{x \in \mathbb{Z}_+^n : -Ax \in \mathbb{Z}^m - b\};$$

$S = \mathbb{Z}^m - b$  lies in a smaller space but the ground set  $\mathbb{Z}_+^n$  is no longer convex, so sublinear CGF's are now ruled out. Instead, CGF's in this context are subadditive, periodic, and satisfy a certain symmetry condition (Gomory and Johnson [15]).




FIGURE 11. Not all cuts are obtained from a CGF.

*Question 5.* Perhaps the most crucial question is whether CGF’s do generate all possible cuts, i.e., whether (8) is able to produce all possible  $c$ ’s satisfying (2). This turns out to be a tough nut to crack, we conclude the paper with some considerations for future research concerning it.

The following counterexample shows that the answer to question 5 is no in general.

**EXAMPLE 6.1 (CGF’S NEED NOT GENERATE ALL CUTS).** In  $\mathbb{R}^2$ , take  $S = (0, 1) \cup \{(\mathbb{Z}, -1)\}$ . The left part of Figure 11, drawn in the  $S$ -space, clearly shows that, if the unit-vector  $(1, 0)$  lies in the recession cone of an  $S$ -free set  $V$ , then it lies on the boundary of this cone.

Now take the identity matrix for  $R$ : in the  $x = (\xi, \eta)$ -space,  $X$  reduces to the singleton  $(0, 1)$  in  $\mathbb{R}^2$  (right part of Figure 11). It can be separated from the origin by the cut  $\eta \geq \xi + 1$ , obtained with  $c = (-1, 1)^\top$ . Knowing that the first column of  $R$  is  $r_1 = (1, 0)^\top$ , a CGF  $\rho$  producing this  $c$  must therefore have  $\rho(r_1) = -1$ . In view of Lemma 3.2,  $(1, 0)$  lies in the interior of  $V_\infty$ ; but we have seen that no  $V$  can satisfy this.

Negative  $c_j$ ’s are therefore troublesome, a general sufficiency theorem is out of reach. To eliminate  $c_j < 0$ , we can restrict the class of instances:

**PROPOSITION 6.2.** *If the recession cone of  $\overline{\text{conv}}(X)$  is the whole of  $\mathbb{R}_+^n$ , then every cut  $c$  lies in  $\mathbb{R}_+^n$ .*

**PROOF.** Each basis vector  $e_j$  of  $\mathbb{R}^n$  lies in  $[\overline{\text{conv}}(X)]_\infty$ : picking some  $x \in X$ ,

$$c^\top(x + te_j) = c^\top x + tc_j \geq 1 \quad \text{for all } t \geq 0;$$

let  $t \rightarrow +\infty$  to see that  $c_j \geq 0$ .  $\square$

This result might suggest that the trouble in Example 6.1 is due to the difference between the recession cones of  $\overline{\text{conv}}(X)$  and of the ground set  $\mathbb{R}_+^n$  in (1a). However, the assumption introduced in Proposition 6.2 does not suffice, as even  $c_j = 0$  brings trouble. In fact, make a “more nonlinear” variant of Example 6.1: instead of the horizontal line  $\psi = -1$ , take for  $S$  the curve  $\psi = -1/|\phi|$  ( $\phi \neq 0$ ). This leaves  $X = \{(0, 1)\}$  unchanged;  $c = (0, 1)^\top$  is a cut and a CGF  $\rho$  generating it has  $\rho(r_1) = 0$ ; this  $\rho$  represents a set  $V(\rho)$ , which has  $(\mathbb{R}_+, 0)$  in its recession cone. Being a neighborhood of the origin,  $V(\rho)$  contains  $A := (0, -\varepsilon)$  for small enough  $\varepsilon > 0$ ; also,  $B := (r, 0) \in V(\rho)_\infty \subset V(\rho)$  for all  $r > 0$  (see Figure 12); by convexity, the whole segment  $[A, B]$  lies in  $V(\rho)$ , which therefore cannot be  $S$ -free.

In these two examples, the conical hull of the  $r_j$ ’s does not cover the whole of  $S$ . In fact,  $S$  contains points that can be reached by no  $x \in \mathbb{R}_+^n$ ; these points have nothing to do with the problem, so forcing  $V$  not to contain them is unduly demanding. Then one may ask whether CGF’s are able to describe all possible cuts, for all possible instances such that  $S \subset \text{cone}(r_1, \dots, r_n)$ . This is an open question. Here we limit ourselves to a reasonably simple sufficiency result, proved with the help of a “comfortable” assumption; it motivated the generalization obtained recently in Cornuéjols et al. [11].




FIGURE 12. Trouble appears when  $V_\infty$  is an asymptote of  $S$ .

THEOREM 6.3. Let an instance of (1) be as described by Proposition 6.2 and assume

$$\text{cone}(r_1, \dots, r_n) := \left\{ \sum_{j=1}^n \lambda_j r_j : \lambda_j \geq 0, j = 1, \dots, n \right\} = \mathbb{R}^q.$$

Then every cut can be obtained from a CGF.

PROOF. Let  $c \in \mathbb{R}_+^n$  and set

$$J_+ := \{j \in \{1, \dots, n\} : c_j > 0\}, \quad J_0 := \{j \in \{1, \dots, n\} : c_j = 0\}.$$

Then introduce in  $\mathbb{R}^q$  the vectors

$$r'_j := \frac{r_j}{c_j}, \quad \text{for } j \in J_+$$

and the polyhedron

$$V := G + K, \quad \text{with } \begin{cases} G := \text{conv}\{r'_j : j \in J_+\}, \\ K := \text{cone}\{r_j : j \in J_0\}. \end{cases}$$

Claim 1.  $V$  is a neighborhood of the origin. In fact, our assumption means that  $\mathbb{R}^q = \text{cone}(G) + K$ : every  $\bar{d} \in \mathbb{R}^n$  has the form

$$\bar{d} = \bar{t}\bar{g} + \bar{k}, \quad \text{with } \bar{t} \geq 0, \quad \bar{g} \in G, \quad \bar{k} \in K.$$

Then compute  $\sigma_V(\bar{d})$  for nonzero  $\bar{d}$ .

Case 1.  $\bar{t} = 0$ . Fixing  $g \in G$  so that  $g + t\bar{k} \in V$  for all  $t \geq 0$ , we have

$$\sigma_V(\bar{d}) = \sigma_V(\bar{k}) \geq \bar{k}^\top (g + t\bar{k}) = \bar{k}^\top g + t\|\bar{k}\|^2, \quad \text{for all } t > 0;$$

let  $t \rightarrow +\infty$  to see that  $\sigma_V(\bar{d}) = +\infty$ .

Case 2.  $\bar{t} > 0$ . Scale  $\bar{d}$  to  $\bar{t}^{-1}\bar{d} \in G + K = V$  to obtain  $\sigma_V(\bar{d}) \geq \bar{t}^{-1}\|\bar{d}\|^2 > 0$ .

Altogether, we have proved that  $\sigma_V(\bar{d}) > 0$  for all  $\bar{d} \neq 0$ , i.e.,  $0 \in \text{int}(V)$ .

Claim 2.  $V$  is  $S$ -free. Take  $\bar{r} \in \text{int}(V)$ . For  $\varepsilon > 0$  small enough,  $\bar{r} + \varepsilon\bar{r} \in V$ :

$$(1 + \varepsilon)\bar{r} = \sum_{j \in J_+} \beta_j r'_j + \sum_{j \in J_0} \mu_j r_j, \quad \text{with } \beta_j, \mu_j \geq 0, \quad \sum_{j \in J_+} \beta_j = 1.$$

Divide by  $1 + \varepsilon$  and set  $\alpha_j = \beta_j / (1 + \varepsilon)$ ,  $\lambda_j = \mu_j / (1 + \varepsilon)$  to get

$$\bar{r} = \sum_{j \in J_+} \alpha_j r'_j + \sum_{j \in J_0} \lambda_j r_j, \quad \text{for } \alpha_j, \lambda_j \geq 0, \quad \sum_{j=1}^n \alpha_j < 1.$$

Introduce the vector  $\bar{x} \in \mathbb{R}^n$  whose coordinates are

$$\bar{x}_j := \begin{cases} \frac{\alpha_j}{c_j} & \text{if } j \in J_+, \\ \lambda_j & \text{if } j \in J_0. \end{cases}$$

Observe that  $\bar{x} \geq 0$  and that

$$R\bar{x} = \sum_{j=1}^n \bar{x}_j r_j = \sum_{j \in J_+} \frac{\alpha_j}{c_j} r_j + \sum_{j \in J_0} \lambda_j r_j = \bar{r}.$$

If  $\bar{r} \in S$  then  $x \in X$  by definition (1a); but

$$c^\top \bar{x} = \sum_{j \in J_+} c_j \frac{\alpha_j}{c_j} = \sum_{j \in J_+} \alpha_j \leq \sum_{j=1}^n \alpha_j < 1$$

and  $x$  cannot lie in  $X$  if  $c$  is a cut. We have proved that  $\text{int}(V) \cap S = \emptyset$ , i.e., that  $V$  is  $S$ -free.

Conclusion. We have proved that the gauge  $\gamma_V$  is a CGF; besides

- for  $j \in J_0$ ,  $r_j$  is a direction of recession of  $V$ :  $\gamma_V(r_j) = 0 = c_j$ ; and
- for  $j \in J_+$ , the property  $r'_j \in V$  gives

$$1 \geq \gamma_V(r'_j) = \frac{1}{c_j} \gamma_V(r_j), \quad \text{hence } \gamma_V(r_j) \leq c_j.$$

In summary,  $\gamma_V$  is a CGF dominating the cut  $c$ .  $\square$

To make question 5 less ambitious, one may ask whether CGF's can reproduce the set of cuts “globally.” In fact, the set of  $c$ 's satisfying (2) is a closed convex set: the opposite of the reverse polar  $X^-$ , in the terminology of Balas [3] and Cornuéjols and Lemaréchal [10]. Then consider the set  $\mathcal{R}_S$  of all representations of a given  $S$ -free set. Given  $(n, R)$ , form the set  $\mathcal{C}$  of  $c \in \mathbb{R}^n$  whose coordinates are  $\rho(r_j)$ , where  $\rho$  describes  $\mathcal{R}_S$ . Is it true that  $\overline{\text{conv}}(\mathcal{C}) = -X^-$ ? This question is open. If the answer is yes, one more question occurs: Example 4.5 tells us that  $\mathcal{R}_S$  cannot be reduced to the maximal  $S$ -free sets; then, what sort of maximality can be imposed while preserving “completeness” of  $\mathcal{R}_S$ ? An answer should need answering question 3 first.

**Acknowledgments.** The discontinuous support function constructed in Remark 5.2 was suggested by R.T. Rockafellar. The authors are indebted to the referees, whose attentive and intelligent reading was very helpful to improve an earlier version of this paper. This work was supported in part by NSF [Grant CMMI1263239], ONR [Grant N00014-09-1-0033], MICINN [Grant MTM2011-29064-C03-01] (Spain) and FONDECYT [Regular Grant 1130176] (Chile).

## References

- [1] Andersen K, Louveaux Q, Weismantel R, Wolsey L (2007) Cutting planes from two rows of a simplex tableau. Fischetti M, Williamson D, eds. *Integer Programming and Combinatorial Optimization*, Lecture Notes in Computer Science, Vol. 4513 (Springer, Berlin), 1–15.
- [2] Averkov G (2013) On maximal  $S$ -free sets and the Helly number of the family of  $S$ -convex sets. *SIAM J. Discrete Math.* 27:1610–1624.
- [3] Balas E (1979) Disjunctive programming. *Ann. Discrete Math.* 5:3–51.
- [4] Basu A, Cornuéjols G, Zambelli G (2011) Convex sets and minimal sublinear functions. *J. Convex Anal.* 18(2):427–432.
- [5] Basu A, Conforti M, Cornuéjols G, Zambelli G (2010) Maximal lattice-free convex sets in linear subspaces. *Math. Oper. Res.* 35(3):704–720.
- [6] Basu A, Conforti M, Cornuéjols G, Zambelli G (2010) Minimal inequalities for an infinite relaxation of integer programs. *SIAM J. Discrete Math.* 24(1):158–168.
- [7] Bixby R, Rothberg E (2007) Progress in computational mixed integer programming—A look back from the other side of the tipping point. *Ann. Oper. Res.* 149(1):37–41.
- [8] Borozan V, Cornuéjols G (2009) Minimal valid inequalities for integer constraints. *Math. Oper. Res.* 34(3):538–546.
- [9] Caprari E, Zaffaroni A (2010) Conically equivalent convex sets and applications. *Pacific J. Optim.* 6:281–303.
- [10] Cornuéjols G, Lemaréchal C (2005) A convex-analysis perspective on disjunctive cuts. *Math. Programming* 106(3):567–586.
- [11] Cornuéjols G, Yildiz S, Wolsey LA (2014) Sufficiency of cut-generating functions. *Math. Programming, Ser. A*, ePub ahead of print April 19, <http://link.springer.com/article/10.1007/s10107-014-0780-2>.
- [12] Dey SS, Wolsey LA (2010) Constrained infinite group relaxations of MIPs. *SIAM J. Optim.* 20(6):2890–2912.
- [13] Gomory RE (1963) An algorithm for integer solutions to linear programs. Graves RL, Wolfe P, eds. *Recent Advances in Mathematical Programming* (McGraw-Hill, New York), 269–302.
- [14] Gomory RE (1969) Some polyhedra related to combinatorial problems. *Linear Alg. Appl.* 2(4):451–558.
- [15] Gomory RE, Johnson EL (1972) Some continuous functions related to corner polyhedra i. *Math. Programming* 3:23–85.
- [16] Hiriart-Urruty J-B, Lemaréchal C (1993) *Convex Analysis and Minimization Algorithms* (Springer, Berlin).
- [17] Hiriart-Urruty J-B, Lemaréchal C (2001) *Fundamentals of Convex Analysis* (Springer, Berlin).
- [18] Johnson EL (1981) Characterization of facets for multiple right-hand side choice linear programs. *Math. Programming Study* 14:112–142.
- [19] Júdice JJ, Sherali H, Ribeiro IM, Faustino AM (2006) A complementarity-based partitioning and disjunctive cut algorithm for mathematical programming problems with equilibrium constraints. *J. Global Optim.* 136:89–114.
- [20] Kiliç-Karzan F (2013) On minimal valid inequalities for mixed integer conic programs. Working Paper 2013-E20, GSIA, Carnegie Mellon University, Pittsburgh.
- [21] Lovász L (1989) Geometry of numbers and integer programming. Iri M, Tanabe K, eds. *Mathematical Programming: Recent Developments and Applications* (Kluwer Academic Publishers, Amsterdam), 177–210.
- [22] Morán R, Dey SS (2011) On maximal  $S$ -free convex sets. *SIAM J. Discrete Math.* 25:379–393.
- [23] Rockafellar RT (1970) *Convex Analysis* (Princeton University Press, Princeton, NJ).
- [24] Zaffaroni A (2013) Convex radiant costarshaped sets and the least sublinear gauge. *J. Convex Anal.* 20(2):307–328.