

# Sleep and awake bruxism in adults and its relationship with temporomandibular disorders: A systematic review from 2003 to 2014

Antonio Jiménez-Silva, Consuelo Peña-Durán, Julio Tobar-Reyes & Raúl Frugone-Zambra

To cite this article: Antonio Jiménez-Silva, Consuelo Peña-Durán, Julio Tobar-Reyes & Raúl Frugone-Zambra (2017) Sleep and awake bruxism in adults and its relationship with temporomandibular disorders: A systematic review from 2003 to 2014, *Acta Odontologica Scandinavica*, 75:1, 36-58, DOI: [10.1080/00016357.2016.1247465](https://doi.org/10.1080/00016357.2016.1247465)

To link to this article: <https://doi.org/10.1080/00016357.2016.1247465>



Published online: 31 Oct 2016.



Submit your article to this journal



Article views: 1097



View related articles



View Crossmark data



Citing articles: 27 View citing articles

REVIEW ARTICLE

## Sleep and awake bruxism in adults and its relationship with temporomandibular disorders: A systematic review from 2003 to 2014

Antonio Jiménez-Silva<sup>a,b</sup>, Consuelo Peña-Durán<sup>c</sup>, Julio Tobar-Reyes<sup>c</sup> and Raúl Frugone-Zambrano<sup>d,e</sup>

<sup>a</sup>Facultad de Ciencias de la Salud, Universidad Autónoma de Chile, Temuco, Chile; <sup>b</sup>Ortodoncia y Ortopedia dentomaxilofacial, Facultad de Odontología, Universidad Andrés Bello, Santiago, Chile; <sup>c</sup>Department of Prosthodontics, Faculty of Dentistry, University of Chile, Santiago, Chile; <sup>d</sup>Faculty of Dentistry, University of the Andes, Santiago, Chile; <sup>e</sup>University of La Serena, La Serena, Chile

### ABSTRACT

**Objective:** In order to establish a relationship between bruxism and temporomandibular disorders (TMDs), a systematic review was performed.

**Materials and methods:** A systematic research was performed based on PubMed, Cochrane Library, Medline, Embase, BIREME, Lilacs and Scielo data bases, between 2003 and 2014 including all languages. Descriptive clinical cases were identified. Two independent authors selected the articles. PICO format was used to analyse the studies and the Newcastle-Ottawa Scale (NOS) was used to verify the quality of the evidence.

**Results:** Thirty-nine studies ( $n=39$ ) were analysed in this review. According to bruxism diagnosis, articles were grouped as follows: polysomnographic diagnosis (PSG) ( $n=7$ ), clinical diagnosis ( $n=11$ ) and survey/self-report ( $n=21$ ). Thirty-three articles ( $n=33$ ) established a positive relation between bruxism and TMD and six ( $n=6$ ) did not. Quality of evidence was low to moderate. In general, the most part of the studies showed shortcomings on their design with bias risk, and also had a low sensitivity on bruxism diagnosis.

**Conclusions:** The evidence based on PSG was not as conclusive as the studies that used surveys and clinical exam to diagnosis bruxism, when bruxism was related to TMD. Sleep bruxism could be associated with myofascial pain, arthralgia and joint pathology as disc displacement and joint noises. Although the evidence at present is inconclusive and does not provide information according to the type of bruxism (bruxism sleep and wakefulness), it is possible to suggest that bruxism would be associated with TMD.

### ARTICLE HISTORY

Received 7 June 2016  
Accepted 8 October 2016

### KEYWORDS

Bruxism; temporomandibular disorders;  
temporomandibular joint;  
sleep bruxism

### Introduction

Bruxism is defined as 'a repetitive jaw-muscle activity characterized by clenching or grinding of the teeth and/or by bracing or thrusting of the mandible'.<sup>[1]</sup> Also, it has been demonstrated that bruxism has two circadian manifestations because it can occur during sleep (sleep bruxism), or during wakefulness (awake bruxism). Moreover, it has been proposed that sleep or awake bruxism can be classified as possible (self-report), probable (self-report plus clinical examination) or definite (self-report plus clinical examination, plus polysomnographic recording).<sup>[1]</sup>

According to the American Academy of Orofacial Pain (AAOP), temporomandibular disorders (TMDs) are defined as 'a group of disorders involving the masticatory muscles, the temporomandibular joint (TMJ), and associated structures'.<sup>[2]</sup> TMDs have a multifactorial etiology and among them, some conditions cause trauma, as the overloading generated because of bruxism.

Muscle overloading due to tooth clenching could be associated with local blood flow and microcirculation disorders,

and pain derived from an ischemia,<sup>[3]</sup> the latter related to substances that sensitize muscles nociceptors.<sup>[4]</sup>

Based on evidence, the relationship between bruxism and TMD is still controversial in the literature due to the complexity of the etiology and diagnostic of both disorders.<sup>[5]</sup> In fact, both the presence and the cause-effect relationship are still undefined [5–7] although there are some studies that demonstrate a positive relation between bruxism and TMD.<sup>[5]</sup> The lack of evidence is based essentially on methodological differences, as data interpretation, operational definitions, methodological designs and diagnostic instrumentation for both disorders.

Looking out the diversity and the scarce reviews that analysed bruxism with a specific diagnostic modality, without focus in evidence level and the insufficiency of inclusion of other languages out of English, this study aims to review the literature between 2003 and 2014. The aim was to find out if bruxism is associated with TMDs but considering the types of bruxism (sleep and awake), TMD diagnostic method, quality of evidence and all languages.

## Materials and methods

To establish the relationship between TMD and bruxism, on 31 January 2015, a systematic review of the literature was conducted between 2003 and 2014. Databases used were as follows: PubMed, Cochrane Library, Embase, Medline, BIREME, Lilacs and Scielo.

### Types of studies

Observational studies, analytical case-control or cohort who have clearly defined aims to determine the relationship between bruxism and TMD.

### Language studies

The search was conducted without limitation of language.

### Types of participants

The studies selected for this review included subjects older than 19 years old of both genders.

### Intervention type

Studies without intervention in order to relate TMD and bruxism.

### Types of results

Primary outcomes: to determine the relationship between sleep and awake bruxism and TMD.

Secondary outcomes: to determine the relationship (in-person or cause and effect) between bruxism and TMD by gender. To determine the association between bruxism and TMD according to age group.

### Data collection

For TMD: Data were collected from studies that showed diagnosis of TMD not limited to any method, with a clear reference to the concept and diagnosis of temporomandibular pathology.

For bruxism: Data were collected from studies that showed a diagnosis of bruxism, defining the type explicitly, according to the circadian expression (sleep, awake or both). The accepted diagnostic studies were as follows: (a) polysomnography diagnosis (PSG); (b) clinical diagnosis with or without self-report and (c) self-report and/or questionnaire. The following diagnostic modalities were accepted: possible, probable and definitive.

For the identification and selection of the number of potentially eligible studies for this review (N), a specific and individualized search strategy for each database was developed. A semantic field was determined for the term 'bruxism' and another semantic field related to the term 'TMD' (Table 1).

### Databases used

1. PubMed database. Filters used were: publication dates: 10 years, 1 January 2003–31 December 2014; Age; Adult +19 years and Text Availability: Abstract.

2. The Cochrane Library. Filters: publication dates: 10 years, 1 January 2003–31 December 2014; Database: Trials.
3. Embase: publication dates: 2003–2014
4. Medline: publication dates: 2003–2014
5. BIREME: date of publication: 2003–2014
6. Lilacs: date of publication: 2003–2014
7. Scielo: date of publication: 2003–2014

### Study selection and data collection

For article selection or first approach, two researchers independently selected potentially eligible articles by title and abstract. When no agreement was found, it was discussed with a third researcher about including it or not within the sample. The titles of the selected articles were transferred to an Excel table. Full articles were read, considering three questions of elimination based on the inclusion criteria for the selection of studies that would complete text analysis: (1) Is it a clinical trial? (2) Are subjects over 19 years? (3) Is the relationship between bruxism and TMD established? Articles that met these criteria were included in the review for the final analysis. The reasons why some studies were excluded were recorded in an adjacent column and presented in the results (Table 2).

### Extracting data from the studies

The PICO criteria (population, intervention, control groups and outcome) were used to make the tables of analysed articles, where we defined in detail the analysis variables: population (sample size, distribution of subjects by gender and age range), intervention (main variables to compare, related to the topic, statistical analysis and confidence intervals and type of method used for the diagnosis of bruxism and TMD), comparison criteria or control (presence of any control group) and outcomes (included the answer to the hypothesis, the presence or causal relationship between bruxism and TMD).

### Presentation of results and quality of evidence

The tables were developed with the summary of the main results of the studies analysed. The quality of evidence was determined by the Newcastle-Ottawa Scale (NOS),<sup>[8]</sup> which measures the quality of the evidence for case-control and cohort studies, assigning a score ranging from 0 to 9 points. For case-control studies, there are three categories: (1) selection (4 points), (2) comparability (2 points) and (3) exposure (3 points). To determine the quality of cohort studies, there were also three categories with a score of level of evidence ranging from 0 to 9 points. The categories were as follows: (1) Selection (4 points), (2) Comparability (2 points) and (3) Outcome (3 points). The quality was determined by the same two researchers in charge of search, where the highest quality achieved is obtained by those items that reached a maximum score of 9.

**Table 1.** Search strategy and terms used for the search.

Database and limits	Search strategy and terms
PubMed ( $n = 1733$ ) Publication dates: 10 years, 1 January 2003–31 December 2014; Age: Adult +19 years and Text Availability: Abstract.	Bruxism [tiab] OR sleep bruxism [tiab] OR nocturnal bruxism [tiab] OR teeth grinding [tiab] OR teeth wear [tiab] OR occlusal wear [tiab] OR tooth clenching [tiab] OR tooth abrasion [tiab] OR parafunctional habits [tiab] OR teeth grinding disorder [tiab] OR nocturnal teeth grinding disorder [tiab] AND Temporomandibular joint disorders [tiab] OR Temporomandibular disorders [tiab] OR Temporomandibular joint Dysfunction Syndrome [tiab] OR Temporomandibular Joint Syndrome [tiab] OR TMJ disorders [tiab] OR TMJ diseases [tiab] OR myofascial pain [tiab] OR myofascial pain syndrome [tiab] OR craniomandibular pain [tiab] OR orofacial pain [tiab] OR arthralgia Temporomandibular joint [tiab] OR osteoarthritis temporomandibular joint [tiab] OR osteoarthritis temporomandibular joint [tiab] OR disc displacement [tiab] OR disc displacement reduction [tiab] OR disc displacement with reduction [tiab] OR disc displacement without reduction [tiab]
The Cochrane Library ( $n = 1735$ ) Publication dates: 2003–2014; Database: Trials.	Bruxism OR sleep bruxism OR nocturnal bruxism OR teeth grinding OR teeth wear OR occlusal wear OR tooth clenching OR tooth abrasion OR parafunctional habits OR teeth grinding disorder OR nocturnal teeth grinding disorder AND Temporomandibular joint disorders OR Temporomandibular disorders OR Temporomandibular joint Dysfunction Syndrome OR Temporomandibular Joint Syndrome OR TMJ disorders OR TMJ diseases OR myofascial pain OR myofascial pain syndrome OR craniomandibular pain OR orofacial pain OR arthralgia Temporomandibular joint OR osteoarthritis temporomandibular joint OR osteoarthritis temporomandibular joint OR disc displacement OR disc displacement reduction OR disc displacement with reduction OR disc displacement without reduction
Lilacs ( $n = 38$ ) Publication year: 2003–2014	Bruxism OR sleep bruxism OR nocturnal bruxism OR teeth grinding disorder AND Temporomandibular disorders OR craniomandibular disorders
Scielo ( $n = 20$ )	Bruxism OR sleep bruxism OR nocturnal bruxism AND Temporomandibular disorders
Bireme ( $n = 212$ )	Bruxism OR sleep bruxism OR nocturnal bruxism OR tooth clenching OR parafunctional habits OR teeth grinding disorder OR nocturnal teeth grinding disorder AND Temporomandibular joint disorders OR Temporomandibular disorders OR Temporomandibular joint Dysfunction Syndrome OR Temporomandibular Joint Syndrome OR TMJ disorders OR TMJ diseases OR craniomandibular disorders OR craniomandibular dysfunction
Año publicación: 2003–2014	
Embase and Medline ( $n = 1008$ ) Date Publication: 2003–2014	'bruxism'/exp OR bruxism OR 'sleep'/exp OR sleep AND ('bruxism'/exp OR bruxism) OR nocturnal AND ('bruxism'/exp OR bruxism) OR parafunctional AND ('habits'/exp OR habits) AND temporomandibular AND ('joint'/exp OR joint) AND disorders OR temporomandibular AND disorders OR temporomandibular AND ('joint'/exp OR joint) AND dysfunction AND ('syndrome'/exp OR syndrome) OR tmj AND disorders AND [2003–2014]/py

## Results

In all, 4746 potentially eligible articles were identified in the first approach in the 7 databases used (Table 1). Based on the selection by title and abstract, 127 studies were selected. Once tabulated in Excel table, 55 articles were eliminated for being repeated. Of the remaining 72 articles, 33 were eliminated in the reading of the full text for not meeting the inclusion criteria for this review (Table 2). Finally, 39 studies were analysed. Figure 1 summarizes the results described.

### Included studies

In all, 39 articles were analysed in this review. The analysis tables were prepared according to the PICO criteria (Tables 3–5). The articles analysed were summarized in: (a) bruxism diagnosed by polysomnography ( $n = 7$ ); (b) bruxism diagnosed clinically with or without self-report ( $n = 11$ ) and (c) bruxism diagnosed by questionnaire and/or self-report ( $n = 21$ ).

### Characteristics of participants

In relation to gender, five studies ( $n = 5$ ) included only women in their sample.[47,48,59,61,78] Four studies ( $n = 4$ ) did not specify the gender of the participants.[13,60,39,76]

### Quality assessment

None of the reviewed articles obtained the highest score based on NOS. The range of scores was between 2 and 7.

## Discussion

In this review, a high relation was founded between bruxism and TMD, but with a lower to moderate level of evidence.

All studies that were analysed showed a high bias risk in their design due to the lack of randomization and blinding, and sensibility for bruxism diagnostic as well.

Almost half of the included articles based their bruxism diagnostic on the self-report and/or surveys, and less than one-third of them based such diagnostic on self-report and/or clinical findings. Although methods had a low sensibility, 20 from 21 and 9 from 11 studies, respectively, founded a positive association between bruxism and TMD. On the other side, 4 studies from 7 that used PSG for bruxism diagnosis, demonstrated a positive association (Table 6).

The evidence, in an effort to evaluate and validate the sleep bruxism self-report in patients with TMD (predominating myofascial pain), indicate that this method would not be a good clinical predictor in case of a moderate or severe bruxism due to a high rate of false positives.[80] Such authors concluded that self-report is not a good indicator for sleep bruxism in clinical and research activities, and they do not recommended to substitute it by the PSG.[80]


In general, the studies presented a high variability in age range; however, they did not determine the risk to have bruxism with a TMD. Also, they did not demonstrate an association between gender and bruxism although there was a higher prevalence of TMD in women, suggesting a higher prevalence of bruxism.

Considering circadian manifestation of bruxism and its relation with TMD, most of the analysed studies ( $n = 18$ ) did

**Table 2.** Studies retrieved in full text and excluded from the review.

First author and year	Reason for exclusion
Fernandes, 2014 [9]	Relationship between sleep bruxism and tinnitus.
Glaros, 2014 [10]	Headache associated with oral parafunctions (tooth contact time, intensity of tooth contact).
Fernandes, 2013 [11]	Changes in EMG activity and masticatory muscle pain.
Köhler, 2013 [12]	Same data as Köhler [13].
Brandini, 2012 [14]	Analysis is based on the significant association of parafunctional habits with non-carious lesions, there is no correlation with TMD.
Glaros, 2012 [15]	Evaluates only tooth clenching and not bruxism (undiagnosed).
Ommerborn, 2012 [16]	Relationship between sleep bruxism and functional occlusal parameters.
Shedden Mora, 2012 [17]	Related levels of somatization, depression, anxiety and NMMA in patients with chronic TMD.
Chandwani, 2011 [18]	Prevalence study shows no association.
González, 2011 [19]	Case series showing no association.
Lucchesi, 2010 [20]	Sleep disturbances associated with headache.
Meeder, 2010 [21]	Unable to extract data from subjects older than 19 years.
van der Meulen, 2010 [22]	Application of a questionnaire if patients think or believe that bruxism is associated with TMD.
Jesus, 2009 [23]	It was not possible to obtain the article.
Santos, 2009 [24]	TMD and tooth wear, no statistical association.
Van Selms, 2009 [25]	Relate multiple parafunctions and did not use the term bruxism.
Branco, 2008 [26]	Prevalence study shows no association.
Kanehira, 2008 [27]	TMD and bruxism with stress is associated separately (did not study relationship between TMD and bruxism).
Marklund, 2008 [28]	Same data as Marklund 2010 [29].
Mundt, 2008 [30]	Same data as Marklund 2005 [31].
Nagamatsu-Sakaguchi, 2008 [32]	Study in adolescents younger than 19 years.
Oginni, 2007 [33]	Occlusal wear and relationship with TMD. Do not define bruxism or wear cause.
Pizolato, 2007 [34]	Evaluates maximum bite force in TMD and bruxism.
Cassanova-Rosado, 2006 [35]	Unable to extract data from subjects older than 19 years.
Branco, 2006 [36]	Descriptive study. Shows frequencies and establishes no association.
Winocur, 2006 [37]	Study in adolescents younger than 19 years.
Glaros, 2005 [38]	Same data as Glaros 2005 [39].
Kino, 2005 [40]	Questionnaire applied only in subjects over 12 years, it is not possible to extract data.
Matheus, 2005 [41]	It was not possible to obtain the article.
Aydin, 2004 [42]	Turkish language is not in the inclusion criteria.
Johansson, 2004 [43]	Same data as Johansson 2003 [44] and Johansson 2006 [45].
Johansson, 2003 [44]	Same data as Johansson, 2004 [43] and Johansson 2006 [45].
Molina, 2003 [46]	No TMD-related bruxism, but characterized and compared signs and symptoms of TMD between the study groups.

EMG: electromyography; NMMA: nocturnal masseter muscle activity; TMD: temporomandibular disorders.

**Figure 1.** Search method, identification, choice and inclusion of articles.

**Table 3.** Summary of bruxism diagnosed based polysomnography (PSG).

Author and year	Type of study	Population	Intervention	Comparison (control group)	Outcome	Conclusions	Comments
Raphael, 2013 [47]	Cases and controls study	124 patients with myofascial pain. 46 controls. Age 39.2 (14.6), a.r.: 19–78 years, All women.	TMD: RDC/TMD Bruxism: self-report, individual interview and PSG $\times$ 2 nights. 6 channels. Phasic episodes: 3 or more explosions (burst) EMG (>0.25 sec and <2.0 sec), Tonics episodes: >2.0 sec.  RDC/SB criteria. More than 4 SB episodes p/h or more than 25 explosions p/h sleep.  <i>t</i> -test and Pearson's correlation test, $p < .05$ .	<i>Case:</i> Group I or myofascial pain according to RDC/TMD. <i>Control:</i> No report of facial pain >1 week in the last 2 years or >1 site tenderness according to RDC/TMD.	TMD higher EMG activity compared to controls. Pain before and after sleeping inversely proportional to RMMA number p/sleepy hour. Increased pain before or after sleep associated with fewer muscular events. High levels of RMS EMG activity during sleep associated with reports of pain before and after sleep. Pain assessed before sleeping or waking is associated with fewer SB.	These data are not able to determine whether the EMG activity during sleep is a risk factor for developing myofascial pain, but supports the hypothesis that a high EMG activity in the dream would be a risk factor for the course of myofascial pain.	No sample size calculation. Without randomization groups. Only women. Blinded and trained examiners in data collection PSG. Validated diagnostic methods.
Raphael, 2012 [48]	Cases and controls study	124 patients with myofascial pain. 46 controls. Age 39.2 (14.6), a.r.: 19–78 years, All women.	TMD: RDC/TMD Bruxism: self-report, individual interview and PSG $\times$ 2 nights. 6 channels. Phasic episodes: 3 or more explosions (burst) EMG (>0.25 sec and <2.0 sec), Tonics episodes: >2.0 sec  RDC/SB criteria. More than 4 SB episodes p/h or more than 25 explosions p/h sleep.  <i>t</i> -test and Pearson's correlation test, $p < .05$ .	<i>Case:</i> Group I or myofascial pain according to RDC/TMD. <i>Control:</i> No report of facial pain in the last 2 years or >1 site tenderness according to RDC/TMD.	Similar pain duration in TMD patients with and without evidence of bruxism PSG. High SB levels (RDC/SB): 10.9% in control and 9.7% in cases. Significantly higher SB in cases (55.3%) than in controls (15.2%).	There would be no relationship between SB and course of myofascial pain in TMD. Treatment of SB should not be considered to maintain or exacerbate TMD myofascial pain.	No sample size calculation. Without randomization groups. Only women. Blinded and trained examiners in data collection PSG. There is no clarity on statistics used, they are expressed in terms of percentages. Validated diagnostic methods.
Rossetti, 2008 [49]	Cases and controls study	26 patients, 17–40 years.	VAS for muscle and joint palpation based on the RDC/TMD scale. One night of PSG application. Calibration on EMG amplification. Episodes of EMG activity >20% of maximum voluntary contraction (MVC) were recorded.  <i>Chi-square test and Fisher's exact test,</i> $p < .05$ .	<i>Active group:</i> 14 patients with TMD (F:8; M:6; Age 27.1 (SD 7.4 years), 17–40 years) without sleep disorders <i>Control group:</i> TMD absence ( $n = 12$ ); 6F, 6M; 27.4 SD 5.2 years old, range 22–40 years old.	No association between bruxism and TMD with $p = .976$ and between bruxism and tenderness, $p = 1.000$ . No difference between groups for variables of sleep. No difference between RMMA in bruxers with and without tenderness ( $p > .05$ ).	Sleep bruxism is not associated with TMD or with tenderness. Pain associated only in some individuals with SB. No information on shielding and calibration of examiners. It applies only 1 night PSG. PSG blind data analysis. Validated diagnostic methods.	

(continued)



Table 3. Continued

Author and year	Type of study	Population	Intervention	Comparison (control group)	Outcome	Conclusions	Comments
Rossetti, 2008 [50]	Cases and controls study	60 individuals; F:48; M:12 (range 19–42 years)	TMD: RDC/TMD myofascial pain (MP) Pain: questionnaire and VAS. Sleep bruxism: PSG, examiner blind to diagnosis. Self-reported tooth clenching during the day was recorded.	Group 1: 30 individuals MP Group 2: 30 healthy individuals.	Significant association between SB and MP ( $p < .04$ ; OR = 3.45, 95% CI 1.07–11.19) Significant association between tooth clenching and MP ( $p < .001$ , OR = 12.0; 95% CI 3.26–44.15)	SB and tooth clenching significantly associated with MP. Tooth clenching more power than SB as a risk factor.	No sample size calculation. Without randomization groups. Blind examiner, without information on calibration. Validated diagnostic methods.
Rompré, 2007 [51]	Cases and controls study	100 subjects with sleep bruxism (SB) without TMD. Age 26.5 ± 0.6 years. 60% women.	$p < .05$ . SB subjects, chosen based on criteria of the AASM. Study PSG for 2 nights (first night for adaptation). Questionnaires: SB, anxiety, sleep habits, stress, fatigue, facial pain, jaw pain upon awakening, masticatory muscles fatigue on awakening. Chi-square test, nonparametric and cluster analysis.	Control group: 43 subjects with no history of SB. 68% women. Age 24.5 ± 0.9 years. Some subjects were excluded after the second night with PSG. Comparison was made between 54 SB versus 34 without SB.	Excluded bruxism was more likely than those included to complain of jaw pain upon awakening and fatigue of masticatory muscles (OR 3.9–4.9). The pain of the excluded bruxism was slightly larger than those included ( $p = .06$ ).	SB-RDC has a high level of discrimination between subjects with sleep bruxism and controls. The pain is often reported among subjects with low frequency SB mandibular muscle contractions.	No calculation of sample size. Without randomization. PSG analysis blind score. No information on examiners calibration. Validated bruxism diagnostic method and TMD not validated.
Camparis, 2006 [52]	Cases and controls study	40 patients (32F, 8M). Age 36.1 ± 11.3 years. a.r.: 17–54 years.	$p < .05$ . Preliminary interview. TMD: RDC/TMD Sleep bruxism (SB): PSG, RDC/TMD axis I and II Fisher's exact test and Mann–Whitney test, $p < .05$ analysis.	Group A (GA): bruxism and TMD ( $n = 20$ ) Group B (GB): bruxism without TMD ( $n = 20$ )	GA: 100% myofascial pain, 10% displacement disk and 85% arthralgia without difference between two groups. 85% of episodes of bruxism related to micro awakenings. Patients without pain 20% versus pain episodes.	There is no conclusive evidence that relationship TMD and sleep bruxism.	No calculation of sample size. Higher proportion of women. Without randomization group. Trained examiner, without information on blind. Validated diagnostic methods.
Baba, 2005 [53]	Cases and controls study	103 subjects: 51F (age 23.7 ± 2.6 years) and 52M (age 24.7 ± years)	Questionnaire 0–4 points for 'symptoms upon awakening' and 0–52 points for 'pain/mandibular function'. Four	No control group.	There was a relationship between EMG activity and joint sounds ( $p < .05$ ). There was no relationship between EMG activity and	Joint noises significantly related to duration of the EMG activity of the masseter muscle when sleeping.	No calculation of sample size. Without randomization group. Examiners blinded and trained for EMG data analysis.

(continued)

**Table 3.** Continued

Author and year	Type of study	Population	Intervention	Comparison (control group)	Outcome	Conclusions	Comments
			trained and calibrated examiners: Maximum mouth opening(score muscle sensitivity (0–24 points) and joint sensitivity, (0–12 points)/joint noise. EMG register of right masseter; EMG device use. Bruxism potential episodes: $>20\%$ MVC (sleep bruxism). Multiple linear regression, $p < .05$ .		questionnaire data, data articular sensitivity and maximum mouth opening ( $p > .05$ ).		Validated bruxism diagnostic method and TMD non-validated diagnostic method.

AASM: American Academy of Sleep Medicine; a/r: age range; CI: confidence interval; EMG: electromyography; F: female; M: male; MP: myofascial pain; MVC: maximal voluntary clenching; OR: odds ratio; PSG: polysomnography; RDC/TMD: research criteria for temporomandibular disorders; RMMA: rhythmic masticatory muscle activity; RMS: root mean square; SB: sleep bruxism; sec: seconds; TMD: temporomandibular disorders; VAS: visual analogue scale.

not distinguish between sleep and awaken bruxism. Also, they did not evidence any tendency for having a specific temporomandibular pathology ( $n = 13$ ) (Table 7). On the contrary, nine studies evidenced a clear relation between sleep bruxism with myofascial pain, followed by arthralgia and intracapsular disorders (Table 7). All studies that used PSG ( $n = 4$ ), two studies that used self-report and/or clinic, and three studies that used self/report and survey founded an association. Five articles differentiated the types of bruxism according to circadian manifestation and they demonstrated an association between bruxism, and joint pathology (noise and pain), and muscle pathology. Bortolotto et al., in 2013, concluded that sleep bruxism should be associated with muscle pain, and awaken bruxism to joint pain.[66] This study had a low evidence rate and a low quality on instruments, so its results could be overestimated. Only one study (Michelotti et al.), using self-report, determined a relation between awaken bruxism and TMD (myofascial pain and disc displacement) with a moderate level of evidence.[69]

Few studies did not distinguish the type of bruxism. This situation is perhaps related to the definition of bruxism that is associated with sleep disorders, and defined by the International Classification of Sleep Disorders or ICDS-2.[81] A contribution to that was published by Lobbezoo et al. who established a bruxism definition based on the circadian manifestation, giving a clear, and an operative definition for this disorder.[1]

Although 33 from 39 studies demonstrated a positive relation between bruxism and TMD, it is not possible to confirm causality or perpetuating relationship with them. There are many other factors involved, and that were not necessarily isolated from the samples of the studies. Current evidence shows facilitating conditions involved in the development of TMD as follows: gender,[82] anxiety,[83] unilateral chewing [84] and facial skeletal structures.[85]

### **Relationship between bruxism and TMDs based on analysis of selected studies**

The seven articles that used PSG for bruxism diagnosis were case and control studies. Also they had a low NOS punctuation scale. In three of them, it was not possible to determine an association between bruxism and TMD, but in four of them a positive relation was established. Raphael et al. diagnosed TMD through RDC/TMD, and demonstrated that the increase in EMG activity during sleep could be a risk factor for myofascial pain.[47] Rossetti et al. [50] related sleep bruxism and awaking clenching by self-report to myofascial pain according to RDC/TMD. Rompré et al. concluded that subjects with sleep bruxism could have pain related to a low frequency of mandibular muscle contractions.[51] Finally, Baba et al. demonstrated a relation between presence of joint noises and an increase in EMG activity during sleep bruxism.[53]

Nine of the studies that used clinical method to diagnose bruxism ( $n = 11$ ) demonstrated a positive association. According to type of pathology, most of them established a relation with the TMJ, including arthralgia,[55,56]

**Table 4.** Summary of studies with clinical diagnosis of bruxism with or without self-report.

Author and year	Type of study	Population	Intervention	Comparison (control group)	Outcome	Conclusions	Comments
Alves, 2013 [54]	Cases and controls study.	80 subjects (73F, 7M).	TMD: clinical examination of signs and symptoms of dysfunction. Bruxism (day and night); clinical examination and questionnaire used in previous study. In addition, a questionnaire developed by researchers about mastication and bruxism implications. Student's t-test, Chi-square test, $p < .05$ .	Group 1 (G1): Bruxism, F:37; M:3 (age 33.4 ± 11.5 years) Group 2 (G2): No Bruxism M: 4; F: 36 (age 36.3 ± 10.1 years).	In G1 observed: limited eating mandibular movement ( $p \leq .01$ ), facial pain when chewing ( $p \leq 0.007$ ), facial muscles fatigue ( $p \leq 0.004$ ), headache ( $p \leq 0.02$ ) and articular noise ( $p \leq 0.001$ ).	Masticatory function was reduced in G1, it may be the result of hyperactivity of the masticatory muscles caused by increased muscle tension.	No sample size calculation. Without randomization groups. Examiners trained, without information on blind. No validated diagnostic methods. Analysis does not differentiate types of bruxism.
Fernandes, 2012 [55]	Cases and controls study.	272 participants. F:87.5%; 36.9 m.a.	TMD: Clinical Protocol orofacial pain (questionnaire AAOP) and RDC/TMD (Axis I and II). Sleep bruxism: AASM Chi-square test, 95%, $p < .05$ .	4 groups study: G1: No TMD pain without SB G2: No TMD pain and SB G3: With TMD pain and SB G4: With TMD pain without SB.	Risk of myofascial pain and arthralgia increases in subjects reporting nocturnal bruxism ( $p < .0001$ and $p < .001$ , respectively).	SB patients showed increased myofascial pain and arthralgia.	No sample size calculation. Without randomization groups. Higher proportion women. Trained examiner, without information on shielding, Validated diagnostic methods.
Manfredini, 2010 [56]	Cases and controls study.	276 patients; F: 193; M: 83. Age 32.2 ± 5.7 years, a/r: 25–44 years old.	Analysis of multiple occlusal factors. TMD: RDC/TMD. Bruxism: clinical examination. Multiple logistic regression Chi-square test, $p < .05$ .	G1: Patients diagnosed IIIa group arthralgia. G2: Diagnosis group IIIb osteoarthritis patients.	Univariate analysis showed an association between the presence of TMJ and bruxism ( $p = .025$ ). Multivariate analysis shows no significant association. T significant factors, bruxism and overjet $> = 5$ mm, specificity was 48.2% and 91.5% and sensitivity 65.4% and 18.2%, respectively.	Overbite greater than or equal to 4 mm combined with clinical diagnosis of bruxism (OR = 4.62), greater than or equal 5mm overjet (OR = 2.83) and asymmetric molar ratio combined with clinically diagnosed bruxism (OR = 2.77) have higher chance of TMJ IIIa and IIIb group.	No sample size calculation. Without randomization groups. Trained examiner, without information on blind. No validated diagnostic method bruxism, TMD validated.
Li, 2009 [57]	Cases and controls study.	40 subjects, 24F, 16M.	Bruxism: International Classification of Sleep Disorders (ICSD).	Bruxers: n = 24, 14F, 10M, age 24.38 ± 3.85 years. Divided into three groups.	No significant difference for all parameters TMJ vibration between MB and	No sample size calculation. In the TMJ vibration analysis, it was concluded that bruxism induces abnormal	(continued)

Table 4. Continued

Author and year	Type of study	Population	Intervention	Comparison (control group)	Outcome	Conclusions	Comments
Mehulic, 2009 [58]	Cases and controls study.	Bruxers were grouped according to Molina criteria. TMD: Questionnaire and clinical examination (status of TMJ), maximum opening and closing pattern examination. TMJ evaluation using Bio JVA system to assess damage, internal degenerative conditions. TMJ surface vibration was analysed by two piezoelectric accelerometers. Mann-Whitney analysis and Chi-square test, $p < .05$ . TMD: questionnaires and clinical examination. Index Helkimo. Bumann manual functional diagnosis (clinical examination, joint and muscular palpation) Bruxism: clinical examination and study in articulator. Analysis t-test, Chi-square test, univariate and multivariate regression, $p < .05$ .	Mild bruxism (MB): $n = 9$ (37.5%) Moderate bruxers: $n = 13$ (54.2%) Severe bruxism: $n = 2$ (8.3%) Moderate and severe bruxism same group (MSB) No bruxers: $n = 16$ . 10F, 6M. $23.89 \pm 1.98$ years.	control group ( $p > .05$ ), The total integral, integral $> 300$ Hz, and the peak amplitude of bruxism group was significantly higher in the MB group ( $p < .05$ ). Vibratory radio TMJ: asymptomatic (75%), MB (77.8%) and MSB (100%). The value of MSB group much larger.	vibrations in the TMJ. Moreover, alterations in the TMJ produced by bruxism may be related to the pathogenesis of TMD	shielding and calibration of examiners. Validated diagnostic method bruxism and TMD not validated. Analysis does not differentiate types of bruxism.	
Janal, 2007 [59]	Cases and controls study.	42 subjects, 18–65 years old. F: 72%; M: 28%.	Bruxism subjects ( $n = 22$ ) and patients without bruxism ( $n = 20$ ). Manual functional examination, joint and muscular palpation	No difference in muscle palpation, deviation and deflection opening. Most common neuromuscular dysfunction in bruxism subjects. No difference in muscle condition groups ( $p = .795$ ) with a trend to be more frequent in patients with bruxism. Difference in anterior-medial dislocation with disc replacement in patients without bruxism ( $p = .049$ ).	Bruxers with more common muscle disorders (neuromuscular incoordination). Patients without bruxism have disorders in disk-condyle complex. There are differences in TMD symptoms between the two study groups.	No calculation of sample size. Without randomization groups. Higher proportion women. No information on blind and calibration of examiners. No validated diagnostic method bruxism, TMD validated. Analysis does not differentiate types of bruxism.	
		51F with myofascial pain (ma: $34.5 \pm 11.0$ years old)	Bruxism: 2 weeks to assess changes in tooth wear (TW). Every night questionnaire SCL-90R applied, stressful life, self-reported bruxism, intensity of spontaneous and start pain.	Control group: 12 women	Study fails to support the model in which tooth wear keeps pain. Without demonstrating that tooth grinding or tightening start pain.	Similar proportion of new/old features in 2 weeks (TW). Major TW patients with myofascial pain controls (1.34 versus 1.23)	No sample size calculation. Without randomization groups. Only women. Blinded and trained examiners.

(continued)

Table 4. Continued

Author and year	Type of study	Population	Intervention	Comparison (control group)	Outcome	Conclusions	Comments
Schierz, 2007 [60]	Cases and controls study.	646 participants. a.r: 35–44 years old.	Bruxism: the anterior teeth wear of maxillary and mandibular scale of 0–4 points was considered as an indicator. TMD: self-reported pain through RDC/TMD (German version) or Helkimo index. Chi square test, multiple logistic regression, $p < .05$ .	TMD group versus healthy group.	TW high levels associated with fewer reports of muscle pain or tenderness. Grinding teeth is related to pain palpation report only unilateral pain.	TMJ pain risk increased 11% as increased tooth wear category, without being significant ( $p = .66$ ).	No validated bruxism diagnostic method, TMD validated.
Storm, 2007 [61]	Cases and controls study.	68F. a.r: 21–70 years, m.a: 13.1 ± 49.7 years	TMD: RDC/TMD and Helkimo index. Bruxism: parafunctions determined by questionnaire (tooth wear evaluated). Load test, MVC for 30 sec to cause pain and fatigue. Chi-square test, analysis parametric and nonparametric, $p < .05$ .	Controls: 46 women without TMD Cases: 22 women with TMD.	45% of cases developed symptoms compared with 15% of controls $p = .007$ for the load test. Joint pain in 26% cases with $p = .002$	The engine of the jaw, especially 'tooth clenching' behaviour is significant in patients with TMD.	No calculation of sample size. Without randomization groups. Only women. Blind examiner without calibration information.
Guiler, 2003 [62]	Cases and controls study.	64 patients with clinical diagnosis bruxism behaviour and TMD degenerative disease (52F, 12M; 29 years, a.r: 13–63)	Clinical findings: pre-auricular pain and chewing muscles VAS scale. MNR 1.5 T (128 TMJ). Evaluated position disc, degenerative changes Bruxism: clinical diagnosis (unspecified). Chi-square analysis	Control group: 30 patients without bruxism and TMD. M: 4; F: 26 (26; 14–50 years old). Study group (SG): 64 patients with bruxism and degenerative disease.	High prevalence condyle changes in patients with bruxism. Significant unilateral and bilateral degenerative disease versus control group ( $p < .05$ ). SG relationship between condylar changes and severity of degenerative TMD. DDWR 55%	High prevalence condyle changes in patients with bruxism.	(continued)



Table 4. Continued

Author and year	Type of study	Population	Intervention	Comparison (control group)	Outcome	Conclusions	Comments
Manfredini, 2003 [6]	Cases and controls study.	289 subjects. M: 93; F: 196. Average 34.4 years old; SD: 13.8 years.	<i>Bruxism:</i> (AAOP). No difference between awake and sleep bruxism. <i>Bruxism:</i> when there was one positive aspect for clinical examination and the examination of one positive anamnesis. <i>TMD:</i> questionnaire (subjective description of signs and symptoms) and RDC/TMD, conducted by another examiner. Fisher's exact test and Chi-square test, $p < .05$ .	Subjects without TMD ( $n = 77$ ) Without bruxism and with bruxism. TMD subjects ( $n = 212$ ) with bruxism and without bruxism.	Bruxism has a greater relationship with muscle disorders than with joint pathology.	versus 38% in bruxism control. Compared to 86% DDwR TMD study group versus 24% control. Difference in prevalence of bruxism between patients with and without TMD ( $p < .05$ ). By pathology group RDC/TMD, there were different subjects without TMD and prevalence bruxism with myofascial pain ( $p < .05$ ), myofascial pain + displacement disc ( $p < .01$ ), myofascial pain + displacement disc, $p < .05$ .	No calculation of sample size. Without randomization group. Trained examiner, without information on blind. Bruxism and TMD diagnosis method validated. Analysis does not differentiate types of bruxism.
Pergamalian, 2003 [63]	Cases and controls study.	84 TMD patients (84% F-16% M; $29.1 \pm 8.1$ years old)	TMD: RDC/TMD, TW using models Calibrating two examiners <i>Bruxism:</i> Questionnaire, presence/frequency ANCOVA, MANOVA, Pearson's correlation test, $p < .05$ .	Independent variable bruxism No bruxism, $n = 10$ Occasional bruxism, $n = 27$ Frequent bruxism, $n = 40$ Seven subjects excluded for inconsistencies in the interview and questionnaire.	Numbers joint pain sites varies according to classification of bruxism ( $p < .05$ ), muscle pain of sites does not change ( $p = .52$ ).	No association between myofascial pain (MP) and TW. Bruxism is related to high levels of muscle pain. Report of minimum bruxism and non-bruxism was associated with high levels of TMJ pain.	No calculation of sample size. Without randomization groups. Examiners blind and calibrated. Diagnosis of bruxism no validated method, TMD validated.

AAOP: American Academy of Orofacial Pain; AASM: American Academy of Sleep Medicine; a.i: age range; DDwR: displacement disc without reduction; DDwR: displacement disc with reduction; F: female; M: male; m.a: mean age; MNR: magnetic nuclear resonance; McN: maximal voluntary clenching; OR: odds ratio; RDC/TMD: research criteria for temporomandibular disorders; SB: sleep bruxism; TMD: temporomandibular disorders; TMJ: temporomandibular joint; TW: tooth wear.

**Table 5.** Summary of bruxism diagnosed based questionnaire/self-report studies.

Author and year	Type of study	Population	Intervention	Comparison (control group)	Outcome	Conclusions	Comments
Blanco Aguilera, 2014 [64]	Cases and controls study	1220 patients. 1020 F (84.4%) y 190 M (15.6%) a.r.: 18–60 years	TMD: RDC/TMD. Calibrated examiners. Previous history of joint blockage 14 question as RDC/TMD Pain intensity (CPI) scale using GPS (0–10) Sleep bruxism (SB) questionnaire based on questions 15c modified RDC/TMD Logistic regression and t-test, $p < .05$ .	G0: No muscle pain (n = 29) G1: muscle pain (MP) (n = 127) G2: Muscle pain and discopathy (n = 237) G3: Muscle pain and arthropathy (n = 350) G4: Muscle pain + discopathy + arthropathy (n = 377) G5: Joint locking.	G4 (OR = 2.04) and G3 (OR = 1.64) significance only with respect to G0. G2 close to statistical significance OR = 1.48). G5 without association with perceived bruxism (OR = 1.17). SB and pain intensity: higher in the group with regard to perception of bruxism without perception of bruxism group ( $p < .0001$ ).	Strong association between SB and the presence of painful symptoms of TMD, especially muscle pathology accompanied by arthralgia. No significant difference in reporting the presence of bruxism and disc displacement.	No sample size calculation. Without randomization group. Higher proportion women. No information on blind and calibration of examiners. SB no validated diagnostic method, and TMD validated.
Ferreira, 2014 [65]	Cases and controls study	201 undergraduate students. a.r.: 17–34 years, m.a.: 20.5 years, 146F, 55M.	TMD: RDC/TMD validated for Brazil. Bruxism: Self-reported parafunctional habits and application of questionnaire 2 questions. Occlusal examination, evaluating discrepancy between MfC and centric relation (CR) with Jig wore. Evaluation interference in laterality and protrusion. Evaluation: overjet, overbite, open bite and crossbite.	TMD subjects (group I or G-MPD-group II or G-DD) or without TMD versus subjects with presence or absence of parafunctional habits (teeth grinding and clenching). Subjects with or without TMD versus occlusal characteristics (overjet, overbite, open bite, crossbite). TMD versus functional alterations.	Tooth clenching: 85 subjects (42.3%), with G-MPD, n = 26 and G-DD with n = 13. Tooth grinding 26 subjects; subjects diagnosed with G-MPD, n = 7 and subjects diagnosed with G-DD, n = 6. Only Tooth clenching association with G-MPD ( $p = .000$ ). No difference between occlusal factors and TMD groups. Tooth clenching + overjet correlate with myofascial pain.	Only tooth clenching and overjet were associated with myofascial pain (G-MPD).	No sample size calculation. Without randomization group. Higher proportion women. Calibrated examiner, without information about blind.
Bortolotto, 2013 [66]	Cases and controls study	172 subjects. 69.19% F, 30.81% F. m.a: 34.82, a.r: 17.70–78 years.	Awake and sleep bruxism: AAOP questionnaire validated, self-administered, nine questions with VAS. TMD: Anamnesis for clinical examination to TMD and wear facets. Calibrated examiners.	TMD subjects with awake and sleep bruxism. TMD patients without awake and sleep bruxism.	Awake bruxism: Joint pain, 22.39% ( $p = .042$ ). Pain maximum aperture 13.48% ( $p = .011$ ). Subjects awake bruxism at risk 2.1 times for joint pain ( $p = .044$ ). Sleep bruxism: facial pain 48.35% ( $p = .003$ ). 2.5-fold risk for muscle pain ( $p = .004$ ).	Awake bruxism habit is the most common and is associated with joint pain, followed sleep bruxism associated with muscle pain, both are risk factor for TMD. The other habits studied did not have the same association.	No calculation of sample size. Without randomization group. Calibrated examiners, without information on shielding. Validated diagnostic method bruxism and TMD no validated method.
Köhler, 2012 [13]	Cases and controls study	Population study in 1983, 1993 and 2003.	Questionnaire: bruxism, facial trauma history and TMD treatment.	Three random samples by year (1704 subjects);	All symptoms strongly associated with tooth wear and	Report bruxism increased during the study period	No sample size calculation.  (continued)



Table 5. Continued

Author and year	Type of study	Population	Intervention	Comparison (control group)	Outcome	Conclusions	Comments
Manfredini, 2012 [67]	Cases and controls study	1704 subjects a.r: 20–70 years.	TMD: interview and Helkimo index. Binary logistic regression. Univariate regression (UR) and multiple regression analysis (MR), $p < .05$ .	1983, 1993 and 2003 grouped by 100 subjects in each age group as 20, 30, 40, 50, 60 and 70.	crepitus). Bruxism symptoms dependent: dislocation OR = 3.9 (UR/MR), $p < .001$ ; difficulty opening UR, OR = 3.1, $p < .0001$ ; MR: OR = 3.9, $p < .001$ ; pain jaw movement, UR: OR = 5.2, $p < .001$ ; MR: OR = 4.9, $p < .0001$ . Bruxism, dependence for dysfunction Index (Ai) UR: OR = 2.6, $p < .0001$ ; MR: OR = 2.4, $p < .0001$ .	clench (except crepitus). Bruxism symptoms mostly associated with TMD symptoms and dysfunction index.	and deterioration of health perception were mostly associated with TMD symptoms and dysfunction index.
Yachida, 2012 [68]	Cases and controls study	115 individuals. 39M. m.a: 36.8 ± 14.0 years. 76F m.a: 32.9 ± 10.2 years.	TMD: RDC/TMD Awake and sleep bruxism: self-report and questionnaire. Chi-square test, $p < .05$ . a.r: 18–81 years. Tel Aviv University: n = 397. F: 79.6%. 35.6 years (±14.7). a.r: 18–84 years.	Group 1: Padova University, Italy patients. F: 74.4%. 42.9 years (±16.1).	Myofascial pain more prevalent in Tel Aviv (36.8%) combined with inflammatory and degenerative disorders in Padova (27.4%), $p < .001$ . At least one positive response questionnaire on bruxism, $p < .001$ . More bruxism in patients with myofascial pain only.	The characteristics of samples studied and the different interpretation of the same pattern of diagnosis may influence the epidemiological reports of bruxism and TMD and relationship between them.	Randomization groups (method not explained). Trained interviewers, without information on blind. No validated diagnostic method bruxism, and TMD method validated.
				CP subjects versus No pain versus TH versus Healthy.	Significant difference in EMG activity of CP subjects with SB and without pain with SB ( $p < .05$ ). EMG positive correlation between activity and number of painful muscles – pain intensity – McGill questionnaire and levels of depression ( $p < .05$ ).	No sample size calculation. Without randomization groups. Formation of groups is indicated. Calibrated examiners. No blind information. Validated diagnostic methods.	(continued)

**Table 5.** Continued

Author and year	Type of study	Population	Intervention	Comparison (control group)	Outcome	Conclusions	Comments
Marklund, 2010 [29]	Cohort study	280 students 98M, 182F.	Bruism: questionnaire TMD: RDC/TMD and frequency questionnaire and symptoms and TMD.	Cases: report signs and symptoms incidence or maintenance TMD Control: no signs and symptoms Without persistence and incidence TMD	Self-reported bruxism associated with signs and symptoms start in the TMJ. Predictive incidence of positive bruxism (0.47) and negative (0.75). Self-reported bruxism OR 2.4. Persistence TMD-Bruxism OR = 3.1.	The self-reported bruxism and crossbite can increase the risk in the incidence and duration of signs and symptoms of TMD	No sample size calculation. No randomization group. Blinded and calibrated examiners. No validated diagnostic method bruxism, TMD validated.
Michelotti, 2010 [69]	Cases and controls study	Total subjects: 668 a.r: 11–79 years. TMD diagnosis 557 subjects 33.2 years; SD 14.4 years. M: 127 (a.r: 11–79 years) 34.5 ± 15.4 years. F: 430 (a.r: 12–76 years); 32.9 ± 14.1 years.	TMD: RDC/TMD Axis I. Bruxism: patient questionnaire, applied with respect to grinding or clenching tooth day according to modified questions RDC/TMD (question 15 Axis 1: during the day you keep the teeth in contact, is this what makes your jaw grinding or clenching? Multinomial logistic regression test, $p < .05$ .	Control group (without pathology TMD). N= 111 subjects (55M, 56F) Group 25–37 age group; n = 34 Group 38–79 years; n = 54 Pathology group: (Group I–II and III RDC/TMD) n = 557 Group II: 408; Group II: 133; Group III: 16	Report of tooth clenching and grinding in the day 2 times more frequent in subjects with TMD compared to the control group and is a risk factor for myofascial pain and disk displacement. No association parafunctional oral and arthralgia, arthritis and osteoarthritis.	Parafunctional daytime activities can be a risk factor for TMD subgroup. More specifically, tooth clenching and grinding of the day was a risk factor for myofascial pain and disk displacement.	No sample size calculation. Without randomization group. Calibrated examiner without blind information. No validated diagnostic method bruxism, TMD validated. No clear definition of awake bruxism.
Österberg, 2007 [70]	Cohort study	Two cohorts of subjects 70 years of age born in 1922 (n = 422); F:232; M:190 1930 (n = 484); F:249, M:235	Two cohorts: subjects examined interval 8 years. TMD symptoms questionnaire index 0–5 (5 questions). Bruxism: questionnaire, one question. Pearson correlation analysis, logistic regression model, $p < .05$ .	Cohort 1 (examination year 1992–1993); n = 422; F: 232; M: 190. Cohort 2 (examination year 2000/2001); n = 484; F: 249; M: 235.	High correlation of TMD and bruxism index, $p < .001$ . Index TMD significant association, $p < .001$ ; OR= 2.23. $p = .942$ , OR = 0.957.	TMD symptoms associated with bruxism and psychosomatic factors and overall health.	No calculation of sample size and randomization group. Only one age group. No information examiners calibration or shielding. No validated diagnostic methods.
Camparis, 2006 [71]	Cases and controls study	100 patients. F: 80; M: 20. a.r: 13–66 years; SD 36.1 years ± 11.3	Group A (GA): bruxism without orofacial pain; n = 30, 17–66 years, mean 33 years, M:18; F:12	Prevalent symptoms in Group B (tooth grinding and clenching daytime, uncomfortable bite,	There are significant differences in long-standing bruxism with and without	No sample size calculation. Without randomization group.	(continued)

Table 5. Continued

Author and year	Type of study	Population	Intervention	Comparison (control group)	Outcome	Conclusions	Comments
Johansson, 2006 [45]	Cases and controls study	Two samples: 50 years ( $n = 12,468$ ) and 60 years ( $n = 6232$ ). M: 50.6%, F: 49.4%.	Orofacial pain questionnaire: 53 questions, 123 items on oral and general health. Logistic regression analysis.	Group B(GB): bruxism with orofacial pain; n=70; 13–59 years, 37.5 years; 62M, 8H	morning stiffness, clicking on TMJ and tinnitus. GA: myofascial pain (10.0%), offset disc (10.0%), arthralgia (6.7%), GB: myofascial pain (95.7%), disk displacement (22.8%) and arthralgia (77.1%). TMD pain 12.1%. Bruxism: 79.9% of TMD pain. Bruxism: odds ratio for TMD pain (OR= 4.2). Difficulty opening mouth (OR= 2.0).	chronic facial pain. The frequency and location of the bilateral pain was statistically different regarding no pain bruxism.	Higher proportion women. Trained examiner without blind information. No validated diagnostic method bruxism, TMD validated.
Sato, 2006 [72]	Cases and controls study	508 TMD patients with more than 1 week.	TMD: questionnaire RDC/TMD Axis I. Questionnaire: (1) Background personal; (2) Subtype TMD Axis I RDC/TMD; (3) Value VAS pain intensity. Bruxism: questionnaire. 25 questions. Asked: do you often let your upper and lower teeth contact continuously during your work or at rest?	Association between TCH and TMD was analysed in 229 chronic pain (CP) patients (85.6% women, mean 32 years). Improvement and no improvement (compared to pain levels presented in the first visit).	Patients wear by Teeth contacting habit (TCH) was 52.4% in patients with chronic pain in TMD. TCH is a risk factor for TMD (OR= 1.994).	Half TMD patients had TCH. TCH could be a prolongation factor TMD pain.	No calculation of sample size. Without randomization group. No information examiners calibration or blind. No validated diagnostic methods.
van der Meulen, 2006 [73]	Cohort study	Two cohorts of patients 'Cohort frequency or CF': 226 patients F: 88.5%; a.r: 13–76 years, m.a: 38.5 ± 13.3 years. 'Cohort degree of stress or CDS' 303 patients. F: 83.8%; a.r: 14–83 years; 37.2 ± 14.2 years.	Analysis, logistic regression, p < .05. Validated questionnaire (12 questions parafunctional habits after 2–3 weeks TMD diagnosis). Difference between two cohorts questionnaire application. Cohort frequency: 12 parafunctions occurrence. Ordinal scale of 5 points to respond. Cohort degree of stress: questions as perceived stress level.	No control group	There is no relationship between the BRUX scale and CPI values.	No clinical relevance related to different types of oral parafunctions with self-report and discomfort for TMD. Causal relationship between TMD and bruxism if exists, is small.	No sample size calculation. Absence control group. No information examiners calibration or blind. Validated diagnostic methods.

(continued)

Table 5. Continued

Author and year	Type of study	Population	Intervention	Comparison (control group)	Outcome	Conclusions	Comments
Ahlberg, 2005 [74]	Cases and controls study	1500 subjects. Group work shifts. M: 45 years ( $\pm 10.6$ ); F: 42.6 years ( $\pm 10.7$ ) Workers day: M: 47.4 years ( $\pm 9.7$ ); F: 45.5 years ( $\pm 10.1$ ). $p < .05$ .	TMD: RDC/TMD axis II CPI was used; 3 questions of intensity of pain. CPI values 0–100. Bruxism scales: BRUX, BITE, SOFT. Regression analysis, $p < .05$ . TMD: RDC/TMD axis II (chronic pain) Bruxism: standardized questionnaire, self-frequency of wear or clench tooth (never, rarely, sometimes, often, continuously) Chi-square test, regression model,	Workers with regular shifts, $n = 750$ turn 8 h × day. Men 56.6%. Workers with irregular shifts ( $n = 750$ ). 46.7% men. $p < .05$ .	Orofacial pain associated with frequent bruxism, $p < .001$ ; female, $p < .001$ ; and sleep disruption, $p < .01$ .	Association between perception of orofacial pain and bruxism report. Bruxism together with sleep disruption may participate simultaneously in developing orofacial pain.	No calculation of sample size. No randomization groups. No information blind and examiners calibration. No validated diagnostic method bruxism. TMD validated. Analysis does not differentiate types of bruxism.
Glaros, 2005 [39]	Cases and controls study	96 subjects with TMD in a central facial pain or from general population.	TMD: RDC/TMD by two calibrated examiners. Quiz time and intensity of tooth contacts. VAS scale for pain and stress. Questionnaire yes/no tooth contacts. Linear regression analysis, $p < .05$ .	Subjects with myofascial pain (MP; $n = 24$ ), myofascial pain + arthralgia (MP + A; $n = 21$ ), disk displacement (DD; $n = 24$ ). Subjects without TMD ( $n = 27$ )	Myofascial pain (MP) + high intensity of arthralgia when teeth are in contact. High levels of MP when teeth are in intense contact versus disk displacement and controls. Correlation between muscle tension and jaw pain.	Parafunctions that increase muscle tension and emotional states are good predictors of levels of mandibular pain in TMD patients and healthy subjects.	No calculation of sample size. No randomization groups. Blind examiners. No validated diagnostic method bruxism. TMD validated. No diagnosis parafunctions neither bruxism is defined.
Kobs, 2005 [75]	Cases and controls study	307 subjects. M:140; F:167; a/r: 20–54 years; m/a: 35.4 years	Signs and symptoms of dysfunction in the stomatognathic system (sensitivity joint noise joint, muscle pain, muscle pain opening (active–passive), opening deviation. Chi-square test, $p < .05$ . 20-year follow-up.	TMD subjects: $n = 114$ . Subjects without TMD: $n = 193$	Group clench teeth: 53.1% bilateral masticatory sensitivity and 31.2% in the 'non-clenchers' group ( $p = .001$ )	There is a relationship between the incidence of dental clench and pathological phenomena in the muscles and joints.	No calculation of sample size. No randomization groups. No information blind and examiners calibration. No validated diagnostic methods.
Magnusson, 2005 [76]	Cohort study	320 subjects aged 7, 11 and 15 years.	TMD: Clinical examination. Clinical dysfunction index (DI). Helkimo index. Parafunctions oral questionnaire. Sleep and awake bruxism.	No control group.	Associated symptoms but weakly correlated to tooth-clenching and tooth-grinding ( $r_s$ values ranging between 0.11 and 0.53, $p$ values	A significant correlation between reported bruxism and TMD symptoms. Baseline report of tooth-grinding at night was a predictor	No calculation of sample size. No randomization groups. Calibrated examiners, without information blind.

(continued)

Table 5. Continued

Author and year	Type of study	Population	Intervention	Comparison (control group)	Outcome	Conclusions	Comments
Mundt, 2005 [31]	Cases and controls study	2963 subjects: F:1493; M:1470 a.r.: 35–74 years	Combining daytime tooth-clenching and/or tooth-grinding at night.  Clinical examination to assess the amount of occlusal wear.  Questionnaire symptoms of masticatory system. Logistic regression, $p < .05$ .	ranging between 0.05 and 0.001). Predictor of symptoms of TMD: 3 independent variables: index occlusal wear ( $OR = 4.3$ , $p = .014$ ), clinically recorded TMJ clicking ( $OR = 3.3$ , $p < .0001$ ), and reported tooth-grinding at night ( $OR = 2.2$ , $p = .023$ ). Bruxism + other oral parafunctions ( $OR = 7.7$ , $p = .0031$ ), and bruxism ( $OR = 5.3$ , $p = .016$ ).	In men and women, the presence of bruxism is associated with tenderness/muscle pain in women ( $p \leq 0.001$ ) and TMJ pain/tenderness in women ( $p = .001$ ) and men ( $p = .028$ ).	No calculation of sample size. Without randomization group. Calibrated examiners without blind information. No validated diagnostic method of bruxism and validated method for TMD.	No validated diagnostic bruxism. TMD method validated. Analysis does not differentiate types of bruxism.
Miyake, 2004 [77]	Cases and controls study	3557 students, 20.4 ( $\pm 2.1$ years) a.r.: 18–26 years M: 2516; F:1041	TMD: 3 questions symptoms questionnaire. According to Onizawa and Yoshida study.  Parafunctions oral questionnaire: nine questions. Two questions for sleep and awake bruxism.  Chi-square analysis. Alfa 5%.	Association between symptoms of TMD and oral parafunctions through questionnaires (no control group)	Tooth clench increases risk noises in the temporomandibular joint ( $OR = 1.86$ , $p < .001$ ), TMJ pain ( $OR = 1.79$ , $p = .001$ ) and impaired mouth opening ( $OR = 1.88$ , $p < .001$ ).	Association between parafunctional activities and symptoms of TMD.	No calculation of sample size. Without randomization group. Blind examiners, without information on calibration. No validated diagnostic methods. No control group

(continued)

**Table 5.** Continued

Author and year	Type of study	Population	Intervention	Comparison (control group)	Outcome	Conclusions	Comments
Fujita, 2003 [78]	Cases and controls study	57 women examined between 1995 and 1998. 12 years and 10 months–42 years and 2 months. m.a.: 23 years and 6 months.	TMD: 'The Craniomandibular Disorders Protocol'. Questions and clinical examination. Bruxism: self-report.	Subjects with TMD (primary symptoms and symptoms) and subjects with habits (bruxism, unilateral chewing, crunch, tongue thrusting, abnormal posture and other habits).	Primary symptoms: 40.4% joint noises, 26.3% sound articulate and pain. Presenting symptoms: Most common noise and joint pain (35.1%), 87.7% had oral habits. Bruxism present in 47.4%. Subjects with bruxism and unilateral chewing showed more complex symptoms.	Comparing primary habits, patients with bruxism and unilateral chewers were more complex symptoms of TMD.	No sample size calculation. Only women. Without randomization group. No information blind or calibration of examiners. Unclear methods and unvalidated diagnoses. Analysis does not differentiate types of bruxism.
Velly, 2003 [79]	Cases and controls study	183 patients (a.r.: 18–60 years)	TMD: myofascial pain (MP) by RDC/TMD Bruxism: self-report questionnaire Two blind examiners. Multivariable logistic regression, $p < .05$ .	G1: 83 patients. Myofascial pain (MP) (M:16, F:67) G2: 100 patients. Control	Only tooth clenching and tooth grinding and clenching were associated with MP. Only grinding was not associated with MP.	Tooth clenching (with or without grinding) are associated with chronic MP.	No calculation of sample size. Without randomization group. Blind examiners, without information on calibration. No validated diagnostic method bruxism, TMD validated.

AAOP: American Academy of Orofacial Pain; a.r.: age range; C.I.: confidence interval; CPI: characteristic pain intensity; EMG: electromyography; F: female; M: male; m.a.: mean age; OR: odds ratio; RDC/TMD: research criteria for temporomandibular disorders; SB: sleep bruxism; TCh: teeth contacting habit; TMD: temporomandibular joint; TMJ: temporomandibular disorders; VAS: visual analogue scale.

osteoarthritis,[56] disc displacement and myofascial pain,[6] degenerative pathology,[62] joint pain,[61] joint noises and joint pathology (with no specification).[57] Also, they observed a relation between bruxism with myofascial pain,[55] muscles disorders,[58] myofascial pain and disc displacement,[6] muscle pain,[63] and facial pain during chewing.[54] These studies had a low to moderate evidence level (NOS Scale) and also, most of them, did not make a distinction between sleep and awaken bruxism.

According to the last international consensus, bruxism diagnosis made by self-report is classified as possible bruxism.[1] This definition of bruxism was used in half of the articles that were analysed. Nevertheless, Raphael et al. concluded that the self-reported bruxism is not a useful

indicator.[80] This is the cause of a high positive relation between oral para-functions and pain that those studies presented, compared with those that used PSG for diagnosing bruxism.

So, the relation between bruxism, TMD and pain could be determined by the diagnostic instruments and also by pre-conceived ideas that patients have on bruxism.[86] For that reason, Paesani et al. performed a correlation study to find out the relation between bruxism diagnostic performed by self-report by a survey and performed by clinical history and clinical exam. They concluded that there is a positive correlation between self-report and clinical findings in diagnosis of awaken bruxism.[86]

In conclusion, the importance of these studies conclusions is to validate that the association between bruxism, TMD and orofacial pain would depend on the diagnostic instruments for all pathologies.

**Table 6.** Summary of studies with different methodologies for the diagnosis of TMD and bruxism [337].

TMD diagnostics	Bruxism's diagnostics		
	Polysomnographic	Clinics/ self-report	Questionnaire/ self-report
RDC/TMD	n = 5	n = 6	n = 10
Helkimo index	n = 0	n = 1	n = 2
AAOP	n = 0	n = 0	n = 1
Signs and symptoms	n = 1	n = 2	n = 2
Questionnaire	n = 1	n = 0	n = 5
Others	n = 0	n = 2	n = 1
Total	7	11	21

AAOP: American Academy of Orofacial Pain; RDC/TMD: research diagnostic criteria for temporomandibular disorders.

**Table 7.** Bruxism association between temporomandibular disorders, according to the classification of bruxism, temporomandibular pathology type and quality of evidence according NOS scale.

Author	Year	Relationship bruxism – TMD			Without relationship bruxism – TMD		
		Bruxism's type	Pathology (TMD)	Score	Author	Year	Score
Raphael [47]	2013	Sleep bruxism	Myofascial pain	5	Raphael [48]	2012	5
Alves [54]	2013	Sleep and awake bruxism	Joint/facial pain	3	Rossetti [49]	2008	5
Blanco Aguilera [64]	2014	Sleep bruxism	Muscular/arthalgia	6	Janal [59]	2007	2
Ferreira [65]	2014	Bruxism	Myofascial pain	2	Schierz [60]	2007	4
Bortolotto [66]	2013	Sleep and awake bruxism	Muscular (SB) – Joint (AB)	3	Camparis [52]	2006	3
Fernandes [55]	2012	Sleep bruxism	Arthralgia – myofascial pain	4	van der Meulen [73]	2006	2
Köhler [13]	2012	Bruxism	Signs and symptoms	5			
Manfredini [67]	2012	Sleep and awake bruxism	Myofascial pain	5			
Yachida [68]	2012	Sleep bruxism	Craniofacial pain	5			
Manfredini [56]	2010	Bruxism	Arthralgia – osteoarthritis	5			
Marklund [29]	2010	Bruxism	Signs and symptoms	4			
Michelotti [69]	2010	Awake bruxism	Myofascial pain – DD	4			
Li [57]	2009	Sleep bruxism	Joint	4			
Mehulić [58]	2009	Bruxism	Muscular	3			
Rossetti [50]	2008	Sleep bruxism	Myofascial pain	5			
Rompré [51]	2007	Sleep bruxism	Pain	4			
Storm [61]	2007	Bruxism	Joint	7			
Osterberg [70]	2007	Bruxism	Signs and symptoms	4			
Camparis [71]	2006	Sleep bruxism	MP/DD/arthalgia	3			
Johansson [45]	2006	Bruxism	Signs and symptoms	4			
Sato [72]	2006	Bruxism	TMJ pain	3			
Ahlberg [74]	2005	Bruxism	Orofacial pain	3			
Baba [53]	2005	Sleep bruxism	Joint noises	3			
Glaros [39]	2005	Bruxism	Mandibular pain	4			
Kobs [75]	2005	Bruxism	Muscular and joint	3			
Magnusson [76]	2005	Sleep and awake bruxism	Signs and symptoms	5			
Mundt [31]	2005	Bruxism	Muscular and joint pain	5			
Miyake [77]	2004	Sleep and awake bruxism	Joint (pain, noises)	3			
Fujita [78]	2003	Bruxism	Joint (pain, noises)	2			
Güler [62]	2003	Bruxism	Joint (degenerative)	4			
Manfredini [6]	2003	Bruxism	Muscular/muscular + joint	4			
Pergamalian[63]	2003	Bruxism	Muscular (pain)	4			
Velly [79]	2003	Bruxism	Myofascial pain	5			

AB: awake bruxism; DD: disk displacement; MP: myofascial pain; SB: sleep bruxism; TMD: temporomandibular disorders; TMJ: temporomandibular joint.

patients with myofascial pain as well, an increasing secretion of peripheral serotonin (5HT) and glutamate on the patients was observed.[88,89] Experimental models based on conscious dental clenching generating isometric contractions by prolonged periods are considered valid and adequate to reproduce pain characteristics of subjects with myofascial pain.[87] Many authors support this method. Svensson et al. concluded that with 15 min of continuing clenching at 25% of the maximal voluntary clenching force (MVCF), patients should feel a moderate level of pain without presence of mechanic hyperalgesia.[90] The same authors also concluded that with 60 min of continuing clenching at 10% of the MVCF, patients should feel a high level of pain.[91] Other studies have also focused on continuous clenching and obtained different results according to the method. Hedenberg-Magnusson et al. induced a repetitive voluntary clenching at 50% of the MVCF for 30 s each 1.5 min during 30 min, and they observed a low pain level.[92] Also, Torisu et al. [93] concluded that continuous clenching at 10% of the MVCF for 30 min should induce a low pain level. Other studies with different method arrived to the same results, that is, with different models of clenching activity, they obtained a low level of pain.[92] Different results were obtained by Von Korff et al. [94] and Farella et al. [95] who induced voluntary clenching at different intensities for 6 times and 5 min each one. They observed moderate levels of pain persisting for 1 day after and also a short-term mechanic hyperalgesia.

Those studies have been performed inducing tooth clenching out of Maximal Intercuspatation (MIC). Takeuchi et al. simulated an incisive clenching at 10% of MVCF during 2 h for 3 consecutive days. They did not obtain muscle or joint pain, suggesting other associated factors.[96]

On the other side, based on finite element modelling, when an overload due to prolonged tooth clenching was simulated as in bruxism, an increase in stress was observed on retrodiscal tissues. According to the authors, its effect could generate a degenerative pathology as osteoarthritis or disc perforation.[97] As well, Comissso et al. using the same method, simulated a rhythmic pattern with a 10% and a 20% of the MVCF.[98] They concluded that stress level over disc tissues is not proportional to muscle activation, but the situation could damage the articular disc.

In general, it is supposed that this situation should be similar to myofascial pain because similar values have been obtained when data are recorded from subjects with myofascial pain. The matter is that etiology and pathogenesis of myofascial pain are not an acute entity and pain results of all the studies are more similar to delayed onset muscle soreness. So there is still the doubt.

There is evidence about the association between masticatory muscular activity of the temporal and masseter muscles during sleep and awaken times, as risk factors for TMD. Despite these findings, the increase in muscle activity cannot be associated with bruxism, due to the low methodological quality of the studies based on non-standardized self-report surveys and poor characterization of bruxism in base to tooth wear, as well.[99]

### **Agreements and disagreements with other reviews**

Two reviews with a similar methodology and objectives with this study were found.[5,100] All agree that there is a lack of a good level of evidence to establish an accurate relationship between bruxism and TMD, and all concluded that there is a low sensibility to establish a correct diagnostic of bruxism.

Discrepancies observed were mainly due to search method, and inclusion-exclusion criteria, as well. Comparing with Manfredini et al.'s article,[5] it can be observed that they did not establish differences between sleep and awaken bruxism. Also, inclusion criteria concerning language, in this review were broader. Moreover, Cunali et al. [100] only evaluated the relationship between sleep bruxism and TMD through a PSG analysis and RDC/TMD instrument, respectively, and concluded that it is not possible to establish a positive relation between both pathologies.

In relation to other studies that did not use similar search method and analysis to this research, they concluded that bruxism could be considered as a risk factor for TMD and headache.[101,102] The main discrepancies between these studies and this one lie on design, analysis method and data recruitment. So their conclusions could be overestimated.

Due to the evidence level, variability on diagnostic methodology for bruxism and TMD, a meta-analysis is not feasible.

### **Conclusion**

Establishing a relationship between bruxism and TMD has been one of the most controversial aspects to demonstrate in literature and although many studies and reviews have been made, it is not yet possible to consistently conclude from high-quality studies a relationship between sleep and awake bruxism and TMD.

In relation to the findings in this review, we can conclude the following:

- Evidence-based diagnosis with PSG was not as conclusive as those performed based on clinical diagnostic survey to determine the bruxism-TMD relationship, and mostly was not established.
- Sleep bruxism would be associated with myofascial pain, arthralgia and joint pathology as disc displacement and joint noises.
- Despite the presence of studies with heterogeneous designs, and based on the studies reviewed, it may be suggested that subjects with sleep and awake bruxism would be associated with the presence of TMD.
- More cohort studies are required, with higher levels of evidence to determine the causal relationship between bruxism and TMD.

### **Acknowledgements**

Thanks to Mr. Juan Fernandez de los Rios from the Language and Translation services of the Faculty of Dentistry, University of Chile for kindly correcting the English spelling and grammar of this paper.

## Disclosure statement

None to declare.

## References

- [1] Lobbezoo F, Ahlberg J, Glaros AG, et al. Bruxism defined and graded: an international consensus. *J Oral Rehabil.* 2013;40:2–4.
- [2] de Leeuw RKG. Orofacial pain: guidelines for assessment, diagnosis and management. Hanover Park (IL): Quintessence Editorial International; 2013.
- [3] Monteiro AAKS. Estimation of blood flow by <sup>133</sup>Xe clearance in human masseter muscle during rest, endurance of isometric contraction, and recovery. *Arch Oral Biol.* 1988;33:561–565.
- [4] Mense S. Nociception from skeletal muscle in relation to clinical muscle pain. *Pain.* 1993;54:241–289.
- [5] Manfredini D, Lobbezoo F. Relationship between bruxism and temporomandibular disorders: a systematic review of literature from 1998 to 2008. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2010;109:e26–e50.
- [6] Manfredini D, Cantini E, Romagnoli M, et al. Prevalence of bruxism in patients with different research diagnostic criteria for temporomandibular disorders (RDC/TMD) diagnoses. *Cranio.* 2003;21:279–285.
- [7] De Meyer MD, De Boever JA. [The role of bruxism in the appearance of temporomandibular joint disorders]. *Rev Belge Med Dent* (1984). 1997;52:124–138.
- [8] Newcastle Ottawa Scale [cited 2015 Oct 12]. Available from: [http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp)
- [9] Fernandes G, Siqueira JT, Godoi Gonçalves DA, et al. Association between painful temporomandibular disorders, sleep bruxism and tinnitus. *Braz Oral Res.* 2014;28:1–7.
- [10] Glaros AG, Hanson AH, Ryen CC. Headache and oral parafunctional behaviors. *Appl Psychophysiol Biofeedback.* 2014; 39:59–66.
- [11] Fernandes G, Franco AL, Gonçalves DA, et al. Temporomandibular disorders, sleep bruxism, and primary headaches are mutually associated. *J Orofac Pain.* 2013;27:14–20.
- [12] Köhler AA, Hugoson A, Magnusson T. Clinical signs indicative of temporomandibular disorders in adults: time trends and associated factors. *Swed Dent J.* 2013;37:1–11.
- [13] Anastassaki Köhler A, Hugoson A, Magnusson T. Prevalence of symptoms indicative of temporomandibular disorders in adults: cross-sectional epidemiological investigations covering two decades. *Acta Odontol Scand.* 2012;70:213–223.
- [14] Brandini DA, Pedrini D, Panzarini SR, et al. Clinical evaluation of the association of noncarious cervical lesions, parafunctional habits, and TMD diagnosis. *Quintessence Int.* 2012;43:255–262.
- [15] Glaros AG, City K, Williams K. Tooth contact versus clenching: oral parafunctions and facial pain. *J Orofac Pain.* 2012;26: 176–180.
- [16] Ommerborn MA, Giraki M, Schneider C, et al. Effects of sleep bruxism on functional and occlusal parameters: a prospective controlled investigation. *Int J Oral Sci.* 2012;4:141–145.
- [17] Shadden Mora M, Weber D, Borkowski S, et al. Nocturnal masseter muscle activity is related to symptoms and somatization in temporomandibular disorders. *J Psychosom Res.* 2012;73: 307–312.
- [18] Chandwani B, Ceneviz C, Mehta N, et al. Incidence of bruxism in TMD population. *N Y State Dent J.* 2011;77:54–57.
- [19] González EJ, Gutiérrez M, Pellitero B. Manifestaciones clínicas de trastornos temporomandibulares en bruxópatas. *Correo Cient Med Holguín.* 2011;15.
- [20] Lucchesi LM, Speciali JG, Santos-Silva R, et al. Nocturnal awakening with headache and its relationship with sleep disorders in a population-based sample of adult inhabitants of São Paulo City, Brazil. *Cephalalgia.* 2010;30:1477–1485.
- [21] Meeder W, Weiss F, Maulén M, et al. Temporomandibular disorders: clinical profile, comorbidity, etiological associations and therapeutic guidelines. *Av Odontoestomatol.* 2010;26:209–216.
- [22] van der Meulen MJ, Ohrbach R, Aartman IH, et al. Temporomandibular disorder patients' illness beliefs and self-efficacy related to bruxism. *J Orofac Pain.* 2010;24:367–372.
- [23] Jesus L, Martins M, Andrade D, et al. Estudo transversal da prevalência de disfunção temporomandibular e bruxismo em estudantes universitários. *Ter Man.* 2009;7:11–14.
- [24] Santos T, Moura A, Piva M, et al. Relação dos desgastes dentários com as disfunções temporomandibulares. *Rev Cir Traumatol Buco-Maxilo-Fac.* 2009;9:73–80.
- [25] van Selms MK, Lobbezoo F, Naeije M. Time courses of myofascial temporomandibular disorder complaints during a 12-month follow-up period. *J Orofac Pain.* 2009;23:345–352.
- [26] Raquel Stumpf B, Carla Stumpf B, Ricardo de Souza T, et al. Frequência de relatos de parafunções nos subgrupos diagnósticos de DTM de acordo com os critérios diagnósticos para pesquisa em disfunções temporomandibulares (RDC/TMD). *Rev Dent Press Ortodon Ortop Facial.* 2008;13:61–69.
- [27] Kanehira H, Agariguchi A, Kato H, et al. Association between stress and temporomandibular disorder. *Nihon Hotetsu Shika Gakkai Zasshi.* 2008;52:375–380.
- [28] Marklund S, Wänman A. Incidence and prevalence of myofascial pain in the jaw-face region. A one-year prospective study on dental students. *Acta Odontol Scand.* 2008;66:113–121.
- [29] Marklund S, Wänman A. Risk factors associated with incidence and persistence of signs and symptoms of temporomandibular disorders. *Acta Odontol Scand.* 2010;68:289–299.
- [30] Mundt T, Mack F, Schwahn C, et al. Association between socio-demographic, behavioral, and medical conditions and signs of temporomandibular disorders across gender: results of the study of health in Pomerania (SHIP-0). *Int J Prosthodont.* 2008;21: 141–148.
- [31] Mundt T, Mack F, Schwahn C, et al. Gender differences in associations between occlusal support and signs of temporomandibular disorders: results of the population-based Study of Health in Pomerania (SHIP). *Int J Prosthodont.* 2005;18:232–239.
- [32] Nagamatsu-Sakaguchi C, Minakuchi H, Clark GT, et al. Relationship between the frequency of sleep bruxism and the prevalence of signs and symptoms of temporomandibular disorders in an adolescent population. *Int J Prosthodont.* 2008;21: 292–298.
- [33] Oginni AO, Oginni FO, Adekoya-Sofowora CA. Signs and symptoms of temporomandibular disorders in Nigerian adult patients with and without occlusal tooth wear. *Community Dent Health.* 2007;24:156–160.
- [34] Pizolato RA, Gavião MBD, Berretin-Felix G, et al. Maximal bite force in young adults with temporomandibular disorders and bruxism. *Braz Oral Res.* 2007;21:278–283.
- [35] Casanova-Rosado JF, Medina-Solís CE, Vallejos-Sánchez AA, et al. Prevalence and associated factors for temporomandibular disorders in a group of Mexican adolescents and youth adults. *Clin Oral Investig.* 2006;10:42–49.
- [36] Stumpf Branco R, Stumpf Branco C, de Souza Tesch R, et al. Frequency of relates of bruxism in patients with temporomandibular disorders. *Rev Bras Med.* 2006;63:464–468.
- [37] Winocur E, Littner D, Adams I, et al. Oral habits and their association with signs and symptoms of temporomandibular disorders in adolescents: a gender comparison. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2006;102:482–487.
- [38] Glaros AG, Williams K, Lausten L, et al. Tooth contact in patients with temporomandibular disorders. *Cranio.* 2005;23:188–193.
- [39] Glaros AG, Williams K, Lausten L. The role of parafunctions, emotions and stress in predicting facial pain. *J Am Dent Assoc.* 2005;136:451–458.
- [40] Kino K, Sugisaki M, Haketa T, et al. The comparison between pains, difficulties in function, and associating factors of patients in subtypes of temporomandibular disorders. *J Oral Rehabil.* 2005;32:315–325.
- [41] Matheus R, Ghelardi I, Vega N, et al. The relationship between parafunctional habits and disc position in symptomatic patients

- with temporomandibular dysfunction. *Rev Bras Odontol.* 2005;62:9–12.
- [42] Aydin G, Keles I, Zog G, et al. [How does the habit of teeth clenching affect the symptoms and signs of patients with temporomandibular disorders?]. *Kulak Burun Bogaz Ihtis Derg* 2004;13:19–24.
- [43] Johansson A, Unell L, Carlsson G, et al. Associations between social and general health factors and symptoms related to temporomandibular disorders and bruxism in a population of 50-year-old subjects. *Acta Odontol Scand.* 2004;62:231–237.
- [44] Johansson A, Unell L, Carlsson GE, et al. Gender difference in symptoms related to temporomandibular disorders in a population of 50-year-old subjects. *J Orofac Pain.* 2003;17:29–35.
- [45] Johansson A, Unell L, Carlsson GE, et al. Risk factors associated with symptoms of temporomandibular disorders in a population of 50- and 60-year-old subjects. *J Oral Rehabil.* 2006;33:473–481.
- [46] Molina OF, dos Santos J, Jr., Nelson S, et al. A clinical comparison of internal joint disorders in patients presenting disk-attachment pain: prevalence, characterization, and severity of bruxing behavior. *Cranio.* 2003;21:17–23.
- [47] Raphael KG, Janal MN, Sirois DA, et al. Masticatory muscle sleep background electromyographic activity is elevated in myofascial temporomandibular disorder patients. *J Oral Rehabil.* 2013;40: 883–891.
- [48] Raphael KG, Sirois DA, Janal MN, et al. Sleep bruxism and myofascial temporomandibular disorders: a laboratory-based polysomnographic investigation. *J Am Dent Assoc.* 2012;143: 1223–1231.
- [49] Rossetti LM, Rossetti PH, Conti PC, et al. Association between sleep bruxism and temporomandibular disorders: a polysomnographic pilot study. *Cranio.* 2008;26:16–24.
- [50] Rossetti LM, de Araujo Cdos R, Rossetti PH, et al. Association between rhythmic masticatory muscle activity during sleep and masticatory myofascial pain: a polysomnographic study. *J Orofac Pain.* 2008;22:190–200.
- [51] Rompré PH, Daigle-Landry D, Guitard F, et al. Identification of a sleep bruxism subgroup with a higher risk of pain. *J Dent Res.* 2007;86:837–842.
- [52] Camparis CM, Formigoni G, Teixeira MJ, et al. Sleep bruxism and temporomandibular disorder: clinical and polysomnographic evaluation. *Arch Oral Biol.* 2006;51:721–728.
- [53] Baba K, Haketa T, Sasaki Y, et al. Association between masseter muscle activity levels recorded during sleep and signs and symptoms of temporomandibular disorders in healthy young adults. *J Orofac Pain.* 2005;19:226–231.
- [54] Alves AC, Alchieri JC, Barbosa GA. Bruxism. Masticatory implications and anxiety. *Acta Odontol Latinoam.* 2013;26:15–22.
- [55] Fernandes G, Franco AL, Siqueira JT, et al. Sleep bruxism increases the risk for painful temporomandibular disorder, depression and non-specific physical symptoms. *J Oral Rehabil.* 2012;39:538–544.
- [56] Manfredini D, Peretta R, Guarda-Nardini L, et al. Predictive value of combined clinically diagnosed bruxism and occlusal features for TMJ pain. *Cranio.* 2010;28:105–113.
- [57] Li X, Lin X, Wang Y. Temporomandibular joint vibration in bruxers. *Cranio.* 2009;27:167–173.
- [58] Mehulic K, Gospic RK, Dundjer A, et al. Optoelectronic pantography diagnostics of temporomandibular disorders in patients with bruxism. *Coll Antropol.* 2009;33:849–856.
- [59] Janal MN, Raphael KG, Klausner J, et al. The role of tooth-grinding in the maintenance of myofascial face pain: a test of alternate models. *Pain Med.* 2007;8:486–496.
- [60] Schierz O, John MT, Schroeder E, et al. Association between anterior tooth wear and temporomandibular disorder pain in a German population. *J Prosthet Dent.* 2007;97:305–309.
- [61] Storm C, Wänman A. A two-year follow-up study of temporomandibular disorders in a female Sami population: validation of cases and controls as predicted by questionnaire. *Acta Odontol Scand.* 2007;65:341–347.
- [62] Güler N, Yatmaz PI, Ataoglu H, et al. Temporomandibular internal derangement: correlation of MRI findings with clinical symptoms of pain and joint sounds in patients with bruxing behaviour. *Dentomaxillofacial Radiol.* 2003;32:304–310.
- [63] Pergamalian A, Rudy TE, Zaki HS, et al. The association between wear facets, bruxism, and severity of facial pain in patients with temporomandibular disorders. *J Prosthet Dent.* 2003;90:194–200.
- [64] Blanco Aguilera A, Gonzalez Lopez L, Blanco Aguilera E, et al. Relationship between self-reported sleep bruxism and pain in patients with temporomandibular disorders. *J Oral Rehabil.* 2014;41:564–572.
- [65] Ferreira FM, Simamoto-júnior PC, Novais VR, et al. Correlation between temporomandibular disorders, occlusal factors and oral parafunction in undergraduate students. *Braz J Oral Sci.* 2014;13:281–287.
- [66] Bortolotto P, Moreira A, Madureira P. Análise dos hábitos parafuncionais e associação com Disfunção das Articulações Temporomandibulares. *Rev Assoc Paul Cir.* 2013;67:216–221.
- [67] Manfredini D, Winocur E, Guarda-Nardini L, et al. Self-reported bruxism and temporomandibular disorders: findings from two specialised centres. *J Oral Rehabil.* 2012;39:319–325.
- [68] Yachida W, Castrillon EE, Baad-Hansen L, et al. Craniofacial pain and jaw-muscle activity during sleep. *J Dent Res.* 2012;91: 562–567.
- [69] Michelotti A, Cioffi I, Festa P, et al. Oral parafunctions as risk factors for diagnostic TMD subgroups. *J Oral Rehabil.* 2010;37: 157–162.
- [70] Osterberg T, Carlsson GE. Relationship between symptoms of temporomandibular disorders and dental status, general health and psychosomatic factors in two cohorts of 70-year-old subjects. *Gerodontology.* 2007;24:129–135.
- [71] Camparis CM, Siqueira JTT. Sleep bruxism: clinical aspects and characteristics in patients with and without chronic orofacial pain. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2006;101:188–193.
- [72] Sato F, Kino K, Sugisaki M, et al. Teeth contacting habit as a contributing factor to chronic pain in patients with temporomandibular disorders. *J Med Dent Sci.* 2006;53:103–109.
- [73] van der Meulen MJ, Lobbezoo F, Aartman IH. a, et al. Self-reported oral parafunctions and pain intensity in temporomandibular disorder patients. *J Orofac Pain.* 2006;20:31–35.
- [74] Ahlberg K, Ahlberg J, Könönen M, et al. Perceived orofacial pain and its associations with reported bruxism and insomnia symptoms in media personnel with or without irregular shift work. *Acta Odontol Scand.* 2005;63:213–217.
- [75] Kobs G, Bernhardt O, Kocher T, et al. Oral parafunctions and positive clinical examination findings. *Stomatologija.* 2005;7: 81–83.
- [76] Magnusson T, Egermark I, Carlsson GE. A prospective investigation over two decades on signs and symptoms of temporomandibular disorders and associated variables. A final summary. *Acta Odontol Scand.* 2005;63:99–109.
- [77] Miyake R, Ohkubo R, Takehara J, et al. Oral parafunctions and association with symptoms of temporomandibular disorders in Japanese university students. *J Oral Rehabil.* 2004;31:518–523.
- [78] Fujita Y, Motegi E, Nomura M, et al. Oral habits of temporomandibular disorder patients with malocclusion. *Bull Tokyo Dent Coll.* 2003;44:201–207.
- [79] Velly AM, Gornitsky M, Philippe P. Contributing factors to chronic myofascial pain: a case-control study. *Pain.* 2003;104:491–499.
- [80] Raphael KG, Janal MN, Sirois DA, et al. Validity of self-reported sleep bruxism among myofascial temporomandibular disorder patients and controls. *J Oral Rehabil.* 2015;42:751–758.
- [81] American Academy of Sleep Medicine. International classification of sleep disorders. 2nd ed. Westchester: American Academy of Sleep Medicine; 2005.
- [82] Bi R-Y, Ding Y, Gan Y-H. A new hypothesis of sex-differences in temporomandibular disorders: estrogen enhances hyperalgesia of inflamed TMJ through modulating voltage-gated sodium

- channel 1.7 in trigeminal ganglion? *Med Hypotheses*. 2015;84:100–103.
- [83] Reissmann DR, John MT, Seedorf H, et al. Temporomandibular disorder pain is related to the general disposition to be anxious. *J Oral Facial Pain Headache*. 2014;28:322–330.
- [84] Santana-Mora U, López-Cedrún J, Mora MJ, et al. Temporomandibular disorders: the habitual chewing side syndrome. *PLoS One*. 2013;8:e59980.
- [85] Manfredini D, Segù M, Arveda N, et al. Temporomandibular joint disorders in patients with different facial morphology. A systematic review of the literature. *J Oral Maxillofac Surg*. 2016;74: 29–46.
- [86] Paesani DA, Lobbezoo F, Gelos C, et al. Correlation between self-reported and clinically based diagnoses of bruxism in temporomandibular disorders patients. *J Oral Rehabil*. 2013;40: 803–809.
- [87] Dawson A. Experimental tooth clenching. A model for studying mechanisms of muscle pain. *Swed Dent J Suppl*. 2013;9–94.
- [88] Ernberg M, Hedenberg-Magnusson B, Alstergren P, et al. The level of serotonin in the superficial masseter muscle in relation to local pain and allodynia. *Life Sci*. 1999;65:313–325.
- [89] Castrillon EE, Ernberg M, Cairns BE, et al. Interstitial glutamate concentration is elevated in the masseter muscle of myofascial temporomandibular disorder patients. *J Orofac Pain*. 2010;24: 350–360.
- [90] Svensson P, Arendt-Nielsen L. Effects of 5 days of repeated submaximal clenching on masticatory muscle pain and tenderness: an experimental study. *J Orofac Pain*. 1996;10:330–338.
- [91] Svensson P, Burgaard A, Schlosser S. Fatigue and pain in human jaw muscles during a sustained, low-intensity clenching task. *Arch Oral Biol*. 2001;46:773–777.
- [92] Hedenberg-Magnusson B, Brodda Jansen G, Ernberg M, et al. Effects of isometric contraction on intramuscular level of neuropeptide Y and local pain perception. *Acta Odontol Scand*. 2006;64:360–367.
- [93] Torisu T, Wang K, Svensson P, et al. Effect of low-level clenching and subsequent muscle pain on exteroceptive suppression and resting muscle activity in human jaw muscles. *Clin Neurophysiol*. 2007;118:999–1009.
- [94] Von Korff M, Dworkin SF, Le Resche L, et al. An epidemiologic comparison of pain complaints. *Pain*. 1988;32:173–183.
- [95] Farella M, Michelotti A, Steenks MH, et al. The diagnostic value of pressure algometry in myofascial pain of the jaw muscles. *J Oral Rehabil*. 2000;27:9–14.
- [96] Takeuchi T, Arima T, Ernberg M, et al. Symptoms and physiological responses to prolonged, repeated, low-level tooth clenching in humans. *Headache*. 2015;55:381–394.
- [97] Abe S, Kawano F, Kohge K, et al. Stress analysis in human temporomandibular joint affected by anterior disc displacement during prolonged clenching. *J Oral Rehabil*. 2013;40:239–246.
- [98] Commisso MS, Martínez-Reina J, Mayo J. A study of the temporomandibular joint during bruxism. *Int J Oral Sci*. 2014;6: 116–123.
- [99] Khawaja SN, McCall W, Dunford R, et al. Infield masticatory muscle activity in subjects with pain-related temporomandibular disorders diagnoses. *Orthod Craniofac Res*. 2015;18:137–145.
- [100] Cunali R, Bonotto D, Machado E, et al. Sleep bruxism and temporomandibular disorders: systematic review. *Rev Dor São Paulo*. 2012;13:360–364.
- [101] Poveda Roda R, Bagan JV, Díaz Fernández JM, et al. Review of temporomandibular joint pathology. Part I: classification, epidemiology and risk factors. *Med Oral Patol Oral Cir Bucal*. 2007;12:292–298.
- [102] Bender SD. Orofacial pain and headache: a review and look at the commonalities. *Curr Pain Headache Rep*. 2014;18:400.