

Body mass index is negatively associated with telomere length: a collaborative cross-sectional meta-analysis of 87 observational studies

Marij Gielen,¹ Geja J Hageman,² Evangelia E Antoniou,³ Katarina Nordfjall,⁴ Massimo Mangino,^{5,6}
 Muthuswamy Balasubramanyam,⁷ Tim de Meyer,⁸ Audrey E Hendricks,^{9,10} Erik J Giltay,¹¹ Steven C Hunt,¹²
 Jennifer A Nettleton,¹³ Klelia D Salpea,¹⁴ Vanessa A Diaz,¹⁵ Ramin Farzaneh-Far,¹⁶ Gil Atzmon,¹⁷ Sarah E Harris,¹⁸
 Lifang Hou,¹⁹ David Gilley,²⁰ Iris Hovatta,^{21,22} Jeremy D Kark,²³ Hisham Nassar,²⁴ David J Kurz,²⁵ Karen A Mather,²⁶
 Peter Willeit,²⁷ Yun-Ling Zheng,²⁸ Sofia Pavanello,²⁹ Ellen W Demerath,³⁰ Line Rode,³¹ Daniel Bunout,³²
 Andrew Steptoe,³³ Lisa Boardman,³⁴ Amelia Marti,^{35,36,37} Belinda Needham,³⁸ Wei Zheng,³⁹ Rosalind Ramsey-Goldman,⁴⁰
 Andrew J Pellatt,⁴¹ Jaakko Kaprio,^{42,43} Jonathan N Hofmann,⁴⁴ Christian Gieger,⁴⁵ Giuseppe Paolisso,⁴⁶
 Jacob BH Hjelmberg,⁴⁷ Lisa Mirabello,⁴⁸ Teresa Seeman,⁴⁸ Jason Wong,⁴⁹ Pim van der Harst,⁵⁰ Linda Broer,⁵¹
 Florian Kronenberg,⁵² Barbara Kollerits,⁵² Timo Strandberg,⁵³ Dan TA Eisenberg,⁵⁴ Catherine Duggan,⁵⁵
 Josine E Verhoeven,⁵⁶ Roxanne Schaakxs,⁵⁶ Raffaella Zannolli,⁵⁷ Rosana MR dos Reis,⁵⁸ Fadi J Charchar,⁵⁹
 Maciej Tomaszewski,^{60,61} Ute Mons,^{62,63} Ilja Demuth,⁶⁴ Andrea Elena Iglesias Molli,⁶⁵ Guo Cheng,⁶⁶ Dmytro Krasnienkov,⁶⁷
 Bianca D'Antono,⁶⁸ Marek Kasielski,⁶⁹ Barry J McDonnell,⁷⁰ Richard Paul Ebstein,⁷¹ Kristina Sundquist,⁷²
 Guillaume Pare,⁷³ Michael Chong,⁷³ and Maurice P Zeegers^{1,74} on behalf of the TELOMAAS group

¹Departments of Complex Genetics; ²Toxicology, NUTRIM School of Nutrition and Translational Research in Metabolism, Maastricht University Medical Center, Maastricht University, Netherlands; ³Department of Clinical Psychological Science, Faculty of Psychology and Neuroscience, Maastricht University, Netherlands; ⁴Department of Medicine, Östersund Hospital, Östersund, Sweden; ⁵Twin Research and Genetic Epidemiology, King's College London, London, United Kingdom; ⁶NIHR Biomedical Research Center at Guy's and St. Thomas' Foundation Trust, London, United Kingdom; ⁷Cell and Molecular Biology, Madras Diabetes Research Foundation, Gopalapuram, Chennai, India; ⁸Department of Mathematical Modeling, Statistics, and Bioinformatics, Ghent University, Ghent, Belgium; ⁹Population Sciences Branch of the National Heart, Lung, and Blood Institute (NHLBI), NIH, NHLBI's Framingham Heart Study, Framingham, MA; ¹⁰Department of Mathematical and Statistical Sciences, University of Colorado-Denver, Denver, CO; ¹¹Department of Psychiatry, Leiden University Medical Center, Leiden, Netherlands; ¹²Cardiovascular Genetics Division, Department of Medicine, University of Utah, Salt Lake City, UT; ¹³Division of Epidemiology, Human Genetics, and Environmental Sciences, University of Texas Health Science Center, Houston, TX; ¹⁴Department of Molecular Biology and Genetics, BSRC "Alexander Fleming," Athens, Greece; ¹⁵Department of Family Medicine, Medical University of South Carolina, Charleston, SC; ¹⁶Division of Cardiology, San Francisco General Hospital, San Francisco, CA; ¹⁷Department of Medicine and Genetics, Albert Einstein College of Medicine, Bronx, NY, and Department of Biology, Faculty of Natural Science, University of Haifa, Haifa, Israel; ¹⁸Center for Cognitive Aging and Cognitive Epidemiology and Medical Genetics Section and Center for Genomics and Experimental Medicine and MRC Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, United Kingdom; ¹⁹Department of Preventive Medicine and Robert H Lurie Comprehensive Cancer Center, Feinberg School of Medicine, Northwestern University, Chicago, IL; ²⁰Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, IN; ²¹Department of Biosciences, University of Helsinki, Helsinki, Finland; ²²Department of Health, National Institute for Health and Welfare, Helsinki, Finland; ²³Epidemiology Unit, Hebrew University-Hadassah School of Public Health and Community Medicine, Jerusalem, Israel; ²⁴Department of Cardiology, Hadassah University Medical Center, Jerusalem, Israel; ²⁵Department of Cardiology, Triemli Hospital, Zurich, Switzerland; ²⁶Centre for Healthy Brain Ageing, Psychiatry, UNSW Australia, Sydney, Australia; ²⁷Department of Neurology, Medical University Innsbruck, Innsbruck, Austria, and Department of Public Health and Primary Care, University of Cambridge, Cambridge, United Kingdom; ²⁸Department of Oncology, Georgetown University Medical Center, Georgetown University, Washington, DC; ²⁹Department of Cardiac, Thoracic, and Vascular Sciences, Unit of Occupational Medicine, University of Padova, Padova, Italy; ³⁰Division of Epidemiology and Community Health, University of Minnesota School of Public Health, Minneapolis, MN; ³¹The Copenhagen General Population Study, Department of Clinical Biochemistry, Copenhagen University Hospital, Herlev and Gentofte Hospital, Copenhagen, Denmark; ³²Institute of Nutrition and Food Technology University of Chile, Santiago, Chile; ³³Department of Epidemiology and Public Health, University College London, London, United Kingdom; ³⁴Division of Gastroenterology and Hepatology, Department of Internal Medicine, Mayo Clinic College of Medicine, Rochester, MN; ³⁵Department of Nutrition, Food Science, and Physiology, University of Navarra, Pamplona, Spain; ³⁶Instituto de Investigación Sanitaria de Navarra, Pamplona, Spain; ³⁷CIBER Fisiopatología de la Obesidad y Nutrición, (CIBERObn), Instituto de Salud Carlos III, Madrid, Spain; ³⁸Department of Epidemiology, University of Michigan, Ann Arbor, MI; ³⁹Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center, Vanderbilt-Ingram Cancer Center, Vanderbilt University Medical Center, Nashville, TN; ⁴⁰Division of Rheumatology, Northwestern University Feinberg School of Medicine, Chicago, IL; ⁴¹Department of Medicine, University of Utah, Salt Lake City, UT; ⁴²Department of Public Health; ⁴³Institute for Molecular Medicine, University of Helsinki, Helsinki, Finland; ⁴⁴Division of Cancer Epidemiology and Genetics, National Cancer Institute, NIH, Bethesda, MD; ⁴⁵Research Unit of Molecular Epidemiology and Institute of Epidemiology II, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany; ⁴⁶Department of Medical, Surgical, Neurological, Metabolic, and Geriatric Sciences, Second University of Naples, Naples, Italy; ⁴⁷Department of Epidemiology, Biostatistics, and Biodemography, Institute of Public Health, University of Southern Denmark, Odense C, Denmark; ⁴⁸Department of Medicine, David Geffen School of Medicine at UCLA, Los Angeles, CA; ⁴⁹Stanford University School of Medicine, Stanford, CA; ⁵⁰Department of Cardiology, University Medical Center Groningen, Groningen, Netherlands; ⁵¹Department of Internal Medicine, Erasmus MC, Rotterdam, Netherlands; ⁵²Division of Genetic

Epidemiology, Department of Medical Genetics, Molecular, and Clinical Pharmacology, Medical University of Innsbruck, Innsbruck, Austria;⁵³University of Helsinki and Helsinki University Central Hospital, Helsinki, Finland; Center for Life Course Epidemiology, University of Oulu, Oulu, Finland;⁵⁴Department of Anthropology and Center for Studies in Demography and Ecology, University of Washington, Seattle, WA;⁵⁵The Fred Hutchinson Cancer Research Center, Seattle, WA;⁵⁶Department of Psychiatry, VU University Medical Center, Amsterdam Public Health Research Institute, Amsterdam, Netherlands;⁵⁷Pediatrics Unit, Azienda Ospedaliera Universitaria, Senese/University of Siena, Policlinico Le Scotte, Siena, Italy;⁵⁸Department of Gynecology and Obstetrics, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, São Paulo, Brazil;⁵⁹School of Science and Technology, Federation University Australia, Department of Physiology, University of Melbourne, Melbourne, Australia, and Department of Cardiovascular Sciences, University of Leicester, Leicester, United Kingdom;⁶⁰Division of Cardiovascular Sciences, Faculty of Medicine, Biology, and Health, University of Manchester, Manchester, United Kingdom;⁶¹Division of Medicine, Manchester University NHS Foundation Trust, Manchester Academic Health Science Centre, Manchester, United Kingdom; Division of ⁶²Clinical Epidemiology and Aging Research; ⁶³Cancer Prevention Unit, German Cancer Research Center (DKFZ), Heidelberg, Germany; ⁶⁴Charité–Universitätsmedizin Berlin (corporate member of Freie Universität Berlin), Humboldt-Universität zu Berlin, and Berlin Institute of Health, Lipid Clinic at the Interdisciplinary Metabolism Center, Berlin, Germany; ⁶⁵CONICET-Universidad de Buenos Aires. Instituto de Inmunología, Genética y Metabolismo (INIGEM). Laboratorio de Diabetes y Metabolismo, Ciudad Autónoma de Buenos Aires, Buenos Aires, Argentina; ⁶⁶Department of Nutrition, Food Safety, and Toxicology, West China School of Public Health, Sichuan University, Chengdu, China; ⁶⁷Department of Epigenetics, DF Chebotarev State Institute of Gerontology NAMS of Ukraine, Kyiv, Ukraine; ⁶⁸Research Center, Montreal Heart Institute, and Psychology Department, University of Montreal, Montreal, Quebec, Canada; ⁶⁹Bases of Clinical Medicine Teaching Center, Medical University of Lodz, Lodz, Poland; ⁷⁰Cardiff School of Sport and Health Sciences, Cardiff Metropolitan University, Cardiff, United Kingdom; ⁷¹Department of Psychology, National University of Singapore, Singapore; ⁷²Center for Primary Health Care Research, Lund University, Region Skåne, Lund, Sweden; ⁷³Population Health Research Institute and McMaster University, Hamilton, Canada; and ⁷⁴CAPHRI School for Public Health and Primary Care, Maastricht University, Maastricht, Netherlands

ABSTRACT

Background: Even before the onset of age-related diseases, obesity might be a contributing factor to the cumulative burden of oxidative stress and chronic inflammation throughout the life course. Obesity may therefore contribute to accelerated shortening of telomeres. Consequently, obese persons are more likely to have shorter telomeres, but the association between body mass index (BMI) and leukocyte telomere length (TL) might differ across the life span and between ethnicities and sexes.

Objective: A collaborative cross-sectional meta-analysis of observational studies was conducted to investigate the associations between BMI and TL across the life span.

Design: Eighty-seven distinct study samples were included in the meta-analysis capturing data from 146,114 individuals. Study-specific age- and sex-adjusted regression coefficients were combined by using a random-effects model in which absolute [base pairs (bp)] and relative telomere to single-copy gene ratio (T/S ratio) TLs were regressed against BMI. Stratified analysis was performed by 3 age categories (“young”: 18–60 y; “middle”: 61–75 y; and “old”: >75 y), sex, and ethnicity.

Results: Each unit increase in BMI corresponded to a -3.99 bp (95% CI: -5.17 , -2.81 bp) difference in TL in the total pooled sample; among young adults, each unit increase in BMI corresponded to a -7.67 bp (95% CI: -10.03 , -5.31 bp) difference. Each unit increase in BMI corresponded to a -1.58×10^{-3} unit T/S ratio (0.16% decrease; 95% CI: -2.14×10^{-3} , -1.01×10^{-3}) difference in age- and sex-adjusted relative TL in the total pooled sample; among young adults, each unit increase in BMI corresponded to a -2.58×10^{-3} unit T/S ratio (0.26% decrease; 95% CI: -3.92×10^{-3} , -1.25×10^{-3}). The associations were predominantly for the white pooled population. No sex differences were observed.

Conclusions: A higher BMI is associated with shorter telomeres, especially in younger individuals. The presently observed difference is not negligible. Meta-analyses of longitudinal studies evaluating change in body weight alongside change in TL are warranted. *Am J Clin Nutr* 2018;108:453–475.

Keywords: BMI, telomere length, obesity, low-grade inflammation, meta-analysis, observational studies

INTRODUCTION

Telomeres, the nucleoprotein structures at the ends of chromosomes, shorten with each cell division in somatic cells (1). When telomere length (TL) reaches a critical value, cells either enter a state of senescence or undergo apoptosis (2). Oxidative stress and chronic inflammation are suggested to play a role in accelerated telomere attrition (3–5). Even before the onset of age-related diseases, obesity might be a contributing factor to the cumulative burden of oxidative stress and chronic inflammation throughout the life course, and obesity may therefore contribute to accelerated shortening of telomeres.

Obesity is a growing health problem, and worldwide its prevalence has more than doubled since 1980 (6). In addition, the burden of diabetes and cardiovascular disease is partly attributable to being overweight and obese (6). Tackling obesity might be a starting point to delay telomere shortening and the onset of age-related diseases. Although obesity is associated with shorter telomeres overall (7), studies in the elderly found no relation between TL and obesity and no relation between TL and mortality (8, 9). We hypothesized that obese persons will have shorter telomeres, compared with those of normal weight of the

The authors reported no funding received for this study.

Present address for AJP: Tulane School of Medicine, New Orleans, LA.

Present address for MG: Maastricht University, Maastricht, the Netherlands.

Supplemental Methods, Supplemental Results, Supplemental Figures 1 and 2, Supplemental Tables 1–3, and Study Protocol for Participating PIs are available from the “Supplementary data” link in the online posting of the article and from the same link in the online table of contents at <https://academic.oup.com/ajcn/>.

Address to correspondence to MG (e-mail: marj.gielen@maastrichtuniversity.nl).

Abbreviations used: bp, base pair; PI, principal investigator; qPCR, quantitative polymerase chain reaction; TL, telomere length; T/S ratio, telomere to single-copy gene ratio.

Received May 15, 2017. Accepted for publication April 27, 2018.

First published online September 11, 2018; doi: <https://doi.org/10.1093/ajcn/nqy107>.

same chronological age, but that the association between obesity and TL will differ across the life span.

Sex and ethnicity may also influence the association between BMI and TL. On average, women have longer telomeres than men (10–12). However, published results on sex differences in associations between BMI and TL are inconsistent (13–15). African Americans and Native Americans have higher rates of obesity (16), and racial differences in TL have frequently been reported in adult African Americans who have longer telomeres than white individuals (17–21), but evidence is lacking with regard to whether the association between BMI and TL differs between ethnicities.

Two recent meta-analyses reported the negative association between BMI and TL on reported summary statistics in the literature, but they did not examine sex differences or the influence of age and ethnicity (7, 22). To further evaluate whether BMI is associated with TL, a large-scale, collaborative, cross-sectional meta-analysis was conducted across observational studies that collected information on BMI and TL of adult individuals. To avoid publication bias and maximize the data in the analyses, a consistent standardized analysis plan across studies was used and principal investigators (PIs) of published studies were contacted and asked to participate in the Telomere Maastricht collaborator (TELOMAAS) group. Because the relation between TL and BMI could be moderated by age, sex, and ethnicity, we completed additional analyses stratifying by these factors.

METHODS

Search strategy

We performed a broad literature search up until 10 November 2017 using PubMed (www.ncbi.nlm.nih.gov/pmc), EMBASE (ovidsp.tx.ovid.com), and the Cochrane (www.cochranelibrary.com) database without restrictions in language or publication date. Numerous studies have measured BMI and TL for purposes other than the association between TL and BMI as an outcome. Therefore, the search was rather broad and not narrowed to TL or BMI. On the basis of the existing relation between obesity, diabetes, and cardiovascular diseases, and because TL is related to aging, we completed a search in which terms related to these conditions were entered. In addition, search items related to study design were entered. The complete search criteria are listed in **Supplemental Methods**. Citation and reference tracking were performed until no new studies were found. One of the authors (MG) performed the literature search and selected potentially relevant publications. Titles and abstracts of potentially relevant studies were screened. In addition, when the abstract indicated that the article reported a study of diabetes or cardiovascular disease, the full text was screened. No additional restrictions for study design were applied.

Eligibility criteria

Studies were included if height and weight or BMI was collected. The corresponding author was invited to participate in the meta-analysis and identified additional unpublished studies. PIs of these unpublished studies were also invited to participate. Cohort studies in healthy individuals at baseline were included,

and if the study design was a case-control study, only controls were included in the meta-analysis. In compiling the database, care was taken to exclude overlapping study cohorts. The study sample (abbreviated as “study”) was taken as the unit for this meta-analysis.

Data extraction

The detailed study protocol can be found in the Supplemental data (“**Study Protocol for Participating PIs**”). The PI of each study completed a questionnaire and additional information was extracted from the manuscript. The following data were collected: study name; study design (cohort or case-control); sample size (cohort size or control group size); presence of the variables age, sex, and ethnicity [when $\geq 70\%$ of the individuals of a sample were of a single ethnicity (e.g., white, African American, Native American, Asian, Hispanic), the sample was classified as a sample of a particular ethnicity; when no ethnicity constituted 70% of the sample, the sample was classified as a mixed sample]; leukocyte TL; and BMI (kg/m^2) and whether BMI was measured or self-reported; white blood cell types from which DNA was extracted for telomere measurements; and method of TL measurement and of DNA storage (Supplemental Material: Study Protocol for Participating PIs). Two metrics were used for TL: absolute TL in base pairs (bp) and relative TL based on telomere to single-copy gene ratio (T/S ratio) (23). A T/S ratio of 0.8 indicates a relative TL, which is 80% of the reference used (100%).

The PI was free to provide the de-identified raw data or to perform linear regression analyses and provide summary statistics. If the PI provided raw data, one of the authors (MG) conducted the linear regression analyses with TL (bp or T/S ratio) as the outcome and BMI as the independent variable to obtain the summary statistics. Three sex groups were defined: men, women, and a combined group of men and women; 4 age groups were defined: “young” (18–60 y), “middle” (61–75 y), and “old” (>76 y) and a combined group of all subjects regardless of their age. For each study, ≤ 12 stratified linear regression analyses were conducted (stratified by the 3 sex groups and 4 age groups). The analyses that included all subjects regardless of sex were adjusted for sex; similarly, the analyses that included all subjects regardless of age were adjusted for age.

If the T/S ratio was used to estimate absolute TL without the use of reference DNA with the known absolute TL, the PI was asked to provide new analyses with the T/S ratio as the outcome. If the PI did not respond to this request, absolute TL based on the T/S ratio was used for analyses and included in the analysis. The regression coefficients (β estimates) and SEs were then used in the meta-analyses. In the case of longitudinal data, one randomly selected measurement of TL along with the corresponding BMI and age for that time point were used in the analysis. The summary statistics thus included the results of 12 linear regression analyses, with TL (bp or T/S ratio) as the outcome and BMI as the independent variable.

Assessment of small study effects

To examine the potential presence of publication bias, visual inspection of funnel plots for asymmetry was performed, followed by the Egger and Begg’s linear regression test for small

study effects (24) and the Duval and Tweedie nonparametric “trim and fill” method (25).

Statistical analysis

Statistical pooling

The primary outcome of the meta-analysis was a pooled estimation of the difference in absolute TL in bp or relative TL (T/S ratio) per unit increase in BMI. Study-specific regression coefficients (β estimates) and SEs were combined by using random-effects pooling in 12 meta-analyses. The assumption of a linear association between BMI and TL was verified by using the raw data provided by the PIs (Supplemental Methods, **Supplemental Results**).

Assessment of heterogeneity

Details are given in Supplemental Methods. Statistical heterogeneity between studies was estimated by using I^2 statistics (26, 27) for each of the 12 meta-analyses. Low heterogeneity was indicated by $I^2 \leq 25\%$, medium heterogeneity by I^2 of 25–50%, and high heterogeneity by $I^2 > 50\%$ (27). To investigate potential effect modification of age and sex, meta-regression analysis was performed with age and sex separately incorporated as covariates. Age was therefore categorized into 3 age categories [“young” (18–60 y), “middle” (61–75 y), and “old” (>75 y)] and also into 2 age categories [“young” (18–60 y) and “other” (>60 y)]. Other potential sources of heterogeneity at the study level (effect modifiers) were also investigated by meta-regression analysis (Supplemental Methods).

Sensitivity analyses

The following sensitivity analyses were performed: 1) outlier analyses by omitting one study at a time, 2) omitting studies with large sample sizes ($n > 5000$), 3) omitting studies that used the relative TL to estimate the absolute TL, 4) stratification by method of measurement of TL [Southern blot compared with quantitative polymerase chain reaction (qPCR)], and 5) using a cutoff of 90% for defining ethnicity (Supplemental Methods).

Statistical analyses were performed with the use of Stata software version 12.0 (StataCorp). All of the statistical tests were 2-sided; P values < 0.05 were considered significant, except where otherwise specified.

RESULTS

Search

The search (PubMed, EMBASE, and Cochrane) yielded 5021 publications, from which 173 potentially relevant publications were identified. Some authors contributed to > 1 publication. As a result, 153 corresponding authors were identified and contacted. Seventy-five corresponding authors responded positively, 69 authors did not respond, 6 declined to participate, and 3 authors did not have the requested data. Because 1 publication could include multiple studies, the PIs (if not the same as corresponding authors) of the studies were contacted. Eight additional studies were identified by the corresponding authors and the PIs of these additional studies were contacted. We decided to exclude 9 studies that used 1) techniques other than Southern blots and




FIGURE 1 Study inclusion flow chart. L/C ratio, telomere/centromere ratio; T/S ratio, telomere to single-copy gene ratio.

qPCR or 2) did not report TL in bp or T/S ratio, because the regression coefficients (β estimates) may not be directly comparable.

In total, 87 unique studies were included in the meta-analyses. Twenty-nine studies measured absolute TL and 58 studies used the T/S ratio. A flow chart of the inclusion procedure is presented in **Figure 1**.

Description of studies

The characteristics of the 87 studies included in this meta-analysis are provided in **Table 1**. Absolute TLs were obtained from 29 studies (3, 5, 13, 14, 17, 28–61; HyperGEN study, unpublished data S. Hunt, A. Aviv, R. Cawthon 2011), of which 4 studies estimated absolute TL on the basis of the T/S ratio (19, 22, 62–68). In 17 studies, Southern blots were used (3, 5, 13, 14, 17, 28–40, 44, 45, 48–52, 55, 56, 59, 61). Fifty-eight studies presented the relative TL (T/S ratio) (4, 15, 69–133; Utah Pedigree Study, unpublished data S. Hunt, A. Aviv, R. Cawthon 2011). One PI provided the data stratified by source of leukocytes (whole blood compared with buffy coat) (104). One PI provided longitudinal data (55).

The total pooled sample of this meta-analysis consisted of 146,114 adults (40% men), the “young” pooled sample (18–60 y) consisted of 81,446 adults (51% men), the “middle” pooled sample (61–75 y) consisted of 42,991 adults (41%

TABLE 1
Characteristics of included study samples¹

Ref	Study name	All, n	Men, n	Women, 18–60 y	Age, n		Cell type	Telomere length measure	DNA	BMI	Data provided	design	Ethnicity (proportion white/black/Asian/Hispanic/Native American)	
					>75 y	>60–75 y								
	Absolute telomere length measured in base pairs													
(3, 28–31)	TwinsUK	3236	286	2950	2630	574	32	Leukocytes	Southern blot RF	Stored	Measured	Summary	Cohort	1/0/0/0/0
(17, 32–34)	Bogalusa	635	635	0	635	0	0	Leukocytes	Southern blot RF	Unknown	Measured	Raw data	Cohort	0.71/0.29/0/0/0
(147)	India CURES Study	40	20	20	37	3	0	Leukocytes	Southern blot RF	Stored	Measured	Raw data	Case-control	0/0/1/0/0
(36)	Campania	528	251	277	320	100	108	Leucocytes	Southern blot RF	Stored	Measured	Raw data	Cohort	1/0/0/0/0
(14, 37–39)	Asklepios	2509	1218	1291	2509	0	0	Leucocytes	Southern blot RF	Stored	Measured	Raw data	Cohort	1/0/0/0/0
(5, 13, 40)	Framingham	1146	557	589	658	444	44	Leukocytes	Southern blot RF	NA	Measured	Summary	Cohort	1/0/0/0/0
(42)	COPD	178	89	89	113	60	5	Leukocytes	Real-time PCR	NA	Measured	Raw data	Case-control	1/0/0/0/0
(43)	Crete	109	109	0	0	0	109	Leukocytes	Real-time PCR	NA	Measured	Raw data	Cohort	1/0/0/0/0
(41)	Zutphen	189	189	0	0	68	121	Leukocytes	Real-time PCR	NA	Measured	Raw data	Cohort	1/0/0/0/0
(17)	Family Heart, African American	625	216	409	459	148	18	Leukocytes	Southern blot RF	NA	Measured	Summary	Cohort	0/1/0/0/0
(17)	Family Heart, white	2603	1170	1433	1419	997	187	Leukocytes	Southern blot RF	NA	Measured	Summary	Cohort	1/0/0/0/0
Not published	HyperGEN, African American	224	108	116	172	51	1	Leukocytes	Southern blot RF	NA	Measured	Summary	Cohort	0/1/0/0/0
Not published	HyperGEN, white	1240	612	628	799	426	15	Leukocytes	Southern blot RF	NA	Measured	Summary	Cohort	1/0/0/0/0
(44, 45)	LSADT	525	171	354	0	82	443	Leukocytes	Southern blot RF	NA	Measured	Summary	Cohort	1/0/0/0/0
(46, 47)	Heart and Soul	954	777	177	274	451	229	Leukocytes	Real-time PCR	Unknown	Measured	Summary	Cohort	0.60/0.116/0.12/0.09/0.03
(57, 58)	Lothian	1057	530	527	0	1057	0	Leucocytes	Real-time PCR	Stored	Measured	Summary	Cohort	1/0/0/0/0
(48, 49)	WarTwins	639	639	0	0	86	553	Leukocytes	Southern blot RF	Stored	Reported	Raw data	Cohort	1/0/0/0/0
(55)	Jerusalem LRC	620	413	207	620	0	0	Leukocytes	Southern blot RF	Buffy coat stored	Measured	Summary	Cohort	1/0/0/0/0
(56)	Jerusalem Palestinians	939	498	441	336	306	0	Leukocytes	Southern blot RF	Buffy coat stored	Measured	Summary	Cross-sectional	1/0/0/0/0

(Continued)

TABLE 1 Continued

Ref	Study name	Age, n			Cell type	Telomere length measure	DNA	BMI	Data provided	design	Ethnicity (proportion white/black/Asian/Hispanic/Native American)		
		All, n	Men, n	Women, 18–60 y									
(50–52)	Helsinki Businessmen Study	487	487	0	250	237	Leukocytes	Southern blot RF	Stored	Reported	Raw data	Cohort	1/0/0/0/0
(53)	Copenhagen General Population Study	45,069	20,422	24,647	14,525	4504	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	1/0/0/0/0
(54)	SOLVABLE	152	0	152	136	16	PBMCs	Real-time PCR	Stored	Measured	Summary	Case-control	0.70/0.22/0.03/0.05/0
(59)	ZTL2008	25	17	8	24	1	PBMCs	Southern blot RF	Stored	Measured	Raw data	Cohort	1/0/0/0/0
(60)	Venado Tuerto 2	401	0	401	325	63	Leukocytes	Real-time PCR	Stored	Measured	Raw data	Cohort	0/0/0/1/0
(61)	NHSC	672	333	339	534	138	Leukocytes	Southern blot RF	Stored	Measured	Summary	Cohort	0/0/1/0/0
Absolute telomere length estimated from T/S ratio													
(19, 62, 66, 67)	South Carolina	323	145	178	305	18	Leukocytes	Real-time PCR	Unknown	Measured	Summary	Cohort	0.57/0.41/0.01/0.01/0
(63, 64, 68)	Bruneck	800	395	405	363	315	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	1/0/0/0/0
(65)	RPCI	174	0	174	111	47	Leukocytes	Real-time PCR	Stored	Reported	Raw data	Case-control	0.93/0.05/0/0/0.02
(22)	ESTHER	3559	1583	1976	1432	2127	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	1*/0/0/0/0
Telomere length based on T/S ratio													
(15)	MONICA	511	183	328	419	92	Leukocytes	Real-time PCR	NA	Measured	Raw data	Cohort	1/0/0/0/0
(15)	MDCC	476	330	146	199	277	Granulocytes	Real-time PCR	NA	Measured	Raw data	Cohort	1/0/0/0/0
Not published	Utah Pedigree Study	964	493	471	725	183	Leukocytes	Real-time PCR	NA	Measured	Summary	Cohort	1/0/0/0/0
(71, 72)	MESA, white	182	89	93	80	80	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	1/0/0/0/0
(71, 72)	MESA, African American	278	125	153	141	109	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	0/1/0/0/0
(71, 72)	MESA, Hispanic	518	252	266	245	231	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	0/0/0/1/0
(73)	EARSII	395	395	0	395	0	Leukocytes	Real-time PCR	Unknown	Measured	Raw data	Case-control	1/0/0/0/0
(74)	UCLA MacArthur	233	115	118	0	144	Leukocytes	Real-time PCR	NA	Reported	Summary	Cohort	1/0/0/0/0

(Continued)

TABLE 1 Continued

Ref	Study name	Age, n				Cell type	Telomere length measure	DNA	BMI	Data provided	design	Ethnicity (proportion white/black/Asian/Hispanic/Native American)		
		All, n	Men, n	Women, 18–60 y	>60–75 y									
(75)	Ashkenazi	359	191	168	50	179	130	Leukocytes	Real-time PCR	Stored	Measured	Raw data	Cohort	1/0/0/0/0
(76)	Warsaw	714	246	468	235	411	68	Leukocytes	Real-time PCR	Stored	Measured	Raw data	Case-control	1/0/0/0/0
(77)	Finland Health 2000 cohort	938	350	588	754	137	47	Leukocytes	Real-time PCR	Stored	Unknown	Summary	Cohort	1/0/0/0/0
(78)	Sister Study I (Vanguard sample)	644	0	644	475	169	0	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	0.83/0.07/0.02/0.02/0.05
(114)	Sister Study II (Genetic Study subcohort)	734	0	734	548	186	0	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	0.92/0.04/0/0.02/0.02
(79)	CAS	183	96	87	112	53	18	Leukocytes	Real-time PCR	Stored	Measured	Raw data	Case-control	1/0/0/0/0
(80)	PATH 40	331	151	180	331	0	0	Leukocytes	Real-time PCR	Stored	Reported	Raw data	Cohort	0.95/0/0.03/0/0.02
(80)	PATH 60	294	157	137	0	294	0	Leukocytes	Real-time PCR	Stored	Reported	Raw data	Cohort	0.97/0/0.02/0/0.01
(81)	Italy alcohol controls	258	258	0	255	3	0	Leukocytes	Real-time PCR	Stored	Reported	Raw data	Case-control	1/0/0/0/0
(82)	Fels Longitudinal Study	257	116	104	196	54	7	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	1/0/0/0/0
(83, 84)	ECRAN	188	38	150	121	41	26	PBMCs	Real-time PCR	NA	Measured	Raw data	Cohort	0/0/0/1/0
(85, 86)	Heart Scan Study	434	206	228	169	259	0	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	1/0/0/0/0
(87, 88)	Boiler workers	104	104	0	97	7	0	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	0.85/0.09/0.02/0.03/0
(89, 90)	Mayo	2886	1470	1416	2001	709	176	Leukocytes	Real-time PCR	Stored	Measured	Raw data	Case-control	0.98/0/0.01/0.01/0
(91, 92, 105, 108–111)	HBCS	1962	911	1051	703	1259	0	Leukocytes	Real-time PCR	NA	Measured	Raw data	Cohort	1/0/0/0/0
(93–95)	PREVEND	7991	3994	3997	6094	1897	0	Leukocytes	Real-time PCR	NA	Measured	Summary	Cohort	0.96/0.01/0.02/0/0.01
(96, 97, 106, 107)	Strong Heart Family Study	3256	1315	1941	2834	340	82	Leukocytes	Real-time PCR	NA	Measured	Summary	Cohort	0/0/0/0/1
(98)	PREDIMED-NAVARRA	521	236	285	81	401	38	Leukocytes	Real-time PCR	Stored	Measured	Summary	RCT	1/0/0/0/0
(99)	NHANES	7349	3542	3807	5034	1564	751	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	0.52/0.18/0/0.30/0
(100)	SWHS	2912	0	2912	1812	1100	0	Leukocytes	Real-time PCR	NA	Measured	Summary	Cohort	0/0/1/0/0
(101)	DHS, white	1073	493	580	821	245	7	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	1/0/0/0/0

(Continued)

TABLE 1 Continued

Ref	Study name	Age, n				Cell type	Telomere length measure	DNA	BMI	Data provided	design	Ethnicity (proportion white/black/Asian/Hispanic/Native American)		
		All, n	Men, n	Women, 18–60 y	>60–75 y									
(101)	DHS, black	1667	606	1061	1348	317	17	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	0/1/0/0/0
(101)	DHS, Hispanic	464	194	270	412	51	1	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	0/0/0/1/0
(102)	DALS	734	401	333	268	366	100	Leukocytes	Real-time PCR	Stored	Measured	Summary	Case-control	0.96/0/0/0.03/0
(105, 113)	FinnTwin study	2096	1101	995	1589	385	122	Leukocytes	Real-time PCR	NA	Measured	Summary	Cohort	1/0/0/0/0
(112)	Former Athletes Study	586	586	0	1	376	209	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	1/0/0/0/0
(104)	USKCS, whole blood	765	442	323	395	320	50	Leukocytes	Real-time PCR	Stored	Measured	Summary	Case-control	0.61/0.39/0/0/0
(104)	USKCS, buffy coat	126	70	56	87	36	3	Leukocytes	Real-time PCR	Stored	Measured	Summary	Case-control	0.66/0.34/0/0/0
(103)	Erasmus Rucphen Study	2449	1082	1367	1900	499	50	Leukocytes	Real-time PCR	Stored	Measured	Summary	Case-control	1/0/0/0/0
(103)	Rotterdam Study	2231	944	1287	556	1272	404	Leukocytes	Real-time PCR	Stored	Measured	Summary	Case-control	1/0/0/0/0
(103)	KORA F3	3113	1509	1604	1768	1051	294	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	1/0/0/0/0
(103)	KORA F4	3014	1457	1557	1824	943	247	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	1/0/0/0/0
(148)	CAVASIC	315	315	155	160	160	0	Leukocytes	Real-time PCR	Stored	Measured	Summary	Case-control	1/0/0/0/0
(103, 149)	SAPHIR	1681	1055	626	1586	95	0	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	1/0/0/0/0
(115)	CLNHS	3467	893	2574	3380	87	0	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	0/0/1/0/0
(116)	NEW Study	437	0	437	304	131	2	Leukocytes	Real-time PCR	Stored	Measured	Raw data	Cohort	0.85/0.08/0.02/0.03/0.02
(117)	NESDO	495	173	322	17	354	124	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	0.95/0.01/0.04/0/0
(118)	NESDA	2936	986	1950	2749	187	0	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	0.97/0.02/0.01/0/0
(119)	PRT	43	0	43	43	0	0	Leukocytes	Real-time PCR	Stored	Measured	Raw data	Case-control	**
(120, 121)	UMS	67	67	0	65	1	1	Leukocytes	Real-time PCR	Stored	Measured	Raw data	Cohort	1/0/0/0/0

(Continued)

TABLE 1 Continued

Ref	Study name	All, n	Men, n	Women, 18–60 y	Age, n		Cell type	Telomere length measure	DNA	BMI	Data provided		design	Ethnicity (proportion white/black/Asian/Hispanic/Native American)
					>75 y	>60–75 y					Raw data	Summary		
(122)	YMCA	1126	1126	0	1126	0	0	Leukocytes	Real-time PCR	Stored	Measured	Raw data	Cohort	1/0/0/0/0
(123–125)	BASE-II	1894	946	948	441	1409	44	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	1**/0/0/0/0
(126)	Kyiv	82	20	62	36	33	13	Leukocytes	Real-time PCR	Stored	Measured	Raw data	Cohort	1/0/0/0/0
(127)	GAHR2	133	53	80	128	5	0	Leukocytes	Real-time PCR	Stored	Measured	Raw data	Cohort	0.93/0.01/0.01/0.05/0
(128)	UMED telomere trial	28	7	21	26	2	0	PBMCs	Real-time PCR	Not stored	Measured	Raw data	Cohort	1/0/0/0/0
(129)	ACCT	904	454	450	469	361	74	Leukocytes	Real-time PCR	Stored	Measured	Summary	Cohort	0.98/0.01/0.01/0/0
(130)	RPE	975	466	509	975	0	0	Leukocytes	Real-time PCR	Stored	Reported	Raw data	Cohort	0/0/1/0/0
(131)	Sweden Mindfulness Study	172	21	151	167	5	0	Leukocytes	Real-time PCR	Stored	Reported	Raw data	RCT	1/0/0/0/0
(132)	EpiDREAM	4205	1565	2640	3363	770	72	Leukocytes	Real-time PCR	Stored	Reported	Summary	—	0.61/0.08/0.21/0.09/0
(133)	INTERHEART	3306	2601	705	2092	997	217	Leukocytes	Real-time PCR	Stored	Reported	Summary	Case-control	0.27/0.08/0.44/0.21/0

*Not measured, but all of Eurasian descent; **not measured. ACCT, Anglo-Cardiff Collaborative Trial; BASE-II, Berlin Aging Study; CAS, calcific aortic valve stenosis study; CAVASIC, Cardiovascular Disease in Intermittent Claudication; CLHNS, Cebu Longitudinal Health and Nutrition Survey; COPD, Chronic Obstructive Lung disease cohort; CURES, Chennai Urban Rural Epidemiology Study; DALIS, Diet, Activity and Lifestyle Study; DHS, Dallas Heart Study; EARSII, European Atherosclerosis Study II; ECRAN, Envejecimiento y Enfermedades Crónicas Asociadas a Nutrición (Aging and Nutrition Associated Chronic Disease); EpiDREAM, Epidemiologic study of the Screenes for DREAM (Diabetes Reduction Assessment with Ramipril and Rosiglitazone Medication); ESTHER, Epidemiological Study on the Chances of Prevention, Early Recognition, and Optimised Treatment of Chronic Diseases in the Older Population; GAHR2, Prospective evaluation of Gender and Age differences in the impact of Hostility and Reactivity on intermediary coronary artery disease risk factors; HBCS, Helsinki Birth Cohort Study; HyperGEN, Hypertension Genetic Epidemiology Network study; KORA F3, Cooperative Health Research in the Region of Augsburg F3 (2004/2005) survey; KORA F4, Cooperative Health Research in the Region of Augsburg F4 (2006/2008) survey; LRC (Jerusalem), Lipid Research Clinic; LSADT, Longitudinal Study of Aging Danish Twins (<https://www.icsr.umich.edu/icsrweb/NACDA/studies/21041>); MDCC, Malmö Diet and Cancer Cohort; MESA, Multi-Ethnic Study of Atherosclerosis (<https://www.mesa-nhlbi.org/>); MONICA, Multinational Monitoring of Trends and Determinants in Cardiovascular Disease; NA, not available (data provided by principal investigator); NESDA, The Netherlands Study of Depression and Anxiety; NESDO, The Netherlands Study of Depression in older persons; NEW, Nutrition and Exercise for Women; NHSC, Nutrition and Health in Southwest China study; PATH 40, Personality and Total Health Through Life Project AGE 40–44; PATH 60, Personality and Total Health Through Life Project AGE 60–64; PBMC, peripheral blood mononuclear cell; PCR, polymerase chain reaction; PREDIMED, Prevención con Dieta Mediterránea; PREVENT, Prevention of Renal and Vascular End-stage Disease; PRT, Progressive Resistance Training; RCT, randomized controlled trial; Ref, reference; RF, restriction fragment; RPCI, Roswell Park Cancer Institute; RPE, Richard Paul Ebstein; SAPHIR, Salzburg Atherosclerosis Prevention program in subjects at High Individual Risk Study; SOLVABLE, Study of Lupus Vascular and Bone Longterm Endpoints; summary, summary statistics; SWHS, Shanghai Women's Health Study; T/S ratio, telomere to single-copy gene ratio; UMED, Uniwersytet MEDyczny w Łodzi (Medical University of Lodz (MUL)); UMS, ultra-marathon study; USKCS, US Kidney Cancer Study; YMCA, Young Men Cardiovascular Association; ZTL2008, Zamolli Telomere Length 2008.

men), and the “old” pooled sample (>75 y) consisted of 8495 adults (65% men). Overall, the majority of the adults were white (including Arab; 83%), followed by Asian (7%), African American (4%), and Hispanic and Native American (both 3%). Six studies provided data of mixed study populations stratified by ethnicity (17, 71, 72, 101, 132, 133; HyperGEN study, unpublished data S. Hunt, A. Aviv, R. Cawthon 2011). Sixty-five studies consisted of >70% white individuals (of which 60 had ≥90% white individuals) (3, 5, 13–15, 17, 22, 28–34, 36–45, 48–59, 63–65, 68, 71–95, 98, 100–103, 105, 108–114, 116–118, 120–129, 131; HyperGEN study, unpublished data S. Hunt, A. Aviv, R. Cawthon 2011; Utah Pedigree Study, unpublished data S. Hunt, A. Aviv, R. Cawthon 2011). Four studies consisted only of African Americans (17, 71, 72, 101; HyperGEN study, unpublished data S. Hunt, A. Aviv, R. Cawthon 2011), 5 only of Asians (35, 61, 100, 115, 130), 1 study only of Native Americans (96, 97, 106, 107), and 4 studies comprised only Hispanics (60, 71, 72, 83, 84, 101). One study could not provide information about ethnicity (119).

Assessment of small study effects

Visual inspection of the funnel plots for absolute TL and for relative TL yielded symmetric plots (Supplemental Figure 1, funnel plots). No publication bias was detected with the use of Egger’s test or Begg’s test. The “trim and fill” method added 1 hypothetical study to the meta-analysis for absolute TL. However, the recalculated summary estimate did not change and was still significant with its inclusion ($\beta = -3.99$; 95% CI: $-5.16, -2.84$; $P < 0.001$).

Statistical pooling

A summary of the β estimates of the meta-analysis is shown in Tables 2 and 3. An overview of the meta-analysis is shown Supplemental Tables 2 and 3 in which the β estimates of all meta-analyses for absolute TL as the outcome (Supplemental Table 2) and of all meta-analyses for relative TL as the outcome (Supplemental Table 3) are presented. The accompanying forest plots are presented in the Supplemental Figure 2.

We confirmed that age was an effect modifier. Because the associations between BMI and TL did not differ significantly between men and women, the results of the sex-specific meta-analyses are shown only in the Supplemental Results.

Overall meta-analysis

Overall, sex- and age-adjusted absolute TL was significantly associated with BMI (Table 2, line 1). Each unit increase in BMI corresponded to a -3.99 bp (95% CI: $-5.17, -2.81$ bp; $I^2 = 0.6\%$) difference in absolute TL (Table 2, line 1; Figure 2). For example, the estimated difference in TL between a normal-weight individual with a BMI of 25 and an obese individual with a BMI >30 is (rounded) >20.0 bp and, if a larger difference is used (BMI: 20 compared with >30), is >39.9 bp. The estimated difference between normal weight and morbid obesity (BMI >40) is >59.9 bp. Overall, each unit increase in BMI corresponded to a -1.58×10^{-3} unit T/S ratio (0.16% decrease; 95% CI: $-2.14 \times 10^{-3}, -1.01 \times 10^{-3}$; $I^2 = 41.1\%$) difference in age- and sex-adjusted relative TL (Table 3, line 1). The estimated difference in relative TL between normal weight and obesity

TABLE 2

Summary of the β estimates (regression coefficients) from the meta-analysis of the association between BMI and telomere length as the outcome and absolute telomere length (in base pairs) as the independent variable¹

Line	Ethnicity	All (total pooled sample) ²			Pooled sample ³												
		n	Estimate	95% CI	I^2 , %	n	Estimate	95% CI	I^2 , %	n	Estimate	95% CI	I^2 , %				
1	Overall	29	-3.99*	-5.17, -2.81*	0.6	23	-7.67*	-10.03, -5.31*	31.2	22	-1.65	-4.41, 1.11	19.7	16	-5.89*	-10.41, -1.37*	5.3
2	White	21	-4.36*	-5.87, -2.85*	11.3	15	-8.77*	-10.42, -7.12*	0.0	15	-2.06*	-4.06, -0.06*	0.0	13	-6.97*	-12.29, -1.64*	15.4
3	African American	2	0.86	-4.75, 6.46	0.0	2	0.960	-5.51, 7.43	1.2	2	4.36	-7.25, 15.97	0.0	1	74.70	-76.02, 225.42	
4	Hispanic	1	5.97	-149.97, 161.91	0.0	1	-45.64	-216.24, 124.95	10.8	1	212.68	-169.98, 595.34	0	0			
5	Asian	2	-7.65	-27.20, 11.91	0.0	2	-48.70	-130.38, 32.99	10.8	1	90.00*	27.28, 152.72*	0	0			

¹ A random-effects model was used. Statistical heterogeneity was estimated by I^2 statistics for each of the meta-analyses. * $P < 0.05$ or $I^2 > 50\%$. “n” indicates number of studies.

² Adjusted for age and sex.

³ Adjusted for sex.

TABLE 3

Summary of the β estimates (regression coefficients) from the meta-analysis of the association between BMI and telomere length as the outcome and relative telomere length (T/S ratio) as the independent variable¹

Line	Ethnicity	All (total pooled sample) ²						Pooled sample ³								
		Estimate	95% CI	I^2 , %	n	Estimate	95% CI	I^2 , %	n	Estimate	95% CI	I^2 , %				
1	Overall	-1.58*	-2.14, -1.01*	32.7	55	-2.58*	-3.92, -1.25*	80.0*	50	-1.08*	-1.76, -0.39*	0.0	29	0.20	-1.40, 1.80	0.0
2	White	-1.87*	-2.44, -1.31*	8.1	40	-2.80*	-4.77, -0.82*	84.1*	37	-1.65*	-2.45, -0.86*	0.0	21	-0.28	-2.29, 1.73	0.0
3	African American	5.66	-6.60, 17.92	80.0*	2	5.21	-5.67, 16.08	68.7	2	0.08	-6.20, 6.36	0.0	1	-0.74	-12.62, 11.14	
4	Hispanic	2.53	-5.18, 10.25	17.7	3	-0.42	-4.19, 3.34	0.0	3	2.31	-2.35, 6.97	0.0	2	27.29	-40.32, 94.61	77.7*
5	Asian	-1.11	-4.23, 2.02	60.4	3	-4.50*	-5.75, -3.25*	0.0	2	2.18	-2.90, 7.27	0.0	0			
6	Native American	-2.64*	-3.60, -1.68*		1	-4.14*	-5.28, -3.00*		1	2.23	-1.00, 5.46		1	4.68	-2.35, 11.71	

¹The unit of the estimates and 95% CI of the T/S ratio is 10^{-3} . A random-effects model was used. Statistical heterogeneity was estimated by I^2 statistics for each of the meta-analyses. * $P < 0.05$ or $I^2 > 50\%$. "n" indicates number of studies. T/S ratio, telomere to single-copy gene ratio.

²Adjusted for age and sex.

³Adjusted for sex.

is a $\geq 7.9 \times 10^{-3}$ unit (0.79% difference) T/S ratio (Table 3, line 1; Figure 3) and between normal weight and morbid obesity is a $\geq 23.7 \times 10^{-3}$ unit T/S ratio (2.37% difference).

Meta-analysis stratified by age categories

Analysis stratified by age category showed that in “young” adults (18–60 y) a unit increase in BMI corresponded to a -7.67 bp (95% CI: $-10.03, -5.31$ bp; $I^2 = 31.2\%$) difference in absolute TL (Table 2, line 1; Figure 4). In “middle”-aged adults (61–75 y) the overall association between BMI and TL was -1.65 bp (95% CI: $-4.41, 1.11$ bp; $I^2 = 19.7$) per unit increase in BMI. In “old” adults (>75 y) the overall association between BMI and TL was -5.89 bp (95% CI: $-10.41, -1.37$ bp; $I^2 = 5.3$) per unit increase in BMI (Table 2, line 1).

For relative TL, each unit increase in BMI corresponded to a -2.58×10^{-3} unit T/S ratio (0.26% decrease; 95% CI: $-3.92 \times 10^{-3}, -1.25 \times 10^{-3}$; $I^2 = 80.0\%$) difference in relative TL in young adults (Table 3, line 1; Figure 5). In “middle”-aged adults, the overall association between BMI and relative TL was found to be a -1.08×10^{-3} unit T/S ratio (0.1% decrease; 95% CI: $-1.76 \times 10^{-3}, -0.39 \times 10^{-3}$; $I^2 = 0.0$) per unit increase in BMI. For “old” adults no significant association [0.20×10^{-3} unit T/S ratio (0.02%); 95% CI: $-1.40 \times 10^{-3}, 1.80 \times 10^{-3}$; $I^2 = 0.0$] was found between BMI and relative TL (Table 3, line 1).

Meta-regression and sources of heterogeneity

Age and ethnicity were effect modifiers in the meta-regression analyses (Supplemental Results). Therefore, all of the analyses were stratified by ethnicity in addition to the originally planned analyses.

With absolute TL as the outcome, stratified analyses showed that all estimates were significant for the white pooled sample (Table 2, line 2) and also for the “middle” pooled Asian sample. However, the latter estimate was an outlier and based on just 1 study (Table 2, line 5).

With relative TL as the outcome, stratified analyses showed that the estimates (except for one estimate for the “old” pooled sample) were significant for the white pooled sample (Table 3, line 2). In addition, the estimates of the overall and “young” pooled Native American samples and the estimate of the “young” pooled Asian sample were significant (Table 3, lines 5 and 6).

Sensitivity analysis

None of the sensitivity analyses resulted in a substantial change in the summary estimate (Supplemental Results).

DISCUSSION

This cross-sectional meta-analysis of 87 observational studies of adult pooled populations confirmed previous observations that BMI is negatively associated with TL. After stratification for age and ethnicity, the negative association between BMI and TL appeared to be stronger in the “young” pooled population (18–60 y) and in the white pooled population. Differences between men and women could not be confirmed.

On the basis of our estimates for absolute TL, an ~ 5 -unit higher BMI appears to be equivalent to a difference in TL of ~ 20 –38 bp or an $\sim 7.9 \times 10^{-3}$ to 13×10^{-3} unit T/S ratio (0.79–1.3%




FIGURE 2 Forest plot of the β estimates (regression coefficients) from the meta-analysis of the association between BMI and absolute telomere length (base pairs) as the outcome in the total pooled population. Random effect model was used and adjusted for age and sex. Statistical heterogeneity was estimated by I^2 statistics for each of the 12 meta-analyses. The shaded boxes indicate the inverse variance weighting of each estimate, and the size of the box indicates the weight. In case no shaded box is visible, the weight is very small. COPD, Chronic Obstructive Lung disease cohort; CURES, Chennai Urban Rural Epidemiology Study; ES, estimate; ESTHER, Epidemiological Study on the Chances of Prevention, Early Recognition, and Optimised Treatment of Chronic Diseases in the Older Population; HyperGEN, Hypertension Genetic Epidemiology Network study; LRC, Lipid Research Clinic; LSADT, Longitudinal Study of Aging Danish Twins (<https://www.icpsr.umich.edu/icpsrweb/NACDA/studies/21041>); NHSC, Nutrition and Health in Southwest China study; RPCI, Roswell Park Cancer Institute; SOLVABLE, Study of Lupus Vascular and Bone Longterm Endpoints; ZTL2008, Zannolli Telomere Length 2008.

difference). Compared with an estimated average yearly decrease (i.e., ~ 25 bp/y or ~ 0.01 T/S ratio/y) in leucocyte TL in adults based on cross-sectional data (3, 32, 134–136), this is equivalent to an increase in biological age of ~ 1 y. A major disadvantage of cross-sectional analysis is the impossibility to infer causation. However, the robust association between higher BMI and lower TL found in this meta-analysis could highlight another potential area of concern for the obesity epidemic.

Because obesity and, more specifically, an increase in leptin and a decrease in adiponectin have been associated with low-grade inflammation and oxidative stress (137), the observed negative association between BMI and leukocyte TL may be due, in part, to the chronic inflammatory state associated with higher leptin. Recently, a negative association was observed between age-related relative TL and serum leptin in 7 cohorts of 11,448 participants, which remained significant after adjustment for BMI (103). These data suggest that, beyond a high BMI, inflammatory conditions, mediated via increase in leptin, likely contribute to telomere shortening. Because a longitudinal intervention study

showed that a reduction in BMI was linked to increasing TL over a 5-y period (98), it is also suggested that a common factor, such as chronic inflammation, is associated both with leptin resistance and with TL.

The negative association between BMI and TL was most apparent in the younger pooled population, in whom a stronger association was found for TL than in the other age groups. Three possible explanations could explain this observation. First, BMI could be a better marker for adiposity in younger individuals aged < 60 y than in older individuals (22). Above 65 y of age BMI may less consistently reflect obesity because of potential loss of muscle and bone mass and height (22). The fact that older men weighed less than the middle-aged men at a given height is attributed to older men having less lean tissue, and a lower BMI can actually reflect a higher fat mass (138). Second, selective survival might be one of the causes of the stronger association found in the younger age category. As Manson et al. (139) stated, “obesity in one’s 40s contributes to the onset of type 2 diabetes in one’s 50s, which leads to myocardial infarction (MI) in one’s




FIGURE 3 Forest plot of the β estimates (regression coefficients) from the meta-analysis of the association between BMI and relative telomere length (T/S ratio) as outcome in the total pooled population. The unit of the estimates and 95% CI is 10^{-3} . A random-effects model was used and adjusted for age and sex. Statistical heterogeneity was estimated by I^2 statistics for each of the 12 meta-analyses. The shaded boxes indicate the inverse variance weighting of each estimate, and the size of the box indicates the weight. In case no shaded box is visible, the weight is very small. ACCT, Anglo-Cardiff Collaborative Trial; BASE-II, Berlin Aging Study; CAS, calcific aortic valve stenosis study; CAVASIC, Cardiovascular Disease in Intermittent Claudication; DALS, Diet, Activity and Lifestyle Study; DHS, Dallas Heart Study; EARSII, European Atherosclerosis Study II; ECRAN, Envejecimiento y Enfermedades Crónicas Asociadas a Nutrición (Aging and Nutrition Associated Chronic Diseases); EPIDREAM, Epidemiologic study of the Screenes for DREAM (Diabetes Reduction Assessment with Ramipril and Rosiglitazone Medication); ES, estimate; GAHR2, Prospective evaluation of Gender and Age differences in the impact of Hostility and Reactivity on intermediary coronary artery disease risk factors; HBCS, Helsinki Birth Cohort Study; KORA F3, Cooperative Health Research in the Region of Augsburg F3 (2004/2005) survey; KORA F4, Cooperative Health Research in the Region of Augsburg F4 (2006/2008) survey; MDCC, Malmö Diet and Cancer Cohort; MESA, Multi-Ethnic Study of Atherosclerosis (<https://www.mesa-nhlbi.org/>); MONICA, Multinational Monitoring of Trends and Determinants in Cardiovascular Disease; NESDA, The Netherlands Study of Depression and Anxiety; NESDO, The Netherlands Study of Depression in older persons; NHANES, National Health and Nutrition Examination Survey; Not publ, not published; PATH 40, Personality and Total Health Through Life Project AGE 40-44; PATH 60, Personality and Total Health Through Life Project AGE 60-64; PREDIMED, Prevención con Dieta Mediterránea; PREVEND, Prevention of Renal and Vascular End-stage Disease; PRT, Progressive Resistance Training; Ref, reference; RPE, Richard Paul Ebstein; SAPHIR, Salzburg Atherosclerosis Prevention program in subjects at High Individual Risk Study; SWHS, Shanghai Women's Health Study; T/S ratio, telomere to single-copy gene ratio; UMED, Uniwersytet MEDyczny w Łodzi (Medical University of Lodz (MUL)); UMS, ultra-marathon study; USKCS, US Kidney Cancer Study; YMCA, Young Men Cardiovascular Association.

60s, heart failure and weight loss due to debilitation and muscle wasting at age 70, and death at age 75". People who suffered from age-related diseases may have died and those who survived may therefore differ from those who died (140). Third, older people are more likely to have chronic diseases that lead to weight

loss and people with chronic diseases are probably less likely to participate in studies (139).

The magnitude of the negative association between BMI and leukocyte TL was found to be largest in the white pooled population. One possible explanation could be that TL




FIGURE 4 Forest plot of the β estimates (regression coefficients) from the meta-analysis of the association between BMI and absolute telomere length (base pairs) as outcome in the “young” pooled population (age ≥ 18 and ≤ 60 y). A random-effects model was used and adjusted for sex. Statistical heterogeneity was estimated by I^2 statistics for each of the 12 meta-analyses. The shaded boxes indicate the inverse variance weighting of each estimate, and the size of the box indicates the weight. In case no shaded box is visible, the weight is very small. COPD, Chronic Obstructive Lung disease cohort; CURES, Chennai Urban Rural Epidemiology Study; ES, estimate; ESTHER, Epidemiological Study on the Chances of Prevention, Early Recognition, and Optimised Treatment of Chronic Diseases in the Older Population; HyperGEN, Hypertension Genetic Epidemiology Network study; LRC, Lipid Research Clinic; LSADT, Longitudinal Study of Aging Danish Twins (<https://www.icpsr.umich.edu/icpsrweb/NACDA/studies/21041>); NHSC, Nutrition and Health in Southwest China study; Not publ, not published; Ref, reference; RPCI, Roswell Park Cancer Institute; SOLVABLE, Study of Lupus Vascular and Bone Longterm Endpoints; ZTL2008, Zannoli Telomere Length 2008.

differs between different cell types (141) and that leukocyte cell subpopulations (142) differ between whites and African Americans. However, because only 4 samples consisted of African Americans, more research is required to resolve whether this observation explains the racial differences in association between TL and BMI for white and African Americans or whether this is a false-positive finding. Second, it was reported that the estimation of visceral adipose tissue, the most relevant tissue that determines the risk of developing chronic metabolic diseases, was different in white and African American adults (143). At a higher BMI or increased waist circumference, white adults had higher amounts of visceral adipose tissue than African-American adults (137). Because the presence of leptin resistance or markers of inflammation was not included in these studies, it remains to be determined whether the relation between BMI, leptin resistance, inflammation, and telomere attrition is different for African Americans compared with whites. In addition, the one study sample consisting of 3256 mostly “young” Native Americans showed similar results as found for the white study population (107). The majority of this study sample was centrally obese, and leukocyte TL was negatively correlated with C-reactive protein.

One of the main strengths of this study is that we did not rely on publications only. Instead, we contacted PIs, which, in turn, pointed us toward important studies we may have missed to obtain the data used in the meta-analysis. In addition, we incorporated potential confounders (age and sex) and sources of heterogeneity (ethnicity and study design). The response rate of the originally contacted PIs was 55%, with a final count of 87 unique studies and $>140,000$ individuals. Although it is impossible to make a direct comparison with the unpublished β estimates of the nonresponders, we assume, also based on the absence of significant publication bias, that the studies in this meta-analysis are a random selection of all studies conducted and that we present a valid representation of the association between BMI and TL. Because of the large variation in adult TL, as well as biological and measurement variation (qPCR), large sample sizes are needed, especially in cross-sectional studies, to detect modest effects (30). In this meta-analysis we were able to detect a significant association of -3.99 bp or a -1.58×10^{-3} unit T/S ratio (0.16%) per unit increase in BMI. Because 36% of our meta-analyses showed a significant association with estimates in the same direction and of the same magnitude (except for 3 estimates of extreme magnitude of one Asian sample in the “middle”-aged




FIGURE 5 Forest plot of the β estimates (regression coefficients) from the meta-analysis of the association between BMI and relative telomere length (T/S ratio) as the outcome in the “young” pooled population (age ≥ 18 y and ≤ 60 y). The unit of the estimates and 95% CI is 10^{-3} . A random-effects model was used and adjusted for sex. Statistical heterogeneity was estimated by I^2 statistics for each of the 12 meta-analyses. The shaded boxes indicate the inverse variance weighting of each estimate, and the size of the box indicates the weight. In case no shaded box is visible, the weight is very small. ACCT, Anglo-Cardiff Collaborative Trial; BASE-II, Berlin Aging Study; CAS, calcific aortic valve stenosis study; CAVASIC, Cardiovascular Disease in Intermittent Claudication; DALS, Diet, Activity and Lifestyle Study; DHS, Dallas Heart Study; EARSII, European Atherosclerosis Study II; ECRAN, Envejecimiento y Enfermedades Crónicas Asociadas a Nutrición (Aging and Nutrition Associated Chronic Diseases); EpiDREAM, Epidemiologic study of the Screenes for DREAM (Diabetes Reduction Assessment with Ramipril and Rosiglitazone Medication); ES, estimate; GAHR2, Prospective evaluation of Gender and Age differences in the impact of Hostility and Reactivity on intermediary coronary artery disease risk factors; HBCS, Helsinki Birth Cohort Study; KORA F3, Cooperative Health Research in the Region of Augsburg F3 (2004/2005) survey; KORA F4, Cooperative Health Research in the Region of Augsburg F4 (2006/2008) survey; MDCC, Malmö Diet and Cancer Cohort; MESA, Multi-Ethnic Study of Atherosclerosis (<https://www.mesa-nhlbi.org/>); MONICA, Multinational Monitoring of Trends and Determinants in Cardiovascular Disease; NESDA, The Netherlands Study of Depression and Anxiety; NESDO, The Netherlands Study of Depression in older persons; Not publ, not published; PATH 40, Personality and Total Health Through Life Project AGE 40–44; PREDIMED, Prevención con Dieta Mediterránea; PREVEND, Prevention of Renal and Vascular End-stage Disease; PRT, Progressive Resistance Training; Ref, reference; RPE, Richard Paul Ebstein; SAPHIR, Salzburg Atherosclerosis Prevention program in subjects at High Individual Risk Study; SWHS, Shanghai Women’s Health Study; T/S ratio, telomere-to-single-copy gene ratio; UMED, Uniwersytet MEDyczny w Łodzi (Medical University of Lodz (MUL)); UMS, ultra-marathon study; USKCS, US Kidney Cancer Study; YMCA, Young Men Cardiovascular Association.

population), we assume that false positive reporting is only of minor concern.

Two recent meta-analyses, which relied on published data, also reported negative associations between BMI and TL. The first smaller scale meta-analysis reported negative regression coefficients on the association between TL and BMI (22), of which 5 studies were also included in this meta-analysis (13, 14, 19, 82, 144). The larger-scale meta-analysis reported a weak

negative correlation, a standardized mean difference of 0.84 (95% CI: 0.22, 1.46) between obese individuals and normal-weight individuals, and an OR of 1.39 (95% CI: 1.15, 1.69) (7). Of the 45 samples that met our inclusion criteria 33 collaborated in our analysis. This shows that the results between the meta-analyses are consistent and very robust.

Although age and ethnicity were taken into account, it should be mentioned that the older study sample was relatively small

(~8400 individuals) and that the majority of the individuals were white (83%). Unfortunately, we did not include smoking in the meta-analysis. Smoking is generally associated with a lower BMI and shorter TL (3, 139, 145), which may have caused an underestimation of the inverse association between BMI and TL. In addition, inflammation was not directly measured. We were also not able to measure telomere attrition because we did not incorporate longitudinal data, and reverse causation cannot be excluded. However, there are very few large-scale studies with repeated measures of TL.

The lengths of telomeres at different ages are highly correlated, and it has been suggested that most of the variation in leukocyte TL in adults is a result of TL at birth and that therefore the impact of environmental and lifestyle factors is rather small (145, 146). In addition, Benetos et al. (145) described that ranking of individuals into deciles according to their TL barely changes across adult life. Our meta-analysis shows that a 5-unit increase in BMI corresponds to a change of ~20 bp or even ~38 bp in the younger pooled population, which is equivalent to 1-y greater biological age, irrespective of ranking.

In summary, a higher BMI is associated with shorter telomeres, especially in the younger pooled population. Being aware of the fact that the association between BMI and TL differs across the life span can lead to further research. Although no causal inference can be drawn and residual confounding may exist, the results were robust across a variety of potential confounders. Given this, we could possibly infer that tackling the obesity epidemic might be a starting point to delay telomere shortening and the onset of age-related diseases, thereby contributing to slower biological aging of the population. However, meta-analyses of longitudinal studies that can evaluate change in body weight alongside change in TL are warranted.

The collaborators in the TELOMAAS group are as follows: Göran Roos (Department of Medical Biosciences, Umeå University, Umeå, Sweden); Birgitta Stegmayr (Department of Public Health and Clinical Medicine, Umeå University, Sweden); Peter Nilsson (Clinical Sciences Medicine, University Hospital, Malmö, Sweden); Mats Eliasson (Department of Medicine, Sunderbyn Hospital, Luleå, Sweden); Olle Melander (Department of Clinical Sciences Medicine, University Hospital, Malmö, Sweden); Tim Spector (Twin Research and Genetic Epidemiology, King's College London, United Kingdom); Abraham Aviv (Center of Human Development and Aging, New Jersey Medical School, University of Medicine and Dentistry of New Jersey, Rutgers The State University of New Jersey, Newark, NJ); Antonysunil Adaikalakoteswari (Division of Metabolic and Vascular Health, University of Warwick, Coventry, United Kingdom); Viswanathan Mohan (Department of Diabetology, Madras Diabetes Research Foundation and Dr Mohans' Diabetes Specialities Center, Gopalapuram, Chennai, India); Sofie Bekaert (Bimera Clinical Research Center Ghent, Ghent University Hospital, Ghent, Belgium); Denil Simon (Department of Mathematical Modeling, Statistics, and Bioinformatics, Ghent University, Ghent, Belgium); Ernst Rietzschel, Marc De Buyzere, Thierry Gillebert, and Caroline Van Daele (Department of Cardiovascular Diseases, Ghent University Hospital, Ghent, Belgium); Michel Langlois (Department of Clinical Chemistry, AZ St-Jan Brugge-Oostende AV, Bruges, Belgium); Patrick Segers (Institute Biomedical Technology, Ghent University, Ghent, Belgium); Gui De Backer and Dirk De Bacquer (Department of Public Health, Ghent University, Ghent, Belgium); Michael Fenech (Department of Animal Food and Health Sciences, CSIRO, Adelaide BC, Australia); David Hughes and Janet Cross (Institute of Food Research, Norwich, United Kingdom); Helen Murphy (Department of Diabetes and Endocrinology, Ipswich Hospital, Ipswich, United Kingdom); Rosemary C Temple (Elsie Bertram Diabetes Center, Norfolk and Norwich University Hospital NHS Trust, Norwich, United

Kingdom); Duncan Fowler (Department of Diabetes and Endocrinology, Ipswich Hospital, Ipswich, United Kingdom); Nicoletta Dozio (Elsie Bertram Diabetes Center, Norfolk and Norwich University Hospital NHS Trust, Norwich, United Kingdom); Jackie Hughes and Mark Winterbone (Institute of Food Research, Norwich, United Kingdom); Shi-Jen Hwang and Daniel Levy [Population Sciences Branch of the National Heart, Lung, and Blood Institute (NHLBI), NIH, NHLBI's Framingham Heart Study, Framingham, MA]; Daan Kromhout (Division of Human Nutrition, Wageningen University, Wageningen, Netherlands); Anthony Kafatos (Department of Preventive Medicine and Nutrition Clinic, Medical School, University of Crete, Heraklion, Crete, Greece); Richard M Cawthon (Eccles Institute of Human Genetics, University of Utah, Salt Lake City, UT); Ana V Diez-Roux (Drexel University School of Public Health, Philadelphia, PA); Steve E Humphries (Department of Cardiovascular Genetics, Institute Cardiovascular Science, University College London, London, United Kingdom); Arch G Mainous III (Department of Health Services Research, Management, and Policy, University of Florida, Gainesville, FL); Marty S Player (Department of Family Medicine, Medical University of South Carolina, Charleston, SC); Charles J Everett (Ralph H Johnson VA Medical Center, Charleston, SC); Nir Barzilai (Department of Medicine and Genetics, Albert Einstein College of Medicine, Bronx, NY); Ian J Deary (Center for Cognitive Ageing and Cognitive Epidemiology and Department of Psychology, University of Edinburgh, Edinburgh, United Kingdom); John M Starr (Center for Cognitive Aging and Cognitive Epidemiology and Alzheimer Scotland Dementia Research Center, University of Edinburgh, Edinburgh, United Kingdom); Thomas von Zglinicki (Institute for Cell and Molecular Biosciences and Newcastle University Institute for Aging, Campus for Aging and Vitality, Newcastle upon Tyne, United Kingdom); Carmen Martin-Ruiz (NIHR Biomedical Research Center at Newcastle upon Tyne, Institute of Neuroscience and Newcastle University Institute for Aging, Campus for Aging and Vitality, Newcastle upon Tyne, United Kingdom); Nazmul Huda (Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, IN); Laura Kananen (Research Programs Unit, Biomedicum-Helsinki, University of Helsinki, Helsinki, Finland); Ronit Sinreich (Epidemiology Unit, Hebrew University-Hadassah School of Public Health and Community Medicine, Jerusalem, Israel); Barbara Kloeckener-Gruissem (Institute of Medical Molecular Genetics, University of Zurich, Schlieren, Switzerland, and Department of Biology, ETHZ, Zurich); Simon Eastaerle and Peter J Milburn (John Curtin School of Medical Research, The Australian National University, Canberra, Australia); Stefan Kiechl and Johann Willeit (Department of Neurology, Medical University Innsbruck, Innsbruck, Austria); Julia Raschenberger (Division of Genetic Epidemiology, Department of Medical Genetics, Molecular and Clinical Pharmacology, Medical University of Innsbruck, Innsbruck, Austria); Susan McCann and Christine Ambrosone (Department of Cancer Prevention and Control, Roswell Park Cancer Institute, Buffalo, NY); Andrea Baccarelli (Department of Environmental Health Sciences, Columbia Mailman School of Public Health, New York, NY); Mirjam Hoxha (Department of Clinical Sciences and Community Health, EPIGET- Epidemiology, Epigenetics and Toxicology Lab, University of Milan, Milan, Italy); Pier Alberto Bertazzi (Department of Clinical Sciences and Community Health, University of Milan and IRCCS Ca Granda Maggiore Hospital Foundation, Milan, Italy); Santo Davide Ferrara and Massimo Montisci (Department of Cardiac, Thoracic, and Vascular Sciences, Unit of Legal Medicine and Forensic Toxicology, University of Padova, Padova, Italy); Miryoung Lee (Community Health, Lifespan Health Research Center, Wright State University Boonshoft School of Medicine, Dayton, OH); Stig E Bojesen and Børge G Nordestgaard (The Copenhagen General Population Study, Department of Clinical Biochemistry, Copenhagen University Hospital, Herlev and Gentofte Hospital, Copenhagen, Denmark); Gladys Barrera, Sandra Hirsch, and María Pía De la Maza (Institute of Nutrition and Food Technology, University of Chile, Chile); Halcyon Skinner (Truven Health, Durham, NC); Kristin Litzelman (National Cancer Institute, Division of Cancer Control and Population Sciences, Rockville, MD); Songwon Seo (Korea Institute of Radiological and Medical Sciences, National Radiation Emergency Medical Center, Seoul, South Korea); Sonia Garcia-Calzon (Department of Nutrition, Food Science, and Physiology, University of Navarra, Pamplona, Spain, and Instituto de Investigación

Sanitaria de Navarra, Pamplona, Spain); Guillermo Zalba (Department of Biochemistry and Genetics, University of Navarra, Pamplona, Spain, and Instituto de Investigación Sanitaria de Navarra, Pamplona, Spain); J Alfredo Martínez and Miguel Angel Martínez-Gonzalez [Department of Nutrition, Food Science, and Physiology, University of Navarra, Pamplona, Spain, and Instituto de Investigación Sanitaria de Navarra, Pamplona, Spain, and CIBER Fisiopatología de la Obesidad y Nutrición, (CIBERObn), Instituto de Salud Carlos III, Madrid, Spain]; David Rehkopf (Department of General Medical Disciplines, Stanford University, Stanford, CA); Jue Lin (Department of Biochemistry and Biophysics, University of California, San Francisco, CA); Yu-Tang Gao (Department of Epidemiology, Shanghai Cancer Institute, Shanghai, China); Xiao-Ou Shu (Department of Medicine, Division of Epidemiology, Vanderbilt University Medical Center, Nashville, TN); Carly Skamra, Alexander Sandhu, and QiQuan Huang (Division of Rheumatology, Northwestern University Feinberg School of Medicine, Chicago, IL); Jungwha Lee (Department of Preventive Medicine, Northwestern University Feinberg School of Medicine, Chicago, IL); Richard Pope (Division of Rheumatology, Northwestern University Feinberg School of Medicine, Chicago, IL); Martha Slattery (Department of Medicine, University of Utah, Salt Lake City, UT); Merja K Laine (Department of General Practice and Primary Health Care, University of Helsinki, Helsinki, Finland, and Vantaa Health Center, Vantaa, Finland); Johan G Eriksson (Department of General Practice and Primary Health Care, University of Helsinki, Helsinki, Finland); Urho M Kujala (Department of Health Sciences, University of Jyväskylä, Jyväskylä, Finland); Rahul Raj (Department of Neurosurgery, Helsinki University Central Hospital, Helsinki, Finland); Heli M Bäckmand (City of Vantaa Health and Social Welfare Department, Vantaa, Finland); Markku Peltonen (Department of Health, National Institute for Health and Welfare, Helsinki, Finland); Seppo Sarna (Department of Public Health, University of Helsinki, Helsinki, Finland); Kendra Schwartz (Department of Family Medicine and Public Health Sciences, Wayne State University, Detroit, MI); Faith Davis (School of Public Health, University of Alberta, Alberta, Canada); Julie Ruterbusch (Department of Family Medicine and Public Health Sciences, Wayne State University, Detroit, MI); Mirjam Hoxha (Department of Occupational and Environmental Health, University of Milan, Milan, Italy); Bridget McCarthy (School of Public Health, University of Illinois at Chicago, Chicago, IL); Nathaniel Rothman and Joanne Colt (Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, MD); Wong-Ho Chow (Department of Epidemiology, MD Anderson Cancer Center, Houston, TX); Mark Purdue (Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, MD); Clemens Baumbach (Research Unit of Molecular Epidemiology and Institute of Epidemiology II, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany); Annette Peters (Institute of Epidemiology II, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany); Martina Müller-Nurasyid (Institute of Genetic Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany, and Department of Medicine I, Ludwig-Maximilians-University Munich, Munich, Germany, and DZHK (German Centre for Cardiovascular Research), partner site Munich Heart Alliance, Munich, Germany); Michelangela Barbieri (Department of Medical, Surgical, Neurological, Metabolic, and Geriatric Sciences, Second University of Naples, Naples, Italy); Kaare C Christensen (Department of Epidemiology, Biostatistics, and Biodemography, Institute of Public Health, University of Southern Denmark, Odense C, Denmark); Anu Loukola (Department of Public Health, University of Helsinki, Helsinki, Finland); Tellervo Korhonen (Department of Public Health, University of Helsinki, Finland, and University of Eastern Finland, Kuopio, Finland); Pamela AF Madden (Washington University School of Medicine, St. Louis, MO); Sharon Merkin (Department of Medicine, David Geffen School of Medicine at UCLA, Los Angeles, CA); Katri Räikkönen (Institute of Behavioural Sciences, University of Helsinki, Helsinki, Finland); Eero Kajantie (Department of General Practice and Primary Health Care, University of Helsinki, Helsinki, Finland); Johan Eriksson (Department of General Practice and Primary Health Care and Helsinki University Hospital, University of Helsinki, Helsinki, Finland); Pascal Arp (Department of Internal Medicine, Erasmus MC, Rotterdam, Netherlands); Albert Hofman and Cornelia M van Duijn (Department of Epidemiology, Erasmus MC, Rotterdam, Netherlands); André G Uitterlinden (Department of Internal Medicine, Erasmus MC, Rotterdam, Netherlands); Lyudmyla Kedenko and Bernhard Paulweber (Department of Internal Medicine I, Paracelsus Medical University, Salzburg, Austria); Margot Haun and (Division of Genetic Epidemiology, Department of Medical Genetics, Molecular and Clinical Pharmacology, Medical University of Innsbruck, Innsbruck, Austria); Barbara Rantner (Division of Genetic Epidemiology, Departments of Medical Genetics, Molecular and Clinical Pharmacology, and Vascular Surgery, Medical University of Innsbruck, Innsbruck, Austria); Angelika Hammerer-Lercher (Central Institute of Medical and Chemical Laboratory Diagnostics, University Hospital Innsbruck, Innsbruck, Austria); Gustav Fraedrich (Department of Vascular Surgery, Innsbruck Medical University, Innsbruck, Austria); Marietta Stadler (Third Medical Department of Metabolic Diseases and Nephrology, Hietzing Hospital, Vienna, Austria); Peter Klein-Weigel (Clinic for Angiology, HELIOS Klinikum Berlin-Buch, Berlin, Germany); M Geoffrey Hayes (Division of Endocrinology, Metabolism, and Molecular Medicine, Department of Medicine, and Center for Genetic Medicine, Feinberg School of Medicine, Northwestern University, Chicago, IL, and Department of Anthropology, Northwestern University, Evanston, IL); Christopher W Kuzawa (Department of Anthropology, Northwestern University, Evanston, IL, and Cells to Society (C2S), The Center on Social Disparities and Health, Institute for Policy Research, Northwestern University, Evanston, IL); Nanette Lee (Office of Population Studies Foundation, University of San Carlos, Cebu, Philippines); Caitlin Mason (Fred Hutchinson Cancer Research Center, Seattle, WA); Rosa-Ana Risques and Peter Rabinovitch (Department of Pathology, University of Washington, Seattle, WA); Ching-Yun Wang (Fred Hutchinson Cancer Research Center, Seattle, WA); Anne McTiernan (Fred Hutchinson Cancer Research Center, Seattle, WA, and Schools of Public Health and Medicine, University of Washington, Seattle, WA); Cristiana LMF Miranda-Furtado (Department of Gynecology and Obstetrics, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, São Paulo, Brazil); Hermann Brenner [Division of Clinical Epidemiology and Aging Research, German Cancer Research Center (DKFZ), Heidelberg, Germany, and Network Aging Research (NAR), University of Heidelberg, Heidelberg, Germany]; Antje Meyer [Center of Stroke Research Berlin (CSB) and Berlin-Brandenburg Center for Regenerative Therapies (BCRT), at Campus Virchow Klinikum, Charité Universitätsmedizin, Berlin, Germany]; Elisabeth Steinhagen-Thiessen (Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Lipid Clinic at the Interdisciplinary Metabolism Center, Germany); Julieta Panero and LaPatricia Carolina Dos Santos [Academia Nacional de Medicina (CONICET), Instituto de Medicina Experimental (IMEX), Laboratorio de Genética de Neoplasias Linfoides, Ciudad Autónoma de Buenos Aires, Argentina]; Claudio Daniel González (Universidad de Buenos Aires, Facultad de Medicina, Departamento de Farmacología, Ciudad Autónoma de Buenos Aires, Argentina); Jorge Vilariño (Hospital FLENI, Departamento de Cardiología, Ciudad Autónoma de Buenos Aires, Argentina); Martha Serey (Hospital Fiorito, Servicio de Endocrinología, Buenos Aires, Argentina); Gloria Edith Cerrone (Universidad de Buenos Aires, Facultad de Farmacia y Bioquímica, Cátedra de Genética y Biología Molecular, Ciudad Autónoma de Buenos Aires, Argentina); Irma Slavutsky [Academia Nacional de Medicina (CONICET), Instituto de Medicina Experimental (IMEX), Laboratorio de Genética de Neoplasias Linfoides, Ciudad Autónoma de Buenos Aires, Argentina]; Gustavo Daniel Frechtel [CONICET-Universidad de Buenos Aires. Instituto de Inmunología, Genética y Metabolismo (INIGEM), Laboratorio de Diabetes y Metabolismo, Ciudad Autónoma de Buenos Aires, Argentina]; Dariusz Nowak (Department of Clinical Physiology, Medical University of Lodz, Lodz, Poland); Makandjou-Ola Eusebio and Mirosława Pietruczuk (Department of Laboratory Diagnostics, II Department of Internal Medicine, Medical University of Lodz, Lodz, Poland); Yasmin (Division of Experimental Medicine and Immunotherapeutics, University of Cambridge, Cambridge, United Kingdom); Lee Butcher (Cardiff School of Sport and Health Sciences, Cardiff Metropolitan University, Cardiff, United Kingdom); John R Cockcroft (Division of Cardiology, New York-Presbyterian Hospital, Columbia University, New York, NY); Ian B

Wilkinson (Division of Experimental Medicine and Immunotherapeutics, University of Cambridge, Cambridge, United Kingdom); Jorge D Erusalimsky (Cardiff School of Sport and Health Sciences, Cardiff Metropolitan University, Cardiff, United Kingdom); Carmel M McEniery (Division of Experimental Medicine and Immunotherapeutics, University of Cambridge, Cambridge, United Kingdom); Onn Siong Yim (Department of Psychology, National University of Singapore, Singapore); Xing Zhang (Singapore-ETH Center for Global Environmental Sustainability, Singapore); Idan Shalev (Department of Biobehavioral Health, Pennsylvania State University, University Park, PA); Mikhail Monakhov (UConn Health, Neuroscience, Farmington, CT); Songfa Zhong (Department of Economics, National University of Singapore, Singapore); Ming Hsu (University of California/Haas School of Business, Berkeley, CA); Soo Hong Chew (Department of Economics, National University of Singapore, Singapore); Poh San Lai (Department of Pediatrics, National University of Singapore, Singapore); Jan Sundquist and Xiao Wang (Center for Primary Health Care Research, Lund University, Region Skåne, Lund, Sweden); Salim Yusuf, Hertzog Gerstein, and Sonia Anand (Population Health Research Institute and McMaster University, Hamilton, Canada).

We thank the principal investigators of the Bogalusa Study, the Sister Study, the Strong Heart Family Study, and the Helsinki Birth Cohort Study (HBCS) for their willingness to share the data but with whom we lost contact despite our efforts.

The TwinsUK study was funded by the Wellcome Trust, European Community's Seventh Framework Programme (FP7/2007–2013). The study also receives support from the National Institute for Health Research (NIHR) BioResource Clinical Research Facility and Biomedical Research Centre based at Guy's and St. Thomas' NHS Foundation Trust and King's College London. The Dallas Heart Study (DHS) research reported in this publication was supported by the National Center for Advancing Translational Sciences of the NIH under award number UL1TR001105. We thank Drs. C Garcia and Kozlitina for sharing the data. The acquisition and analyses of the data from the Multi-Ethnic Study of Atherosclerosis (MESA) was funded by contracts N01-HC-95159 through N01-HC-95165 and N01-HC-95169 from the National Heart, Lung, and Blood Institute, by R01 HL076831 (PI: Diez Roux), and by funding from the MacArthur Foundation. Jennifer A Nettleton was funded by a career development award from the NIH, the National Institute of Diabetes and Digestive and Kidney Diseases (5K01DK082729-04). MESA thanks the investigators, staff, and participants of the MESA study for their valuable contributions. A full list of participating MESA investigators and institutions can be found at <http://www.mesa-nhlbi.org>. SE Humphries is a British Heart Foundation Professor and he and Klelia D Salpea are supported by the British Heart Foundation (RG008/08 and FS/06/053) and by the National Institute for Health Research University College London Hospitals Biomedical Research Centre. The EARSII study was funded by the EC Concerted Action BMH1 CT92-0206. Phenotype collection was supported by Age UK (The Disconnected Mind project). The work was undertaken by The University of Edinburgh Centre for Cognitive Ageing and Cognitive Epidemiology, which is part of the cross-council Lifelong Health and Wellbeing Initiative (MR/K026992/1). Funding from the Biotechnology and Biological Sciences Research Council and Medical Research Council (MRC) is gratefully acknowledged. The Sister Study was supported by the Intramural Research Program of the NIH, National Institute of Environmental Health Sciences (Z01 ES 044005). The Jerusalem LRC study was funded by the Chief Scientist of the Israel Ministry of Health (300000-5352), the Israel Science Foundation (593/01), and the US-Israel Binational Science Foundation (87-00419). The Jerusalem Palestinians study was funded by the USAID MERC program (grant TA-MOU-01-M21-002) and by a research grant from DCURE Israel. The PATH study was supported by the Australian National Health and Medical Research Council program grant 17805 and project grant 418020. The Bruneck Study was supported by a grant from the "Tiroler Wissenschaftsfonds," Austria, and the Dr Johannes and Hertha Tuba Foundation. The Fels Longitudinal Study, Research reported in this manuscript was supported by the NIH grants (R03AG023251, R01HD012252).

The authors thank the Indian Health Service facilities. The views expressed in this article are those of the authors and do not necessarily reflect those

of the Indian Health Service. This study was supported by NIH grants R01DK091369, K01AG034259, R21HL092363, and cooperative agreement grants U01HL65520, U01HL41642, U01HL41652, U01HL41654, and U01HL65521.

SOLVABLE was supported by NIH/NIAMS T32AR07611, K24AR002138, and P60 AR30692, which is now updated as P60 AR064464, Kirkland Scholar Award, and the AFMR Summer Clinical and the Eleanor Wood-Prince grant: A Project of the Woman's Board of Northwestern Memorial Hospital. The Former Athletes study was funded by the Ministry of Education and Culture, the Juho Vainio Foundation, the Finnish Heart Research Foundation, Paavo Nurmi Foundation, the Finnish Cultural Foundation, and by a grant from Medical Society of Finland, Finska Läkaresällskapet. We thank National Institute for Health and Welfare, Department of Public Health, University of Helsinki, Sports and Exercise Medicine Department of Health Sciences, University of Jyväskylä, Paavo Nurmi Centre, Turku, and ORTON Research Institute, Invalid Foundation, Helsinki, for collaboration during the large epidemiological and clinical research program in the year 2008. The Finnish Twin Cohort data come primarily from the Nicotine Addictions Genetics family study of twins and their siblings. Data collection for this work was supported by Academy of Finland grants and NIH grant DA12854 (PAFM), and further support was from ENGAGE–European Network for Genetic and Genomic Epidemiology, FP7-HEALTH-F4-2007, grant agreement 201413; the Academy of Finland Center of Excellence in Complex Disease Genetics (grants 213506 and 129680; to J Kaprio), and the Academy of Finland (grants 141054, 265240, and 263278; to J Kaprio). The US Kidney Cancer Study (USKCS) was supported, in part, by the intramural research program of the Division of Cancer Epidemiology and Genetics, National Cancer Institute, NIH. The KORA study was initiated and financed by the Helmholtz Zentrum München– German Research Center for Environmental Health, which is funded by the German Federal Ministry of Education and Research (BMBF) and by the State of Bavaria. Furthermore, KORA research was supported within the Munich Center of Health Sciences (MC-Health), Ludwig-Maximilians-Universität, as part of LMUinnovativ. The Erasmus Rucphen Family (ERF) study was supported by grants from The Netherlands Organisation for Scientific Research (NWO), Erasmus MC, the Center for Medical Systems Biology (CMSB), the European Community's Seventh Framework Programme (FP7/2007-2013), the ENGAGE Consortium (grant HEALTH-F4-2007- 201413), and the Netherlands Consortium for Healthy Ageing (grant 050-060-810). We are grateful to all general practitioners for their contributions, to Petra Veraart for her help in genealogy, Jeannette Vergeer for the supervision of the laboratory work, and Peter Snijders for his help in data collection. The Rotterdam Study is supported by the Erasmus MC and Erasmus University Rotterdam; the Netherlands Organization for Scientific Research (NWO); the Netherlands Organization for Health Research and Development (ZonMw); the Research Institute for Diseases in the Elderly (RIDE); the Netherlands Genomics Initiative (NGI); the Ministry of Education, Culture, and Sciences; the Ministry of Health, Welfare, and Sports; the European Commission (DG XII); and the Municipality of Rotterdam. The analysis of data from the NHANES was funded by grant R01 AG033592 from the National Institute on Aging (PI: Elissa Epel). The Zutphen Elderly Study was supported by grants from The Netherlands Prevention Foundation (Preventiefonds) and the survey on Crete was financially supported by Danone. The RPCI study was supported by grants from Susan G Komen for the Cure (BCTR 0600562) and the US NIH Cancer Center Support Grant (P30 CA016056). The Heart Scan Study was funded by the British Heart Foundation (RG/10/005/28296) and the Medical Research Council, United Kingdom. A Steptoe is funded by the British Heart Foundation. The Helsinki Businessmen Study (HBS) was initially funded by the Academy of Finland and various foundations during the decades, most recently by the Sohlberg Foundation, the Jahnsen Foundation, and Gustav Voch Victoria Frimurarestiftelse. Funding has also been provided by the Helsinki University Central Hospital, the Oulu University Hospital (VTR/EVO-funding), and the Academy of Finland grants 286294 and 294154. Laboratory analysis of the Cebu Longitudinal Health and Nutrition Survey (CLHNS) was funded by NSF (Doctoral Dissertation Improvement grant BCS-0962282), the Wenner-Gren Foundation (grant

8111), and institutional support from Northwestern University; data and sample collection were funded by the NIH (grants RR20649 and ES10126). DNA extracts were provided by Karen Mohlke. Richard Cawthon, Justin Tackney, Katarina Nordfjäll, Klelia Salpea, Christine Ackerman, and Margrit Urbanek provided laboratory advice. We especially thank the many researchers at the Office of Population Studies, University of San Carlos, Cebu, the Philippines, for their central role in study design and data collection, and the Filipino participants who provided their time and samples for this study. NESDA is funded through the Geestkracht program of the Netherlands Organization for Health Research and Development (Zon-Mw, grant 10-000-1002) and is supported by participating universities and mental health care organizations. JE Verhoeven and telomere length assaying were supported through an NWO-VICI grant (91811602). The authors of PRT thank Gislaine Satyko Kogure and Rafael Costa Silva for personal guidance regarding progressive resistance training. We are grateful to the São Paulo Research Foundation (Fundação de Amparo à Pesquisa do Estado de São Paulo–FAPESP), for financial support with grants 2010/08800-8 and 2012/11566-2 (RMR) and fellowship 2012/11069-9 (CLMF). Fadi Charchar's work was supported by a National Health and Medical Research Council of Australia grant APP1123472. Maciej Tomaszewski's work on the Y chromosome was supported by British Heart Foundation project grants PG/16/49/32176 and PG/12/9/29376. The ESTHER study was funded by the Baden-Württemberg State Ministry of Science, Research, and Arts (Stuttgart, Germany); the Federal Ministry of Education and Research (Berlin, Germany); and the Federal Ministry of Family Affairs, Senior Citizens, Women, and Youth (Berlin, Germany). The Berlin Aging Study II (BASE-II) was supported by the German Federal Ministry of Education and Research (BMBF grants 16SV5536K, 16SV5537, 16SV5538, and 16SV5837; previously 01UW0808). The study "Sex- and Gender-Sensitive Prevention of Cardiovascular and Metabolic Disease in Older Adults in Germany" (GendAge) includes analyses of BASE-II baseline data and follow-up analyses of BASE-II participants with a focus on sex and gender differences in cardiovascular and metabolic diseases. GendAge is also supported by the BMBF (grants 01GL1716A and 01GL1716B). Additional contributions (e.g., financial, equipment, logistics, personnel) were made from each of the other participating sites [i.e., the Max Planck Institute for Human Development (MPIB); Max Planck Institute for Molecular Genetics (MPIMG); Charite-Universitätsmedizin, University Medicine, German; Institute for Economic Research (DIW) Humboldt-Universität zu Berlin; all located in Berlin, Germany, and University of Lübeck in Lübeck, and University of Tübingen, Germany].

This study corresponds to a substudy of "Venado Tuerto 2." The authors thank Hugo Grancelli, Dra Monica Damiano, Daniel Fox, and Carlos Ranalli; the Regional Centre for Development; School of Social Work of Venado Tuerto; Nutrition Society of Venado Tuerto; and the Department of Cardiology Hospital Alejandro Gutierrez of Venado Tuerto for their collaboration.

GAHR2 was supported by grants awarded to Dr. D'Antono by the Canadian Institutes of Health Research (CIHR; MOP 79456 and 111017) and the Montreal Heart Institute Foundation. The RPE study was funded by research grants from the Singapore Ministry of Education, the AXA Research Foundation, and the Templeton World Charity Foundation, Inc. The Sweden Mindfulness Study was supported by the Swedish Research Council and FORTE to Kristina Sundquist and Jan Sundquist and the National Heart, Lung, and Blood Institute of the NIH under award R01HL116381 to Kristina Sundquist. The Ukrainian part of the data of the Kyev study was completed within the framework of the "Research of anthropometric, biochemical and genetic risk factors of development of the glucose regulation disturbances and diabetes mellitus in Ukraine study" conducted at the Komisarenko Institute of Endocrinology and Metabolism (National Academy of Medical Sciences grant 0116 U 002166).

The authors' responsibilities were as follows—MG: wrote the manuscript, researched data, contributed to the design and discussion, and had responsibility for final content; GJH, EEA, and MPZ: contributed to the design and discussion and reviewed and edited the manuscript; KN, MM, MB, TdM, AEH, EJG, SCH, JAN, KDS, VAD, RF-F, GA, SEH, LH, DG, IH, JDK, HN, DJK, KAM, PW, Y-LZ, SP, EWD, LR, DB, AS, LB Boardman, AM, BN, WZ,

RR-G, AJP, JK, JNH, CG, G Paolisso, JBHH, LM, TS, JW, KS, PvdH, LB Broer, FK, BK, TS, DTAE, CD, JEV, RS, RZ, RMRdR, FJC, MTT, UM, ID, AEIM, GC, DK, BD, MK, BJM, RPE, KS, G Pare, and MC: researched data and reviewed and edited the manuscript; and all authors: read and approved the final manuscript. None of the authors declared a conflict of interest.

REFERENCES

1. Allsopp RC, Harley CB. Evidence for a critical telomere length in senescent human fibroblasts. *Exp Cell Res* 1995;219:130–6.
2. Hayflick L. Mortality and immortality at the cellular level: a review *Biochemistry (Mosc)* 1997;62:1180–90.
3. Valdes AM, Andrew T, Gardner JP, Kimura M, Oelsner E, Cherkas LF, Aviv A, Spector TD. Obesity, cigarette smoking, and telomere length in women. *Lancet* 2005;366:662–4.
4. Sampson MJ, Winterbone MS, Hughes JC, Dozio N, Hughes DA. Monocyte telomere shortening and oxidative DNA damage in type 2 diabetes. *Diabetes Care* 2006;29:283–9.
5. Demissie S, Levy D, Benjamin EJ, Cupples LA, Gardner JP, Herbert A, Kimura M, Larson MG, Meigs JB, Keaney JF, et al. Insulin resistance, oxidative stress, hypertension, and leukocyte telomere length in men from the Framingham Heart Study. *Aging Cell* 2006;5:325–30.
6. WHO. Fact sheet 311. Obesity and overweight 2012. Available from: <http://www.who.int/mediacentre/factsheets/fs311/en/index.htm> 2012.
7. Mundstock E, Sarria EE, Zatti H, Mattos Louzada F, Kich Grun L, Herbert Jones M, Guma FT, Mazzola (In Memoriam) J, Epifanio M, Stein RT, et al. Effect of obesity on telomere length: systematic review and meta-analysis. *Obesity (Silver Spring)* 2015;23:2165–74.
8. Njajou OT, Cawthon RM, Blackburn EH, Harris TB, Li R, Sanders JL, Newman AB, Nalls M, Cummings SR, Hsueh WC. Shorter telomeres are associated with obesity and weight gain in the elderly. *Int J Obes (Lond)* 2012;36:1176–9.
9. Martin-Ruiz CM, Gussekloo J, van Heemst D, von Zglinicki T, Westendorp RG. Telomere length in white blood cells is not associated with morbidity or mortality in the oldest old: a population-based study. *Aging Cell* 2005;4:287–90.
10. Benetos A, Okuda K, Lajemi M, Kimura M, Thomas F, Skurnick J, Labat C, Bean K, Aviv A. Telomere length as an indicator of biological aging: the gender effect and relation with pulse pressure and pulse wave velocity. *Hypertension* 2001;37:381–5.
11. Gardner M, Bann D, Wiley L, Cooper R, Hardy R, Nitsch D, Martin-Ruiz C, Shiels P, Sayer AA, Barbieri M, et al. Gender and telomere length: systematic review and meta-analysis. *Exp Gerontol* 2014;51:15–27.
12. McGrath M, Wong JY, Michaud D, Hunter DJ, De Vivo I. Telomere length, cigarette smoking, and bladder cancer risk in men and women. *Cancer Epidemiol Biomarkers Prev* 2007;16:815–9.
13. O'Donnell CJ, Demissie S, Kimura M, Levy D, Gardner JP, White C, D'Agostino RB, Wolf PA, Polak J, Cupples LA, et al. Leukocyte telomere length and carotid artery intimal medial thickness: the Framingham Heart Study. *Arterioscler Thromb Vasc Biol* 2008;28:1165–71.
14. Bekaert S, De Meyer T, Rietzschel ER, De Buyzere ML, De Bacquer D, Langlois M, Segers P, Cooman L, Van Damme P, Cassiman P, et al. Telomere length and cardiovascular risk factors in a middle-aged population free of overt cardiovascular disease. *Aging Cell* 2007;6:639–47.
15. Nordfjäll K, Eliasson M, Stegmayr B, Melander O, Nilsson P, Roos G. Telomere length is associated with obesity parameters but with a gender difference. *Obesity (Silver Spring)* 2008;16:2682–9.
16. Story M, Evans M, Fabsitz RR, Clay TE, Holy Rock B, Broussard B. The epidemic of obesity in American Indian communities and the need for childhood obesity-prevention programs. *Am J Clin Nutr* 1999;69(Suppl):747S–54S.
17. Hunt SC, Chen W, Gardner JP, Kimura M, Srinivasan SR, Eckfeldt JH, Berenson GS, Aviv A. Leukocyte telomeres are longer in African Americans than in whites: the National Heart, Lung, and Blood Institute Family Heart Study and the Bogalusa Heart Study. *Aging Cell* 2008;7:451–8.
18. Chen W, Kimura M, Kim S, Cao X, Srinivasan SR, Berenson GS, Kark JD, Aviv A. Longitudinal versus cross-sectional evaluations of leukocyte telomere length dynamics: age-dependent telomere

- shortening is the rule. *J Gerontol A Biol Sci Med Sci* 2011;66:312–9.
19. Diaz VA, Mainous AG, Player MS, Everett CJ. Telomere length and adiposity in a racially diverse sample. *Int J Obes (Lond)* 2010;34:261–5.
 20. Zhu H, Wang X, Gutin B, Davis CL, Keeton D, Thomas J, Stallmann-Jorgensen I, Mookken G, Bundy V, Snieder H, et al. Leukocyte telomere length in healthy Caucasian and African-American adolescents: relationships with race, sex, adiposity, adipokines, and physical activity. *J Pediatr* 2011;158:215–20.
 21. Fitzpatrick AL, Kronmal RA, Kimura M, Gardner JP, Psaty BM, Jenny NS, Tracy RP, Hardikar S, Aviv A. Leukocyte telomere length and mortality in the Cardiovascular Health Study. *J Gerontol A Biol Sci Med Sci* 2011;66:421–9.
 22. Muezzinler A, Zaineddin AK, Brenner H. Body mass index and leukocyte telomere length in adults: a systematic review and meta-analysis. *Obes Rev* 2014;15:192–201.
 23. Cawthon RM. Telomere measurement by quantitative PCR. *Nucleic Acids Res* 2002;30:e47.
 24. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629–34.
 25. Duval S, Tweedie R. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* 2000;56:455–63.
 26. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539–58.
 27. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557–60.
 28. Aviv A, Valdes A, Gardner JP, Swaminathan R, Kimura M, Spector TD. Menopause modifies the association of leukocyte telomere length with insulin resistance and inflammation. *J Clin Endocrinol Metab* 2006;91:635–40.
 29. Valdes AM, Richards JB, Gardner JP, Swaminathan R, Kimura M, Xiaobin L, Aviv A, Spector TD. Telomere length in leukocytes correlates with bone mineral density and is shorter in women with osteoporosis. *Osteoporos Int* 2007;18:1203–10.
 30. Richards JB, Valdes AM, Gardner JP, Kato BS, Siva A, Kimura M, Lu X, Brown MJ, Aviv A, Spector TD. Homocysteine levels and leukocyte telomere length. *Atherosclerosis* 2008;200:271–7.
 31. Cherkas LF, Hunkin JL, Kato BS, Richards JB, Gardner JP, Surdulescu GL, Kimura M, Lu X, Spector TD, Aviv A. The association between physical activity in leisure time and leukocyte telomere length. *Arch Intern Med* 2008;168:154–8.
 32. Gardner JP, Li S, Srinivasan SR, Chen W, Kimura M, Lu X, Berenson GS, Aviv A. Rise in insulin resistance is associated with escalated telomere attrition. *Circulation* 2005;111:2171–7.
 33. Aviv A, Chen W, Gardner JP, Kimura M, Brimacombe M, Cao X, Srinivasan SR, Berenson GS. Leukocyte telomere dynamics: longitudinal findings among young adults in the Bogalusa Heart Study. *Am J Epidemiol* 2009;169:323–9.
 34. Chen W, Gardner JP, Kimura M, Brimacombe M, Cao X, Srinivasan SR, Berenson GS, Aviv A. Leukocyte telomere length is associated with HDL cholesterol levels: the Bogalusa Heart Study. *Atherosclerosis* 2009;205:620–5.
 35. Adakalakeswari A, Balasubramanyam M, Ravikumar R, Deepa R, Mohan V. Association of telomere shortening with impaired glucose tolerance and diabetic macroangiopathy. *Atherosclerosis* 2007;195:83–9.
 36. Barbieri M, Paolisso G, Kimura M, Gardner JP, Boccardi V, Papa M, Hjelmborg JV, Christensen K, Brimacombe M, Nawrot TS, et al. Higher circulating levels of IGF-1 are associated with longer leukocyte telomere length in healthy subjects. *Mech Ageing Dev* 2009;130:771–6.
 37. De Meyer T, Rietzschel ER, De Buyzere ML, De Bacquer D, Van Criekinge W, De Backer GG, Gillebert TC, Van Oostveldt P, Bekaert S. Paternal age at birth is an important determinant of offspring telomere length. *Hum Mol Genet* 2007;16:3097–102.
 38. De Meyer T, Rietzschel ER, De Buyzere ML, Langlois MR, De Bacquer D, Segers P, Van Damme P, De Backer GG, Van Oostveldt P, Van Criekinge W, et al. Systemic telomere length and preclinical atherosclerosis: the Asklepios Study. *Eur Heart J* 2009;30:3074–81.
 39. Rietzschel ER, De Buyzere ML, Bekaert S, Segers P, De Bacquer D, Cooman L, Van Damme P, Cassiman P, Langlois M, van Oostveldt P, et al. Rationale, design, methods and baseline characteristics of the Asklepios Study. *Eur J Cardiovasc Prev Rehabil* 2007;14:179–91.
 40. Vasan RS, Demissie S, Kimura M, Cupples LA, White C, Gardner JP, Cao X, Levy D, Benjamin EJ, Aviv A. Association of leukocyte telomere length with echocardiographic left ventricular mass: the Framingham Heart Study. *Circulation* 2009;120:1195–202.
 41. Houben JM, Giltay EJ, Rius-Ottenheim N, Hageman GJ, Kromhout D. Telomere length and mortality in elderly men: the Zutphen Elderly Study. *J Gerontol A Biol Sci Med Sci* 2011;66:38–44.
 42. Houben JM, Mercken EM, Ketelslegers HB, Bast A, Wouters EF, Hageman GJ, Schols AM. Telomere shortening in chronic obstructive pulmonary disease. *Respir Med* 2009;103:230–6.
 43. Rius-Ottenheim N, Houben JM, Kromhout D, Kafatos A, van der Mast RC, Zitman FG, Geleijnse JM, Hageman GJ, Giltay EJ. Telomere length and mental well-being in elderly men from the Netherlands and Greece. *Behav Genet* 2012;42:278–86.
 44. Jeanclous E, Schork NJ, Kyvik KO, Kimura M, Skurnick JH, Aviv A. Telomere length inversely correlates with pulse pressure and is highly familial. *Hypertension* 2000;36:195–200.
 45. Hjelmborg JV, Nzietchueng R, Kimura M, Gardner JP, Bladbjerg EM, Christensen K, Aviv A, Benetos A. Leukocyte telomere length is inversely correlated with plasma Von Willebrand factor. *Thromb Res* 2010;125:e339–42.
 46. Farzaneh-Far R, Lin J, Epel E, Lapham K, Blackburn E, Whooley MA. Telomere length trajectory and its determinants in persons with coronary artery disease: longitudinal findings from the heart and soul study. *PLoS One* 2010;5:e8612.
 47. Farzaneh-Far R, Lin J, Epel ES, Harris WS, Blackburn EH, Whooley MA. Association of marine omega-3 fatty acid levels with telomeric aging in patients with coronary heart disease. *JAMA* 2010;303:250–7.
 48. Huda N, Tanaka H, Herbert BS, Reed T, Gilley D. Shared environmental factors associated with telomere length maintenance in elderly male twins. *Aging Cell* 2007;6:709–13.
 49. Fitzpatrick AL, Reed T, Goldberg J, Buchwald D. The association between prolonged fatigue and cardiovascular disease in World War II veteran twins. *Twin Res* 2004;7:571–7.
 50. Strandberg TE, Kovanen PT, Eklund KK. Is the “lipid paradox” in rheumatoid arthritis really a paradox? Comment on the article by Bartels et al. *Arthritis Rheum* 2011;63:3644–5; author reply: 5–6.
 51. Strandberg TE, Saijonmaa O, Fyhrquist F, Tilvis RS, Strandberg AY, Miettinen TA, Pitkala KH, Salomaa V. Telomere length in old age and cholesterol across the life course. *J Am Geriatr Soc* 2011;59:1979–81.
 52. Strandberg TE, Saijonmaa O, Tilvis RS, Pitkala KH, Strandberg AY, Miettinen TA, Fyhrquist F. Association of telomere length in older men with mortality and midlife body mass index and smoking. *J Gerontol A Biol Sci Med Sci* 2011;66:815–20.
 53. Rode L, Nordestgaard BG, Weischer M, Bojesen SE. Increased body mass index, elevated C-reactive protein, and short telomere length. *J Clin Endocrinol Metab* 2014;99:E1671–5.
 54. Skamra C, Romero-Diaz J, Sandhu A, Huang Q, Lee J, Pearce W, McPherson DD, Sutton-Tyrrell K, Pope R, Ramsey-Goldman R. Telomere length in patients with systemic lupus erythematosus and its associations with carotid plaque. *Rheumatology* 2013;52:1101–8.
 55. Kark JD, Goldberger N, Kimura M, Sinnreich R, Aviv A. Energy intake and leukocyte telomere length in young adults. *Am J Clin Nutr* 2012;95:479–87.
 56. Kark JD, Nassar H, Shaham D, Sinnreich R, Goldberger N, Aboudi V, Bogot NR, Kimura M, Aviv A. Leukocyte telomere length and coronary artery calcification in Palestinians. *Atherosclerosis* 2013;229:363–8.
 57. Harris SE, Martin-Ruiz C, von Zglinicki T, Starr JM, Deary IJ. Telomere length and aging biomarkers in 70-year-olds: the Lothian Birth Cohort 1936. *Neurobiol Aging* 2012;33:1486, e3–8.
 58. Deary IJ, Gow AJ, Taylor MD, Corley J, Brett C, Wilson V, Campbell H, Whalley LJ, Visscher PM, Porteous DJ, et al. The Lothian Birth Cohort 1936: a study to examine influences on cognitive ageing from age 11 to age 70 and beyond. *BMC Geriatr* 2007;7:28.
 59. Zannolli R, Mohn A, Buoni S, Pietrobelli A, Messina M, Chiarelli F, Miracco C. Telomere length and obesity. *Acta Paediatr* 2008;97:952–4.
 60. Iglesias Molli AE, Panero J, Dos Santos PC, Gonzalez CD, Vilarino J, Sereday M, Cerrone GE, Slavutsky I, Frechtel GD. Metabolically

- healthy obese women have longer telomere length than obese women with metabolic syndrome. *PLoS One* 2017;12:e0174945.
61. Xue HM, Liu QQ, Tian G, Quan LM, Zhao Y, Cheng G. Television watching and telomere length among adults in southwest China. *Am J Public Health* 2017;107:1425–32.
 62. Diaz VA, Mainous AG III, Everett CJ, Schoepf UJ, Codd V, Samani NJ. Effect of healthy lifestyle behaviors on the association between leukocyte telomere length and coronary artery calcium. *Am J Cardiol* 2010;106:659–63.
 63. Willeit P, Willeit J, Brandstatter A, Ehrlenbach S, Mayr A, Gasperi A, Weger S, Oberhollenzer F, Reindl M, Kronenberg F, et al. Cellular aging reflected by leukocyte telomere length predicts advanced atherosclerosis and cardiovascular disease risk. *Arterioscler Thromb Vasc Biol* 2010;30:1649–56.
 64. Willeit P, Willeit J, Mayr A, Weger S, Oberhollenzer F, Brandstatter A, Kronenberg F, Kiechl S. Telomere length and risk of incident cancer and cancer mortality. *JAMA* 2010;304:69–75.
 65. Zheng YL, Ambrosone C, Byrne C, Davis W, Nesline M, McCann SE. Telomere length in blood cells and breast cancer risk: investigations in two case-control studies. *Breast Cancer Res Treat* 2010;120:769–75.
 66. Mainous AG III, Codd V, Diaz VA, Schoepf UJ, Everett CJ, Player MS, Samani NJ. Leukocyte telomere length and coronary artery calcification. *Atherosclerosis* 2010;210:262–7.
 67. Mainous AG III, Everett CJ, Diaz VA, Baker R, Mangino M, Codd V, Samani NJ. Leukocyte telomere length and marital status among middle-aged adults. *Age Ageing* 2011;40:73–8.
 68. Ehrlenbach S, Willeit P, Kiechl S, Willeit J, Reindl M, Schanda K, Kronenberg F, Brandstatter A. Influences on the reduction of relative telomere length over 10 years in the population-based Bruneck study: introduction of a well-controlled high-throughput assay. *Int J Epidemiol* 2009;38:1725–34.
 69. Bull CF, O'Callaghan NJ, Mayrhofer G, Fenech MF. Telomere length in lymphocytes of older South Australian men may be inversely associated with plasma homocysteine. *Rejuvenation Res* 2009;12:341–9.
 70. Cross JA, Brennan C, Gray T, Temple RC, Dozio N, Hughes JC, Levell NJ, Murphy H, Fowler D, Hughes DA, et al. Absence of telomere shortening and oxidative DNA damage in the young adult offspring of women with pre-gestational type 1 diabetes. *Diabetologia* 2009;52:226–34.
 71. Nettleton JA, Diez-Roux A, Jenny NS, Fitzpatrick AL, Jacobs DR Jr. Dietary patterns, food groups, and telomere length in the Multi-Ethnic Study of Atherosclerosis (MESA). *Am J Clin Nutr* 2008;88:1405–12.
 72. Roux AV, Ranjit N, Jenny NS, Shea S, Cushman M, Fitzpatrick A, Seeman T. Race/ethnicity and telomere length in the Multi-Ethnic Study of Atherosclerosis. *Aging Cell* 2009;8:251–7.
 73. Salpea KD, Nicaud V, Tiret L, Talmud PJ, Humphries SE. The association of telomere length with paternal history of premature myocardial infarction in the European Atherosclerosis Research Study II. *J Mol Med (Berl)* 2008;86:815–24.
 74. Puterman E, Lin J, Blackburn E, O'Donovan A, Adler N, Epel E. The power of exercise: buffering the effect of chronic stress on telomere length. *PLoS One* 2010;5:e10837.
 75. Atzmon G, Cho M, Cawthon RM, Budagov T, Katz M, Yang X, Siegel G, Bergman A, Huffman DM, Schechter CB, et al. Evolution in health and medicine Sackler colloquium: genetic variation in human telomerase is associated with telomere length in Ashkenazi centenarians. *Proc Natl Acad Sci USA* 2010;107(Suppl 1):1710–7.
 76. Hou L, Savage SA, Blaser MJ, Perez-Perez G, Hoxha M, Dioni L, Pegoraro V, Dong LM, Zatonski W, Lissowska J, et al. Telomere length in peripheral leukocyte DNA and gastric cancer risk. *Cancer Epidemiol Biomarkers Prev* 2009;18:3103–9.
 77. Kananen L, Surakka I, Pirkola S, Suvisaari J, Lonnqvist J, Peltonen L, Ripatti S, Hovatta I. Childhood adversities are associated with shorter telomere length at adult age both in individuals with an anxiety disorder and controls. *PLoS One* 2010;5:e10826.
 78. Kim S, Parks CG, DeRoo LA, Chen H, Taylor JA, Cawthon RM, Sandler DP. Obesity and weight gain in adulthood and telomere length. *Cancer Epidemiol Biomarkers Prev* 2009;18:816–20.
 79. Kurz DJ, Kloeckener-Gruissem B, Akhmedov A, Eberli FR, Buhler I, Berger W, Bertel O, Luscher TF. Degenerative aortic valve stenosis, but not coronary disease, is associated with shorter telomere length in the elderly. *Arterioscler Thromb Vasc Biol* 2006;26:e114–7.
 80. Mather KA, Jorm AF, Milburn PJ, Tan X, Eastaer S, Christensen H. No associations between telomere length and age-sensitive indicators of physical function in mid and later life. *J Gerontol A Biol Sci Med Sci* 2010;65:792–9.
 81. Pavanello S, Hoxha M, Dioni L, Bertazzi PA, Snenghi R, Nalesso A, Ferrara SD, Montisci M, Baccarelli A. Shortened telomeres in individuals with abuse in alcohol consumption. *Int J Cancer* 2011;129:983–92.
 82. Lee M, Martin H, Firpo MA, Demerath EW. Inverse association between adiposity and telomere length: the Fels Longitudinal Study. *Am J Hum Biol* 2011;23:100–6.
 83. Barrera G, Cases T, Bunout D, de la Maza MP, Leiva L, Rodriguez JM, Hirsch S. Associations between socioeconomic status, aging and functionality among older women. *Geriatr Nurs* 2017;38(4):347–51
 84. Bunout D, Barrera G, de la Maza MP, Leiva L, Hirsch S. Effect of weight maintenance or gain in a 10 years period over telomere length, sirtuin 1 and 6 expression and carotid intima media thickness. *J Hum Nutr Diet* 2015;28:155–64.
 85. Jackowska M, Hamer M, Carvalho LA, Erusalimsky JD, Butcher L, Steptoe A. Short sleep duration is associated with shorter telomere length in healthy men: findings from the Whitehall II cohort study. *PLoS One* 2012;7:e47292.
 86. Zalli A, Carvalho LA, Lin J, Hamer M, Erusalimsky JD, Blackburn EH, Steptoe A. Shorter telomeres with high telomerase activity are associated with raised allostatic load and impoverished psychosocial resources. *Proc Natl Acad Sci USA* 2014;111:4519–24.
 87. Wong JY, De Vivo I, Lin X, Christiani DC. Cumulative PM(2.5) exposure and telomere length in workers exposed to welding fumes. *J Toxicol Environ Health A* 2014;77:441–55.
 88. Wong JY, De Vivo I, Lin X, Grashow R, Cavallari J, Christiani DC. The association between global DNA methylation and telomere length in a longitudinal study of boilermakers. *Genet Epidemiol* 2014;38:254–64.
 89. Boardman LA, Litzelman K, Seo S, Johnson RA, Vanderboom RJ, Kimmel GW, Cunningham JM, Gangnon RE, Engelman CD, Riegert-Johnson DL, et al. The association of telomere length with colorectal cancer differs by the age of cancer onset. *Clin Transl Gastroenterol* 2014;5:e52.
 90. Skinner HG, Gangnon RE, Litzelman K, Johnson RA, Chari ST, Petersen GM, Boardman LA. Telomere length and pancreatic cancer: a case-control study. *Cancer Epidemiol Biomarkers Prev* 2012;21:2095–100.
 91. Savolainen K, Eriksson JG, Kajantie E, Lahti M, Raikonen K. The history of sleep apnea is associated with shorter leukocyte telomere length: the Helsinki Birth Cohort Study. *Sleep Med* 2014;15:209–12.
 92. Savolainen K, Raikonen K, Kananen L, Kajantie E, Hovatta I, Lahti M, Lahti J, Pesonen AK, Heinonen K, Eriksson JG. History of mental disorders and leukocyte telomere length in late adulthood: the Helsinki Birth Cohort Study (HBCS). *J Psychiatr Res* 2012;46:1346–53.
 93. Hoen PW, Rosmalen JG, Schoevers RA, Huzen J, van der Harst P, de Jonge P. Association between anxiety but not depressive disorders and leukocyte telomere length after 2 years of follow-up in a population-based sample. *Psychol Med* 2013;43:689–97.
 94. Huzen J, Wong LS, van Veldhuisen DJ, Samani NJ, Zwinderman AH, Codd V, Cawthon RM, Benus GF, van der Horst IC, Navis G, et al. Telomere length loss due to smoking and metabolic traits. *J Intern Med* 2014;275:155–63.
 95. Kingma EM, de Jonge P, van der Harst P, Ormel J, Rosmalen JG. The association between intelligence and telomere length: a longitudinal population based study. *PLoS One* 2012;7:e49356.
 96. Zhao J, Zhu Y, Lin J, Matsuguchi T, Blackburn E, Zhang Y, Cole SA, Best LG, Lee ET, Howard BV. Short leukocyte telomere length predicts risk of diabetes in American Indians: the Strong Heart Family Study. *Diabetes* 2014;63:354–62.
 97. Zhu Y, Voruganti VS, Lin J, Matsuguchi T, Blackburn E, Best LG, Lee ET, MacCluer JW, Cole SA, Zhao J. QTL mapping of leukocyte telomere length in American Indians: the Strong Heart Family Study. *Aging* 2013;5:704–16.
 98. Garcia-Calzon S, Gea A, Razquin C, Corella D, Lamuela-Raventos RM, Martinez JA, Martinez-Gonzalez MA, Zalba G, Marti A. Longitudinal association of telomere length and obesity indices in an intervention study with a Mediterranean diet: the PREDIMED-NAVARRA trial. *Int J Obes (Lond)* 2014;38:177–82.

99. Needham BL, Adler N, Gregorich S, Rehkopf D, Lin J, Blackburn EH, Epel ES. Socioeconomic status, health behavior, and leukocyte telomere length in the National Health and Nutrition Examination Survey, 1999–2002. *Soc Sci Med* 2013;85:1–8.
100. Cui Y, Cai Q, Qu S, Chow WH, Wen W, Xiang YB, Wu J, Rothman N, Yang G, Shu XO, et al. Association of leukocyte telomere length with colorectal cancer risk: nested case-control findings from the Shanghai Women's Health Study. *Cancer Epidemiol Biomarkers Prev* 2012;21:1807–13.
101. Kozlitina J, Garcia CK. Red blood cell size is inversely associated with leukocyte telomere length in a large multi-ethnic population. *PLoS One* 2012;7:e51046.
102. Pellatt AJ, Wolff RK, Lundgreen A, Cawthon R, Slattery ML. Genetic and lifestyle influence on telomere length and subsequent risk of colon cancer in a case control study. *Int J Mol Epidemiol Genet* 2012;3:184–94.
103. Broer L, Raschenberger J, Deelen J, Mangino M, Codd V, Pietilainen KH, Albrecht E, Amin N, Beekman M, de Craen AJ, et al. Association of adiponectin and leptin with relative telomere length in seven independent cohorts including 11,448 participants. *Eur J Epidemiol* 2014;29:629–38.
104. Hofmann JN, Baccarelli A, Schwartz K, Davis FG, Ruterbusch JJ, Hoxha M, McCarthy BJ, Savage SA, Wacholder S, Rothman N, et al. Risk of renal cell carcinoma in relation to blood telomere length in a population-based case-control study. *Br J Cancer* 2011;105:1772–5.
105. Kajantie E, Pietilainen KH, Wehkalampi K, Kananen L, Raikonen K, Rissanen A, Hovi P, Kaprio J, Andersson S, Eriksson JG, et al. No association between body size at birth and leucocyte telomere length in adult life—evidence from three cohort studies. *Int J Epidemiol* 2012;41:1400–8.
106. Chen S, Lin J, Matsuguchi T, Blackburn E, Yeh F, Best LG, Devereux RB, Lee ET, Howard BV, Roman MJ, et al. Short leukocyte telomere length predicts incidence and progression of carotid atherosclerosis in American Indians: the Strong Heart Family Study. *Aging* 2014;6:414–27.
107. Chen S, Yeh F, Lin J, Matsuguchi T, Blackburn E, Lee ET, Howard BV, Zhao J. Short leukocyte telomere length is associated with obesity in American Indians: the Strong Heart Family Study. *Aging* 2014;6:380–9.
108. Savolainen K, Eriksson JG, Kajantie E, Lahti J, Raikonen K. Telomere length and hypothalamic-pituitary-adrenal axis response to stress in elderly adults. *Psychoneuroendocrinology* 2015;53:179–84.
109. Savolainen K, Eriksson JG, Kajantie E, Pesonen AK, Raikonen K. Associations between the five-factor model of personality and leukocyte telomere length in elderly men and women: the Helsinki Birth Cohort Study (HBCS). *J Psychosom Res* 2015;79(3):233–8.
110. Savolainen K, Eriksson JG, Kananen L, Kajantie E, Pesonen AK, Heinonen K, Raikonen K. Associations between early life stress, self-reported traumatic experiences across the lifespan and leukocyte telomere length in elderly adults. *Biol Psychol* 2014;97:35–42.
111. Tiainen AM, Mannisto S, Blomstedt PA, Moltchanova E, Perala MM, Kaartinen NE, Kajantie E, Kananen L, Hovatta I, Eriksson JG. Leukocyte telomere length and its relation to food and nutrient intake in an elderly population. *Eur J Clin Nutr* 2012;66:1290–4.
112. Laine MK, Eriksson JG, Kujala UM, Raj R, Kaprio J, Backmand HM, Peltonen M, Sarna S. Effect of intensive exercise in early adult life on telomere length in later life in men. *J Sports Sci Med* 2015;14:239–45.
113. Loukola A, Wedenoja J, Keskitalo-Vuokko K, Broms U, Korhonen T, Ripatti S, Sarin AP, Pitkaniemi J, He L, Hapola A, et al. Genome-wide association study on detailed profiles of smoking behavior and nicotine dependence in a twin sample. *Mol Psychiatry* 2014;19:615–24.
114. Kim S, Sandler DP, Carswell G, De Roo LA, Parks CG, Cawthon R, Weinberg CR, Taylor JA. Telomere length in peripheral blood and breast cancer risk in a prospective case-cohort analysis: results from the Sister Study. *Cancer Causes Control* 2011;22:1061–6.
115. Eisenberg DT, Hayes MG, Kuzawa CW. Delayed paternal age of reproduction in humans is associated with longer telomeres across two generations of descendants. *Proc Natl Acad Sci USA* 2012;109:10251–6.
116. Mason C, Risques RA, Xiao L, Duggan CR, Imayama I, Campbell KL, Kong A, Foster-Schubert KE, Wang CY, Alfano CM, et al. Independent and combined effects of dietary weight loss and exercise on leukocyte telomere length in postmenopausal women. *Obesity (Silver Spring)* 2013;21:E549–54.
117. Comijs HC, van Marwijk HW, van der Mast RC, Naarding P, Oude Voshaar RC, Beekman AT, Boshuisen M, Dekker J, Kok R, de Waal MW, et al. The Netherlands Study of Depression in Older Persons (NESDO): a prospective cohort study. *BMC Res Notes* 2011;4:524.
118. Penninx BW, Beekman AT, Smit JH, Zitman FG, Nolen WA, Spinhoven P, Cuijpers P, De Jong PJ, Van Marwijk HW, Assendelft WJ, et al. The Netherlands Study of Depression and Anxiety (NESDA): rationale, objectives and methods. *Int J Methods Psychiatr Res* 2008;17:121–40.
119. Miranda-Furtado CL, Ramos FK, Kogure GS, Santana-Lemos BA, Ferriani RA, Calado RT, Dos Reis RM. A nonrandomized trial of progressive resistance training intervention in women with polycystic ovary syndrome and its implications in telomere content. *Reprod Sci* 2016;23:644–54.
120. Denham J, Nelson CP, O'Brien BJ, Nankervis SA, Denniff M, Harvey JT, Marques FZ, Codd V, Zukowska-Szczechowska E, Samani NJ, et al. Longer leukocyte telomeres are associated with ultra-endurance exercise independent of cardiovascular risk factors. *PLoS One* 2013;8:e69377.
121. Chilton WL, Marques FZ, West J, Kannourakis G, Berzins SP, O'Brien BJ, Charchar FJ. Acute exercise leads to regulation of telomere-associated genes and microRNA expression in immune cells. *PLoS One* 2014;9:e92088.
122. Bloomer LD, Nelson CP, Denniff M, Christofidou P, Debiec R, Thompson J, Zukowska-Szczechowska E, Samani NJ, Charchar FJ, Tomaszewski M. Coronary artery disease predisposing haplogroup I of the Y chromosome, aggression and sex steroids—genetic association analysis. *Atherosclerosis* 2014;233:160–4.
123. Gerstorff D, Bertram L, Lindenberger U, Pawelec G, Demuth I, Steinhagen-Thiessen E, Wagner GG. Editorial. *Gerontology* 2016;62:311–5.
124. Bertram L, Bockenhoff A, Demuth I, Duzel S, Eckardt R, Li SC, Lindenberger U, Pawelec G, Siedler T, Wagner GG, et al. Cohort profile: The Berlin Aging Study II (BASE-II). *Int J Epidemiol* 2014;43:703–12.
125. Meyer A, Salewsky B, Buchmann N, Steinhagen-Thiessen E, Demuth I. Relative leukocyte telomere length, hematological parameters and anemia—data from the Berlin Aging Study II (BASE-II). *Gerontology* 2016;62:330–6.
126. Khalangot M, Krasnienkov D, Vaiserman A, Avilov I, Kovtun V, Okhrimenko N, Koliada A, Kravchenko V. Leukocyte telomere length is inversely associated with post-load but not with fasting plasma glucose levels. *Exp Biol Med (Maywood)* 2017;242:700–8.
127. Starnino L, Busque L, Tardif JC, D'Antonio B. Psychological profiles in the prediction of leukocyte telomere length in healthy individuals. *PLoS One* 2016;11:e0165482.
128. Kasielski M, Eusebio MO, Pietruczuk M, Nowak D. The relationship between peripheral blood mononuclear cells telomere length and diet - unexpected effect of red meat. *Nutr J* 2016;15:68.
129. McDonnell BJ, Yasmin, Butcher L, Cockcroft JR, Wilkinson IB, Erusalimsky JD, McEniery CM. The age-dependent association between aortic pulse wave velocity and telomere length. *J Physiol* 2017;595:1627–35.
130. Yim OS, Zhang X, Shalev I, Monakhov M, Zhong S, Hsu M, Chew SH, Lai PS, Ebstein RP. Delay discounting, genetic sensitivity, and leukocyte telomere length. *Proc Natl Acad Sci USA* 2016;113:2780–5.
131. Wang X, Sundquist K, Hedelius A, Palmer K, Memon AA, Sundquist J. Leukocyte telomere length and depression, anxiety and stress and adjustment disorders in primary health care patients. *BMC Psychiatry* 2017;17:148.
132. Gerstein HC, Yusuf S, Holman R, Bosch J, Pogue J; Dream Trial Investigators. Rationale, design and recruitment characteristics of a large, simple international trial of diabetes prevention: the DREAM trial. *Diabetologia* 2004;47:1519–27.
133. Yusuf S, Hawken S, Ounpuu S, Dans T, Avezum A, Lanas F, McQueen M, Budaj A, Pais P, Varigos J, et al. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet* 2004;364:937–52.
134. van der Harst P, van der Steege G, de Boer RA, Voors AA, Hall AS, Mulder MJ, van Gilst WH, van Veldhuisen DJ; Merit-Hf Study Group.

- Telomere length of circulating leukocytes is decreased in patients with chronic heart failure. *J Am Coll Cardiol* 2007;49:1459–64.
135. Steenstrup T, Hjelmborg JV, Mortensen LH, Kimura M, Christensen K, Aviv A. Leukocyte telomere dynamics in the elderly. *Eur J Epidemiol* 2013;28:181–7.
 136. Muezzinler A, Zaineddin AK, Brenner H. A systematic review of leukocyte telomere length and age in adults. *Ageing Res Rev* 2013;12:509–19.
 137. de Heredia FP, Gomez-Martinez S, Marcos A. Obesity, inflammation and the immune system. *Proc Nutr Soc* 2012;71:332–8.
 138. Spiegelman D, Israel RG, Bouchard C, Willett WC. Absolute fat mass, percent body fat, and body-fat distribution: which is the real determinant of blood pressure and serum glucose? *Am J Clin Nutr* 1992;55:1033–44.
 139. Manson JE, Bassuk SS, Hu FB, Stampfer MJ, Colditz GA, Willett WC. Estimating the number of deaths due to obesity: can the divergent findings be reconciled? *J Womens Health* 2007;16:168–76.
 140. Aviv A, Valdes AM, Spector TD. Human telomere biology: pitfalls of moving from the laboratory to epidemiology. *Int J Epidemiol* 2006;35:1424–9.
 141. Lin J, Epel E, Cheon J, Kroenke C, Sinclair E, Bigos M, Wolkowitz O, Mellon S, Blackburn E. Analyses and comparisons of telomerase activity and telomere length in human T and B cells: insights for epidemiology of telomere maintenance. *J Immunol Methods* 2010;352:71–80.
 142. Freedman DS, Gates L, Flanders WD, Van Assendelft OW, Barboriak JJ, Joesoef MR, Byers T. Black/white differences in leukocyte subpopulations in men. *Int J Epidemiol* 1997;26:757–64.
 143. Camhi SM, Bray GA, Bouchard C, Greenway FL, Johnson WD, Newton RL, Ravussin E, Ryan DH, Smith SR, Katzmarzyk PT. The relationship of waist circumference and BMI to visceral, subcutaneous, and total body fat: sex and race differences. *Obesity (Silver Spring)* 2011;19:402–8.
 144. Fitzpatrick AL, Kronmal RA, Gardner JP, Psaty BM, Jenny NS, Tracy RP, Walston J, Kimura M, Aviv A. Leukocyte telomere length and cardiovascular disease in the cardiovascular health study. *Am J Epidemiol* 2007;165:14–21.
 145. Benetos A, Kark JD, Susser E, Kimura M, Sinnreich R, Chen W, Steenstrup T, Christensen K, Herbig U, von Bornemann Hjelmborg J, et al. Tracking and fixed ranking of leukocyte telomere length across the adult life course. *Aging Cell* 2013;12: 615–21.
 146. Broer L, Codd V, Nyholt DR, Deelen J, Mangino M, Willemsen G, Albrecht E, Amin N, Beekman M, de Geus EJ, et al. Meta-analysis of telomere length in 19,713 subjects reveals high heritability, stronger maternal inheritance and a paternal age effect. *Eur J Hum Genet* 2013;21:1163–8.
 147. Adaikalakoteswari A, Balasubramanyam M, Mohan V. Telomere shortening occurs in Asian Indian type 2 diabetic patients. *Diabet Med* 2005;22:1151–6.
 148. Raschenberger J, Kollerits B, Hammerer-Lercher A, Rantner B, Stadler M, Haun M, Klein-Weigel P, Fraedrich G, Kronenberg F. The association of relative telomere length with symptomatic peripheral arterial disease: results from the CAVASIC study. *Atherosclerosis* 2013;229:469–74.
 149. Heid IM, Wagner SA, Gohlke H, Iglseder B, Mueller JC, Cip P, Ladurner G, Reiter R, Stadlmayr A, Mackevics V, et al. Genetic architecture of the APM1 gene and its influence on adiponectin plasma levels and parameters of the metabolic syndrome in 1,727 healthy Caucasians. *Diabetes* 2006;55:375–84.