



Mapping the neuroanatomy of functional decline in Alzheimer's disease from basic to advanced activities of daily living

Andrea Slachevsky^{1,2,3,5,17} · Gonzalo Forno^{1,2,3} · Paulo Barraza⁴ · Eneida Mioshi⁶ · Carolina Delgado^{7,17} · Patricia Lillo^{1,16,17} · Fernando Henriquez^{1,2,3,4} · Eduardo Bravo⁸ · Mauricio Farias⁸ · Carlos Muñoz-Neira³ · Agustín Ibañez^{10,11,12,13,14} · Mario A. Parra^{9,10} · Michael Hornberger^{6,15}

Received: 27 July 2018 / Revised: 15 January 2019 / Accepted: 26 February 2019 / Published online: 4 April 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

Background Impairments in activities of daily living (ADL) are a criterion for Alzheimer's disease (AD) dementia. However, ADL gradually decline in AD, impacting on advanced (a-ADL, complex interpersonal or social functioning), instrumental (IADL, maintaining life in community), and finally basic functions (BADL, activities related to physiological and self-maintenance needs). Information and communication technologies (ICT) have become an increasingly important aspect of daily functioning. Yet, the links of ADL, ICT, and neuropathology of AD dementia are poorly understood. Such knowledge is critical as it can provide biomarker evidence of functional decline in AD.

Methods ADL were evaluated with the Technology–Activities of Daily Living Questionnaire (T-ADLQ) in 33 patients with AD and 30 controls. ADL were divided in BADL, IADL, and a-ADL. The three domain subscores were covaried against gray matter atrophy via voxel-based morphometry.

Results Our results showed that three domain subscores of ADL correlate with several brain structures, with a varying degree of overlap between them. BADL score correlated mostly with frontal atrophy, IADL with more widespread frontal, temporal and occipital atrophy and a-ADL with occipital and temporal atrophy. Finally, ICT subscale was associated with atrophy in the precuneus.

Conclusions The association between ADL domains and neurodegeneration in AD follows a traceable neuropathological pathway which involves different neural networks. This the first evidence of ADL phenotypes in AD characterised by specific patterns of functional decline and well-defined neuropathological changes. The identification of such phenotypes can yield functional biomarkers for dementias such as AD.

Keywords Alzheimer's disease · Functional impairment · Activities of daily living · Technology–activities of daily living questionnaire

Background

Alzheimer's disease (AD) is one of the most common form of age-related dementia, affecting more than 25 million people worldwide, with the number of new cases raising continuously, both in developed and developing countries [1, 2]. The diagnosis of dementia due to AD is based on the

presence of a gradual onset of cognitive impairment, mainly an episodic memory impairment with evidence of cognitive dysfunction in at least one other cognitive domain, whose severity has led to a significant functional decline in activities of daily living (AD), interfere with the ability to function at work or at usual activities [3].

The confirmation of the presence and severity of impairment in ADL is critical for the diagnosis of dementia [3]. Commonly, ADL have been divided in basic ADL (BADL) and instrumental ADL (IADL). BADL are defined as activities related to basic physiological and self-maintenance needs, including tasks such as eating, toileting or getting dressed.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00415-019-09260-w>) contains supplementary material, which is available to authorized users.

✉ Andrea Slachevsky
andrea.slachevsky@uhile.cl

Extended author information available on the last page of the article

IADL include activities, essential to maintain independent living and maintaining life in community, such as managing finances, shopping, handle medications or using the public transport [4, 5]. Recently, advanced ADL (a-ADL) has emerged as an additional important category in ADL [5, 6]. a-ADL are defined as more complex activities, not being essential to maintain an independent life, are considered voluntary [7] and include activities necessary for complex interpersonal or social functioning such as using household technology, going on holidays, practice hobbies, etc. [4, 6, 8]. a-ADL that require higher levels of cognitive, physical, and social functions, are very sensitive to subtle cognitive impairment and could contribute to early diagnosis of dementia [4]. Nonetheless, the definition of a-ADL and its division from IADL is complex and need to consider cultural variability, since ADL performance is influenced by cultural events [9–12]. Moreover, as culture evolves, any scale that is sensitive to early ADL deficits must also evolve to measure newly relevant activities.

In the last decades, information and communication technologies (ICT) have become an increasingly important aspect of daily functioning and the use of electronic devices are essential in different everyday life tasks, such as communication, work or recreational activities. The use of everyday technology may be of particular concern in people with dementia because most patients typically continue to live at home, in the same social context as before the disability and, as a result, they are expected to manage the everyday technology that is common in that context [13]. ICT could include either IADL or a-ADL depending on the complexity of the technology and sociocultural factors shaping technology use [6, 14].

Despite recent advances in the development of ADL scales, the relationship between those outcomes and structural brain changes in AD is poorly understood, especially considering the neural correlates of IADL or BADL. BADL dysfunction in AD was associated with atrophy in the temporal, cingulate, hippocampus, caudate, frontal, and parietal regions, whereas IADL dysfunction was linked to atrophy in the frontal, temporal, parietal, insula, and caudate regions [15]. Hippocampal and cortical gray matter volume loss was associated with rapid IADL decline in AD [16]. Parietal and temporal lobe atrophy at baseline predict further IADL impairment over time [17]. Complementary, PET studies have reported an association between greater rate of IADL impairment over time and middle frontal, orbitofrontal and posterior cingulate hypometabolism in AD [18].

However, to our knowledge, there is no study investigating the neural correlates of a-ADL. Nor is their evidence that such correlates differ from those reported for BADL and IADL. Moreover, no study has incorporated ICT as an important aspect of functional assessment. The aim of this study was to investigate the neural correlates of the global

score and subscores of the Technology–Activities of Daily Living Questionnaire (T-ADLQ) in patients with AD in comparison to healthy controls (HC). Specifically, we examined which brain areas were associated with a-ADL impairment in AD in comparison with BADL and IADL scores. In a first step, total T-ADLQ scores, BADL, IADL and a-ADL subscores were regressed against gray matter atrophy via voxel-based morphometry (VBM). In a second step, we performed an inclusive masking analysis to verify which areas of brain atrophy would overlap between BADL and a-ADL, and between IADL and a-ADL. Finally, we performed an exclusive masking analysis to verify areas of brain atrophy displaying no overlap between BADL, a-ADL, and IADL. We hypothesized that these three ADL domains would exhibit shared and segregated neuroanatomical substrates and that a-ADL would be associated to regions involved in more complex cognitive tasks.

Methods

Participants

A cohort comprising 63 participants was recruited for the study. This cohort was divided into two groups matched according to sex, age, and years of education: 33 subjects with a clinical diagnosis of AD and 30 healthy controls (HC). Patients were recruited from the Memory and Neuropsychiatric Clinic at Hospital del Salvador, and the Neurology and Neurosurgery Department at Hospital Clínico Universidad de Chile (HCUCH), both located in Santiago, Chile. HC were recruited from a variety of sources, including spouses or relatives of patients with dementia. The inclusion criteria considered Spanish-speaking participants older than 60 years of age. All participants required a reliable proxy who had known them for at least 5 years. Specifically, a proxy was someone who was able to provide information about ADL performance, behavioral changes, and patients' general medical history. The exclusion criteria included illiteracy, underlying neurological or psychiatric illness that could affect cognition (except AD), physical disability, and sensory disturbance that could interfere with the neuropsychological assessment. All AD patients met the NINCDS-ADRDA criteria for probable AD [3]. Diagnosis was made by consensus between senior neurologists (AS and CD) based on extensive clinical protocol, interviews with a reliable proxy, laboratory tests and global cognitive functioning. Briefly, AD patients displayed a history of significant episodic memory loss, within the context of preserved behavioral and personality, and scored above 0.5 on the Clinical Dementia Rating scale (CDR) [3]. HC did not report memory complaints, had a score of 0 on the CDR [3], and their cognitive performance was considered normal

according to local normative data for the Addenbrooke's Cognitive Examination—Revised Chilean Version (ACE-R-Ch) (> 76) [19]. Scores of the T-ADLQ were not considered to establish the diagnosis. Ethical approval for this study was obtained from the Ethical and Scientific Committees of the East Metropolitan Health Service and the HCUCH. All the participants, and their caregivers, provided informed consent in accordance with the Declaration of Helsinki.

Clinical and neuropsychological examination

All proxies and participants were interviewed separately to obtain the CDR scores. The T-ADLQ was completed by proxies as we have previously described [20]. Experienced clinical psychologists trained in the administration of our neuropsychological protocol and blinded to the diagnosis of each subject carried out the neuropsychological assessment. In addition to the MMSE [21], and the ACE-R-Ch [19] to assess global cognitive functioning, the neuropsychological protocol included The Boston Naming Test as an index of naming abilities. The Rey-Osterrieth Complex Figure Test was used to measure visuospatial constructional abilities [22]. Forward and backward digit-span tasks provided an index of working memory while the Word Free and Cued Selective Reminding Test (FCSRT) was used to assess episodic memory. The Frontal Assessment Battery (FAB), which is a screening test for executive dysfunction that assesses conceptualization, mental flexibility, motor programming, resistance to interference, inhibitory control, and environmental autonomy, was also applied [23]. Other tests of executive functions (EF) included the Modified Version of the Wisconsin Card Sorting Test (MCST) [24], which informs on cognitive flexibility, verbal fluency tests including both Phonemic Verbal Fluency test (i.e., words beginning with letters F, A, and S in 1 min) and Semantic Fluency test (i.e., animals in 1 min) and the Trail Making Test A and B [25, 26].

T-ADLQ

The T-ADLQ [20] consists of seven subscales: self-care, household care, employment and recreation, shopping and money, travel, communication and ICT. Each item is rated on a four-point scale. For each activity, a rating is provided for instances in which the patient may have never performed that activity in the past, stopped the activity prior to the onset of dementia, or for which the proxy did not have information [27]. The overall functional impairment (FI) was calculated for each domain as well as for the global questionnaire as follows: (sum of all ratings not rated ND/DK) / (3 × total number of items not rated ND/DK). The denominator represents the score that would have been obtained if the most severe level of impairment had been indicated for all items

rated not ND/DK [27]. This equation ensures that the functional impairment score was based on the actual functioning of the patients relative to their own premorbid functioning. Higher percentage scores indicate greater deterioration.

An expert panel (two neurologists, three psychologists, one occupational therapist) gathered the activities of the T-ADLQ in three domains (BADL—IADL—a-ADL). To ensure consistency of the division, each expert classified each activity independently and then a consensus was reached to harmonize the different classifications. The outcomes from the consensus classification are presented in Table 1.

Statistical analyses for demographic and neuropsychological data

The Statistical Package for the Social Sciences (SPSS) version 20 for Windows (IBM Corp., Armonk, NY, USA) was used to analyze the demographic and neuropsychological data. We obtained descriptive statistics for such data, used Chi-squared for the categorical variables, and performed two-tailed independent-sample *t* tests for the comparisons between AD and HC. Differences with a $p < 0.05$

Table 1 Division of the three domains of the Technology—Activities of Daily Living Questionnaire (T-ADLQ)

| Basic ADL | Instrumental ADL | Advanced ADL |
|---------------------------------|--|-----------------|
| Eating | Taking pills or medicine | Employment |
| Dressing | Handling cash | Recreation |
| Bathing | Managing finances | Organization |
| Elimination | Public transportation | Travel |
| Interest in personal appearance | Driving | Internet access |
| | Mobility around the neighborhood | Email access |
| | Traveling outside familiar environment | Computer use |
| | Preparing meals, cooking | |
| | Setting the table | |
| | Housekeeping | |
| | Home maintenance | |
| | Home repair | |
| | Laundry | |
| | Food shopping | |
| | Using the telephone | |
| | Talking | |
| | Understanding | |
| | Reading | |
| | Writing | |
| | Cell phone use | |
| | ATM use | |

The T-ADLQ scale is presented in supplementary files

were considered significant. Additionally, the effect sizes (Cohen's-*d* statistic) were calculated to determine the magnitude of the group differences. According to Cohen, effect sizes between 0.2 and 0.49 are considered small; those between 0.5 and 0.79, moderate; and those 0.8, large [28].

MRI acquisition

MRI acquisition was performed in two 1.5 T MRI scanners, a Philips Intera Nova Dual gradient system (45mT/m), and a Siemens Symphony Maestro Class (Erlangen, Germany) with 20 mT/m gradient system. High resolution anatomical scans were obtained using a T1-weighted three dimensional gradient recalled echo acquisition: 3D T1 fast field echo sequence on Philips scanner, and 3D T1 fast low angle shot on Siemens scanner, both with the same acquisition parameters (TE = 4.6 ms, TR = 25 ms; flip angle = 30°, field of view on frequency = 250 mm, 256 × 256 matrix, isotropic voxel size 1 × 1 × 1 mm).

VBM analysis

MRI data were analyzed with FSL-VBM, a (Voxel-based Morphometry) VBM analysis [29, 30] that is part of the FSL software package (<http://www.fmrib.ox.ac.uk/fsl/fslvb/index.html>) [31]. First, tissue segmentation was carried out using FMRIB's automatic segmentation tool (FAST) from brain-extracted images [32]. The resulting grey matter partial volume maps were then aligned to the Montreal Neurological Institute standard space (MNI152) using the non-linear registration approach using FNIRT [33] which uses a b-spline representation of the registration warp field [34]. A study-specific template was created, combining AD and HC images, to which the native gray matter images were re-registered nonlinearly. The registered partial volume maps were then modulated (to correct for local expansion or contraction), by dividing them by the Jacobian of the warp field. The modulated images were then smoothed with an isotropic Gaussian kernel with a sigma of 3 mm (FWHM: 8 mm).

The statistical analysis was performed via a voxel wise general linear model (GLM) to investigate gray matter intensity differences. Permutation-based nonparametric testing (with 5000 permutations per contrast) [35] was used to form clusters with the threshold-free cluster enhancement (TFCE) method [31]. The significance threshold was $p < 0.05$ and tests were corrected for multiple comparisons via Family-wise Error (FWE) correction across space, unless otherwise stated. For uncorrected results, a threshold of 100 contiguous voxels was used, at $p < 0.001$ to reduce the likelihood of significant clusters. Regions of significant atrophy were superimposed on the MNI standard brain, with maximum coordinates provided in MNI space. Areas of significant gray

matter loss were localized with reference to the Harvard-Oxford probabilistic cortical and subcortical atlas.

In a first step, differences in gray matter intensities between AD patients and HC were assessed. To control for a possible scanner site effect, we introduced scanner site as a nuisance covariate for the group contrasts. Next, correlations between gray matter atrophy and T-ADLQ total score and the scores of the three domains of the T-ADLQ, i.e., BADL, IADL and a-ADL subscores, were entered as covariates in the design matrix of the VBM analysis for AD patients combined with HC. This procedure improves the statistical power to detect brain-behavior relationships [36]. In a third step, we explored the overlap of brain atrophy between the BADL, IADL and a-ADL subscores performing an inclusive masking analysis. For statistical power, a covariate-only statistical model with a t-contrast was used, providing an index of association between brain atrophy and scores on the functional scales. The statistical maps generated from the contrast using BADL, IADL and a-ADL subscores as covariate, were scaled using a threshold of $p < 0.001$, following which, the scaled contrasts were multiplied to create an inclusive, or overlap, mask across groups. In a fourth step, we performed a contrast analysis between the three subscores BADL, IADL and a-ADL subscores of the T-ADLQ to study the existence of significant anatomical differences between the different domains. For the exclusive masks, the same procedure described above was adopted. However, the scaled images were subsequently subtracted from each other, to create an exclusive mask for each condition.

Result

Demographic and neuropsychological data

Demographic and neuropsychological scores are shown in Table 2. AD and HC groups did not differ in terms of sex, age, or education (all $p > 0.05$). In brief, AD patients exhibited scores significantly higher on assessments of severity of the disease (CDR) and lower on measures of global cognitive efficiency (ACE-R-Ch and MMSE) and episodic memory (FCSRT) relative to HC. Compared to the HC group, the AD group was impaired on the global scores of T-ADLQ [$F(1,67) = 70.981, p < 0.001$]; the three ADL domains and the ICT subscores (see Table 2) The details of the neuropsychological battery in HC and AD subjects are shown in Supplementary Table 1.

VBM: groups comparison analysis

Results are shown in Table 3 and Fig. 1. The AD group was contrasted with HC group to reveal patterns of brain atrophy. The AD group showed significant gray matter atrophy

Table 2 Demographic, global cognitive, and functional characteristics of AD and HC

| | AD | Control | <i>t</i> test/ χ^2 | Effect size (Cohen's <i>d</i>) | 95% CI | |
|--|---------------------|---------------------|-------------------------|------------------------------------|---------|---------|
| | | | | | Lower | Upper |
| <i>N</i> | 33 | 29 | | | | |
| Sex (m:f) | 13:20 | 9:20 | 0.492 | | | |
| Age (years) | 73.09 ± 6.96 | 72.03 ± 5.99 | 0.636 | 0.163 | −2.265 | 4.378 |
| Education (years) | 12.21 ± 4.45 | 12.59 ± 3.73 | −0.356 | −0.092 | −2.476 | 1.728 |
| MMSE | 20.79 ± 4.88 (30) | 28.07 ± 1.67 (30) | −7.648** | −1.996 | −9.185 | −5.377 |
| ACE-R | 62.24 ± 15.64 (100) | 92.55 ± 5.60 (100) | −9.887** | −2.580 | −36.441 | −24.177 |
| CDR | 1.66 ± 0.65 (3) | 0 ± 00 (3) | 13.648** | 3.611 | 1.413 | 1.899 |
| CDR-SB | 5.84 ± 2.82 (18) | 0 ± 00 (18) | 11.157** | 2.928 | 4.796 | 6.892 |
| CDR-AIG | 1.13 ± 0.83 (3) | 0 ± 00 (3) | 7.269** | 1.925 | 0.815 | 1.435 |
| T-ADLQ (%) Total | 38.00 ± 16.96 (100) | 8.41 ± 8.90 (100) | 8.425** | 2.184 | 22.562 | 36.611 |
| BADL domain subscore of the T-ADLQ (%) | 5.70 ± 11.56 (100) | 0.93 ± 2.94 (100) | 2.158* | 0.565 | 0.347 | 9.184 |
| IADL domain subscore of the T-ADLQ (%) | 43.73 ± 19.39 (100) | 7.79 ± 10.23 (100) | 8.939** | 2.318 | 27.893 | 43.975 |
| a-ADL domain of the T-ADLQ (%) | 52.33 ± 22.00 (100) | 19.76 ± 19.85 (100) | 6.087** | 1.554 | 21.869 | 43.280 |

Data are presented in mean ± standard deviation (total score)

AD Alzheimer's diseases, CDR clinical dementia rating, CDR-SB clinical dementia rating—sum of box, CDR-AIG clinical dementia rating—algorithm, MMSE mini-mental state examination, ACE-R Addenbrooke's cognitive examination revised, MoCA Montreal cognitive assessment, T-ADLQ technology of daily living questionnaire, BADL basic activities of daily life, IADL instrumental ADL, a-ADL advanced activities of daily life

* $p < 0.05$, ** $p < 0.001$

Table 3 VBM showing significant gray matter intensity decrease in AD in contrast HC uncorrected by scanner

| Regions | Hemisphere | MNI coordinates | | | Number of voxel |
|--|------------|-----------------|----------|----------|-----------------|
| | | <i>x</i> | <i>y</i> | <i>z</i> | |
| Hippocampus | Left | −24 | −34 | −10 | 1513 |
| Hippocampus | Right | 24 | −36 | −8 | 1155 |
| Precentral gyrus | Left | −38 | 0 | 28 | 122 |
| Precuneus cortex | Right | 16 | −64 | 32 | 119 |
| Inferior frontal gyrus, par opercularis/precentral gyrus | Right | 36 | 8 | 26 | 117 |
| Inferior temporal gyrus/temporal fusiform cortex, posterior division | Right | 44 | −14 | −30 | 111 |

All results corrected for multiple comparisons (family-wise error) at $p < 0.05$; only cluster with at least 100 contiguous voxels included

MNI Montreal Neurological Institute

in bilateral hippocampal brain regions, bilateral precentral gyrus, and a right lateralized atrophy in the precuneus cortex, inferior frontal gyrus (par opercularis), inferior temporal gyrus, and temporal fusiform cortex (posterior division) ($P_{\text{fwe corr}} < 0.05$). Similar results were obtained in the analysis covarying for scanner site (see Supplementary Table 2 and Supplementary Figure 1).

Correlations with T-ADLQ subscores

VBM correlations with T-ADLQ total score are presented in supplementary files (see Supplementary Table 3 and Supplementary Figure 2). In brief, T-ADLQ total score covaried

with bilateral atrophy in the parahippocampal gyrus (anterior division) and the inferior temporal gyrus (posterior division and temporo-occipital region), and a right lateralized atrophy in the lateral occipital cortex (inferior and superior division) ($p_{\text{unorr}} < 0.001$).

BADL, IADL and a-ADL subscores of the T-ADLQ were entered as covariate in the design matrix of the VBM analysis. Results are shown in Table 4 and Fig. 2. For the AD group, the score on the BADL domain covaried with atrophy in the left supplementary motor cortex and right frontal regions (orbital cortex and superior/middle frontal gyrus) ($p_{\text{unorr}} < 0.001$). The IADL subscore covaried with atrophy in several areas widely distributed, highlighting the left




Fig. 1 VBM showing significant gray matter intensity decrease in AD in contrast HC uncorrected by scanner. VBM analysis showing brain areas of decreased gray matter intensity in AD patients in comparison with controls (MNI coordinates $X = -38$; $Y = -36$; $Z = -8$). Colored

voxel show regions that were significant in the analysis with $p < 0.05$ corrected for multiple comparisons (family-wise error), with a cluster threshold of 100 contiguous voxels. Clusters are overlaid on the MNI standard brain


Table 4 VBM correlation with T-ADLQ subscores in AD in comparison with HC uncorrected by scanner

| Regions | Hemisphere | MNI coordinates | | | Number of voxel |
|--|------------|-----------------|-----------|------------|-----------------|
| | | x BADL | y IADL | z a-ADL | |
| BADL domain subscore | | | | | |
| Supplementary motor cortex | Left | -14 | 6 | 50 | 274 |
| Frontal orbital cortex | Right | 32 | 28 | 0 | 260 |
| Superior frontal gyrus/middle frontal gyrus | Right | 24 | 26 | 44 | 139 |
| IADL domain subscore | | | | | |
| Para cingulate gyrus | Left | -4 | 34 | 26 | 3363 |
| Temporal fusiform cortex, anterior division | Right | 32 | -2 | -40 | 1936 |
| Temporal fusiform cortex/parahippocampal gyrus, anterior division | Left | -28 | -10 | -38 | 1418 |
| Intracalcarine cortex | Right | 8 | -66 | 12 | 836 |
| Frontal operculum cortex/central opercular cortex | Right | 42 | 10 | 10 | 512 |
| Parahippocampal gyrus, posterior division/temporal fusiform cortex, posterior division | Left | -32 | -34 | -16 | 379 |
| Intracalcarine cortex | Left | -12 | -70 | 10 | 191 |
| Precentral gyrus | Left | -38 | 0 | 22 | 185 |
| Lateral occipital cortex, inferior division | Right | 34 | -78 | 10 | 146 |
| Lingual gyrus/occipital fusiform gyrus | Right | 20 | -74 | -6 | 107 |
| Superior parietal lobule/angular gyrus | Right | 32 | -50 | 42 | 101 |
| a-ADL domain subscore | | | | | |
| Lingual gyrus/intracalcarine cortex | Left | -14 | -62 | 4 | 1379 |
| Parahippocampal gyrus, anterior division | Left | -24 | -10 | -38 | 1169 |
| Parahippocampal gyrus, anterior division | Right | 30 | -14 | -28 | 732 |
| Paracingulate gyrus | Left | -12 | 28 | 32 | 324 |
| Superior frontal gyrus | Right | 12 | 10 | 58 | 175 |
| Superior frontal gyrus | Left | -18 | 8 | 46 | 131 |
| Paracingulate gyrus/superior frontal gyrus | Right | 12 | 30 | 40 | 107 |
| Paracingulate gyrus | Right | 4 | 26 | 40 | 106 |

BADL basic ADL, *IADL* instrumental ADL, *a-ADL* advanced ADL, *MNI* Montreal Neurological Institute

All results uncorrected at $p < 0.001$. Only cluster with at least 100 contiguous voxels included

Fig. 2 VBM correlations with T-ADLQ subscores in AD in comparison with HC uncorrected by scanner. VBM analysis showing brain areas in which gray matter intensity correlates significantly with T-ADLQ subscores respectively: **a** BADL (MNI coordinates $X=32; Y=6; Z=44$), **b** IADL (MNI coordinates $X=-4; Y=-70; Z=-16$) and **c** a-ADL subscores (MNI coordinates $X=30; Y=-62; Z=46$) in AD in comparison with HC. Colored voxel show regions that were significant in the analysis with $p < 0.001$ uncorrected. For all analysis, a cluster threshold of 100 contiguous voxels was used. Clusters are overlaid on the MNI standard brain



paracingulate gyrus, bilateral temporal fusiform cortex, left parahippocampal gyrus (anterior division), and right intracalcarine cortex ($p_{\text{unorr}} < 0.001$). Finally, the a-ADL score covaried mainly with atrophy in the left lingual gyrus, intracalcarine cortex, and bilateral parahippocampal gyrus (anterior division) ($p_{\text{unorr}} < 0.001$). We present the VBM results showing regions of significant gray matter intensity decrease that covary with the ICT subscale in supplementary material (see Supplementary Table 4 and Supplementary Figure 3).

Overlap analysis

We performed an inclusive masking analysis to verify which areas of brain atrophy overlap when accounting for functional impairment in AD as measured respectively by IADL domain and a-ADL domain, BADL domain, and a-ADL domain. Results show that the BADL domain overlap with neither the IADL nor the a-ADL domain subscores

of the T-ADLQ. We found an overlap between the IADL and a-ADL domain subscores of the T-ADLQ (see Table 5 and Fig. 3).

Contrast analysis

In a final step, we performed a contrast analysis between the BADL, IADL and a-ADL domain subscores of the T-ADLQ to study the existence of significant anatomical differences between the different domains. Results shows that a-ADL were exclusively associated with one cluster in the left lingual gyrus and the anterior division of the parahippocampal gyrus. The score of the IADL covaried exclusively with atrophy in several areas, mainly the left paracingulate and cingulate gyrus (anterior division), the right and left temporal fusiform cortex (anterior division) and the right middle frontal gyrus ($p_{\text{uncorr}} < 0.001$) (see Table 6 and Fig. 3).

Table 5 VBM overlap between IADL and a-ADL subscores in AD compared with HC uncorrected by scanner

| Regions | Hemisphere | MNI coordinates | | | Number of voxel |
|---|------------|-----------------|-----|-----|-----------------|
| | | x | y | z | |
| Regions of overlap | | | | | |
| Parahippocampal gyrus, anterior division | Right | 30 | -14 | -28 | 471 |
| Precuneus cortex | Right | 14 | -56 | 12 | 463 |
| Parahippocampal gyrus, anterior division | Left | -28 | -12 | -36 | 373 |
| Parahippocampal gyrus, posterior division | Left | -28 | -36 | -14 | 232 |
| Paracingulate gyrus | Left | -10 | 28 | 36 | 191 |
| Intracalcarine cortex | Left | -12 | -68 | 10 | 108 |

All results uncorrected at $p < 0.001$. Only cluster with at least 100 contiguous voxels included
MNI Montreal Neurological Institute




Fig. 3 VBM overlap and exclusive mask analysis for IADL and a-ADL in AD compared with HC. VBM analysis showing overlap brain regions between T-ADLQ IADL and a-ADL domain subscores (green), and regions that correlate exclusively with IADL (red) and a-ADL (blue) in AD in comparison with HC (MNI coordinates

$X=30; Y=-62; Z=40$). Colored voxel show regions that were significant in the analysis with $p < 0.001$ uncorrected. For all analysis, a cluster threshold of 100 contiguous voxels was used. Clusters are overlaid on the MNI standard brain

Table 6 VBM exclusive regions that correlate with IADL and a-ADL subscores in AD compared with HC

| Regions IADL domain subscores | Hemisphere | MNI coordinates | | | Number of voxel |
|--|------------|-----------------|-----|-----|-----------------|
| | | x | y | z | |
| Paracingulate gyrus/cingulate gyrus, anterior division | Left | -8 | 36 | 22 | 1930 |
| Temporal fusiform cortex, anterior division | Right | 30 | -2 | -42 | 1258 |
| Temporal fusiform cortex, anterior division | Left | -30 | 0 | -40 | 862 |
| Middle frontal gyrus | Right | 26 | 26 | 40 | 731 |
| Central operculum cortex | Right | 42 | 8 | 10 | 482 |
| Precuneus cortex | Right | 18 | -58 | 20 | 242 |
| Central operculum cortex | Left | -36 | 0 | 20 | 185 |
| Lateral occipital cortex, inferior division | Right | 36 | -82 | 8 | 146 |
| Occipital fusiform gyrus | Right | 26 | -66 | -8 | 107 |
| a-ADL domain subscores | | | | | |
| Lingual gyrus | Left | -16 | -62 | 2 | 646 |
| Parahippocampal gyrus, anterior division | Left | -20 | -10 | -40 | 340 |

All results uncorrected at $p < 0.001$. Only cluster with at least 100 contiguous voxels included
IADL instrumental ADL, *a-ADL* advanced ADL, *MNI* Montreal Neurological Institute

Discussion

To the best of our knowledge, this is the first study assessing the neural correlates of a-ADL and ICT in AD. Our results showed that T-ADLQ subscores correlate with several brain structures, with a varying degree of overlap when the three domains of activities of the questionnaire were considered. The BADL score correlated mostly with frontal atrophy, IADL with more widespread frontal, temporal and occipital atrophy and a-ADL with occipital and temporal atrophy. The inclusive masking analysis did not show areas that overlap between BADL, IADL and a-ADL. However, the bilateral parahippocampal region (PHR) and the precuneus cortex are implicated in both, the IADL and a-ADL. In the exclusive masking analysis, we found an association between IADL and the paracingulate gyrus and the temporal fusiform cortex, while a-ADL correlated more with lingual gyrus atrophy.

T-ADQL total score correlated with a large cluster centered in the temporal fusiform cortex (anterior division) and parahippocampal gyrus (anterior division). Concerning the association with the temporal cortex, our results are in line with studies in both AD and other neurological disorders. Medial temporal lobe atrophy has been associated with functional impairment in both subjects with stroke [37] and mild cognitive impairment (MCI) [38]. The association with the PHR is not surprising due to the involvement of the PHR in multiple cognitive process relevant to everyday functioning as visuospatial processing, episodic memory, and contextual associative processing [39]. Moreover, PHR atrophy has been reported as an early biomarker of AD [40], and hypometabolism in this region has been associated with greater decline in IADL performance [18]. Concerning BADL, AD patients assessed in our study were in mild to moderate stages of the disease and presented mild impairment in some BADL activities (average of 5% of impairment). BADL correlated with two small clusters in left supplementary motor cortex and right frontal regions (orbital cortex and superior/middle frontal gyrus). Although we acknowledge that these results should be interpreted with caution due to the characteristics of our samples, they are in agreement with several studies which have reported that motor abilities are crucial for BADL [41–43] and tend to decline as the disease progresses [44].

IADL correlated with the left paracingulate gyrus, bilateral temporal fusiform cortex, left parahippocampal gyrus (anterior division), and right intracalcarine cortex, among others. Our results are in agreement with previous studies on the neural correlates of IADL [15]. IADL impairment has been associated with inferior temporal and lateral parietal (supramarginal) atrophy [17], decreased gray matter volume in the medial frontal and temporo-parietal cortices

in early stages of AD [45], and white matter lesion [46]. The multiple brain areas related with IADL have been attributed to the complex nature of these activities and their increased demand [15].

a-ADL correlated with the left lingual gyrus, intracalcarine cortex, and bilateral parahippocampal gyrus (anterior division), areas that has been associated with higher-order cognitive functions [39], such as explicit memory [47]. As Braak and Braak described, the brain atrophy of AD patients progresses following a hierarchical model, with early atrophy in the hippocampus and parahippocampus gyrus [48], linked to memory and visuospatial impairments in the early stages of the disease, progressing to a generalized pattern of brain atrophy linked to a wide deterioration of cognitive domains [49].

In the overlap analysis, we found that the bilateral atrophy of the parahippocampal gyrus, the paracingulate gyrus, and the intracalcarine cortex are associated to both IADL and a-ADL impairments. Atrophy of these areas has been associated with impairment in episodic memory, language, praxis and visual perception. These symptoms appear in the early stages of AD [47, 50], corresponding to the typical progression of the disease, starting with impairment in complex cognitive domains, including IADL, to difficulties in BADL in the most advanced stages [51].

In the exclusive analysis, we found that left parahippocampal gyrus correlated exclusively with a-ADL. This area has been extensively associated with topographical learning and spatial navigation [52–54], and is crucial for normal adaptive behaviors in work, traveling, or during other activities that comprise a-ADL.

Finally, we found that the ICT subscale is associated with atrophy in the precuneus, area that has been related with visuospatial functioning, attentional shift, and processing speed [55, 56]. It has also been considered a higher-order area that is generally involved in controlling spatial aspects of motor behavior, and episodic memory retrieval [57]. Evidence shows that attentional and visuospatial abilities are necessary for internet searching [58]. Attentional engagement has also been described in motor-cognitive skills when working with touch-screen terminal [59]. Recent evidence suggests that hypometabolism in the precuneus may be a biomarker of potential progression to AD [60]. From this perspective, functional changes associated to this region may reflect the earliest manifestations of the impact of AD on ADL, suggesting that these novel functional assessment tools could be considered sensitivity tools to aid in the early diagnosis of AD. Longitudinal studies in AD and other dementias are mandatory to support this hypothesis.

Our division of the T-ADLQ items in BADL, IADL and a-ADL should be interpreted with caution as it was performed based on an expert panel in clinical diagnosis of dementia and research in functional assessment. Although

experts hold experience in cognitive neurology (AS and CD), neuropsychology (CMN, FH and GF) and occupational therapy (EM), some of such decisions can be influenced by sociocultural factors. For example, classification of computer use as an a-ADL and use of mobile phones as an IADL may not be representative of every sociocultural or generational context. The study here presented was carried out in 2015 when only 10% of the targeted elderly population reported use of internet, 80% of them had access to the internet via landline connectivity and only 20% via mobile phones [61, 62]. Moreover, elderly people in Chile, as in other countries such as Portugal, report that they mainly use mobile phones for basic functions such as answering and calling [63]. Also, in a recent paper by our group [64], we provide additional evidence on the validity of our proposed division of the T-ADLQ in the BADL, IADL and a-ADL categories. Nevertheless, further studies need to address the subdomains characterization of evolving T-ADLQ, which would need to be updated considering intra-country specificity and sociocultural factors [5, 65].

Some methodological issues warrant consideration. First, our neuroimaging results regarding the correlation between VBM and subscore domain did not survive conservative corrections for multiple comparisons and were therefore reported uncorrected at $p < 0.001$. However, we reduced the likelihood of false positive results, by applying cluster extent thresholds of 100 contiguous voxels in the analysis. Importantly, Monte Carlo simulations and experimental data demonstrate that cluster thresholding is an effective tool to reduce the probability of false positive findings without compromising the statistical power of the study [66]. Given our sample size, the application of stringent cluster extended thresholds, and our a priori assumptions, we are confident that our results do not represent false positive findings. However, it will be important to replicate these findings in larger patient cohorts using corrected neuroimaging approaches. Second, the diagnosis of AD was established on clinical grounds without any neuropathological confirmation for the diagnoses. Nevertheless, AD patients presented bilateral atrophy in the hippocampal region, a structural biomarker of AD. Moreover, clinical pathological studies suggested that NINCDS-ADRDA criteria are reliable for the diagnosis of AD. Finally, patients included in the study also fulfilled criteria for Alzheimer's clinical syndrome according to the latest NIA-AA Research Framework [67].

In conclusion, our study suggests that combining a domain specific approach to ADL with neuropathological data drawn from MRI, specific functional phenotypes can be identified. We have demonstrated that in the sample of AD investigated here such a phenotype is characterized by widespread atrophy of the prefrontal, temporal and occipital brain regions significantly associated with functional impairments that follow a gradient of deterioration in the

diseases continuum. Such functional phenotypes seemingly inform about such a continuum. Our data also suggest that ADL assessment via such an approach can be very sensitive to neurodegenerative processes, and that the association of types of functional decline and AD progression does follow a traceable neuropathological pathway that involves different neural network. Our results are consistent with the existence of a specific pattern of functional loss in the activities of daily living, namely functional phenotypes, beginning with impairment in the a-ADL, followed by losses in IADL and finally progressing to BADL [4].

Further studies need to address the contribution of white matter lesion to functional impairment [68]. Finally, generalization of our results to other neurodegenerative diseases should be made with cautious. Nevertheless, futures studies are needed to investigate if other neurodegenerative disease leads to functional phenotypes different to those seen in AD. Indeed, ADL dysfunction in FTD is associated with atrophy in network different to that reported in AD and the left superior frontal gyrus is the only region implicated in IADL dysfunction in both FTD and AD [15].

Acknowledgements Funding from CONICYT/FONDECYT/11404223 (to A.S, C.D, E.B, F.H, C.M., and P.B); FONDAF Program Grant 15150012 (to AS, PL, GF and AI); CONICYT/FONDECYT (Regular 1170010), CONICYT/FONDECYT regular 1160940 (to AS and PL), the INECO Foundation (to AI), the Global Brain Health Institute (GBHI-UCSF) (to AI), the Inter-American Development Bank (IADB) (to AI and AS), PIA-CONICYT Basal Funds for Centers of Excellence Project FB0003 (to A.S; PB and FH) and the Alzheimer's Society (Grant AS-SF-14-008) (to MAP).

Compliance with ethical standards

Conflicts of interest On behalf of all authors, the corresponding author states that there is no conflict of interest.

Ethical approval All the procedures conducted with the participants of this study were carried out according to the Declaration of Helsinki. Ethical approvals were provided by the Ethical and Scientific Committees of the East Metropolitan Health Service and Hospital Clínico Universidad de Chile from Santiago, Chile.

References

1. Ferri CP, Prince M, Brayne C, Brodaty H, Fratiglioni L, Ganguli M, Hall K, Hasegawa K, Hendrie H, Huang Y, Jorm A, Mathers C, Menezes PR, Rimmer E, Scazufca M (2005) Global prevalence of dementia: a Delphi consensus study. *Lancet* 366:2112–2117
2. Qiu C, Kivipelto M, von Strauss E (2009) Epidemiology of Alzheimer's disease: occurrence, determinants, and strategies toward intervention. *Dialogues Clin Neurosci* 11:111–128
3. McKhanna GM, Knopman DS, Chertkow H, Hyman BT, Jack CRJ, Kawas CH, Klunk WE, Koroshetz WJ, Manly JJ, Mayeux R, Mohs RC, Morris JC, Rossor MN, Scheltens P, Carrillo MC, Thies B, Weintraub S, Phelps CH (2011) The diagnosis of dementia due to Alzheimer's disease: recommendations from the National

- Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 7:263–269
4. De Vriendt P, Gorus E, Cornelis E, Bautmans I, Petrovic M, Mets T (2013) The Advanced activities of daily living: a tool allowing the evaluation of subtle functional decline in mild cognitive impairment. *J Nutr Health Aging* 17:64–71
 5. De Vriendt P, Gorus E, Cornelis E, Velghe A, Petrovic M, Mets T (2012) The process of decline in advanced activities of daily living: a qualitative explorative study in mild cognitive impairment. *Int Psychogeriatr* 24:974–986
 6. Dias EG, de Andrade FB, de Oliveira Duarte YA, Ferreira Santos JL, Lúcia Lebrão M (2015) Advanced activities of daily living and incidence of cognitive decline in the elderly: the SABE Study. *Cad Saúde Pública* 8:1–13
 7. Reuben DB, Laliberte L, Hiris J, Mor V (1990) A hierarchical exercise scale to measure function at the advanced activities of daily living (AADL) level. *J Am Geriatr Soc* 38:855–861
 8. De Vriendt P, Mets T, Petrovic M, Gorus E (2015) Discriminative power of the advanced activities of daily living (a-ADL) tool in the diagnosis of mild cognitive impairment in an older population. *Int Psychogeriatr* 27:1419–1427
 9. Femia EE, Zarit SH, Johansson B (1997) Predicting change in activities of daily living: a longitudinal study of the oldest old in Sweden. *J Gerontol Psychol Sci* 52B:294–302
 10. Goldberg TE, Koppel J, Keehlisen L, Christen E, Dreses-Werringloer U, Conejero-Goldberg C, Gordon ML, Davies P (2010) Performance-based measures of everyday function in mild cognitive impairment. *Am J Psychiatry* 167:845–853
 11. Reisberg B, Finkel S, Overall J, Schmidt-Gollas N, Kanowski S, Lehfeld H, Hulla F, Sclan S, Wilms H, Heininger I, Stemmler M, Poon L, Kluger A, Cooler C, Bergener M, Hugonot-Diener L, Robert P, Antipolis S, Erzigkeit H (2001) The Alzheimer's disease activities of daily living international scale (ADL-IS). *Int Psychogeriatr* 13:163–181
 12. Kalisch T, Richter J, Lenz M, Kattenstroth J-C, Kolankowska I, Tegenthoff M, Dinse HR (2011) Questionnaire-based evaluation of everyday competence in older adults. *Clin Interv Aging* 6:37–46
 13. Rosenberg L, Kottorp A, Winblad B, Nygard L (2009) Perceived difficulty in everyday technology use among older adults with or without cognitive deficits. *Scand J Occup Ther* 16:216–226
 14. Luborsky MR (1993) Sociocultural factors shaping technology usage: fulfilling the promise. *Technol Disabil* 2:71–78
 15. Mioshi E, Hodges JR, Hornberger M (2013) Neural correlates of activities of daily living in frontotemporal dementia. *J Geriatr Psychiatry Neurol* 26:51–57
 16. Cahn-Weiner D, Farias S, Julian L, Harvey DJ, Kramer JH, Reed BR, Mungas D, Wetzel M, Chui H (2007) Cognitive and neuroimaging predictors of instrumental activities of daily living. *J Int Neuropsychol Soc* 13:747–757
 17. Marshall GA, Lorus N, Locascio JJ, Hyman BT, Rentz DM, Johnson KA, Sperling RA (2014) Regional cortical thinning and cerebrospinal biomarkers predict worsening daily functioning across the Alzheimer's disease spectrum. *J Alzheimers Dis* 41:719–728
 18. Roy K, Pepin LC, Philiossaint M, Lorus N, Becker JA, Locascio JJ, Rentz DM, Sperling RA, Johnson KA, Marshall GA (2014) Regional fluorodeoxyglucose metabolism and instrumental activities of daily living across the Alzheimer's disease spectrum. *J Alzheimers Dis* 42:291–300
 19. Muñoz-Neira C, Henriquez F, Ihnen J, Sánchez-Cea M, Flores P, Slachevsky A (2012) Propiedades Psicométricas y Utilidad Diagnóstica del Addenbrooke's Cognitive Examination—Revised (ACE-R) en una muestra de ancianos chilenos. *Rev Med Chil* 140:1006–1013
 20. Muñoz-Neira C, López L, Riveros O, Núñez-Huasaf R, Flores J, Slachevsky P A (2012) The technology—activities of daily living questionnaire: a version with a technology-related subscale. *Dement Geriatr Cogn Disord* 33:361–371
 21. Folstein MF, Folstein SE, McHugh PR (1975) Mini-mental state: a practical method for grading the cognitive state of patients for clinician. *J Psychiatr Res* 12:189–198
 22. Rey A (1959) Test de copie d'une figure complexe. Centre de Psychologie Appliquée, Paris
 23. Dubois B, Slachevsky A, Litvan I, Pillon B (2000) The FAB: a frontal assessment battery at bedside. *Neurology* 55:1621–1626
 24. Nelson HE (1976) A modified card sorting test sensitive to frontal lobe defects. *Cortex* 12:313–324
 25. Reitan R (1958) Validity of the trail making test as an indication of organic brain damage. *Percept Mot Skills* 8:271–276
 26. Arango-Lasprilla JC, Rivera D, Aguayo A, Rodriguez W, Garza M, Saracho C, Rodriguez-Agudelo Y, Aliaga A, Weiler G, Luna M, Longoni M, Ocampo-Barba N, Galarza-del-Angel J, Panyavin I, Guerra A, Esenarro L, Garcia de la Cadena P, Martinez C, Perrin P (2015) Trail making test: normative data for the Latin American Spanish speaking adult population. *NeuroRehabilitation* 37:639–661
 27. Johnson N, Barion A, Rademaker A, Rehkemper G, Weintraub S (2004) The activities of daily living questionnaire: a validation study in patients with dementia. *Alzheimer Dis Assoc Disord* 18:223–230
 28. Cohen J (1988) Statistical power analysis for the behavioral sciences. L. Erlbaum Associates, Hillsdale
 29. Ashburner J, Friston KJ (2000) Voxel-based morphometry—the methods. *Neuroimage* 11:805–821
 30. Good CD, Johnsrude IS, Ashburner J, Henson RN, Friston KJ, Frackowiak RS (2001) A voxel-based morphometric study of ageing in 465 normal adult human brains. *Neuroimage* 14:21–36
 31. Smith SM, Jenkinson M, Woolrich MW, Beckmann CF, Behrens TE, Johansen-Berg H, Bannister PR, De Luca M, Drobnjak I, Flitney DE, Niazy RK, Saunders J, Vickers J, Zhang Y, De Stefano N, Brady JM, Matthews PM (2004) Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage* 23(Suppl 1):S208–S219
 32. Zhang Y, Brady M, Smith S (2001) Segmentation of brain MR images through a hidden Markov random field model and the expectation-maximization algorithm. *IEEE Trans Med Imaging* 20:45–57
 33. Andersson JLR, Jenkinson M, Smith S (2007) Non-linear optimisation. FMRIB technical report. In: FMRIB Analysis Group technical reports. FMRIB technical report
 34. Rueckert D, Sonoda LI, Hayes C, Hill DLG, Leach MO, Hawkes DJ (1999) Non-rigid registration using free-form deformations: application to breast MR images. *IEEE Trans Med Imaging* 18:712–721
 35. Nichols TE, Holmes AP (2002) Nonparametric permutation tests for functional neuroimaging: a primer with examples. *Hum Brain Mapp* 15:1–25
 36. Irish M, Piguet O, Hodges JR, Hornberger M (2014) Common and unique gray matter correlates of episodic memory dysfunction in frontotemporal dementia and Alzheimer's disease. *Hum Brain Mapp* 35:1422–1435
 37. Henon H, Pasquier F, Durieu I, Pruvo JP, Leys D (1998) Medial temporal lobe atrophy in stroke patients: relation to pre-existing dementia. *J Neurol Neurosurg Psychiatry* 65:641–647
 38. Yoshida D, Shimada H, Makizako H, Doi T, Ito K, Kato T, Shimokata H, Washimi Y, Endo H, Suzuki T (2012) The relationship between atrophy of the medial temporal area and daily activities in older adults with mild cognitive impairment. *Aging Clin Exp Res* 24:423–429

39. Aminoff EM, Kveraga K, Bar M (2013) The role of the parahippocampal cortex in cognition. *Trends Cogn Sci* 17:379–390
40. Echavarrri C, Aalten P, Uylings HB, Jacobs HI, Visser PJ, Gronenschild EH, Verhey FR, Burgmans S (2011) Atrophy in the parahippocampal gyrus as an early biomarker of Alzheimer's disease. *Brain Struct Funct* 215:265–271
41. Nijboer T, van de Port I, Schepers V, Post M, Visser-Meily A (2013) Predicting functional outcome after stroke: the influence of neglect on basic activities in daily living. *Front Hum Neurosci* 7:182
42. Mercier L, Audet T, Hébert R, Rochette A, Dubois MF (2001) Impact of motor, cognitive, and perceptual disorders on ability to perform activities of daily living after stroke. *Stroke* 32:2602–2608
43. Gialanella B, Santoro R, Ferlucchi C (2013) Predicting outcome after stroke: the role of basic activities of daily living predicting outcome after stroke. *Eur J Phys Rehabil Med* 49:629–637
44. Scarmeas N, Hadjigeorgioi GM, Papadimitriou A, Dubois B, Sarazin M, Brandt J, Albert M, Marder K, Bell K, Honig LS, Wegesin D, Stern Y (2004) Motor signs during the course of Alzheimer disease. *Neurology* 63:975–982
45. Vidoni ED, Honea RA, Burns JM (2010) Neural correlates of impaired functional independence in early Alzheimer's disease. *J Alzheimers Dis* 19:517–527
46. Baune BT, Schmidt WP, Roesler A, Berger K (2009) Functional consequences of subcortical white matter lesions and MRI-defined brain infarct in an elderly general population. *J Geriatr Psychiatry Neurol* 22:266–273
47. Golby A, Silverberg G, Race E, Gabrieli S, O'Shea J, Knierim K, Stebbins G, Gabrieli J (2005) Memory encoding in Alzheimer's disease: an fMRI study of explicit and implicit memory. *Brain* 128:773–787
48. Braak H, Braak E (1997) Diagnostic criteria for neuropathologic assessment of Alzheimer's disease. *Neurobiol Aging* 18: S85–S88
49. Almkvist O (1996) Neuropsychological features of early Alzheimer's disease: preclinical and clinical stages. *Acta Neurol Scand* 94:63–71
50. Crosson B, Sadek JR, Bobholz JA, Gokcay D, Mohr CM, Leonard CM, Maron L, Auerbach EJ, Browd SR, Freeman AJ, Briggs RW (1999) Activity in the paracingulate and cingulate sulci during word generation: an fMRI study of functional anatomy. *Cereb Cortex* 9:307–316
51. Desai AK, Grossberg GT, Sheth DN (2004) Activities of daily living in patients with dementia: clinical relevance, methods of assessment and effects of treatment. *CNS Drugs* 18:853–875
52. Maguire EA, Burgess N, Donnett JG, Frackowiak RS, Frith CD, O'Keefe J (1998) Knowing where and getting there: a human navigation network. *Science* 280:921–924
53. Aguirre GK, Detre JA, Alsop DC, D'Esposito M (1996) The parahippocampus subserves topographical learning in man. *Cereb Cortex* 6:823–829
54. Mellet E, Briscogne S, Tzourio-Mazoyer N, Ghaem O, Petit L, Zago L, Etard O, Berthoz A, Mazoyer B, Denis M (2000) Neural correlates of topographic mental exploration: the impact of route versus survey perspective learning. *NeuroImage* 12:588–600
55. Wenderoth N, Debaere F, Sunaert S, Swinnen SP (2005) The role of anterior cingulate cortex and precuneus in the coordination of motor behaviour. *Eur J Neurosci* 22:235–246
56. Karas G, Scheltens P, Rombouts S, van Schijndel R, Klein M, Jones B, van der Flier W, Vrenken H, Barkhof F (2007) Precuneus atrophy in early-onset Alzheimer's disease: a morphometric structural MRI study. *Neuroradiology* 49:967–976
57. Cavanna AE, Trimble MR (2006) The precuneus: a review of its functional anatomy and behavioural correlates. *Brain* 129:564–583
58. Small GW, Moody TD, Siddarth P, Bookheimer SY (2009) Your brain on Google: patterns of cerebral activation during internet searching. *Am J Geriatr Psychiatry* 17:116–126
59. Sagari A, Iso N, Moriuchi T, Ogahara K, Kitajima E, Tanaka K, Tabira T, Higashi T (2015) Changes in cerebral hemodynamics during complex motor learning by character entry into touchscreen terminals. *PLoS One* 10(10):e0140552
60. Counts SE, Ikonovic MD, Mercado N, Vega IE, Mufson EJ (2017) Biomarkers for the early detection and progression of Alzheimer's disease. *Neurotherapeutics* 14:35–53
61. Subsecretaría de Telecomunicaciones del Gobierno de Chile (2017) Estudio Octava Encuesta sobre Acceso, Usos y Usuarios de Internet en Chile. In: Subsecretaría de Telecomunicaciones (Subtel). Santiago, Chile
62. Subsecretaría de Telecomunicaciones de Chile (2018) IX Encuesta de Acceso y Usos de Internet—Informe Final. In: Subsecretaría de Telecomunicaciones, Subtel, Santiago, Chile
63. Neves BB, Amaro F (2012) Too old for technology? How the elderly of Lisbon use and perceive ICT. *J Commu Inform* 8(1):1–12
64. Delgado C, Vergara RC, Martinez M, Musa G, Henriquez F, Slachevsky A (2019) Neuropsychiatric symptoms in Alzheimer's disease are the main determinants of functional impairment in advanced everyday activities. *J Alzheimers Dis* 67:381–392
65. Fillenbaum GG, Chandra V, Ganguli M, Pandav R, Gilby JE, Seaberg EC, Belle S, Baker C, Echement DA, Nath LM (1999) Development of an activities of daily living scale to screen for dementia in an illiterate rural older population in India. *Age Ageing* 28:161–168
66. Forman SD, Cohen JD, Fitzgerald M, Eddy WF, Mintun MA, Noll DC (1995) Improved assessment of significant activation in functional magnetic resonance imaging (fMRI): use of a cluster-size threshold. *Magn Reson Med* 33:636–647
67. Jack CR Jr, Bennett DA, Blennow K, Carrillo MC, Dunn B, Haeberlein SB, Holtzman DM, Jagust W, Jessen F, Karlawish J, Liu E, Molinuevo JL, Montine T, Phelps C, Rankin KP, Rowe CC, Scheltens P, Siemers E, Snyder HM, Sperling R, Contributors (2018) NIA-AA research framework: toward a biological definition of Alzheimer's disease. *Alzheimers Dement* 14:535–562
68. Farias ST, Mungas D, Reed B, Haan MN, Jagust WJ (2004) Everyday functioning in relation to cognitive functioning and neuroimaging in community-dwelling Hispanic and non-Hispanic older adults. *J Int Neuropsychol Soc* 10:342–354

Affiliations

Andrea Slachevsky^{1,2,3,5,17} · **Gonzalo Forno**^{1,2,3} · **Paulo Barraza**⁴ · **Eneida Mioshi**⁶ · **Carolina Delgado**^{7,17} · **Patricia Lillo**^{1,16,17} · **Fernando Henriquez**^{1,2,3,4} · **Eduardo Bravo**⁸ · **Mauricio Farias**⁸ · **Carlos Muñoz-Neira**³ · **Agustin Ibañez**^{10,11,12,13,14} · **Mario A. Parra**^{9,10} · **Michael Hornberger**^{6,15}

¹ Geroscience Center for Brain Health and Metabolism (GERO), Faculty of Medicine, University of Chile, Avenida Salvador 486, Providencia, Santiago, Chile

² Neuropsychology and Clinical Neuroscience Laboratory (LANNEC), Physiopathology Department - ICBM, Neurociencia and East Neuroscience Departments, Faculty of Medicine, University of Chile, Avenida Salvador 486, Providencia, Santiago, Chile

³ Memory and Neuropsychiatric Clinic (CMYN) Neurology Department, Hospital del Salvador and Faculty of Medicine, University of Chile, Av. Salvador 364, Providencia, Santiago, Chile

⁴ Center for Advanced Research in Education (CIAE), University of Chile, 8330014 Santiago, Chile

⁵ Servicio de Neurología, Departamento de Medicina, Clínica Alemana-Universidad del Desarrollo, Santiago, Chile

⁶ School of Health Sciences, University of East Anglia, Norwich, UK

⁷ Neurology and Neurosurgery Department, Clinical Hospital, University of Chile, Santiago, Chile

⁸ Neuroradiologic Department, Institute of Neurosurgery Asenjo, Santiago, Chile

⁹ Psychology Department, School of Psychological Sciences & Health, University of Strathclyde, Glasgow, UK

¹⁰ Universidad Autonoma del Caribe, Barranquilla, Colombia

¹¹ Institute of Cognitive and Translational Neuroscience (INCYT), INECO Foundation, Favaloro University, Buenos Aires, Argentina

¹² National Scientific and Technical Research Council (CONICET), Av. Rivadavia 1917, C1033AAJ Buenos Aires, Argentina

¹³ Center for Social and Cognitive Neuroscience (CSCN), School of Psychology, Universidad Adolfo Ibañez, Diagonal Las Torres 2640, Santiago, Chile

¹⁴ Centre of Excellence in Cognition and its Disorders, Australian Research Council (ACR), Sydney, Australia

¹⁵ Norfolk and Suffolk Foundation Trust, Norwich, UK

¹⁶ Departamento de Neurociencia y Neurología Sur, Facultad de Medicina, Universidad de Chile, Santiago, Chile

¹⁷ Departamento de Neurociencia, Facultad de Medicina, Universidad de Chile, Santiago, Chile