

Safety Monitoring Boards at the UTSW site (Beverley Adams-Huet, MS, Sherwood Brown, MD, Kevin C. Kelly, PharmD, and Robert F. Reilly, MD) and the Central VA (Clinical Science Research and Development, Central Data Monitoring Committee, Hines CSPCC); psychiatry consultants (Benji Kurian, MD [UTSW] and Collin Vas, MBBS [Dallas VA]); nephrology faculty (Peter Van Buren, MD, MSc [UTSW]); and UTSW nephrology residents and fellows (Masoud Afshar, MD, Lei Chen, MD, Michael Concepcion, MD, Vishal Jaikaransingh, MD, Naseem Sunnoqrot, MD, Venkata Yalamanchili, MBBS). We also acknowledge the UTSW research personnel who were compensated for their role: Anuoluwapo Adelodun, MBBS, MPH, Patricia Alvarez, MSW, Mieshia Beamon, MS, Susamei Khamphong, BA, Ammar Nassri, MD, Michael Phan, PharmD, MBA, MHSM, David Rezaei, PharmD (Dallas VA site), Staci Schwartz, BA, Francisco Sanchez, BS, Kyle West, MS, and the UTSW Nephrology Clinical and Translational Research Center.

Disclaimer: The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIDDK, the National Institutes of Health, or the Department of Veterans Affairs.

Prior Presentation: Parts of these results were presented in abstract form at the American Society of Nephrology Kidney Week meeting; October 25, 2018; San Diego, CA.

Peer Review: Received December 4, 2018. Evaluated by 2 external peer reviewers, with direct editorial input from a Statistics/Methods Editor, an Associate Editor, and the Editor-in-Chief. Accepted in revised form September 10, 2019.

Publication Information: © 2019 by the National Kidney Foundation, Inc. Published online January 23, 2020 with doi [10.1053/j.ajkd.2019.09.007](https://doi.org/10.1053/j.ajkd.2019.09.007)

References

- Hedayati SS, Minhajuddin AT, Toto RD, Morris DW, Rush AJ. Prevalence of major depressive episode in CKD. *Am J Kidney Dis.* 2009;54(3):424-432.
- Hedayati SS, Minhajuddin AT, Afshar M, Toto RD, Trivedi MH, Rush AJ. Association between major depressive episodes in patients with chronic kidney disease and initiation of dialysis, hospitalization, or death. *JAMA.* 2010;303(19):1946-1953.
- Palmer SC, Vecchio M, Craig JC, et al. Association between depression and death in people with CKD: a meta-analysis of cohort studies. *Am J Kidney Dis.* 2013;62(3):493-505.
- Hedayati SS, Gregg LP, Carmody T, et al. Effect of sertraline on depressive symptoms in patients with chronic kidney disease without dialysis dependence: the CAST randomized clinical trial. *JAMA.* 2017;318(19):1876-1890.
- Leonard BE. Inflammation and depression: a causal or coincidental link to the pathophysiology? *Acta Neuropsychiatr.* 2018;30(1):1-16.
- Rutledge T, Reis VA, Linke SE, Greenberg BH, Mills PJ. Depression in heart failure a meta-analytic review of prevalence, intervention effects, and associations with clinical outcomes. *J Am Coll Cardiol.* 2006;48(8):1527-1537.
- Akchurin OM, Kaskel F. Update on inflammation in chronic kidney disease. *Blood Purif.* 2015;39(1-3):84-92.
- Tynan RJ, Weidenhofer J, Hinwood M, Cairns MJ, Day TA, Walker FR. A comparative examination of the anti-inflammatory effects of SSRI and SNRI antidepressants on LPS stimulated microglia. *Brain Behav Immun.* 2012;26(3):469-479.
- Wiedlocha M, Marcinowicz P, Krupa R, et al. Effect of antidepressant treatment on peripheral inflammation markers - a meta-analysis. *Prog Neuropsychopharmacol Biol Psychiatry.* 2018;80(pt C):217-226.
- Strawbridge R, Arnone D, Danese A, Papadopoulos A, Herane Vives A, Cleare AJ. Inflammation and clinical response to treatment in depression: a meta-analysis. *Eur Neuro-psychopharmacol.* 2015;25(10):1532-1543.

Maintenance Peritoneal Dialysis in Children With Autosomal Recessive Polycystic Kidney Disease: A Comparative Cohort Study of the International Pediatric Peritoneal Dialysis Network Registry



To the Editor:

ARPKD is a rare disorder but an important cause of early-onset pediatric kidney failure.¹⁻³ PD has been recommended as the method of choice for initiating dialysis in infants and small children,⁴⁻⁶ but there are concerns regarding the feasibility of maintenance PD in ARPKD patients with their massively enlarged kidneys.^{2,7} Some centers perform uni- or bilateral nephrectomies of native ARPKD kidneys.⁸

To systematically evaluate maintenance PD characteristics, residual kidney function, and patient and PD technique survival in pediatric ARPKD patients, we compared data from the International Pediatric Peritoneal Dialysis Network (IPPN) registry from children with ARPKD (n = 79), CNS (n = 79), and CAKUT (n = 158; [Tables 1 and S1](#)). Groups were matched for age and time on dialysis ([Fig S1](#)). PD modalities included CAPD and APD (~80% of patients). Patients with highly individualized regimens were excluded from the analysis of PD prescription. Differences in PD prescription over observation time were examined by linear mixed regression models ([Item S1](#); [Table S2](#)).

Overall peritoneal fill volumes of all PD modalities combined were ~15% lower in ARPKD patients (704 ± 24 [SEM] vs 841 ± 24 and 822 ± 20 mL/m²/d for CNS and CAKUT, respectively). This tendency was confirmed in models accounting for PD modalities ([Fig 1A](#); [Table S3](#)) or oligoanuria ([Fig S2](#); [Table S4](#)). Among CAPD patients, those with ARPKD received significantly more cycles ([Fig 1A](#)). Among APD patients, number of cycles and time spent on cycle were similar in all groups. Due to lower fill volumes, ARPKD patients on APD achieved lower PD fluid turnover than CNS and CAKUT patients ([Fig 1A](#)). Lower PD fluid glucose concentrations were prescribed in ARPKD patients on APD, resulting in significantly lower peritoneal glucose exposure. Despite this, ARPKD patients on APD achieved similar ultrafiltration volumes and thus had a higher ultrafiltration per glucose exposure ratio ([Fig 1A](#)). It is tempting to speculate that portal hypertension may add nonosmotic ultrafiltration in ARPKD patients.

At the baseline visit, 35 ARPKD, 39 CNS, and 23 CAKUT patients were already anuric ([Table 1](#)). Urine output at baseline was highest in CAKUT patients and lowest in CNS

Table 1. Patient and PD Characteristics at Enrollment, Incidence of PD Complications

	ARPKD	CNS	CAKUT	P for ARPKD vs	
				CNS	CAKUT
Characteristics					
No. of patients	79	79	158		
Age at BL visit, y	2.42 [0.84-6.77]	2.40 [0.78-5.56]	2.42 [0.80-6.76]	0.8	0.9
Age at start of current PD, y	1.28 [0.08-4.45]	1.53 [0.49-4.44]	1.27 [0.08-4.31]	0.1	0.9
Age at start of first KRT, ^a y	0.58 [0.05-3.46]	1.50 [0.49-4.16]	1.08 [0.07-4.16]	0.02	0.3
PD duration at BL visit, mo	4.0 [0.9-16.2]	3.4 [1.0-12.5]	3.8 [1.1-14.8]	0.7	0.9
F/U time, mo	13.2 [5.6-25.3]	8.4 [3.3-17.8]	12.7 [4.7-24.1]	0.1	0.9
Male sex	37 (47%)	43 (54%)	124 (78.5%)	0.3	<0.001
Anthropometric data at BL visit					
Height, SDS	-2.69 ± 1.48	-2.38 ± 1.76	-2.86 ± 1.74	0.2	0.4
Body mass index, SDS	0.34 ± 1.64	-0.04 ± 1.55	-0.03 ± 1.41	0.1	0.08
Urine output at BL visit ^b					
Urine output, mL/m ² /d	145 [0-1,071]	37 [0-586]	848 [378-1,516]	0.07	<0.001
Pts with oligoanuria	35 (49%)	39 (56%)	23 (16.7%)	0.4	<0.001
PD modality at BL visit					
				0.3	0.9
CAPD	17 (22%)	15 (19%)	32 (20.3%)		
APD (NIPD)	40 (51%)	32 (41%)	80 (50.6%)		
APD (CCPD)	21 (27%)	29 (37%)	43 (27.2%)		
Other	1 (1%)	3 (4%)	3 (1.9%)		
PD fluids at BL visit					
				0.8	0.1
Acidic lactate	40 (51%)	38 (48%)	96 (60.8%)		
pH-neutral fluid	39 (49%)	41 (52%)	62 (39.2%)		
Complications During F/U^c					
	n = 37	n = 38	n = 75		
Exit-site and tunnel infections					
Rate per patient-y of F/U ^d	0.06	0.15	0.23		
Pts with ≥1 infection	2 (5%)	5 (13%)	15 (20%)	0.2	0.04
Pts with >1 infection	1 (3%)	2 (5%)	5 (7%)	0.6	0.4
Peritonitis episodes					
Rate per patient-y of F/U ^d	0.48	0.66	0.56		
Pts with ≥1 episode	12 (32%)	11 (29%)	25 (33%)	0.7	0.9
Pts with >1 episode	4 (11%)	8 (21%)	14 (19%)	0.2	0.3
Access revisions					
Rate per patient-y of F/U ^d	0.27	0.09	0.23		
Pts with ≥1 revision	8 (22%)	3 (8%)	18 (24%)	0.09	0.8
Pts with >1 revision	4 (11%)	1 (3%)	3 (4%)	0.2	0.2

Note: Unless otherwise indicated, data are count (percent), median [interquartile range], or mean ± SD. P values are based on χ^2 , Mann-Whitney U, or t tests. Abbreviations: APD, automated peritoneal dialysis; BL, baseline; CAPD, continuous ambulatory peritoneal dialysis; CAKUT, congenital anomalies of the kidneys and urinary tract; CCPD, continuous cycling peritoneal dialysis (APD with daytime dwell[s]); CNS, congenital nephrotic syndrome; F/U, follow-up; KRT, kidney replacement therapy; NIPD, nocturnal intermittent peritoneal dialysis (APD without daytime dwell); PD, peritoneal dialysis; SD, standard deviation; SDS, standard deviation score.

^aAge at earliest initiation of KRT and includes KRT before current PD; ARPKD, n = 78; CAKUT, n = 154.

^bAvailable urine data at BL: ARPKD, n = 72; CNS, n = 70; CAKUT, n = 138; oligoanuria defined as urine output < 100 mL/m²/d.

^cOnly includes pts followed up from start of PD, pts with first visit >3 mo after PD starting date excluded.

^dPatient-years of F/U defined as sum of F/U years per pt; total patient-years of F/U: ARPKD, 48; CNS, 46; CAKUT, 95.

patients, and the latter showed more frequent and earlier onset of oligoanuria over the course of PD (Fig S3). Patients with and without oligoanuria differed in key PD parameters, though mostly independent of the renal diagnosis (Fig S2; Table S4).

In incident maintenance PD patients (Table S5), exit-site and tunnel infections were reported less frequently in ARPKD than CAKUT patients (0.06 vs 0.23 episodes/year), whereas peritonitis frequency did not differ (Table 1). The need for PD access revisions in ARPKD patients was higher than in CNS patients but similar to CAKUT patients (Table 1). There were no significant differences in

hospitalizations (Table S6) or anthropometric or metabolic parameters (Table S7).

Death on dialysis was observed in 13 ARPKD, 13 CNS, and 8 CAKUT patients (Table S8) with cumulative survival rates after 4 years on PD of 78% in ARPKD, 73% in CNS, and 95% in CAKUT (Fig S4). Patient age and the CAKUT diagnosis were inversely correlated with mortality (Fig S4; Table S9). While mortality was generally higher in patients who commenced PD in the first year of life, survival in this age group did not differ between ARPKD and CNS (Table S10).

PD technique survival rate was ~80% after 4 years, without differences between disease groups (Fig 1B;




Figure 1. (A) PD prescription in patients. (B) Kaplan-Meier PD technique survival curves. (A) Data shown as model parameter estimates (mean and 95% confidence interval) and are based on linear mixed regression models (LMMs) for the entire observation time with an interaction term between diagnosis (ARPKD, CNS, CAKUT) and PD modality (CAPD, APD) (Total N: ARPKD, 72; CNS, 72; CAKUT, 144; Tables S2, S3); P values for CAPD are based on LMM with ARPKD/CAPD as reference group; P values for APD are based on LMM with ARPKD/APD as reference group. UF/GLU-ratio: ratio between ultrafiltration volume (L/m²/d) and glucose exposure (g/kg/d). (B) PD technique failure defined as switch to hemodialysis, death, or termination due to PD complications (infectious or noninfectious).

Table S11). Younger patients had higher risk for PD technique failure (Table S11).

Our study has some limitations. Because IPPN focuses on PD-related information, genotypes, hepatic involvement, gastrostomy tube insertion, vesicostomies, or nephrectomies were not systematically documented. Such data are currently collected in disease-specific cohort studies.^{9,10} Because the median age at baseline was 2.4 years, we may have missed early-onset disease-specific aspects. Furthermore, reporting bias cannot be excluded due to the voluntary nature of the registry. Potential regional practice specificities cannot be sufficiently addressed.

In summary, maintenance PD can be performed successfully in children with ARPKD, with good patient survival and comparable technique outcomes as observed in other early-onset kidney diseases. Minor adaptations of PD prescription are usually required, probably to comply with the large kidney size. Remarkably, higher ultrafiltration per glucose ratios are achieved in children with ARPKD on APD, possibly related to portal hypertension.

Abdelaziz Akarkach, MD, Kathrin Burgmaier, MD, Anja Sander, MSc, Nakysa Hooman, MD, Lale Sever, MD, Francisco Cano, MD, Pedro Zambrano, MD, Ilay Bilge, MD, Joseph T. Flynn, MD, Onder Yavascan, MD, Patricia G. Vallés, MD, PhD, Reyner Loza Munarriz, MD, Hireen P. Patel, MD, Erkin Serdaroglu, MD, Vera H. Koch, MD, Angela del Carmen Suarez, MD, Monica Galanti, MD, Claudia Gonzalez Celedon, MD, Anabella Rébora, MD, Jameela A. Kari, MD, FRCP, Cynthia J. Wong, MD, Ewa Elenberg, MD, Luisa F. Rojas, MD, Bradley A. Warady, MD, Max C. Liebaw, MD, Franz Schaefer, MD, on behalf of the IPPN Registry

Argentina (PGV); Cayetano Heredia Hospital, Lima, Peru (RLM); Department of Pediatrics, Ohio State University College of Medicine and Nationwide Children's Hospital, Columbus, OH (HPP); Department of Pediatric Nephrology, Dr. Behçet Uz Children's Hospital, Izmir, Turkey (ES); Pediatric Nephrology Unit, Instituto da Criança Hospital das Clinicas, University of Sao Paulo Medical School, Sao Paulo, Brazil (VHK); Department of Nephrology, Hospital de Niños Sor María Ludovica La Plata, Buenos Aires, Argentina (AdCS); Pediatric Nephrology, Roberto del Río Hospital (MG); Hospital Sótero del Río, Santiago, Chile (CGC); Pediatric Dialysis Unit, Senniad, Hospital Evangelico, Montevideo, Uruguay (AR); Pediatric Nephrology Center of Excellence and Pediatric Department, Faculty of Medicine, King Abdulaziz University, Jeddah, Kingdom of Saudi Arabia (JAK); Division of Nephrology, Department of Pediatrics, Stanford University School of Medicine, Palo Alto, CA (CJW); Texas Children's Hospital, Houston, TX (EE); Baxter Servicio al Cliente Colombia, Medellin-Antioquia, Colombia (LFR); Division of Nephrology, Children's Mercy Kansas City, University of Missouri-Kansas City School of Medicine, Kansas City, MO (BAW); Center for Molecular Medicine Cologne, University of Cologne, Faculty of Medicine and University Hospital Cologne, Cologne, Germany (MCL); and Division of Pediatric Nephrology, Center for Pediatrics and Adolescent Medicine, University Hospital of Heidelberg, Heidelberg, Germany (FS).

Address for Correspondence: Max C. Liebaw, MD, Department of Pediatrics and Center for Molecular Medicine, University Hospital of Cologne, Kerpener Str. 62, 50937 Cologne, Germany (e-mail: max.liebaw@uk-koeln.de) or Franz Schaefer, MD, Division of Pediatric Nephrology, Center for Pediatrics and Adolescent Medicine, Heidelberg University Medical Center, Im Neuenheimer Feld 430, 69120 Heidelberg, Germany (e-mail: franz.schaefer@med.uni.heidelberg.de)

Authors' Contributions: Research idea and study design: AA*, KB*, AS, BAW, MCL, FS; data acquisition: NH, LS, FC, PZ, IB, JTF, OY, PGV, RLM, HPP, ES, VHK, ACS, MG, CGC, AR, JAK, CJW, EE, LFR, BAW, FS; data analysis/interpretation: AA, KB, AS, MCL, FS; statistical analysis: AA, KB, AS, MCL, FS; study supervision: BAW, MCL, FS; joint senior authors: MCL and FS. Each author contributed important intellectual content during manuscript drafting or revision. Each author agrees to be personally accountable for the individual's own contributions and to ensure that questions pertaining to the accuracy or integrity of any portion of the work, even one in which the author was not directly involved, are appropriately investigated and resolved, including with documentation in the literature if appropriate. *Contributed equally to this work.

Support: The authors gratefully acknowledge the support by the International Society for Peritoneal Dialysis, Baxter Health Care, Fresenius Medical Care, Germany, and Ipsen. MCL and KB were supported by grants of the Medical Faculty of University of Cologne and the Marga and Walter Boll-Foundation. MCL and FS are supported by the German Federal Ministry of Research and Education (BMBF grant 01GM1515). FS was supported by the European Rare Kidney Disease Reference Network (ERKNet), and several authors of this work are ERKNet members. The funders had no role in the design and conduct of the study; collection, management, analysis, or interpretation of the data; or preparation, review, or approval of the manuscript.

Financial Disclosure: MCL has received honoraria for scientific lectures from Pfizer, and, representing the University Hospital of Cologne, has served on an advisory board for Otsuka. The other authors declare that they have no other relevant financial interests.

Peer Review: Received April 19, 2019. Evaluated by 2 external peer reviewers, with direct editorial input from a Statistics/Methods Editor, an Associate Editor, and the Editor-in-Chief. Accepted in revised form October 6, 2019.

Supplementary Material

Supplementary File (PDF)

Figure S1-S4, Items S1-S2, Tables S1-S11.

Article Information

IPPN Registry: A list of the registry's principal investigators is provided in [Item S2](#).

Authors' Affiliations: University of Cologne, Faculty of Medicine and University Hospital Cologne, Department of Pediatrics, Cologne, Germany (AA, KB, MCL); Institute of Medical Biometry and Informatics, University of Heidelberg, Heidelberg, Germany (AS); Ali-Asghar Clinical Research Development Center, Iran University of Medical Sciences, Tehran, Iran (NH); Pediatric Nephrology, Cerrahpaşa Faculty of Medicine, Istanbul University-Cerrahpaşa, Istanbul, Turkey (LS); Division of Pediatrics, Luis Calvo Mackenna Children's Hospital, Faculty of Medicine, University of Chile (FC); Hospital Dr Exequiel Gonzalez Cortes, Santiago, Chile (PZ); Department of Pediatric Nephrology, Istanbul University Medical Faculty, Istanbul, Turkey (IB); Department of Pediatrics, University of Washington School of Medicine (JTF); Division of Nephrology, Seattle Children's Hospital, Seattle, WA (JTF); Department of Pediatric Nephrology, Tepecik Teaching and Research Hospital, Izmir, Turkey (OY); Hospital Pediatrico Humberto Notti, Mendoza,

Publication Information: © 2019 by the National Kidney Foundation, Inc. Published online January 23, 2020 with doi 10.1053/j.ajkd.2019.10.009

References

1. Guay-Woodford LM, Desmond RA. Autosomal recessive polycystic kidney disease: the clinical experience in North America. *Pediatrics*. 2003;111(5, pt 1):1072-1080.
2. Hoyer PF. Clinical manifestations of autosomal recessive polycystic kidney disease. *Curr Opin Pediatr*. 2015;27(2):186-192.
3. Harris PC, Torres VE. Polycystic kidney disease. *Annu Rev Med*. 2009;60:321-337.
4. Zurowska AM, Fischbach M, Watson AR, Edefonti A, Stefanidis CJ; European Paediatric Dialysis Working Group. Clinical practice recommendations for the care of infants with stage 5 chronic kidney disease (CKD5). *Pediatr Nephrol*. 2013;28(9):1739-1748.
5. Guay-Woodford LM, Bissler JJ, Braun MC, et al. Consensus expert recommendations for the diagnosis and management of autosomal recessive polycystic kidney disease: report of an international conference. *J Pediatr*. 2014;165(3):611-617.
6. Gimpel C, Avni FE, Bergmann C, et al. Perinatal diagnosis, management, and follow-up of cystic renal diseases: a clinical practice recommendation with systematic literature reviews. *JAMA Pediatr*. 2018;172(1):74-86.
7. Beil S, Drube J, Gluer S, Lehner F, Ehrich JHH, Pape L. End-stage renal disease due to ARPKD in the first months of life: transplantation or dialysis?—two case reports. *Pediatr Transplant*. 2010;14(6):E75-E78.
8. Mallett TM, O'Hagan E, McKeever KG. Early bilateral nephrectomy in infantile autosomal recessive polycystic kidney disease. *BMJ Case Rep*. 2015;2015:bcr2015211106.
9. Ebner K, Feldkoetter M, Ariceta G, et al. Rationale, design and objectives of ARegPKD, a European ARPKD registry study. *BMC Nephrol*. 2015;16:22.
10. Alzarka B, Morizono H, Bollman JW, Kim D, Guay-Woodford LM. Design and implementation of the Hepatorenal Fibrocystic Disease Core Center Clinical Database: a centralized resource for characterizing autosomal recessive polycystic kidney disease and other hepatorenal fibrocystic diseases. *Front Pediatr*. 2017;5:80.