

Inflammation-associated suppression of metabolic gene networks in acute and chronic liver disease

Por: [Campos, G](#) (Campos, Gisela)^[1]; [Schmidt-Heck, W](#) (Schmidt-Heck, Wolfgang)^[2]; [De Smedt, J](#) (De Smedt, Jonathan)^[3]; [Widera, A](#) (Widera, Agata)^[1]; [Ghallab, A](#) (Ghallab, Ahmed)^[4]; [Puetter, L](#) (Puetter, Larissa)^[1]; [González, D](#) (Gonzalez, Daniela)^[1]; [Edlund, K](#) (Edlund, Karolina)^[1]; [Cadenas, C](#) (Cadenas, Cristina)^[1]; [Marchan, R](#) (Marchan, Rosemarie)^[1] ...[Más](#)
[Ver número de ResearchID y ORCID de Web of Science](#)

ARCHIVES OF TOXICOLOGY

Volumen: 94

Número: 1

Páginas: 205-217

DOI: 10.1007/s00204-019-02630-3

Fecha de publicación: JAN 2020

Tipo de documento: Article

[Ver impacto de la revista](#)

Abstract

Inflammation has been recognized as essential for restorative regeneration. Here, we analyzed the sequential processes during onset of liver injury and subsequent regeneration based on time-resolved transcriptional regulatory networks (TRNs) to understand the relationship between inflammation, mature organ function, and regeneration. Genome-wide expression and TRN analysis were performed time dependently in mouse liver after acute injury by CCl₄ (2 h, 8 h, 1, 2, 4, 6, 8, 16 days), as well as lipopolysaccharide (LPS, 24 h) and compared to publicly available data after tunicamycin exposure (mouse, 6 h), hepatocellular carcinoma (HCC, mouse), and human chronic liver disease (non-alcoholic fatty liver, HBV infection and HCC). Spatiotemporal investigation differentiated lobular zones for signaling and transcription factor expression. Acute CCl₄ intoxication induced expression of gene clusters enriched for inflammation and stress signaling that peaked between 2 and 24 h, accompanied by a decrease of mature liver functions, particularly metabolic genes. Metabolism decreased not only in pericentral hepatocytes that underwent CCl₄-induced necrosis, but extended to the surviving periportal hepatocytes. Proliferation and tissue restorative TRNs occurred only later reaching a maximum at 48 h. The same upstream regulators (e.g. inhibited RXR function) were implicated in increased inflammation and suppressed metabolism. The concomitant inflammation/metabolism TRN occurred similarly after acute LPS and tunicamycin challenges, in chronic mouse models and also in human liver diseases. Downregulation of metabolic genes occurs concomitantly to induce inflammation-associated genes as an early response and appears to be initiated by similar upstream regulators in acute and chronic liver diseases in humans and mice. In the acute setting, proliferation and restorative regeneration associated TRNs peak only later when metabolism is already suppressed.

Palabras clave

Palabras clave de autor: [Liver injury](#); [Regeneration](#); [Transcriptomics](#); [Gene networks](#)

KeyWords Plus:[NUCLEAR FACTOR 4-ALPHA](#); [EXPRESSION PROFILES](#); [MOUSE-LIVER](#); [HEPATOCYTE](#); [REGENERATION](#); [AUTOPHAGY](#); [CELLS](#); [PATHOPHYSIOLOGY](#); [HEPA TOTOXICITY](#); [IRE1-ALPHA](#)

Información del autor

Dirección para petición de copias: Hengstler, JG; Godoy, P (autor para petición de copias)

- + Tech Univ Dortmund, IfADo Leibniz Res Ctr Working Environm & Human Fa, Ardeystr 67, D-44139 Dortmund, Germany.

Direcciones:

- + [1] Tech Univ Dortmund, IfADo Leibniz Res Ctr Working Environm & Human Fa, Ardeystr 67, D-44139 Dortmund, Germany
- + [2] Nat Product Res & Infect Biol eV, Leibniz Inst, Hans Knoll Inst, Jena, Germany
- + [3] Katholieke Univ Leuven, Stem Cell Inst, Leuven, Belgium
- + [4] South Valley Univ, Fac Vet Med, Dept Forens & Vet Toxicol, Qena, Egypt
- + [5] Univ Chile, Fac Med, Biomed Neurosci Inst, Santiago, Chile
- + [6] Ctr Gerosci Brain Hlth & Metab GERO, Santiago, Chile
- + [7] Buck Inst Res Aging, Novato, CA 94945 USA
- + [8] Harvard Sch Publ Hlth, Dept Immunol & Infect Dis, Boston, MA USA
- + [9] Univ Cologne, Inst Neurophysiol, Med Fac, Cologne, Germany
- + [10] Univ Tübingen, Dept Expt & Clin Pharmacol & Toxicol, Tübingen, Germany
- + [11] Fed Inst Risk Assessment, Dept Food Safety, Max Dohrn Str 8-10, D-10589 Berlin, Germany
- + [12] Univ Regensburg Hosp, Ctr Liver Cell Res, Dept Pediat & Juvenile Med, Regensburg, Germany
- + [13] Thomas Jefferson Univ, Dept Pathol Anat & Cell Biol, Philadelphia, PA 19107 USA
- + [14] Vertex Pharmaceut, 3215 Merryfield Row, San Diego, CA 92121 USA
- + [15] Eli Lilly & Co, Lilly Corp Ctr, Lilly Res Labs, Indianapolis, IN 46285 USA
- + [16] Univ Edinburgh, MRC Ctr Regenerat Med, Edinburgh E16 4UU, Midlothian, Scotland

Direcciones de correo electrónico:hengstler@ifado.de; godoy@ifado.de

Financiación

Entidad financiadora Mostrar más información	Número de concesión
Comision Nacional de Investigacion Cientifica y Tecnologica (CONICYT) CONICYT FONDECYT	1140549

Helmholtz Virtuelles Institut Multifunktionale Biomaterialien für die Medizin (DE)	0313854
FP7 Health	266838
Nottingham City Primary Care Trust (GB)	267038

Editorial

SPRINGER HEIDELBERG, TIERGARTENSTRASSE 17, D-69121 HEIDELBERG, GERMANY

Información de la revista

- **Impact Factor:** [Journal Citation Reports](#)

Categorías / Clasificación

Áreas de investigación:Toxicology

Categorías de Web of Science:Toxicology

Información del documento

Idioma:English

Número de acceso: WOS:000511435900014

ID de PubMed: 31919559

ISSN: 0340-5761

eISSN: 1432-0738