

Visual Dermatology: Eruptive Vulvar Syringoma

Journal of Cutaneous Medicine and Surgery
2020, Vol. 24(2) 189–189
© The Author(s) 2019
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1203475419888868
jcms.sagepub.com

Daniel Morgado-Carrasco¹ and Pablo Vargas-Mora² 



Syringomas are benign adnexal tumors which mainly occur on the periorbital area of young women, although can be observed on the neck, chest, extremities, and genitalia. In some cases, syringomas can be eruptive or even generalized, and multiple syringomas can be observed in familial syringomatosis. Syringomas in the context of tumor syndromes have been rarely reported. Vulvar syringomas have been scarcely described. The majority of patients present with pruritic skin-colored papules on the labia majora.¹

ORCID iD

Pablo Vargas-Mora  <https://orcid.org/0000-0002-9388-2940>

Reference

1. Huang Y-H, Chuang Y-H, Kuo T-tong, Yang L-C, Hong H-S, Kuo T. Vulvar syringoma: a clinicopathologic and immunohistologic study of 18 patients and results of treatment. *J Am Acad Dermatol.* 2003;48(5):735-739.

¹Department of Dermatology, Hospital Clínic de Barcelona, Universitat de Barcelona, Spain ²Department of Dermatology, Faculty of Medicine, Universidad de Chile, Santiago, Chile. **Corresponding Author:** Daniel Morgado-Carrasco, Department of Dermatology, Hospital Clínic de Barcelona, Universitat de Barcelona, Calle Villarroel 170, Barcelona 08036, Spain; Email: morgadodaniel8@gmail.com. The authors have no conflicts of interest regarding this article, and have received no funding for this work.