



The 13th European Nutrition Conference, FENS 2019, was held at the Dublin Convention Centre, 15–18 October 2019

Ultra-processed food consumption and breast cancer risk

Inge Huybrechts¹, Isabelle Romieu¹, Neha Kandpur², Katerina Katsikari¹,
Gabriela Torres-Mejia³, Gloria I. Sanchez⁴, Maria Luisa Garmendia⁵, Carolina Porras⁶,
Carine Biessy¹, Marc J. Gunter¹, Carlos Augusto Monteiro² and Sabina Rinaldi¹

¹International Agency for Research on Cancer, Lyon, France,

²Universidade de São Paulo, Sao Paulo, Brazil,

³National Institute of Public Health, Cuernavaca, Morelos, Mexico,

⁴University of Antioquia, Medellín, Colombia,

⁵Institute of Nutrition and Food Technology, Universidad de Chile, Santiago de Chile, Chile and

⁶Agencia Costarricense de Investigaciones Biomédicas-Fundación INCIENSA (ACIB-FUNIN), San José, Costa Rica

Abstract

Objectives: The consumption of ultra-processed foods (UPF) has been linked to an increased risk of cancer in western populations, particularly for breast cancer (BC). Although the consumption of UPF is increasing rapidly in Latin American (LA) populations too, there has been no evaluation of its association with BC among LA women, where 20% of the BC cases present at ages younger than 45 years. We therefore evaluated the role of UPF intake on BC risk in young women participating in the Latin American PRECAMA study.

Methods: The PRECAMA study is an ongoing study coordinated by the International Agency for Research on Cancer (IARC) and involves scientific teams in four LA countries (Chile, Colombia, Costa Rica and Mexico); 406 incident BC cases and population based controls aged 20–45 years are recruited. Trained nurses administered a lifestyle and food frequency questionnaires (FFQ), conducted anthropometric measurements and collected biological samples. Tumour receptor status are determined in a centralized laboratory. UPF consumption was estimated by applying the NOVA classification to the dietary consumption data obtained via the FFQs. The association of the consumption of UPF and BC was determined using conditional logistic regression analysis adjusting for potential confounding factors.

Results: The median age at recruitment was 40 years. The median percentage of calories from UPF was 24% (10th to 90th percentile: 9 to 43%). UPF intake was positively associated with BC risk (OR for tertiles 2 and 3 in comparison with tertile 1 were 1.84 (95% CI 1.21–2.81) and 1.69 (95% CI 0.99–2.89) respectively). Among the UPF most strongly associated with BC, we identified the intake of industrial bread, packaged sweet and savoury snacks, breakfast cereals, cakes and desserts, and ready-eat/fast food. Comparing the 3rd with the 1st tertile of energy intake from the UPF group “fast food” the OR for BC risk was 1.93 (95% CI 1.23–3.04).

Discussion: Our results show positive associations between the consumption of UPF and BC risk in young women in LA. Therefore, global actions to decrease UPF intakes are urgently needed to address the cancer burden. In addition, further studies are needed to disentangle mechanisms relating UPF intake and carcinogenic processes in the breast. The European Prospective Investigation into Cancer and Nutrition (EPIC) study, a large-scale cohort including ~500,000 participants and 18,814 BC cases will be used to unravel the mechanistic pathways underlying these positive associations between the consumption of UPF and BC risk.

Conflict of Interest

There is no conflict of interest