

PAPER • OPEN ACCESS

Perspective on topological states of non-Hermitian lattices

To cite this article: Luis E F Foa Torres 2019 *J. Phys. Mater.* **3** 014002

View the [article online](#) for updates and enhancements.

Recent citations

- [Emergent PT symmetry in a double-quantum-dot circuit QED setup](#)
Archak Purkayastha *et al*
- [Floquet topological phase in a generalized PT -symmetric lattice](#)
Elizabeth Noelle Blose
- [Correspondence between Winding Numbers and Skin Modes in Non-Hermitian Systems](#)
Kai Zhang *et al*

INTERNATIONAL OPEN ACCESS WEEK
OCTOBER 19-26, 2020

ALL ECS ARTICLES. ALL FREE. ALL WEEK.
www.ecsdl.org

**NOW
AVAILABLE**



PAPER

Perspective on topological states of non-Hermitian lattices

OPEN ACCESS

RECEIVED
20 May 2019REVISED
11 August 2019ACCEPTED FOR PUBLICATION
2 September 2019PUBLISHED
8 November 2019

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Luis E F Foa Torres

Departamento de Física, Facultad de Ciencias Físicas y Matemáticas, Universidad de Chile, Santiago, Chile

E-mail: luis.foatorres@uchile.cl

Keywords: topological states, non-Hermitian lattices, PT-symmetry, Floquet systems

Abstract

The search of topological states in non-Hermitian systems has gained a strong momentum over the last two years climbing to the level of an emergent research front. In this perspective we give an overview with a focus on connecting this topic to others like Floquet systems. Furthermore, using a simple scattering picture we discuss an interpretation of concepts like the Hamiltonian's defectiveness, i.e. the lack of a full basis of eigenstates, crucial in many discussions of topological phases of non-Hermitian Hamiltonians.

1. Introduction

More than one hundred years after Edwin Hall discovered the Hall effect, the finding of the integer quantum Hall effect [1] and its topological origin [2] was followed by a sequence of breakthroughs [3–5] that sprouted into the discovery of topological insulators in two [6] and three dimensions [7]. Unlike conventional phases of matter, the new topological phases [1–7] cannot be described by a local order parameter [8–10]. Now, the frontiers in this field keep expanding at a rapid pace in ever more interesting directions, like the search for gapless but topological phases such as Weyl semimetals [11, 12] and topological states in Floquet systems [13–15].

One of the latest research fronts in this area is the search for topological states of non-Hermitian lattices [16–22]. The interest in non-Hermitian Hamiltonians was originally focused in \mathcal{PT} -symmetric Hamiltonians [23] as a generalization of quantum mechanics where the Hermiticity constraint could be removed while keeping a real spectra. Today, this has shifted to non-Hermitian Hamiltonians regarded as an effective description of, for example, open quantum systems [24, 25], where the finite lifetime introduced by electron–electron or electron–phonon interactions [26–28], or disorder [29], is modeled through a non-Hermitian term, or in the physics of lasing [30–34]. An additional source of momentum in this field comes from the study of systems where the quantum mechanical description is used after mapping to a Schrödinger-like equation, as in systems with gain and loss (as found in optics and photonics [35–38]), surface Maxwell waves [39], and topoelectrical circuits [40, 41].

Topological effects are intrinsic to features unique to non-Hermitian Hamiltonians such as the topological structure of exceptional points, points in parameter space where the eigenvalues and eigenvectors coalesce (the non-Hermitian counterpart of Hermitian degeneracies), and have been studied for a few decades [42–46]. The focus of the present new wave of interest is, however, on a different aspect: given a non-Hermitian lattice, a system where a *motif* is periodically repeated in space, what is the phenomenology and how do we classify the topological nature of the resulting states? Here, by topological one means that there is underlying bulk–boundary correspondence relating a bulk topological invariant to the existence (or absence) of states localized at a boundary (between topologically different phases).

The search of topological states of non-Hermitian lattices has become one of the most exciting emergent fronts at the crossroads of condensed matter physics, optics and photonics, acoustics and quantum physics. Together with Floquet systems, non-Hermitian lattices represent our first strong steps on the land of non-equilibrium, dynamical, topological phases. The departure from the Hermitian paradigm brings many surprises, like the fact that because of the extreme sensitivity to boundary conditions, a pristine non-Hermitian lattice may become devoid of extended states [47, 48], an effect termed the non-Hermitian skin effect [49, 50].

On the other hand, there is an intense activity in the search for a consistent classification [51–55], the definition of the topological invariants [56–58], and the search for a bulk-boundary correspondence [47, 48, 59, 60].

Here we provide an overview of this fascinating topic. We start by discussing how to define a gap for a complex spectrum. Later on, we discuss the information encoded in the richer spectrum of non-Hermitian systems and how to make sense of the non-Hermitian terms. In section 5, we discuss non-Hermitian degeneracies (exceptional points) and later on in section 6, by resorting to a scattering picture, we put forward an interpretation of the Hamiltonian defectiveness (the lack of a full basis of eigenvectors). In section 7 we give a flavor of the many paths proposed for building a bulk-boundary correspondence. Finally, in section 8, we present the connections with Floquet systems.

2. Defining a gap for a complex spectrum

Unlike Hermitian Hamiltonians, non-Hermitian systems have eigenvalues with a real and an imaginary part [61]. Thus, since complex numbers do not have an order relation, we cannot recur to the usual (Hermitian) definition of a gap [50].

To fix ideas let us consider a non-Hermitian lattice with Bloch-type eigenstates and eigenenergies $E_n(\mathbf{k})$, where $\hbar\mathbf{k}$ is the quasimomentum and n is the band index. There are many possible extensions of the gap concept. [51] defines a band (say n) as *separable* if $E_n(\mathbf{k}) \neq E_m(\mathbf{k})$ for all $m \neq n$ and all \mathbf{k} ; *isolated* if $E_n(\mathbf{k}) \neq E_m(\mathbf{k}')$ for all $m \neq n$ and all \mathbf{k}, \mathbf{k}' ; *inseparable* when for some quasimomentum the (complex) energy becomes degenerate with that of another band.

Alternatively one can generalize the concept of a bandgap as *the prohibition of touching a base energy* [52], also called the *point gap*. This base energy is most typically set to zero but it can be complex in the general case. In the Hermitian case we can continuously deform the spectrum in such a way that each energy band is mapped to a point along the (real) energy axis, all without touching a base energy set inside the gap (and therefore closing it). In the non-Hermitian case, when we sweep over the Brillouin zone the bands describe loops in the complex plane. If the base point lies inside the loop, the loop cannot be deformed onto a point without touching the base.

Thus, the prohibition of touching a base energy [52] is one physically reasonable generalization of the gap concept. Having a spectrum which describes a loop in a two-dimensional space allows us to define a topological invariant as the winding number of that loop, without any reference to the eigenstates¹. This winding number appears in many proposals for classifying non-Hermitian lattices including that by Gong *et al* [52], and Shen *et al* [51] (where it is dubbed *vorticity*). But the reader might have realized that this opens up the possibility of having a non-trivial topological invariant even in a system with a *single* band! Indeed, a single loop (and hence a single band) enclosing the base energy point gives a non-vanishing winding number. This contrasts with the Hermitian case, where at least two bands are required to have a non-zero Chern number². Here, having a different invariant allows for the rules to change.


Finally, one can also define the *line gap*: if the energy bands do not cross a line in the complex energy plane, the system is said to have a line gap [54, 60]. This concept accommodates the idea of a secluded region in the complex energy plane between two bands. The point and line gap definitions are complementary in the sense that one might be more relevant than the other in a particular situation. The existence of these two types of gaps is also a consequence of moving from a one-dimensional energy line for Hermitian systems to a plane for the case with a complex spectrum: in the Hermitian case, the secluded regions of energy can only be zero-dimensional (e.g. a segment of the line as in figure 1(a)), while in the non-Hermitian one can either be zero-dimensional (point gap, as in figure 1(b)) or one-dimensional (line gap, as in figure 1(c)).

3. The intriguing imaginary part

Up till now we have seen how taking the energy spectrum to the complex plane brings some surprises. But what does it mean to have an imaginary part of the eigenenergies and what information does it encode? The very first point is that non-Hermitian systems can be regarded as non-equilibrium systems. From a dynamical viewpoint, the states may even be unstable as the time-dependent Schrödinger equation is non-unitary. For long enough times, only the state(s) with the largest imaginary part of the eigenenergy survives.

¹ Note that this is in stark contrast with the topological invariants in Hermitian systems, which depend only on the eigenstates and not the eigenenergies.

² Recall that the sum of the Berry curvature over all available bands is zero at each point in parameter space.



Non-Hermiticity however, does not necessarily lead to a complex spectrum [23, 50]. If a Hamiltonian has \mathcal{PT} symmetry³ and that symmetry is preserved by the eigenstates then the corresponding eigenvalues are real [23]. But even in that case, the vanishing imaginary part stems from a dynamical situation where gains and losses are compensated.

In spite of the problems brought by non-Hermiticity, the imaginary part of the spectrum brings opportunities to clearly define things that prove difficult within standard quantum mechanics such as, for example, time scales. Being not a dynamical variable but rather a parameter, time does not have a corresponding operator in quantum mechanics, thereby complicating the determination of time scales such as the tunneling time [62]. Many authors (see [24, 25] and references therein) have pointed out that from a non-Hermitian perspective, the imaginary part of the energies provide precisely those evasive time scales, almost for free.

4. Making sense of gains, losses and asymmetric couplings

The conventional Hermitian quantum mechanics is so strongly built in our thinking that to many of us it might seem devoid of physical meaning to speak about non-Hermitian Hamiltonians. However, non-Hermitian Hamiltonians arise as a natural effective description of different systems. This includes photonic [35, 36, 38] and acoustic systems where gains and losses are naturally present, and also open quantum systems. In the latter case, non-Hermiticity arises when tracing out over the infinite degrees of freedom of what we consider as the *environment* acting on the *sample* of interest. Although defining the line that separates what belongs to the sample and what remains in the environment is arbitrary, the response obtained, for example through Green function techniques, is exact and independent of where we decided to draw this line.

The reader at this point might be puzzled about the fact that the effective Hamiltonians typically encountered in electronic systems contain only losses, because of the eventual leak into the environment, but no gains. How to make sense of gains when all the physical properties seem to be well described by the retarded self-energy corrections? The answer lies in the boundary conditions. The information concerning the boundary conditions in a scattering problem, e.g. the particle is coming from the left or right, is absent in the effective Hamiltonian containing the usual retarded self-energy corrections. This information has to be supplemented somehow if we want to compute the scattering state projected on the sample. One possible path is the Lippmann–Schwinger equation, another is akin to flipping the sign of the imaginary part of the self-energy correction to the signal where the source is; the latter provides a non-Hermitian Hamiltonian with losses *and* gains.

In most models studied in the context of topological states, gains and losses might be distributed over the system, thus allowing for a lattice structure, or they might have some symmetry with respect to an interface, thereby forming a boundary between two regions [63–65].

Besides gains and losses, a non-Hermitian Hamiltonian can also have asymmetric couplings between states, i.e. $\langle i|\mathcal{H}|j\rangle \neq \langle j|\mathcal{H}|i\rangle^*$. These sort of asymmetric or non-reciprocal hoppings appear in many models

³ \mathcal{H} is said to be \mathcal{PT} -symmetric if the combined action of the parity \mathcal{P} and time-reversal \mathcal{T} operators leave it invariant, $\mathcal{H} = (\mathcal{PT})^* \mathcal{H} (\mathcal{PT})$.

[56, 66, 67], and though they can be more difficult to assimilate in condensed matter systems (unless one thinks of them as an effective model or just a mapping between a model with gain and losses and another one with non-reciprocal hoppings) they are common in photonics.

5. Exceptional points and defectiveness

Having a non-Hermitian Hamiltonian $\mathcal{H} = \mathcal{H}_0 + \lambda\mathcal{H}'$ does not lead by itself to new physics, for example, if the non-Hermitian term \mathcal{H}' commutes with the Hermitian \mathcal{H}_0 , the eigenstates do not change. One of the key features unique to non-Hermitian Hamiltonians is defectiveness. This is, the eigenvectors of a non-Hermitian matrix can have an incomplete set of eigenvectors. Take for example the simplest Jordan block, a 2×2 matrix with all elements equal to 1 except for the lower left element which is set to zero. It has the eigenvalue $\lambda = 1$ with degeneracy two but has a single eigenvector. This matrix is said to be *defective*, a situation that is truly unique to non-Hermitian Hamiltonians.

Defectiveness typically emerges at the so-called exceptional points. Let us think of a generic situation with a Hamiltonian of the form $\mathcal{H} = \mathcal{H}_0 + \lambda\mathcal{H}_1$, where λ is a parameter controlling the strength of the second term. Assuming Hermitian \mathcal{H}_0 and \mathcal{H}_1 , the levels repel each other and avoid crossing when λ changes [68], as long as λ is real. But when λ is allowed to adopt complex values, the levels can *coalesce*, i.e. they can merge while rendering the matrix defective. This coalescence appears as a square root singularity of the eigenenergies as a function of the parameter λ and is called an exceptional point [69] (for a recent review on these points in photonics see [46]). The importance of these points was emphasized by Berry [70, 71]. Because of the complex shape of the energy Riemann surfaces in the vicinity of an exceptional point (see figure 2), which also lead to a breakdown of the adiabatic theorem, there are many counter-intuitive properties such as a strong dependence on the parameter λ close to those points, or the phenomenon dubbed chiral state conversion [72–74]⁴. This has fueled fascinating experiments with microwaves [42–45, 74–76], optical waveguides [77], nuclear magnetic resonance [78], and also in optomechanical systems [79]. Exceptional points lead to intriguing phenomena such as unidirectional invisibility [80], single-mode lasers [81, 82], or enhanced sensitivity in optics [83–85]. Many of these phenomena can be understood in terms of an environment-mediated interaction [25, 78, 86–89].

Exceptional points may also acquire different flavors by becoming anisotropic [90], packing onto exceptional surfaces as reported by Okugawa and Yokoyama [91], exceptional lines driven by disorder [92], or exceptional rings [93–95]. They can also be engineered [96] or used as a means to interpret the physics of surface states in three-dimensional systems [97].

For \mathcal{PT} symmetric Hamiltonians, exceptional points mark the onset of a transition from a region with real eigenvalues (and thus oscillatory solutions) to one with complex (conjugate) eigenvalues (and thus unstable growing or decaying solutions), all in spite of the evolution respecting always \mathcal{PT} symmetry (this corresponds to a cut along γ_{diff} in figures 2(a)–(b))⁵.



6. Interpreting defectiveness

In spite of the pivotal role of defectiveness in phenomena unique to non-Hermitian systems, no interpretation is usually given in the context of the search for topological states. Here, we use the simple scattering picture mentioned in previous sections to make sense of defectiveness. The main idea is that a lattice with at least one translationally invariant direction (in d –spatial dimensions) under a non-equilibrium or dynamic condition where, for example, the particles are bound to come from the left (e.g. a boundary condition) can be mapped to a non-Hermitian lattice in dimension $d - 1$ spanning a portion of the original Hilbert space, this is represented in figure 3. The effective Hamiltonian for the reduced system is such as to provide the scattering states of the ancestor or parent Hermitian lattice projected onto the chosen Hilbert space. This path has been explored earlier in tight-binding lattices while trying to make sense of non-Hermitian Hamiltonians in a series of studies by Jin and Song [99, 100] and others [101].

We start by mentioning that this approach has an immediate weakness: there is no guarantee that a non-Hermitian Hamiltonian can always be assimilated to the effective description of a scattering situation (typically stemming from a Hermitian Hamiltonian in larger Hilbert space plus a boundary condition, e.g. incidence


⁴ Note that there has been an intense recent discussion on the need (or lack of it) to encircle the exceptional point to obtain chiral state conversion [72, 73] and also on the dependence on the initial point [74]. We urge the reader to get updated on these issues by reading these references.

⁵ Note that the transition from \mathcal{PT} preserved to \mathcal{PT} broken eigenstates may be shifted away from the exceptional points due to nonlinearities, as reported in [98].



direction of the particles)⁶. That said, a key point to note is the realization that the effective Hamiltonian with gains and losses also encodes information on the boundary condition (which is typically enforced separately!). As such, defectiveness, the lack of a full set of linearly-independent eigenvectors, can be understood as the lack of enough states which are compatible with the boundary condition.

⁶ Furthermore, in many cases, even when this parent or ancestor Hermitian Hamiltonian exists, the effective description usually involves an energy-dependent Hamiltonian (and so the link is restricted to a discrete set of energies (resonances)).



As an example of the above arguments, let us think of a ribbon made of a Chern insulator with particles incident from the left (our boundary condition as typically used to compute transmission coefficients). Now we decide that we are interested only in the projection of those scattering states compatible with the boundary condition over a stripe perpendicular to the translationally invariant direction. In this case, the effective one-dimensional system will sustain an edge state at a single edge and not both, thus reducing the number of linearly-independent eigenvector(s).

Based on the scattering picture it is thus reasonable to expect that a non-Hermitian lattice in dimension d is an effective description of an Hermitian lattice in dimension $d + 1$ plus boundary conditions, when such a parent lattice exists. This *dimensionality link* is consistent with several findings in the literature including: the observation in [52] that a simple one-dimensional chain with asymmetric hoppings is the non-Hermitian proxy of the Chern insulator (as a topological phase not requiring a symmetry), and the anomalous localization transitions in non-Hermitian systems found in low dimensions [102]. In this sense, the non-Hermitian effective description is very much like a shadow (in the sense of a projection) of the higher-dimensional Hermitian ancestor, in a situation reminiscent of what happens in quasicrystals [103, 104].

6.1. A path to obtain a Hermitian parent of a non-Hermitian Hamiltonian

At this point the reader might wonder how to proceed to obtain the Hermitian parent of a given non-Hermitian Hamiltonian. Here we will sketch the procedure for a simple case following Jin and Song [99, 100]. The first point is to refine our starting question. Specifically, we want to obtain a Hermitian parent of a given non-Hermitian Hamiltonian so that a subset of eigenstates of both Hamiltonians share the same projection of the corresponding eigenstates over the common part of the Hilbert space.

In many cases an answer to the previous question can be obtained by referring to a scattering picture. Figure 4(a) represents a tight-binding network containing an absorbing on-site term ($-i\gamma$) at one site. H_{sub} is an arbitrary network with Hermitian terms. We know that such a non-Hermitian term can serve as an effective description of the interaction with the outer world, see for example [105] where this Hamiltonian approach was put forward in the context of quantum transport in the presence of decoherence effects. Therefore one can seek a Hermitian parent in the network of figure 4(b), which is the same network as the one in (a) with the non-Hermitian term replaced by the semi-infinite lead acting as an outer world.

To connect both systems one needs to impose that the solutions projected over the common Hilbert space coincide. The solutions can be chosen as a superposition of plane waves with unknown amplitudes, as indicated in the figure. Replacing this ansatz into the Schrödinger equation will lead to equations that need to be fulfilled so that the requirement is met. Jin and Song [99] showed that the conditions lead to an interesting connection:

real-energy eigenstates of PT-symmetric non-Hermitian tight-binding Hamiltonians have a counterpart (with equal projection over the common Hilbert space): the resonant transmission states of the Hermitian lattice⁷.

If we consider the same system but now invert the sign of the imaginary on-site potential (thus representing a source rather than a sink) as shown in figure 4(c), one can show that the connection with the Hermitian parent Hamiltonian also occurs but this time at a different state: a resonant state (shown in figure 4(d)) which is the time-reversed solution of the one in figure 4(b). Therefore, we see that the full Hermitian parent Hamiltonian has two linearly-independent solutions related by time-reversal symmetry (TRS), but each of them corresponds to a *different* non-Hermitian child and each of them is defective at that energy. We can see that the boundary condition on the Hermitian parent necessary to define the scattering states, affects the non-Hermitian child.

6.2. Extreme defectiveness from higher-order exceptional points and the non-Hermitian skin effect

Finally, we comment on an extreme case of defectiveness found in non-Hermitian lattices: a pristine (without disorder) finite system may become devoid of extended states having all eigenstates localized close to a boundary [47, 48]; the same systems typically show an extreme sensitivity of the spectrum to a change in the boundary conditions (from open to periodic) [47]. This anomalous localization was attributed to the proximity (in parameter space) to exceptional points with an order scaling with the system size in [48], an effect which was interpreted as an environment-mediated interaction effect. The same effect was confirmed analytically in [49] for a non-Hermitian SSH model with asymmetric hoppings and dubbed the *non-Hermitian skin effect*. Recently, Lee and Thomale [66] presented a characterization of these peculiar boundary modes and provided conditions for their existence. The non-Hermitian skin effect presents a conceptual obstacle for building a bulk-boundary correspondence as typically the topological invariants are based on the bulk eigenstates of the system, which in these cases may have a completely different character than the eigenstates of the finite system, no matter how large [47]. We note that, after submission of this article, different experiments have put forward realizations of this anomalous localization in topoelectrical circuits [106, 107], quantum walks [108] mechanical systems [109] and magnons [110].

7. The many paths to a bulk-boundary correspondence

The term topological is used in connection to a given state to mean that there is an underlying bulk-boundary correspondence. Establishing a bulk-boundary correspondence in the most usual sense requires: (i) characterizing the bulk states through a topological invariant, typically an integral over the Brillouin zone of a kernel that depends only on the bulk (Bloch-type) eigenstates; (ii) characterizing the edge or boundary states when translational symmetry is lost in at least one space direction, and (iii) linking the topological invariant with the boundary modes. Since the pioneering TKNN paper [2], this has been a crucial issue where progress is still underway even in Hermitian Hamiltonians [111, 112]; the interested reader can find more extensive accounts on this topic in [8, 10, 113, 114]. Different paths are currently being intensively explored in the quest for a bulk-boundary correspondence that may allow for classification of the topological phases of non-Hermitian lattices [22, 115–118]. These forking paths arise because of the many possible options; the very first one is the use of the Hamiltonian or Green's functions as a starting point. In any of the two cases, one might use the initial non-Hermitian single-particle Hamiltonian or a modified one. More flavors are added by the different possible definitions of a gap and the fact that symmetries are enriched by the lifting of the Hermiticity constraint. In the following we mention only a few of the related works.

Early attempts to complete this program include [51], where the *vorticity*, defined earlier in section 2, appears as a topological invariant. We emphasize that this is at odds with what happens in Hermitian topological systems, as the invariant is always computed from the eigenstates not the eigenvalues. One must also note that in [51] one of the underlying assumptions is that the bandstructure is separable, thereby ruling out cases with non-Hermitian degeneracies.

Another proposal for a classification was presented in [52] for the case of the point gaps introduced in section 2. The resulting classification is analogous with the Hermitian periodic table of topological invariants [119] in the Altland Zirnbauer symmetry classes [120]. This was more recently complemented with a study of cases comprising line gaps [54]. The general strategy is based on reducing the original problem to a Hermitian one where the classification is known. One of the routes allowing for such a reduction is considering the doubled Hamiltonian [52, 60]:

⁷ Further work by the same authors generalized this result to a situation where the PT-symmetry requirement is relaxed [100].

$$\mathcal{H}_{\text{doubled}} = \begin{pmatrix} 0 & \mathcal{H} \\ \mathcal{H}^\dagger & 0 \end{pmatrix},$$

which by construction is Hermitian. From the scattering perspective used earlier to interpret defectiveness, one can also interpret the benefit of doubling the Hamiltonian: in this the information which is missing in the non-Hermitian Hamiltonian is restored, thus avoiding the difficulties introduced by defectiveness and the loss of linearly-independent eigenvectors.

In most of these works the symmetries of the effective Hamiltonian play a crucial role. As compared with usual Hermitian systems, the symmetries are also much enriched in the non-Hermitian case [116]. Indeed, symmetries such as particle hole symmetry fork into two since complex conjugation and transposition become distinct [54], while others get unified, as antiunitary symmetries which are distinct in the Hermitian case and now can be mapped onto each other [121]. There are also new symmetries which appear for the non-Hermitian case, such as pseudo-Hermiticity [122], while others such as non-Hermitian chiral symmetry [123, 124] may remain hidden [125].

In cases where the definition of the topological invariants is attempted directly from the non-Hermitian Hamiltonian, the presence of exceptional points may bring issues such as the non-Hermitian skin effect mentioned in the previous section. Such anomalous localization jeopardizes a direct path connecting a bulk topological invariant to the boundary modes; the sensitivity to the boundary conditions is extreme in this case and the bulk spectrum differs from that of the finite system and, furthermore, the character of the eigenstates might be disparate (one might have extended states while the other may show only localized states). This motivated other authors to choose non-Bloch invariants [56, 126], i.e. invariants based on a finite system rather than bulk states, or invariants defined in real space [127]. A similar approach, without reference to Bloch states but rather a system with open boundaries, was presented by Kunst and collaborators [59].

Another path is the use of Green's functions as put forward by Zirnstein *et al* [128]. This is motivated by the fact that as one special and useful case of a response function, Green's functions describe observables and could therefore be used as an alternative to the Hamiltonian as the starting point for a topological characterization [129, 130]. This has been particularly useful for interacting systems [131]. In the case of non-Hermitian systems, [128] focuses on the one-dimensional case where they find that although there might be no correspondence between topological invariants obtained with periodic boundary conditions and the boundary eigenstates with open boundary conditions, the winding number signals a topological phase transition in the bulk where there is a spatial growth of the Green function.

Borgnia and collaborators [60] combined local Green's functions with the doubled Hamiltonian construction to bring together skin effects and bridge the gap for a bulk-boundary correspondence. On a different path, but with the same unifying aim, Kunst and collaborators [132] put forward a transfer matrix approach. [115] focused on lattices in one dimension with losses added as on-site terms identifying a winding number. Finally, the authors in [133] proposed a path based on singular value decomposition aimed at restoring the correspondence between open and periodic boundary conditions (the reader may also enjoy the discussion on the connection with the doubled Hamiltonian approach in that reference).

8. Connection with Floquet systems

In this section we examine a few connections with the physics of so-called Floquet systems. During the last few years there has been an intense activity in the area of driven systems, exploring ways of harnessing light or time-dependent potentials to substantially alter the states of a material in useful ways. A prominent example is inducing chiral edge states in an otherwise normal material by shining a laser on it [13, 14, 134].

Since the prevalent theory for time-periodic systems is the Floquet theory [135, 136], laser-induced states are referred to in the literature as *Floquet edge states* [13, 137, 138] or, more generally, Floquet–Bloch states [139] to mean states not necessarily of topological origin. Along this new avenue for topological states, studies have focused on diverse materials and systems including normal insulators, [14], graphene [140–143], other two-dimensional materials [144–146], Rashba wires [147], ultracold matter [148], and topological insulators [139, 149–151]. Issues such as thermalization [152] or even the spontaneous generation of magnetism [153] in laser-illuminated systems are some of the latest fronts explored in this area.

Like non-Hermitian systems, Floquet systems can also be classified as non-equilibrium systems. Indeed, the states can even be unstable and typically one needs active sources or sinks. However, besides this first fact there are many more subtle connecting points. A problem of this type involving a time-dependent term is usually solved in Floquet space, which is the direct product between the usual Hilbert space and the space of time-periodic functions with the period of the Hamiltonian; the latter adds a replica index which can be assimilated to different ‘photon’ channels [154]. Many observables require projecting over a single replica (like the time-averaged density of states or in a transport calculation where one needs to assume a reference incident channel

for scattering), and thus the associated properties correspond to those of the effective Hamiltonian for that portion of Floquet space. This effective Hamiltonian might also be non-Hermitian, but even when it is not, the mere partition of the space leads to effects similar to those observed in non-Hermitian lattices. For example, because of the dissimilar weight on different replicas, there could be Floquet edge states that do not contribute to the transport response and remain silent [155]; the coexistence of edge states with a continuum of states belonging to other replicas may introduce a lifetime to those states (see [156–158]), so the spectrum can be considered as effectively complex!

Based on these similarities, many tools used for Floquet systems find a way to non-Hermitian Hamiltonians. This is the case for the doubled Hamiltonian approach mentioned earlier for non-Hermitian systems which has been used earlier for Floquet symmetry-protected topological phases [159].

To close this section we note that there are a few studies that examine the interplay between non-Hermiticity and driving in models with gains and losses [160–163], or others with non-reciprocal interactions [164].

9. Final remarks

Besides providing an overview of the literature in this rapidly growing field, here we have discussed a few interpretative issues related to concepts like defectiveness, the lack of a full basis of eigenstates for a given non-Hermitian operator. The chosen viewpoint corresponds to the scattering theory which provides a starting point to guide our search. The main idea is to think of a non-Hermitian Hamiltonian as an effective description for a parent Hermitian model (on a restricted Hilbert space) plus a boundary condition. The effective system is much like the shadow (in the sense of a projection) of the higher-dimensional parent. Within this picture, defectiveness can be seen as the existence of eigenstates of the parent Hamiltonian which are incompatible with the boundary condition.

The quest for new phenomena now shows many promising possibilities, like the counterpart of higher-order topological states explored by Edvardsson and others [165–167], or the use of non-Hermiticity to model the finite lifetime introduced by electron–electron or electron–phonon interactions [26] and the search for gapless topological phases in the non-Hermitian arena [168–171].

From the experimental side, there have been many more experiments aimed at exceptional points in small systems (see discussion in section 5) than in lattices, as discussed here. One noteworthy exception is the experimental observation of bulk Fermi arcs originating from radiative losses in photonic crystal slabs [172]. The tipping point is expected to come when the loop closes by feeding new experiments, confirming or refuting existing theory. These are likely to come first in photonics and acoustics or perhaps something that our dear reader has just unveiled. In any case, as the writer said, ‘*may your trails be crooked, winding, lonesome, dangerous, leading to the most amazing view*’.⁸

Acknowledgments

LEFFT acknowledges support from FONDECYT (Chile) under grant number 1170917.

ORCID iDs

Luis E Foa Torres  <https://orcid.org/0000-0002-6319-9593>

References

- [1] von Klitzing K, Dorda G and Pepper M 1980 New method for high-accuracy determination of the fine-structure constant based on quantized Hall resistance *Phys. Rev. Lett.* **45** 494–7
- [2] Thouless D J, Kohmoto M, Nightingale M P and den Nijs M 1982 Quantized Hall conductance in a two-dimensional periodic potential *Phys. Rev. Lett.* **49** 405–8
- [3] Haldane F D M 1988 Model for a quantum Hall effect without Landau levels: condensed-matter realization of the ‘Parity Anomaly’ *Phys. Rev. Lett.* **61** 2015–8
- [4] Kane C L and Mele E J 2005 Quantum spin Hall effect in graphene *Phys. Rev. Lett.* **95** 226801
- [5] Bernevig B A, Hughes T L and Zhang S-C 2006 Quantum spin Hall effect and topological phase transition in HgTe quantum wells *Science* **314** 1757–61
- [6] König M *et al* 2007 Quantum spin Hall insulator state in HgTe quantum wells *Science* **318** 766–70
- [7] Hsieh D *et al* 2008 A topological Dirac insulator in a quantum spin Hall phase *Nature* **452** 970–4
- [8] Bernevig B A and Hughes T L 2013 *Topological Insulators and Topological Superconductors* (Princeton, NJ: Princeton University Press)

⁸ Quote from Edward Abbey’s *Desert Solitaire*, New York, McGraw-Hill, 1968.

- [9] Frank Ortmann S O V, Roche S and Molenkamp L W (ed) 2015 *Topological Insulators: Fundamentals and Perspectives* (New York: Wiley)
- [10] Asbóth J K, Oroszlány L and Pályi A 2016 *A Short Course on Topological Insulators, vol 919 of Lecture Notes in Physics* (Cham: Springer International Publishing)
- [11] Vafeek O and Vishwanath A 2014 Dirac fermions in solids: from high-T_c cuprates and graphene to topological insulators and Weyl semimetals *Annual Review of Condensed Matter Physics* **5** 83–112
- [12] Hasan M Z, Xu S-Y, Belopolski I and Huang S-M 2017 Discovery of Weyl fermion semimetals and topological Fermi arc states *Annual Review of Condensed Matter Physics* **8** 289–309
- [13] Oka T and Aoki H 2009 Photovoltaic Hall effect in graphene *Phys. Rev. B* **79** 081406
- [14] Lindner N H, Refael G and Galitski V 2011 Floquet topological insulator in semiconductor quantum wells *Nat. Phys.* **7** 490–5
- [15] McClver J W *et al* 2018 Light-induced anomalous Hall effect in graphene arXiv:1811.03522 [cond-mat]
- [16] Rudner M S and Levitov L S 2009 Topological transition in a non-Hermitian quantum walk *Phys. Rev. Lett.* **102** 065703
- [17] Esaki K, Sato M, Hasebe K and Kohmoto K 2011 Edge states and topological phases in non-Hermitian systems *Phys. Rev. B* **84** 205128
- [18] Diehl S, Rico E, Baranov M A and Zoller P 2011 Topology by dissipation in atomic quantum wires *Nat. Phys.* **7** 971–7
- [19] Yuce C 2015 Topological phase in a non-Hermitian symmetric system *Phys. Lett. A* **379** 1213–8
- [20] San-Jose P, Cayao J, Prada E and Aguado R 2016 Majorana bound states from exceptional points in non-topological superconductors *Sci. Rep.* **6** 21427
- [21] Lee T E 2016 Anomalous edge state in a non-Hermitian lattice *Phys. Rev. Lett.* **116** 133903
- [22] Leykam D, Bliokh K Y, Huang C, Chong Y and Nori F 2017 Edge modes, degeneracies, and topological numbers in non-Hermitian systems *Phys. Rev. Lett.* **118** 040401
- [23] Bender C M and Boettcher S 1998 Real spectra in non-Hermitian Hamiltonians having \mathcal{PT} symmetry *Phys. Rev. Lett.* **80** 5243–6
- [24] Rotter I 2009 A non-Hermitian Hamilton operator and the physics of open quantum systems *J. Phys. A: Math. Theor.* **42** 153001
- [25] Rotter I and Bird J P 2015 A review of progress in the physics of open quantum systems: theory and experiment *Rep. Prog. Phys.* **78** 114001
- [26] Kozii V and Fu L 2017 Non-Hermitian topological theory of finite-lifetime quasiparticles: prediction of bulk Fermi arc due to exceptional point arXiv:1708.05841 [cond-mat]
- [27] Yoshida T, Peters R and Kawakami N 2018 Non-Hermitian perspective of the band structure in heavy-fermion systems arXiv:1805.01172 [cond-mat]
- [28] Yamamoto K, Nakagawa M, Adachi K, Takasan K, Ueda M and Kawakami N 2019 Theory of non-Hermitian fermionic superfluidity with a complex-valued interaction *Phys. Rev. Lett.* **123** 123601
- [29] Zyuzin A A and Zyuzin A Y 2018 Flat band in disorder-driven non-Hermitian Weyl semimetals *Phys. Rev. B* **97** 041203
- [30] Liertzer M *et al* 2012 Pump-induced exceptional points in lasers *Phys. Rev. Lett.* **108** 173901
- [31] Peng B *et al* 2016 Chiral modes and directional lasing at exceptional points *Proc. Natl Acad. Sci.* **113** 6845–50
- [32] Harari G *et al* 2018 Topological insulator laser: theory *Science* **359** eaar4003
- [33] Bandres M A *et al* 2018 Topological insulator laser: experiments *Science* **359** eaar4005
- [34] Zhao H *et al* 2018 Topological hybrid silicon microlasers *Nat. Commun.* **9** 981
- [35] Rüter C E *et al* 2010 Observation of parity-time symmetry in optics *Nat. Phys.* **6** 192–5
- [36] Longhi S 2017 Parity-time symmetry meets photonics: a new twist in non-Hermitian optics *EPL* **120** 64001
- [37] Feng L, El-Ganainy R and Ge L 2017 Non-Hermitian photonics based on parity-time symmetry *Nat. Photonics* **11** 752
- [38] Ozawa T *et al* 2019 Topological photonics *Rev. Mod. Phys.* **91** 015006
- [39] Bliokh K Y, Leykam D, Lein M and Nori F 2019 Topological non-Hermitian origin of surface Maxwell waves *Nat. Commun.* **10** 580
- [40] Lee C H *et al* 2018 Topoelectrical circuits *Communications Physics* **1** 39
- [41] Ezawa M 2019 Non-Hermitian higher-order topological states in nonreciprocal and reciprocal systems with their electric-circuit realization *Phys. Rev. B* **99** 201411
- [42] Dembowski C *et al* 2001 Experimental observation of the topological structure of exceptional points *Phys. Rev. Lett.* **86** 787–90
- [43] Dembowski C *et al* 2004 Encircling an exceptional point *Phys. Rev. E* **69** 056216
- [44] Stehmann T, Heiss W D and Scholtz F G 2004 Observation of exceptional points in electronic circuits *J. Phys. A: Math. Gen.* **37** 7813–9
- [45] Hu W, Wang H, Shum P P and Chong Y D 2017 Exceptional points in a non-Hermitian topological pump *Phys. Rev. B* **95** 184306
- [46] Miri M-A and Alú A 2019 Exceptional points in optics and photonics *Science* **363** eaar7709
- [47] Xiong Y 2018 Why does bulk boundary correspondence fail in some non-Hermitian topological models *Journal of Physics Communications* **2** 035043
- [48] Martínez Alvarez V M, Barrios Vargas J E and Foa Torres L E F 2018 Non-Hermitian robust edge states in one dimension: anomalous localization and eigenspace condensation at exceptional points *Phys. Rev. B* **97** 121401
- [49] Yao S and Wang Z 2018 Edge states and topological invariants of non-Hermitian systems *Phys. Rev. Lett.* **121** 086803
- [50] Martínez Alvarez V M, Barrios Vargas J E, Berdakin M and Foa Torres L E F 2018 Topological states of non-Hermitian systems *The European Physical Journal Special Topics* **227** 1295–308
- [51] Shen H, Zhen B and Fu L 2018 Topological band theory for non-Hermitian Hamiltonians *Phys. Rev. Lett.* **120** 146402
- [52] Gong Z *et al* 2018 Topological phases of non-Hermitian systems *Phys. Rev. X* **8** 031079
- [53] Yin C, Jiang H, Li L, Lü R and Chen S 2018 Geometrical meaning of winding number and its characterization of topological phases in one-dimensional chiral non-Hermitian systems *Phys. Rev. A* **97** 052115
- [54] Kawabata K, Shiozaki K, Ueda M and Sato M 2018 Symmetry and topology in non-Hermitian physics arXiv:1812.09133 [cond-mat, physics:math-ph, physics:physics, physics:quant-ph]
- [55] Li L, Lee C H and Gong J 2019 Geometric classification of non-Hermitian topological systems through the singularity ring arXiv:1905.04965 [cond-mat, physics:quant-ph]
- [56] Yao S, Song F and Wang Z 2018 Non-Hermitian Chern bands and Chern numbers arXiv:1804.04672 [cond-mat, physics:physics]
- [57] Ghatak A and Das T 2019 New topological invariants in non-Hermitian systems *J. Phys. Condens. Matter* **31** 263001
- [58] Jiang H, Yang C and Chen S 2018 Topological invariants and phase diagrams for one-dimensional two-band non-Hermitian systems without chiral symmetry *Phys. Rev. A* **98** 052116
- [59] Kunst F K, Edvardsson E, Budich J C and Bergholtz E J 2018 Biorthogonal bulk-boundary correspondence in non-Hermitian systems *Phys. Rev. Lett.* **121** 026808
- [60] Borgnia D S, Kruchkov A J and Slager R-J 2019 Non-Hermitian boundary modes arXiv:1902.07217 [cond-mat, physics:quant-ph]
- [61] Moiseyev N 2011 *Non-Hermitian Quantum Mechanics* (Non-Hermitian Quantum Mechanics) (Cambridge: Cambridge University Press)

- [62] Landauer R and Martin T 1994 Barrier interaction time in tunneling *Rev. Mod. Phys.* **66** 217–28
- [63] Ni X *et al* 2018 \mathcal{PT} phase transitions of edge states at \mathcal{PT} symmetric interfaces in non-Hermitian topological insulators *Phys. Rev. B* **98** 165129
- [64] Yuce C 2018 Edge states at the interface of non-Hermitian systems *Phys. Rev. A* **97** 042118
- [65] Ryu J-W *et al* 2019 Emergent localized states at the interface of a twofold \mathcal{PT} -symmetric lattice arXiv:1904.00444 [cond-mat]
- [66] Lee C H and Thomaie R 2019 Anatomy of skin modes and topology in non-Hermitian systems *Phys. Rev. B* **99** 201103
- [67] Lee C H, Li L and Gong J 2019 Hybrid higher-order skin-topological modes in nonreciprocal systems *Phys. Rev. Lett.* **123** 016805
- [68] von Neuman J and Wigner E 1929 Über merkwürdige diskrete eigenwerte. Über das Verhalten von eigenwerten bei adiabatischen Prozessen *Physikalische Zeitschrift* **30** 467–70
- [69] Kato T 1995 Perturbation theory for linear operators *Classics in Mathematics* 2nd edn (Berlin: Springer)
- [70] Berry M 1994 Pancharatnam, virtuoso of the Poincaré sphere: an appreciation *Curr. Sci.* **67** 220–3
- [71] Berry M 2010 Geometric Phase Memories *Nat. Phys.* **6** 148–150
- [72] Hassan A U *et al* 2017 Chiral state conversion without encircling an exceptional point *Phys. Rev. A* **96** 052129
- [73] Hassan A U *et al* 2017 Chiral state conversion without encircling an exceptional point *Phys. Rev. A* **96** 069908 erratum Hassan A U *et al* 2017 *Phys. Rev. A* **96** 052129
- [74] Zhang X-L, Wang S, Hou B and Chan C 2018 Dynamically encircling exceptional points: *in situ* control of encircling loops and the role of the starting point *Phys. Rev. X* **8** 021066
- [75] Bittner S *et al* 2012 \mathcal{PT} symmetry and spontaneous symmetry breaking in a microwave billiard *Phys. Rev. Lett.* **108** 024101
- [76] Doppler J *et al* 2016 Dynamically encircling an exceptional point for asymmetric mode switching *Nature* **537** 76–9
- [77] Ghosh S N and Chong Y D 2016 Exceptional points and asymmetric mode conversion in quasi-guided dual-mode optical waveguides *Sci. Rep.* **6** 19837
- [78] Övarez G A, Danieli E P, Levstein P R and Pastawski H M 2006 Environmentally induced quantum dynamical phase transition in the spin swapping operation *J. Chem. Phys.* **124** 194507
- [79] Xu H, Mason D, Jiang L and Harris J G E 2016 Topological energy transfer in an optomechanical system with exceptional points *Nature* **537** 80–3
- [80] Lin Z *et al* 2011 Unidirectional invisibility induced by \mathcal{PT} -symmetric periodic structures *Phys. Rev. Lett.* **106** 213901
- [81] Feng L, Wong Z J, Ma R-M, Wang Y and Zhang X 2014 Single-mode laser by parity-time symmetry breaking *Science* **346** 972–5
- [82] Hodaie H, Miri M-A, Heinrich M, Christodoulides D N and Khajavikhan M 2014 Parity-time-symmetric microring lasers *Science* **346** 975–8
- [83] Chen W, Kaya Özdemir Ö, Zhao G, Wiersig J and Yang L 2017 Exceptional points enhance sensing in an optical microcavity *Nature* **548** 192–6
- [84] Hodaie H *et al* 2017 Enhanced sensitivity at higher-order exceptional points *Nature* **548** 187–91
- [85] Rechtsman M C 2017 Applied physics: optical sensing gets exceptional *Nature* **548** 161–2
- [86] Bird J P and Ochiai Y 2004 Electron spin polarization in nanoscale constrictions *Science* **303** 1621–2
- [87] Yoon Y *et al* 2012 Coupling quantum states through a continuum: a mesoscopic multistate Fano resonance *Phys. Rev. X* **2** 021003
- [88] Auerbach N and Zelevinsky V 2011 Super-radiant dynamics, doorways and resonances in nuclei and other open mesoscopic systems *Rep. Prog. Phys.* **74** 106301
- [89] Eleuch H and Rotter I 2017 Resonances in open quantum systems *Phys. Rev. A* **95** 022117
- [90] Xiao Y-X, Zhang Z-Q, Hang Z H and Chan C T 2019 Anisotropic exceptional points of arbitrary order arXiv:1903.01737 [cond-mat, physics:quant-ph]
- [91] Okugawa R and Yokoyama T 2019 Topological exceptional surfaces in non-Hermitian systems with parity-time and parity-particle-hole symmetries *Phys. Rev. B* **99** 041202
- [92] Moors K, Zyuzin A A, Zyuzin A Y, Tiwari R P and Schmidt T L 2019 Disorder-driven exceptional lines and Fermi ribbons in tilted nodal-line semimetals *Phys. Rev. B* **99** 041116
- [93] Xu Y, Wang Sh-T and Duan L-M 2017 Weyl exceptional rings in a three-dimensional dissipative cold atomic gas *Phys. Rev. Lett.* **118** 045701
- [94] Yoshida T and Hatsugai Y 2019 Exceptional rings protected by emergent symmetry for mechanical systems arXiv:1904.10764 [cond-mat]
- [95] Yoshida T, Peters R, Kawakami N and Hatsugai Y 2019 Symmetry-protected exceptional rings in two-dimensional correlated systems with chiral symmetry *Phys. Rev. B* **99** 121101
- [96] Molina R A and González J 2018 Surface and 3d quantum Hall effects from engineering of exceptional points in nodal-line semimetals *Phys. Rev. Lett.* **120** 146601
- [97] González J and Molina R A 2017 Topological protection from exceptional points in Weyl and nodal-line semimetals *Phys. Rev. B* **96** 045437
- [98] Ge L 2016 Anomalous parity-time-symmetry transition away from an exceptional point *Phys. Rev. A* **94** 013837
- [99] Jin L and Song Z 2010 Physics counterpart of the \mathcal{PT} non-Hermitian tight-binding chain *Phys. Rev. A* **81** 032109
- [100] Jin L and Song Z 2011 A physical interpretation for the non-Hermitian Hamiltonian *J. Phys. A: Math. Theor.* **44** 375304
- [101] Hernandez-Coronado H, Krejčířik D and Siegl P 2011 Perfect transmission scattering as a \mathcal{PT} -symmetric spectral problem *Phys. Lett. A* **375** 2149–52
- [102] Hatano N and Nelson D R 1996 Localization transitions in non-Hermitian quantum mechanics *Phys. Rev. Lett.* **77** 570–3
- [103] Kraus Y E, Lahini Y, Ringel Z, Verbin M and Zilberberg O 2012 Topological states and adiabatic pumping in quasicrystals *Phys. Rev. Lett.* **109** 106402
- [104] Martinez Alvarez V M and Coutinho-Filho M D 2019 Edge states in trimer lattices *Phys. Rev. A* **99** 013833
- [105] D'Amato J L and Pastawski H M 1990 Conductance of a disordered linear chain including inelastic scattering events *Phys. Rev. B* **41** 7411–20
- [106] Helbig T *et al* 2019 Observation of bulk boundary correspondence breakdown in topoelectrical circuits arXiv:1907.11562 [cond-mat]
- [107] Hoffman T *et al* 2019 Reciprocal skin effect and its realization in a topoelectrical circuit arXiv:1908.02759 [cond-mat]
- [108] Xiao L *et al* 2019 Observation of non-Hermitian bulk-boundary correspondence in quantum dynamics arXiv:1907.12566 [cond-mat]
- [109] Ghatak A, Brandenbourger M, van Wezel J and Coullais C 2019 Observation of non-Hermitian topology and its bulk-edge correspondence arXiv:1907.11619 [cond-mat]
- [110] Yu T *et al* 2019 Magnon accumulation in chirally coupled magnets arXiv:1909.00953 [cond-mat]
- [111] Rhim J-W, Behrends J and Bardarson J H 2017 Bulk-boundary correspondence from the intercellular Zak phase *Phys. Rev. B* **95** 035421

- [112] Rhim J-W, Bardarson J H and Slager R-J 2018 Unified bulk-boundary correspondence for band insulators *Phys. Rev. B* **97** 115143
- [113] Ortmann F *et al* (ed) 2015 *Topological Insulators: Fundamentals and Perspectives* (New York: Wiley)
- [114] Vanderbilt D 2018 *Berry Phases in Electronic Structure Theory: Electric Polarization, Orbital Magnetization and Topological Insulators* (Cambridge: Cambridge University Press)
- [115] Rudner M S, Levin M and Levitov L S 2016 Survival, decay, and topological protection in non-Hermitian quantum transport arXiv:1605.07652 [cond-mat]
- [116] Zhou H and Lee J Y 2018 Periodic table for topological bands with non-Hermitian Bernard-LeClair symmetries arXiv:1812.10490 [cond-mat, physics:physics, physics:quant-ph]
- [117] Ge Z-Y *et al* 2019 Topological band theory for non-Hermitian systems from a quantum field viewpoint arXiv:1903.09985 [cond-mat, physics:hep-th, physics:quant-ph]
- [118] Jin L and Song Z 2019 Bulk-boundary correspondence in a non-Hermitian system in one dimension with chiral inversion symmetry *Phys. Rev. B* **99** 081103(R)
- [119] Schnyder A P, Ryu S, Furusaki A and Ludwig A W W 2008 Classification of topological insulators and superconductors in three spatial dimensions *Phys. Rev. B* **78** 195125
- [120] Altland A and Zirnbauer M R 1997 Nonstandard symmetry classes in mesoscopic normal-superconducting hybrid structures *Phys. Rev. B* **55** 1142–61
- [121] Kawabata K, Higashikawa S, Gong Z, Ashida Y and Ueda M 2019 Topological unification of time-reversal and particle-hole symmetries in non-Hermitian physics *Nat. Commun.* **10** 297
- [122] Mostafazadeh A 2002 Pseudo-Hermiticity versus PT-symmetry. II. A complete characterization of non-Hermitian Hamiltonians with a real spectrum *J. Math. Phys.* **43** 2814–6
- [123] Ge L 2017 Symmetry-protected zero-mode laser with a tunable spatial profile *Phys. Rev. A* **95** 023812
- [124] Rivero J D H and Ge L 2019 Chiral symmetry in non-Hermitian systems: product rule, Clifford algebra and pseudo-chirality arXiv:1903.02231 [cond-mat, physics:physics, physics:quant-ph]
- [125] Qi B, Zhang L and Ge L 2018 Defect states emerging from a non-Hermitian flatband of photonic zero modes *Phys. Rev. Lett.* **120** 093901
- [126] Yang X, Cao Y and Zhai Y 2019 Non-Hermitian Weyl semimetals: non-Hermitian skin effect and non-Bloch bulk-boundary correspondence arXiv:1904.02492 [cond-mat]
- [127] Song F, Yao S and Wang Z 2019 Non-Hermitian topological invariants in real space arXiv:1905.02211 [cond-mat, physics:physics]
- [128] Zirnstern H-G, Refael G and Rosenow B 2019 Bulk-boundary correspondence for non-Hermitian Hamiltonians via Green functions arXiv:1901.11241 [cond-mat, physics:physics, physics:quant-ph]
- [129] Volovik G E and Yakovenko V M 1989 Fractional charge, spin and statistics of solitons in superfluid ^3He film *J. Phys. Condens. Matter* **1** 5263–74
- [130] Volovik G E 2009 *The Universe in a Helium Droplet* (Oxford: Oxford University Press)
- [131] Gurarie V 2011 Single-particle Green's functions and interacting topological insulators *Phys. Rev. B* **83** 085426
- [132] Kunst F K and Dwivedi V 2018 Non-Hermitian systems and topology: a transfer matrix perspective arXiv:1812.02186 [cond-mat, physics:quant-ph]
- [133] Herviou L, Bardarson J H and Regnault N 2019 Defining a bulk-edge correspondence for non-Hermitian Hamiltonians via singular-value decomposition *Phys. Rev. A* **99** 052118
- [134] Kitagawa T, Oka T, Brataas A, Fu L and Demler E 2011 Transport properties of nonequilibrium systems under the application of light: photoinduced quantum Hall insulators without Landau levels *Phys. Rev. B* **84** 235108
- [135] Kohler S, Lehmann J and Hänggi P 2005 Driven quantum transport on the nanoscale *Phys. Rep.* **406** 379–443
- [136] Moskalets M and Büttiker M 2002 Floquet scattering theory of quantum pumps *Phys. Rev. B* **66** 205320
- [137] Gomez-Leon A and Platero G 2013 Floquet–Bloch theory and topology in periodically driven lattices *Phys. Rev. Lett.* **110** 200403
- [138] Perez-Piskunow P M, Usaj G, Balseiro C A and Foa Torres L E F 2014 Floquet chiral edge states in graphene *Phys. Rev. B* **89** 121401(R)
- [139] Wang Y H, Steinberg H, Jarillo-Herrero P and Gedik N 2013 Observation of Floquet–Bloch states on the surface of a topological insulator *Science* **342** 453–7
- [140] Kibis O V 2010 Metal-insulator transition in graphene induced by circularly polarized photons *Phys. Rev. B* **81** 165433
- [141] Calvo H L, Pastawski H M, Roche S and Foa Torres L E F 2011 Tuning laser-induced band gaps in graphene *Appl. Phys. Lett.* **98** 232103
- [142] Zhou Y and Wu M W 2011 Optical response of graphene under intense terahertz fields *Phys. Rev. B* **83** 245436
- [143] Sentef M *et al* 2015 Theory of Floquet band formation and local pseudospin textures in pump-probe photoemission of graphene *Nat. Commun.* **6** 7047
- [144] Sie E J *et al* 2015 Valley-selective optical Stark effect in monolayer WS_2 *Nat. Mater.* **14** 290–4
- [145] López A, Scholz A, Santos B and Schliemann J 2015 Photoinduced pseudospin effects in silicene beyond the off-resonant condition *Phys. Rev. B* **91** 125105
- [146] Huamán A and Usaj G 2019 Floquet spectrum and two-terminal conductance of a transition-metal dichalcogenide ribbon under a circularly polarized laser field *Phys. Rev. B* **99** 075423
- [147] Klinovaja J, Stano P and Loss D 2016 Topological Floquet phases in driven coupled Rashba nanowires *Phys. Rev. Lett.* **116** 176401
- [148] Eckardt A 2017 Colloquium: atomic quantum gases in periodically driven optical lattices *Rev. Mod. Phys.* **89** 011004
- [149] Dóra B, Cayssol J, Simon F and Moessner R 2012 Optically engineering the topological properties of a spin Hall insulator *Phys. Rev. Lett.* **108** 056602
- [150] Dal Lago V, Atala M and Foa Torres L E F 2015 Floquet topological transitions in a driven one-dimensional topological insulator *Phys. Rev. A* **92** 023624
- [151] González J and Molina R A 2016 Macroscopic degeneracy of zero-mode rotating surface states in 3D Dirac and Weyl semimetals under radiation *Phys. Rev. Lett.* **116** 156803
- [152] Moessner R and Sondhi S L 2017 Equilibration and order in quantum Floquet matter *Nat. Phys.* **13** 424–8
- [153] Rudner M S and Song J C W 2018 Berryogenesis: self-induced Berry flux and spontaneous non-equilibrium magnetism
- [154] Shirley J H 1965 Solution of the Schrödinger equation with a Hamiltonian periodic in time *Phys. Rev.* **138** B979–87
- [155] Foa Torres L E F, Perez-Piskunow P M, Balseiro C A and Usaj G 2014 Multiterminal conductance of a Floquet topological insulator *Phys. Rev. Lett.* **113** 266801
- [156] Rudner M S, Lindner N H, Berg E and Levin M 2013 Anomalous edge states and the bulk-edge correspondence for periodically-driven two dimensional systems *Phys. Rev. X* **3** 031005
- [157] Usaj G, Perez-Piskunow P M, Foa Torres L E F and Balseiro C A 2014 Irradiated graphene as a tunable Floquet topological insulator *Phys. Rev. B* **90** 115423

- [158] Perez-Piskunow P M, Foa Torres L E F and Usaj G 2015 Hierarchy of Floquet gaps and edge states for driven honeycomb lattices *Phys. Rev. A* **91** 043625
- [159] Roy R and Harper F 2017 Periodic table for Floquet topological insulators *Phys. Rev. B* **96** 155118
- [160] Yuce C 2015 PT symmetric Floquet topological phase *Eur. Phys. J. D* **69** 184
- [161] Turker Z, Tombuloglu S and Yuce C 2018 PT symmetric Floquet topological phase in SSH model *Phys. Lett. A* **382** 2013–6
- [162] Zhou L and Gong J 2018 Non-Hermitian Floquet topological phases with arbitrarily many real-quasienergy edge states arXiv:1807.00988 [cond-mat, physics:quant-ph]
- [163] Zhou L and Pan J 2019 Non-Hermitian Floquet topological phases in the double-kicked rotor arXiv:1908.02066 [cond-mat, physics:quant-ph]
- [164] Li H, Kottos T and Shapiro B 2018 Floquet-network theory of nonreciprocal transport *Physical Review Applied* **9** 044031
- [165] Edvardsson E, Kunst F K and Bergholtz E J 2019 Non-Hermitian extensions of higher-order topological phases and their biorthogonal bulk-boundary correspondence *Phys. Rev. B* **99** 081302
- [166] Liu T *et al* 2019 Second-order topological phases in non-Hermitian systems *Phys. Rev. Lett.* **122** 076801
- [167] Ezawa M 2018 Electric-circuit realization of non-Hermitian higher-order topological systems arXiv:1810.04527 [cond-mat]
- [168] Budich J C, Carlström J, Kunst F K and Bergholtz E J 2019 Symmetry-protected nodal phases in non-Hermitian systems *Phys. Rev. B* **99** 041406
- [169] Carlström J, Stålhammar M, Budich J C and Bergholtz E J 2019 Knotted non-Hermitian metals *Phys. Rev. B* **99** 161115
- [170] Bergholtz E J and Budich J C 2019 Non-Hermitian Weyl physics in topological insulator ferromagnet junctions *Physical Review Research* **1** 012003
- [171] Yang X M, Wang P, Jin L and Song Z 2019 Visualizing topology of real-energy gapless phase arising from exceptional point arXiv:1905.07109 [cond-mat, physics:physics, physics:quant-ph]
- [172] Zhou H *et al* 2018 Observation of bulk Fermi arc and polarization half charge from paired exceptional points *Science* **359** 1009–12