

DIRECTIONAL DYNAMICAL CUBES FOR MINIMAL \mathbb{Z}^d -SYSTEMS

CHRISTOPHER CABEZAS, SEBASTIÁN DONOSO, AND ALEJANDRO MAASS

ABSTRACT. We introduce the notions of *directional dynamical cubes* and *directional regionally proximal relation* defined via these cubes for a minimal \mathbb{Z}^d -system (X, T_1, \dots, T_d) . We study the structural properties of systems that satisfy the so called *unique closing parallelepiped property* and we characterize them in several ways. In the distal case, we build the maximal factor of a \mathbb{Z}^d -system (X, T_1, \dots, T_d) that satisfies this property by taking the quotient with respect to the directional regionally proximal relation. Finally, we completely describe distal \mathbb{Z}^d -systems that enjoy the unique closing parallelepiped property and provide explicit examples.

1. INTRODUCTION

The study of *cubical structures* or *cubespaces* was initiated by Host and Kra in [18] as a fundamental tool to show the L^2 convergence of Furstenberg multiple averages for a single transformation. Later, Host, Kra and Maass in [19] introduced cube structures in topological dynamics given by a single homeomorphism proving a structure theorem for transitive systems with the *unique completion property* (this condition is referred in this work as the *unique closing parallelepiped property*). The main result is that this property characterizes inverse limits of minimal rotations on nilmanifolds.

Soon after, the notion of cubespaces became a major theme of research in the development of *higher order Fourier analysis*, starting from Antolín-Camarena and Szegedys's axiomatic formulation of them [1]. Further description of such theory was given by Candela in [5, 6] and another point of view was developed by Gutman, Manners and Varjú in [14, 15, 16]. In all these works a key feature is the study of the unique closing parallelepiped property and the conclusion (roughly speaking) is that a compact space whose cubes have the unique closing parallelepiped property has a *nilstructure*.

Interestingly, the study of *dynamical cubes*, *i.e.*, those defined in [18] and [19], has reached an independent interest, mainly because of his various applications that range from the construction of nilfactors, recognizing patterns in number theory, the study of recurrence in topological dynamics and even the study of pointwise convergence of averages in ergodic theory [8, 9, 10, 16, 20]. Also, extensions to general group actions have been recently carried out in [13].

In this work, we consider a slightly different (but related) notion that we call *directional dynamical cubes*. Such notion first appeared (without a name) in the work of Host [17], where he gave a proof of Tao's L^2 convergence of Furstenberg multiple averages for commuting transformations [24]. Motivated by those works, the study of directional cubes in topological dynamics started in [7], where the authors introduced for a minimal \mathbb{Z}^2 -system (X, T_1, T_2) a space of cubes denoted by $\mathbf{Q}_{T_1, T_2}(X)$ and proved a structure theorem for systems with the unique closing parallelepiped property (here X is a compact metric space and $T_1, T_2 : X \rightarrow X$ are commuting homeomorphisms). Additionally, the authors introduced the notion of (T_1, T_2) -regionally proximal relation $\mathcal{R}_{T_1, T_2}(X)$ associated to the cube structure $\mathbf{Q}_{T_1, T_2}(X)$. This is a strong variant of the classical regionally proximal relation of order one $\mathbf{RP}^{[1]}(X, \langle T_1, T_2 \rangle)$ for \mathbb{Z}^2 -systems. In the distal case they proved that $\mathcal{R}_{T_1, T_2}(X)$ is an equivalence relation and that $(X/\mathcal{R}_{T_1, T_2}(X), T_1, T_2)$ is the maximal factor with the unique closing parallelepiped property.

The purpose of this article is to extend the program of research started in [7] to arbitrary \mathbb{Z}^d -systems, *i.e.*, d commuting homeomorphisms $T_1, \dots, T_d : X \rightarrow X$ of a compact metric space X with $d \geq 2$.

We define the *space of directional cubes* $\mathbf{Q}_{T_1, \dots, T_d}(X)$ as the closure in X^{2^d} of the points

$$(T_1^{n_1 \varepsilon_1} \dots T_d^{n_d \varepsilon_d} x)_{\varepsilon \in \{0,1\}^d},$$

where $x \in X$ and $\mathbf{n} = (n_1, \dots, n_d) \in \mathbb{Z}^d$. Here we point out that since the transformations T_j could be different, the space of directional cubes $\mathbf{Q}_{T_1, \dots, T_d}(X)$ has much less symmetries than its relative ones defined for a single homeomorphism.

As in previous works we are interested in \mathbb{Z}^d -systems with the unique closing parallelepiped property, as a way of understanding the topological counterpart of the characteristic factors introduced by Host in

This work was funded by Proyecto/Grant PIA AFB-170001 and Math-AmSud DYSTIL.

The second author is also supported by Fondecyt Iniciación Grant 11160061.

[17] for an arbitrary number of commuting transformations. We also introduce the (T_1, \dots, T_d) -regionally proximal relation $\mathcal{R}_{T_1, \dots, T_d}(X)$ of (X, T_1, \dots, T_d) associated to these cube structures in order to give a complete description of systems with the unique closing parallelepiped property and define the maximal factor with this property for any \mathbb{Z}^d -system. We defer the precise definitions of the relation to Section 4.

We show the following structure theorem for minimal distal systems.

Theorem 1. *Let (X, T_1, \dots, T_d) be a minimal distal \mathbb{Z}^d -system. The following statements are equivalent:*

- (1) *(X, T_1, \dots, T_d) has the unique closing parallelepiped property, i.e., if $\mathbf{x}, \mathbf{y} \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ have $2^d - 1$ coordinates in common, then $\mathbf{x} = \mathbf{y}$.*
- (2) *$\mathcal{R}_{T_1, \dots, T_d}(X) = \Delta_X$.*
- (3) *The structure of (X, T_1, \dots, T_d) can be described as follows: (i) it is a factor of a minimal distal \mathbb{Z}^d -system (Y, T_1, \dots, T_d) which is a joining of \mathbb{Z}^d -systems $(Y_1, T_1, \dots, T_d), \dots, (Y_d, T_1, \dots, T_d)$, where for each $i \in \{1, \dots, d\}$ the action of T_i on Y_i is the identity; (ii) for each $i, j \in \{1, \dots, d\}$, $i < j$, there exists a \mathbb{Z}^d -system $(Y_{i,j}, T_1, \dots, T_d)$ which is a common factor of (Y_i, T_1, \dots, T_d) and (Y_j, T_1, \dots, T_d) and where T_i and T_j act as the identity; and (iii) Y is jointly relatively independent with respect to the systems $((Y_{i,j}, T_1, \dots, T_d) : i, j \in \{1, \dots, d\}, i < j)$.*

We will see in Section 7 that Y , the Y_j 's and the $Y_{i,j}$'s are given explicitly by the cube structure.

When $d = 2$ we have that $(Y_{1,2}, T_1, T_2)$ is the trivial system (because it is minimal and the actions are identities). Then, relative independence implies that (Y, T_1, T_2) can be seen as the direct product of the systems (Y_1, T_1, T_2) and (Y_2, T_1, T_2) , so Theorem 1 generalizes the structure theorem proved by Donoso and Sun for minimal distal systems in [7] when $d = 2$. Also, the structure in the previous theorem can be thought of as a topological version of *pleasant extensions* developed in a measure theoretical setting by Austin in [3].

We also prove,

Theorem 2. *Let (X, T_1, \dots, T_d) be a minimal distal \mathbb{Z}^d -system. Then, the relation $\mathcal{R}_{T_1, \dots, T_d}(X)$ is an equivalence relation and $(X/\mathcal{R}_{T_1, \dots, T_d}(X), T_1, \dots, T_d)$ is the maximal factor of (X, T_1, \dots, T_d) with the unique closing parallelepiped property. This means that $(X/\mathcal{R}_{T_1, \dots, T_d}(X), T_1, \dots, T_d)$ has the unique closing parallelepiped property and that any factor of (X, T_1, \dots, T_d) that enjoys that property is necessarily a factor of $(X/\mathcal{R}_{T_1, \dots, T_d}(X), T_1, \dots, T_d)$.*

Finally we develop interesting applications of the directional dynamical cubes to study recurrence properties of systems with the unique closing parallelepiped property. It turns out that this property can be characterized studying the sets of return times of a \mathbb{Z}^d -system.

In the previous results, we do not know if the assumption of distality is essential or not. The main feature we use about distal systems is that they enjoy the *gluing property* (we start its discussion in Definition 10) and most of proofs only use this property. We ask then

Problem. *How much of our theory can be extended for general minimal \mathbb{Z}^d -systems? In particular, is the unique closing parallelepiped property preserved under factor maps? We show in this article that this is true for factor maps between minimal distal systems, but we do not know if this holds for general minimal \mathbb{Z}^d -systems.*

1.1. Organization of the paper. In Section 2 we give some basic material used throughout the paper. In Section 3 we present directional dynamical cubes for a \mathbb{Z}^d -system (X, T_1, \dots, T_d) , introduce the main terminology and give their general properties. Next, in Section 4 we introduce the associated (T_1, \dots, T_d) -regionally proximal relation and establish some general results. In Section 5 we give a characterization of \mathbb{Z}^d -systems with the unique closing parallelepiped property in the distal case and in Section 6 we show the structure of a minimal distal system with the unique closing parallelepiped property. Section 7 is devoted to the proof of Theorem 1. Then, in Section 8 we study the sets of return times for minimal distal systems with the unique closing parallelepiped property and give a theorem that characterizes these systems using their return time sets. In Section 9 we provide a family of explicit examples of minimal distal \mathbb{Z}^d -systems with the unique closing parallelepiped property.

Acknowledgement. We thank Wenbo Sun for many helpful discussions. We also thank the anonymous referee for valuable remarks.

2. PRELIMINARIES

2.1. Basic notions from topological dynamics. A *topological dynamical system* is a pair (X, G) , where X is a compact metric space and G is a group of homeomorphisms of the space X into itself. We

also refer to X as a G -system. We use $\rho_X(\cdot, \cdot)$ to denote the metric of X and we let $\Delta_X = \{(x, x) : x \in X\}$ denote the diagonal of $X \times X$.

If $d \geq 1$ is an integer and $T_1, \dots, T_d : X \rightarrow X$ are d commuting homeomorphisms of X , we write (X, T_1, \dots, T_d) to denote the topological dynamical system $(X, \langle T_1, \dots, T_d \rangle)$, where $\langle T_1, \dots, T_d \rangle$ is the group spanned by T_1, \dots, T_d . Throughout this paper those systems are called \mathbb{Z}^d -systems.

A *factor map* between the dynamical systems (Y, G) and (X, G) is an onto and continuous map $\pi : Y \rightarrow X$ such that $\pi \circ g = g \circ \pi$ for every $g \in G$. We say that (Y, G) is an *extension* of (X, G) or that (X, G) is a *factor* of (Y, G) . When π is bijective we say that π is an isomorphism and that (Y, G) and (X, G) are isomorphic or conjugate.

Let $(X_1, G), \dots, (X_k, G)$ be k topological dynamical systems. A *joining* between them is a closed subset $Z \subseteq X_1 \times \dots \times X_k$ which is invariant under the diagonal action $g \times \dots \times g$ (k times) for all $g \in G$ and projects onto each factor.

Suppose the systems $(X_1, G), \dots, (X_k, G)$ are extensions of a common system (W, G) and for $i \in \{1, \dots, k\}$ denote by $\pi_i : X_i \rightarrow W$ the associated factor map. We say that a joining $Z \subseteq X_1 \times \dots \times X_k$ is *relatively independent* with respect to W if for every $(x_1, \dots, x_k) \in Z$, for every $i \in \{1, \dots, k\}$ and for every $x'_i \in X_i$ with $\pi_i(x'_i) = \pi_i(x_i)$ we have

$$(x_1, \dots, x_{i-1}, x'_i, x_{i+1}, \dots, x_k) \in Z.$$

That is, we can freely change coordinates of points in the joining when the corresponding projection to the factor W coincide.

Given a topological dynamical system (X, G) and a point $x \in X$ we denote by $\mathcal{O}_G(x) = \{gx : g \in G\}$ the orbit of x . We say that (X, G) is *transitive* if there exists a point in X whose orbit is dense. We say that (X, G) is *minimal* if the orbit of any point is dense in X . A system (X, G) is *pointwise almost periodic* if for any $x \in X$ the system $(\overline{\mathcal{O}_G(x)}, G)$ is minimal.

A topological dynamical system (X, G) is *distal* if for every $x, y \in X$ such that $x \neq y$ we have

$$\inf_{g \in G} d(gx, gy) > 0.$$

Distal systems have many interesting properties that we will use in the sequel (see [2], chapters 5 and 7). We recall some of them:

Theorem 3.

- (1) *The Cartesian product of distal systems is distal.*
- (2) *Distality is preserved by taking factors and subsystems.*
- (3) *A distal system is minimal if and only if it is transitive, i.e., a distal system is pointwise almost periodic.*
- (4) *If (X, G) is distal and G' is a subgroup of G , then (X, G') is distal.*
- (5) *Factor maps between distal systems are open maps.*

If (X, G) is a topological dynamical system, we can see G as a subset of X^X . We define the *enveloping semigroup* $E(X, G)$ of (X, G) as the closure (for the pointwise convergence) of G in X^X . We have that $E(X, G)$ is compact and Hausdorff, usually non-metrizable. Algebraic properties of the enveloping semigroup have a precise translation into dynamical properties of the system. For example, a topological dynamical system is distal if and only if its enveloping semigroup is a group.

A *left ideal* in a enveloping semigroup $E(X, G)$ is a non-empty subset $I \subseteq E(X, G)$ such that $E(X, G)I \subseteq I$. A *minimal left ideal* is one which does not properly contain a left ideal. An *idempotent* in a enveloping semigroup $E(X, G)$ is an element $u \in E(X, G)$ such that $u^2 = u$. We denote by $J(E(X, G))$ the set of idempotents in $E(X, G)$. We can introduce a *quasi-order* $<$ on the set $J(E(X, G))$ by defining $v < u$ if and only if $vu = v$. If $v < u$ and $u < v$ we say that u and v are *equivalent* and we write $u \sim v$. An idempotent $u \in J(E(X, G))$ is *minimal* if whenever $v \in J(E(X, G))$ and $v < u$ then $u < v$ (or $u \sim v$).

We recall some results about the enveloping semigroup that we use later.

Theorem 4. *Let (Y, G) and (X, G) be topological dynamical systems.*

- (1) *If $\pi : Y \rightarrow X$ is a factor map, then there exists a unique continuous semigroup homomorphism $\theta : E(Y, G) \rightarrow E(X, G)$ such that $\pi(py) = \theta(p)\pi(y)$ for all $y \in Y$ and $p \in E(Y, G)$. If $u \in E(Y, G)$ is a minimal idempotent then $\theta(u) \in E(X, G)$ is a minimal idempotent too.*
- (2) *$x, x' \in X$ are proximal if and only if there exists $p \in E(X, G)$ such that $px = px'$.*
- (3) *$x \in X$ is an almost periodic point (i.e., its orbit is minimal) if and only if there exists a minimal idempotent $u \in E(X, G)$ such that $ux = x$.*

2.2. Cubes and faces. For an integer $d \geq 1$ we write $[d] = \{1, \dots, d\}$. We view $\{0, 1\}^d$ in one of two ways, either as a sequence $\varepsilon = \varepsilon_1 \dots \varepsilon_d$ of 0's and 1's written without commas or parentheses, or as a subset of $[d]$. A subset $\varepsilon \subseteq [d]$ corresponds to the sequence $\varepsilon_1 \dots \varepsilon_d \in \{0, 1\}^d$ such that for each $i \in [d]$, $i \in \varepsilon$ if and only if $\varepsilon_i = 1$, *i.e.*, seen as a vector it is the sum of the canonical vectors indexed by the set ε .

For any set X we write $X^{[d]} = X^{2^d}$. A point $\mathbf{x} \in X^{[d]}$ can be described in one of two equivalent ways, depending on the context and convenience:

$$\mathbf{x} = (x_\varepsilon : \varepsilon \in \{0, 1\}^d) = (x_\varepsilon : \varepsilon \subseteq [d]).$$

For $x \in X$ we write $x^{[d]} = (x, \dots, x) \in X^{[d]}$ and the diagonal of $X^{[d]}$ is denoted by $\Delta_X^{[d]} = \{x^{[d]} : x \in X\}$.

A point $\mathbf{x} \in X^{[d]}$ can be decomposed as $\mathbf{x} = (\mathbf{x}', \mathbf{x}'')$ with $\mathbf{x}', \mathbf{x}'' \in X^{[d-1]}$, where $\mathbf{x}' = (x_{\eta_0} : \eta \in \{0, 1\}^{d-1})$ and $\mathbf{x}'' = (x_{\eta_1} : \eta \in \{0, 1\}^{d-1})$. We also isolate the first coordinate writing $X_*^{[d]} = X^{2^{d-1}}$ and a point $\mathbf{x} \in X^{[d]}$ as $\mathbf{x} = (x_\emptyset, \mathbf{x}_*)$, where $x_\emptyset \in X$ and $\mathbf{x}_* = (x_\varepsilon : \varepsilon \subseteq [d], \varepsilon \neq \emptyset) \in X_*^{[d]}$.

For $r \leq d$, let $J \subseteq [d]$ with $|J| = d - r$ and $\xi \in \{0, 1\}^{d-r}$. A *face of dimension r* of \mathbf{x} is an element of the form $(x_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_J = \xi) \in X^{[r]}$, where $\varepsilon_J = (\varepsilon_i : i \in J)$.

Let (X, T_1, \dots, T_d) be a \mathbb{Z}^d -system. For $j \in [d]$, the *j -th face transformation* $T_j^{[d]} : X^{[d]} \rightarrow X^{[d]}$ is defined for every $\mathbf{x} \in X^{[d]}$ and every $\varepsilon \subseteq [d]$ by:

$$(T_j^{[d]} \mathbf{x})_\varepsilon = \begin{cases} T_j x_\varepsilon & \text{if } j \in \varepsilon, \\ x_\varepsilon & \text{if } j \notin \varepsilon. \end{cases}$$

The *face group of dimension d* is the group $\mathcal{F}_{T_1, \dots, T_d}$ of transformations of $X^{[d]}$ generated by the face transformations. Let G be the group spanned by T_1, \dots, T_d and set $G_{[d]}^\Delta = \{g^{[d]} : g \in G\}$. We denote by $\mathcal{G}_{T_1, \dots, T_d}$ the subgroup of $G_{[d]}^\Delta$ generated by $\mathcal{F}_{T_1, \dots, T_d}$ and $G_{[d]}^\Delta$.

3. DIRECTIONAL DYNAMICAL CUBES FOR \mathbb{Z}^d -SYSTEMS

In this section we present the notion of directional dynamical cubes for \mathbb{Z}^d -systems (X, T_1, \dots, T_d) together with its main properties. This is a generalization of the dynamical cubes introduced by Donoso and Sun in [7] when $d = 2$.

Definition 5. Let (X, T_1, \dots, T_d) be a \mathbb{Z}^d -system. The set of *directional dynamical cubes* associated to (X, T_1, \dots, T_d) is defined by,

$$\mathbf{Q}_{T_1, \dots, T_d}(X) = \overline{\{(T_1^{n_1 \varepsilon_1} \dots T_d^{n_d \varepsilon_d} x)_{\varepsilon \in \{0, 1\}^d} : x \in X, \mathbf{n} = (n_1, \dots, n_d) \in \mathbb{Z}^d\}} \subseteq X^{[d]}.$$

Adittionally, given $x_0 \in X$ we consider the following restriction of theses cubes to $X_*^{[d]}$,

$$\mathbf{K}_{T_1, \dots, T_d}^{x_0}(X) = \overline{\{(T_1^{n_1 \varepsilon_1} \dots T_d^{n_d \varepsilon_d} x_0)_{\varepsilon \in \{0, 1\}^d \setminus \{\emptyset\}} : \mathbf{n} = (n_1, \dots, n_d) \in \mathbb{Z}^d\}} \subseteq X_*^{[d]}.$$

Given $\{j_1, \dots, j_k\} \subseteq [d]$ and $x_0 \in X$, since $(X, T_{j_1}, \dots, T_{j_k})$ is a \mathbb{Z}^k -system, then we will often consider the set of cubes $\mathbf{Q}_{T_{j_1}, \dots, T_{j_k}}(X) \subseteq X^{[k]}$ and $\mathbf{K}_{T_{j_1}, \dots, T_{j_k}}^{x_0}(X) \subseteq X_*^{[k]}$. For example, if $k = 3$ and $j_1, j_2, j_3 \in [d]$ then $\mathbf{Q}_{T_{j_1}, T_{j_2}, T_{j_3}}(X)$ is the closure of the set of points

$$(x, T_{j_1}^{n_1} x, T_{j_2}^{n_2} x, T_{j_1}^{n_1} T_{j_2}^{n_2} x, T_{j_3}^{n_3} x, T_{j_1}^{n_1} T_{j_3}^{n_3} x, T_{j_2}^{n_2} T_{j_3}^{n_3} x, T_{j_1}^{n_1} T_{j_2}^{n_2} T_{j_3}^{n_3} x),$$

for $x \in X$ and $n_1, n_2, n_3 \in \mathbb{Z}$.

Remark that the generators of \mathbb{Z}^d are given with a precise order. A different choice of this order produces a different set of dynamical cubes. This is the reason why we call them directional cubes.

3.1. Basic structural properties.

Proposition 6. Let (X, T_1, \dots, T_d) be a \mathbb{Z}^d -system and take $k \in [d]$. We have,

- (1) $x^{[d]} \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ for every $x \in X$.
- (2) For all $x, y \in X$ and for all $j \in [d]$, $(x, y) \in \mathbf{Q}_{T_j}(X)$ if and only if $(y, x) \in \mathbf{Q}_{T_j}(X)$.
- (3) The set $\mathbf{Q}_{T_1, \dots, T_d}(X)$ is invariant under $\mathcal{G}_{T_1, \dots, T_d}$ and $\mathbf{K}_{T_1, \dots, T_d}^{x_0}(X)$ is invariant under $\mathcal{F}_{T_1, \dots, T_d}^{x_0}$ ¹.
Therefore $(\mathbf{Q}_{T_1, \dots, T_d}(X), \mathcal{G}_{T_1, \dots, T_d})$ and $(\mathbf{K}_{T_1, \dots, T_d}^{x_0}(X), \mathcal{F}_{T_1, \dots, T_d}^{x_0})$ are topological dynamical systems.
- (4) (Projection) Let $\mathbf{x} \in \mathbf{Q}_{T_1, \dots, T_d}(X)$, $\{j_1, \dots, j_k\} \subseteq [d]$ and $\xi \in \{0, 1\}^{d-k}$. Then,

$$(x_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_{[d] \setminus \{j_1, \dots, j_k\}} = \xi) \in \mathbf{Q}_{T_{j_1}, \dots, T_{j_k}}(X).$$

¹For convenience, we let $\mathcal{F}_{T_1, \dots, T_d}^{x_0}$ denote the restriction of $\mathcal{F}_{T_1, \dots, T_d}$ to $\mathbf{K}_{T_1, \dots, T_d}^{x_0}(X)$

(5) (Duplication) Let $\mathbf{x} = (x_\eta : \eta \in \{0, 1\}^k) \in \mathbf{Q}_{T_{j_1}, \dots, T_{j_k}}(X)$ with $\{j_1, \dots, j_k\} \subseteq [d]$. If $\mathbf{y} \in X^{[d]}$ is defined such that for every $\varepsilon \in \{0, 1\}^d$,

$$y_\varepsilon = x_\eta \iff \forall \ell \in \{1, \dots, k\}, \varepsilon_{j_\ell} = \eta_\ell,$$

then $\mathbf{y} \in \mathbf{Q}_{T_1, \dots, T_d}(X)$.

Proof. Properties (1) to (3) follow directly from definitions. We only prove properties (4) and (5). For property (4) let $\mathbf{x} \in \mathbf{Q}_{T_1, \dots, T_d}(X)$, $\{j_1, \dots, j_k\} \subseteq [d]$ and $\xi \in \{0, 1\}^{d-k}$. By definition, there exist $(x^i)_{i \in \mathbb{N}} \subseteq X$ and $(\mathbf{n}(i))_{i \in \mathbb{N}} \subseteq \mathbb{Z}^d$ such that

$$\forall \varepsilon \in \{0, 1\}^d, \quad x_\varepsilon = \lim_{i \rightarrow \infty} T_1^{n_1(i)\varepsilon_1} \dots T_d^{n_d(i)\varepsilon_d} x^i.$$

Now, consider $\eta \in \{0, 1\}^d$ such that

$$\eta_\ell = \begin{cases} 0 & \ell \in \{j_1, \dots, j_k\}, \\ \xi_\ell & \ell \in [d] \setminus \{j_1, \dots, j_k\}, \end{cases}$$

and set $y^i = T_1^{n_1(i)\eta_1} \dots T_d^{n_d(i)\eta_d} x^i$. Then, for $\varepsilon \in \{0, 1\}^d$ such that $\varepsilon_{[d] \setminus \{j_1, \dots, j_k\}} = \xi$ we have

$$\begin{aligned} x_\varepsilon &= \lim_{i \rightarrow \infty} T_1^{n_1(i)\varepsilon_1} \dots T_d^{n_d(i)\varepsilon_d} x^i, \\ &= \lim_{i \rightarrow \infty} T_{j_1}^{n_{j_1}(i)\varepsilon_{j_1}} \dots T_{j_k}^{n_{j_k}(i)\varepsilon_{j_k}} y^i. \end{aligned}$$

This implies, by definition, that $(x_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_{[d] \setminus \{j_1, \dots, j_k\}} = \xi) \in \mathbf{Q}_{T_{j_1}, \dots, T_{j_k}}(X)$.

Now we prove property (5). Let $\mathbf{x} = (x_\eta : \eta \in \{0, 1\}^k) \in \mathbf{Q}_{T_{j_1}, \dots, T_{j_k}}(X)$ with $\{j_1, \dots, j_k\} \subseteq [d]$. By definition, there exist $(x^i)_{i \in \mathbb{N}} \subseteq X$ and $(\mathbf{m}(i))_{i \in \mathbb{N}} \subseteq \mathbb{Z}^k$ such that

$$\forall \eta \in \{0, 1\}^k, \quad x_\eta = \lim_{i \rightarrow \infty} T_{j_1}^{m_1(i)\eta_1} \dots T_{j_k}^{m_k(i)\eta_k} x^i.$$

Define $\mathbf{n}(i) \in \mathbb{Z}^d$ by

$$n_p(i) = \begin{cases} m_\ell(i) & \text{if } p = j_\ell \text{ for some } j_\ell \in \{j_1, \dots, j_k\}, \\ 0 & \text{if } p \in [d] \setminus \{j_1, \dots, j_k\}. \end{cases}$$

Then, for $\varepsilon \in \{0, 1\}^d$ and $\eta \in \{0, 1\}^k$ such that $\varepsilon_{j_\ell} = \eta_\ell$ for all $\ell \in \{1, \dots, k\}$ we have

$$\begin{aligned} y_\varepsilon = x_\eta &= \lim_{i \rightarrow \infty} T_{j_1}^{m_1(i)\eta_1} \dots T_{j_k}^{m_k(i)\eta_k} x^i, \\ &= \lim_{i \rightarrow \infty} T_1^{n_1(i)\varepsilon_1} \dots T_d^{n_d(i)\varepsilon_d} x^i, \end{aligned}$$

which implies (5). \square

Proposition 6 (3) tells us that $(\mathbf{Q}_{T_1, \dots, T_d}(X), \mathcal{G}_{T_1, \dots, T_d})$ is a topological dynamical system. The following proposition states a little more. Its proof requires the technology of the enveloping semigroup. Basic properties of enveloping semigroups were considered in the preliminaries, for a complete exposition see [2].

Proposition 7. *Let (X, T_1, \dots, T_d) be a minimal \mathbb{Z}^d -system. Then, $(\mathbf{Q}_{T_1, \dots, T_d}(X), \mathcal{G}_{T_1, \dots, T_d})$ is a minimal system. Furthermore, if (X, T_1, \dots, T_d) is distal, then $(\mathbf{Q}_{T_1, \dots, T_d}(X), \mathcal{G}_{T_1, \dots, T_d})$ is distal too.*

Proof. Let $E(\mathbf{Q}_{T_1, \dots, T_d}(X), \mathcal{G}_{T_1, \dots, T_d})$ be the enveloping semigroup of the system $(\mathbf{Q}_{T_1, \dots, T_d}(X), \mathcal{G}_{T_1, \dots, T_d})$. For every $\varepsilon \in \{0, 1\}^d$ let $\pi_\varepsilon : \mathbf{Q}_{T_1, \dots, T_d}(X) \rightarrow X$ be the projection onto the ε -th coordinate and let $\pi_\varepsilon^* : E(\mathbf{Q}_{T_1, \dots, T_d}(X), \mathcal{G}_{T_1, \dots, T_d}) \rightarrow E(X, T_1, \dots, T_d)$ be the corresponding semigroups homomorphism.

Let $u \in E(\mathbf{Q}_{T_1, \dots, T_d}(X), G_{[d]}^\Delta)$ denote a minimal idempotent for the system $(\mathbf{Q}_{T_1, \dots, T_d}(X), G_{[d]}^\Delta)$. We show that u is also a minimal idempotent in $E(\mathbf{Q}_{T_1, \dots, T_d}(X), \mathcal{G}_{T_1, \dots, T_d})$. It suffices to show that if $v \in E(\mathbf{Q}_{T_1, \dots, T_d}(X), \mathcal{G}_{T_1, \dots, T_d})$ with $vu = v$, then $uv = u$. Projecting onto coordinates we deduce that $\pi_\varepsilon^*(vu) = \pi_\varepsilon^*(v)\pi_\varepsilon^*(u) = \pi_\varepsilon^*(v)$ for every $\varepsilon \in \{0, 1\}^d$. Since the action of $G_{[d]}^\Delta$ is diagonal, then the projection of a minimal idempotent of $E(\mathbf{Q}_{T_1, \dots, T_d}(X), G_{[d]}^\Delta)$ is a minimal idempotent in $E(X, T_1, \dots, T_d)$. Then, we have that $\pi_\varepsilon^*(u)\pi_\varepsilon^*(v) = \pi_\varepsilon^*(u)$ for every $\varepsilon \in \{0, 1\}^d$. Since $\mathbf{Q}_{T_1, \dots, T_d}(X) \subseteq X^{[d]}$, we can view the elements of $E(\mathbf{Q}_{T_1, \dots, T_d}(X), \mathcal{G}_{T_1, \dots, T_d})$ as vectors of dimension 2^d . Thus the projections to the coordinates determine an element of $E(\mathbf{Q}_{T_1, \dots, T_d}(X), \mathcal{G}_{T_1, \dots, T_d})$. We have deduced that $uv = u$, which implies that u is a minimal idempotent in $E(\mathbf{Q}_{T_1, \dots, T_d}(X), \mathcal{G}_{T_1, \dots, T_d})$.

Let $x \in X$. Since (X, T_1, \dots, T_d) is minimal, there exists a minimal idempotent $u \in E(X, T_1, \dots, T_d)$ such that $ux = x$. Consider $u^{[d]} \in E(\mathbf{Q}_{T_1, \dots, T_d}(X), G_{[d]}^\Delta)$. We have that $u^{[d]}x^{[d]} = x^{[d]}$, so $x^{[d]}$ is an almost periodic point in $X^{[d]}$ (so in $\mathbf{Q}_{T_1, \dots, T_d}(X)$) under the action of $G_{[d]}^\Delta$. We observe that the

point $x^{[d]}$ is minimal under the action of $\mathcal{G}_{T_1, \dots, T_d}$ since, by the property proved before, $u^{[d]}$ is also a minimal idempotent in $E(\mathbf{Q}_{T_1, \dots, T_d}(X), \mathcal{G}_{T_1, \dots, T_d})$. As $\mathcal{O}(x^{[d]}, \mathcal{G}_{T_1, \dots, T_d}) = \mathbf{Q}_{T_1, \dots, T_d}(X)$, we conclude that $(\mathbf{Q}_{T_1, \dots, T_d}(X), \mathcal{G}_{T_1, \dots, T_d})$ is a minimal system.

Now, if (X, T_1, \dots, T_d) is distal, then $(X^{[d]}, G^{[d]})$ is also distal by Theorem 3 (1), which implies that $(X^{[d]}, \mathcal{G}_{T_1, \dots, T_d})$ is distal by Theorem 3 (4). Additionally, since $\mathbf{Q}_{T_1, \dots, T_d}(X)$ is invariant under $\mathcal{G}_{T_1, \dots, T_d}$ we also get that $(\mathbf{Q}_{T_1, \dots, T_d}(X), \mathcal{G}_{T_1, \dots, T_d})$ is distal. \square

The following proposition states that directional cubes pass through factors.

Proposition 8. *Let $\pi: Y \rightarrow X$ be a factor map between the \mathbb{Z}^d -systems (Y, T_1, \dots, T_d) and (X, T_1, \dots, T_d) . Then,*

$$\pi^{[d]}(\mathbf{Q}_{T_1, \dots, T_d}(Y)) = \mathbf{Q}_{T_1, \dots, T_d}(X),$$

where $\pi^{[d]}: Y^{[d]} \rightarrow X^{[d]}$ is defined from π coordinatewise.

Proof. It follows directly from definition that $\pi^{[d]}(\mathbf{Q}_{T_1, \dots, T_d}(Y)) \subseteq \mathbf{Q}_{T_1, \dots, T_d}(X)$. Then, if (X, T_1, \dots, T_d) is minimal, by Proposition 7 the proof follows directly. If not, let $\mathbf{x} \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ and take sequences $(x^i)_{i \in \mathbb{N}} \subseteq X$ and $(\mathbf{n}(i))_{i \in \mathbb{N}} \subseteq \mathbb{Z}^d$ such that

$$\forall \varepsilon \in \{0, 1\}^d, \quad x_\varepsilon = \lim_{i \rightarrow \infty} T_1^{n_1(i)\varepsilon_1} \dots T_d^{n_d(i)\varepsilon_d} x^i.$$

For each $i \in \mathbb{N}$ take $y^i \in \pi^{-1}(\{x^i\})$. By compactness, we can assume that $\lim_{i \rightarrow \infty} y^i = y$ and

$$\lim_{i \rightarrow \infty} T_1^{n_1(i)\varepsilon_1} \dots T_d^{n_d(i)\varepsilon_d} y^i = y_\varepsilon,$$

for all $\varepsilon \in \{0, 1\}^d$. Now, by continuity of π , we have that $\lim_{i \rightarrow \infty} \pi(y^i) = \pi(y) = x$ and

$$\pi(y_\varepsilon) = \pi \left(\lim_{i \rightarrow \infty} T_1^{n_1(i)\varepsilon_1} \dots T_d^{n_d(i)\varepsilon_d} y^i \right) = \lim_{i \rightarrow \infty} T_1^{n_1(i)\varepsilon_1} \dots T_d^{n_d(i)\varepsilon_d} \pi(y^i) = x_\varepsilon,$$

for all $\varepsilon \in \{0, 1\}^d$. Then, $\mathbf{y} \in \mathbf{Q}_{T_1, \dots, T_d}(Y)$ and $\pi^{[d]}(\mathbf{y}) = \mathbf{x} \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. This proves the lemma. \square

We finish with two crucial properties that appear in the different cube theories developed in topological dynamics [19, 1, 7, 14, 15, 16, 13]. We need to introduce some terminology.

Definition 9. Let $\mathbf{x} = (x_\varepsilon : \varepsilon \in \{0, 1\}^d)$, $\mathbf{y} = (y_\varepsilon : \varepsilon \in \{0, 1\}^d)$ in $X^{[d]}$ and $j \in [d]$. We say that \mathbf{x} and \mathbf{y} coincide in the j -upper face if

$$(x_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_j = 1) = (y_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_j = 1).$$

Similarly, we define when \mathbf{x} and \mathbf{y} coincide in their j -lower face. Also, we say that the j -upper face of \mathbf{x} coincides with the j -lower face of \mathbf{y} if

$$(x_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_j = 1) = (y_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_j = 0).$$

Definition 10 (Gluing operation). Let $\mathbf{x} = (x_\varepsilon : \varepsilon \in \{0, 1\}^d)$ and $\mathbf{y} = (y_\varepsilon : \varepsilon \in \{0, 1\}^d)$ in $X^{[d]}$. Suppose that the j -upper face of \mathbf{x} coincides with the j -lower face of \mathbf{y} . We define the gluing of \mathbf{x} and \mathbf{y} along the j -face as the point $\mathbf{z} = (z_\varepsilon : \varepsilon \in \{0, 1\}^d) \in X^{[d]}$ such that

$$z_\varepsilon = \begin{cases} x_\varepsilon & \varepsilon_j = 0, \\ y_\varepsilon & \varepsilon_j = 1. \end{cases}$$

Definition 11 (Crucial properties). Let (X, T_1, \dots, T_d) be a \mathbb{Z}^d -system.

- (1) *Unique closing parallelepiped property.* We say that (X, T_1, \dots, T_d) has the *unique closing parallelepiped property* if whenever $\mathbf{x}, \mathbf{y} \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ have $2^d - 1$ coordinates in common then $\mathbf{x} = \mathbf{y}$.
- (2) *Gluing property.* We say that (X, T_1, \dots, T_d) has the *gluing property* if for every $j \in [d]$, if $\mathbf{x}, \mathbf{y} \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ are such that the j -upper face of \mathbf{x} coincides with the j -lower face of \mathbf{y} , then the point given by the gluing of \mathbf{x} and \mathbf{y} along their j -face belongs to $\mathbf{Q}_{T_1, \dots, T_d}(X)$.

In the measure theoretical category the *unique closing parallelepiped property* is exactly the property that characteristic factors for multiple averages for systems with commuting transformations satisfy [17, Section 2.6]. In the minimal distal case the gluing property always holds.

Lemma 12 (Gluing Lemma). *Let (X, T_1, \dots, T_d) be a minimal distal \mathbb{Z}^d -system. Then, (X, T_1, \dots, T_d) has the gluing property.*

Proof. We write the proof for $j = d$. The general proof is the same modulo a small change of notation. Let $\mathbf{x}', \mathbf{x}'', \mathbf{y}'' \in X^{[d-1]}$ be such that $\mathbf{x} = (\mathbf{x}', \mathbf{x}''), \mathbf{y} = (\mathbf{x}'', \mathbf{y}'') \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. By definition, the point given by the gluing of \mathbf{x} and \mathbf{y} along their d -face is $\mathbf{z} = (\mathbf{x}', \mathbf{y}'')$.

Pick any $a \in X$. Then, by Proposition 6 (1), $(a^{[d-1]}, a^{[d-1]}) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. By Proposition 7, there exists a sequence $(g_n)_{n \in \mathbb{N}} = ((g'_n, g''_n))_{n \in \mathbb{N}} \in \mathcal{G}_{T_1, \dots, T_d}$ such that $g_n(\mathbf{x}', \mathbf{x}'') = (g'_n \mathbf{x}', g''_n \mathbf{x}'') \rightarrow (a^{[d-1]}, a^{[d-1]})$. We can assume, by compactness, that $g''_n \mathbf{y}'' \rightarrow \mathbf{u}$ and thus $(g'_n \mathbf{x}', g''_n \mathbf{y}'') \rightarrow (a^{[d-1]}, \mathbf{u}) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. Now, we have that $g_n(\mathbf{x}', \mathbf{y}'') = (g'_n \mathbf{x}', g''_n \mathbf{y}'') \rightarrow (a^{[d-1]}, \mathbf{u})$, and this point belongs to the closed orbit of $(\mathbf{x}', \mathbf{y}'')$ under $\mathcal{G}_{T_1, \dots, T_d}$. By distality (see Proposition 7), this orbit is minimal and thus it follows that $(\mathbf{x}', \mathbf{y}'')$ is in the closed orbit of $(a^{[d-1]}, \mathbf{u})$, which implies that $\mathbf{z} = (\mathbf{x}', \mathbf{y}'') \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. \square

Remark 13. Remark that the same property holds for a subset of the transformations. That is, if (X, T_1, \dots, T_d) is a minimal distal \mathbb{Z}^d -system, then $(X, T_{i_1}, T_{i_2}, \dots, T_{i_k})$, $1 \leq k \leq d$, $1 \leq i_1 < i_2 < \dots < i_k \leq d$, also has the gluing property in its space of cubes $\mathbf{Q}_{T_{i_1}, T_{i_2}, \dots, T_{i_k}}(X)$. The proof of this is exactly the same as in Lemma 12, but considering the system $\mathbf{Q}_{T_{i_1}, T_{i_2}, \dots, T_{i_k}}(X)$ endowed with the action generated by $\mathcal{G}_{T_{i_1}, \dots, T_{i_k}}$ and $G_{[k]}^\Delta$ (instead of just $\mathcal{G}_{T_{i_1}, \dots, T_{i_k}}$). We point out that this joint action is minimal, since it is the projection of $(\mathbf{Q}_{T_1, T_2, \dots, T_d}(X), \mathcal{G}_{T_1, \dots, T_d})$ onto a k dimensional face (recall Proposition 6 (4)).

3.2. Symmetries of the euclidean cube and directional cubes. As was mentioned above, a different choice of the order of the transformations in a \mathbb{Z}^d -system produces a different set of dynamical cubes. Thus, in order to simplify some proofs in the next sections we introduce the terminology proposed in [18] about Euclidean permutations of the d -dimensional cube $\{0, 1\}^d$.

3.2.1. Digit permutations. Let $d \geq 1$ be an integer. Given a permutation $\sigma: [d] \rightarrow [d]$ we consider the symmetry that it induces in the euclidean cube $\{0, 1\}^d$, which is defined by setting $\sigma(\varepsilon)_i = \varepsilon_{\sigma(i)}$ for $i \in [d]$. Here we slightly abuse of the notation and continue calling this transformation σ . Let $\sigma_*: X^{[d]} \rightarrow X^{[d]}$ denote the transformation

$$\sigma_*((x_\varepsilon: \varepsilon \in \{0, 1\}^d)) = (x_{\sigma(\varepsilon)}: \varepsilon \in \{0, 1\}^d).$$

We refer to it as a *digit permutation* of $X^{[d]}$.

Lemma 14. *Let (X, T_1, \dots, T_d) be a \mathbb{Z}^d -system and $\sigma: [d] \rightarrow [d]$ be a permutation. Then,*

$$\sigma_*(\mathbf{Q}_{T_{\sigma(1)}, \dots, T_{\sigma(d)}}(X)) = \mathbf{Q}_{T_1, \dots, T_d}(X)$$

Proof. First we prove that for $x \in X$ and $(n_1, \dots, n_d) \in \mathbb{Z}^d$ the point $\sigma_*((T_{\sigma(1)}^{n_1 \varepsilon_1} \dots T_{\sigma(d)}^{n_d \varepsilon_d} x : \varepsilon \in \{0, 1\}^d))$ belongs to $\mathbf{Q}_{T_1, \dots, T_d}(X)$. This will prove that $\sigma_*(\mathbf{Q}_{T_{\sigma(1)}, \dots, T_{\sigma(d)}}(X)) \subseteq \mathbf{Q}_{T_1, \dots, T_d}(X)$. The other inclusion follows analogously by considering the inverse of σ .

Let $\eta \in \{0, 1\}^d$. Since $\sigma(\eta) = \eta_{\sigma(1)} \dots \eta_{\sigma(d)}$ we have that

$$\begin{aligned} \sigma_*((T_{\sigma(1)}^{n_1 \varepsilon_1} \dots T_{\sigma(d)}^{n_d \varepsilon_d} x : \varepsilon \in \{0, 1\}^d))_\eta &= (T_1^{n_1 \varepsilon_1} \dots T_d^{n_d \varepsilon_d} x : \varepsilon \in \{0, 1\}^d)_{\sigma(\eta)}, \\ &= T_{\sigma(1)}^{n_1 \eta_{\sigma(1)}} \dots T_{\sigma(d)}^{n_d \eta_{\sigma(d)}} x, \\ &= T_1^{m_1 \eta_1} \dots T_d^{m_d \eta_d} x, \end{aligned}$$

where $m_i = n_{\sigma^{-1}(i)}$ for all $i \in [d]$. It is then clear that $\sigma_*((T_{\sigma(1)}^{n_1 \varepsilon_1} \dots T_{\sigma(d)}^{n_d \varepsilon_d} x : \varepsilon \in \{0, 1\}^d))$ belongs to $\mathbf{Q}_{T_1, \dots, T_d}(X)$. \square

We remark that $\mathbf{Q}_{T_1, \dots, T_d}(X)$ and $\mathbf{Q}_{T_{\sigma(1)}, \dots, T_{\sigma(d)}}(X)$ do not need to (and usually do not) coincide. This lack of symmetry is one of the main differences with the case where all transformations are powers of a given transformation T , *i.e.*, when $T_i = T^i$ for $i \in [d]$. In the latter case, all digit symmetries hold. We think this is one of the reasons why we cannot expect the unique closing parallelepiped property to imply the strong algebraic consequences that one obtains in the single transformation case [18, 19], or more generally, when the whole group is acting in all directions [1, 16, 13]. Nevertheless, we still have a reduced number of symmetries that leave invariant $\mathbf{Q}_{T_1, \dots, T_d}(X)$ that we describe in the next section.

3.2.2. Reflections. For every $j \in [d]$ we consider the following Euclidean permutation:

$$\begin{aligned} \Phi_j: \quad \{0, 1\}^d &\rightarrow \{0, 1\}^d \\ \varepsilon &\mapsto \Phi_j(\varepsilon) = \begin{cases} \varepsilon_k & \text{if } k \neq j, \\ 1 - \varepsilon_k & \text{if } k = j. \end{cases} \end{aligned}$$

The transformation Φ_j is the one that takes the upper j -face and turn it into the lower j -face and vice versa. We naturally define the induced action $\Phi_{j*}: X^{[d]} \rightarrow X^{[d]}$ by setting

$$(\Phi_{j*}(\mathbf{x}))_\varepsilon = x_{\Phi_j(\varepsilon)}, \text{ for all } \varepsilon \in \{0, 1\}^d,$$

where $\mathbf{x} = (x_\varepsilon : \varepsilon \in \{0, 1\}^d)$. We have,

Lemma 15 (Face permutation invariance). *For every $j \in [d]$ the transformation Φ_{j*} leaves invariant $\mathbf{Q}_{T_1, \dots, T_d}(X)$.*

Proof. Since Φ_{j*} is clearly continuous, it suffices to show that for every $x \in X$ and $(n_1, \dots, n_d) \in \mathbb{Z}^d$ the image of $(T_1^{n_1 \varepsilon_1} \dots T_d^{n_d \varepsilon_d} x : \varepsilon \in \{0, 1\}^d)$ under Φ_{j*} belongs to $\mathbf{Q}_{T_1, \dots, T_d}(X)$. We have that

$$\begin{aligned} \Phi_{j*}((T_1^{n_1 \varepsilon_1} \dots T_d^{n_d \varepsilon_d} x : \varepsilon \in \{0, 1\}^d)) &= (T_1^{n_1 \varepsilon_1} \dots T_j^{n_j(1-\varepsilon_j)} \dots T_d^{n_d \varepsilon_d} x : \varepsilon \in \{0, 1\}^d) \\ &= (T_1^{n_1 \varepsilon_1} \dots T_j^{(-n_j)\varepsilon_j} \dots T_d^{n_d \varepsilon_d} (T^{n_j} x) : \varepsilon \in \{0, 1\}^d) \end{aligned}$$

and this point clearly belongs to $\mathbf{Q}_{T_1, \dots, T_d}(X)$ since $T^{n_j} x \in X$ and $(n_1, \dots, -n_j, \dots, n_d) \in \mathbb{Z}^d$. \square

Definition 16 (The insertion procedure). Let $\mathbf{x} = (x_\varepsilon : \varepsilon \in \{0, 1\}^d)$ and $\mathbf{y} = (y_\varepsilon : \varepsilon \in \{0, 1\}^d)$ be points in $X^{[d]}$. We say that the point $\mathbf{z} = (z_\varepsilon : \varepsilon \in \{0, 1\}^d) \in X^{[d]}$ is the result of inserting the j -upper face of \mathbf{x} into the j -lower face of \mathbf{y} if

$$z_\varepsilon = \begin{cases} y_\varepsilon & \text{if } \varepsilon_j = 1, \\ x_{\Phi_j(\varepsilon)} & \text{if } \varepsilon_j = 0. \end{cases}$$

Similarly one defines the insertion of the j -lower face of \mathbf{x} into the j -upper face of \mathbf{y}

From Lemma 15 and the gluing property we get,

Lemma 17. *Let (X, T_1, \dots, T_d) be a \mathbb{Z}^d -system where the gluing property holds (in particular when the system is minimal and distal). If $\mathbf{x}, \mathbf{y} \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ coincide in their j -upper face, then the point obtained by inserting the j -lower face of \mathbf{x} into the j -upper face of \mathbf{y} also belongs to $\mathbf{Q}_{T_1, \dots, T_d}(X)$.*

4. THE (T_1, \dots, T_d) -REGIONALLY PROXIMAL RELATION

Let (X, T_1, \dots, T_d) be a \mathbb{Z}^d -system. We define a relation in X associated with the cube structure $\mathbf{Q}_{T_1, \dots, T_d}(X)$ established in Section 3. This relation generalizes the definition given by Donoso and Sun for two commuting transformations in [7]. The motivation is to find precisely the relations producing the maximal factors with the unique closing parallelepiped property. We start defining for $j \in [d]$ the following transformations:

$$\begin{aligned} \Psi_j^0: \quad \{0, 1\}^{d-1} &\rightarrow \{0, 1\}^d \\ \varepsilon &\mapsto \Psi_j^0(\varepsilon) = \varepsilon_1 \varepsilon_2 \dots \varepsilon_{j-1} 0 \varepsilon_j \dots \varepsilon_d, \\ \Psi_j^1: \quad \{0, 1\}^{d-1} &\rightarrow \{0, 1\}^d \\ \varepsilon &\mapsto \Psi_j^1(\varepsilon) = \varepsilon_1 \varepsilon_2 \dots \varepsilon_{j-1} 1 \varepsilon_j \dots \varepsilon_d. \end{aligned}$$

The maps Ψ_j^0 and Ψ_j^1 are used to replicate a face. Let $j \in [d]$ and $\mathbf{x} \in X^{[d-1]}$. If we define $\mathbf{y} \in X^{[d]}$ by $y_\varepsilon = x_\eta$ if $\varepsilon = \Psi_j^0(\eta) \vee \varepsilon = \Psi_j^1(\eta)$, we have that both the j -lower face of \mathbf{y} and the j -upper face of \mathbf{y} are equal to \mathbf{x} , *i.e.*,

$$(y_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_j = 0) = (y_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_j = 1) = \mathbf{x} \in X^{[d-1]}.$$

Definition 18. For $x, y \in X$, $\mathbf{a}_* \in X_*^{[d-1]}$ and $j \in [d]$, define $\mathbf{z}(x, y, \mathbf{a}_*, j) \in X^{[d]}$ with coordinates

$$(\mathbf{z}(x, y, \mathbf{a}_*, j))_\varepsilon = \begin{cases} x & \text{if } \varepsilon = \emptyset, \\ y & \text{if } \varepsilon = \{j\}, \\ (\mathbf{a}_*)_\eta & \text{if } \varepsilon = \Psi_j^0(\eta) \vee \varepsilon = \Psi_j^1(\eta). \end{cases}$$

Definition 19. For each $j \in [d]$ we define the T_j -regionally proximal relation as

$$\mathcal{R}_{T_j}(X) = \left\{ (x, y) \in X \times X : \exists \mathbf{a}_* \in X_*^{[d-1]}, \mathbf{z}(x, y, \mathbf{a}_*, j) \in \mathbf{Q}_{T_1, \dots, T_d}(X) \right\}.$$

For $j \in [d]$ let $\sigma: [d] \rightarrow [d]$ be the permutation that interchanges j and 1 (and is the identity everywhere else). By Lemma 14, we have that σ_* maps $\mathbf{Q}_{T_j, \dots, T_1, \dots, T_d}(X)$ onto $\mathbf{Q}_{T_1, \dots, T_d}(X)$. It then follows that $(x, y) \in \mathcal{R}'_{T_j}$ if and only if $(x, y) \in \mathcal{R}_{T_j}$, where \mathcal{R}'_{T_j} is the same relation but in the cube $\mathbf{Q}_{T_j, \dots, T_1, \dots, T_d}(X)$ (so the order of the transformations is changed, interchanging T_1 and T_j). The advantage of doing so is that (modulo changing the order of the transformations) we can assume that the points (x, y) are in R_{T_1} , where T_1 is the first transformation. This point of view is only esthetical, but it will lighten some proofs notations in the next sections.

Finally, we define the (T_1, \dots, T_d) -regionally proximal relation as,

$$\mathcal{R}_{T_1, \dots, T_d}(X) = \bigcap_{j=1}^d \mathcal{R}_{T_j}(X).$$

As an example, for $d = 3$ the relations are defined as follows:

$$\begin{aligned} \mathcal{R}_{T_1}(X) &= \{(x, y) \in X \times X : (x, y, a, a, b, b, c, c) \in \mathbf{Q}_{T_1, T_2, T_3}(X) \text{ for some } a, b, c \in X\}, \\ \mathcal{R}_{T_2}(X) &= \{(x, y) \in X \times X : (x, a, y, a, b, c, b, c) \in \mathbf{Q}_{T_1, T_2, T_3}(X) \text{ for some } a, b, c \in X\}, \\ \mathcal{R}_{T_3}(X) &= \{(x, y) \in X \times X : (x, a, b, c, y, a, b, c) \in \mathbf{Q}_{T_1, T_2, T_3}(X) \text{ for some } a, b, c \in X\}. \end{aligned}$$

Graphically we represent these relations as follows:




FIGURE 1. Representation of the relations $\mathcal{R}_{T_1}(X)$, $\mathcal{R}_{T_2}(X)$ and $\mathcal{R}_{T_3}(X)$ in the case $d = 3$.

It is easy to see that these relations are reflexive, symmetric, closed and invariant under the action of $\langle T_1, \dots, T_d \rangle$. In the next section we prove that these relations are transitive in the minimal distal case, but we do not know yet if they are transitive in the general minimal case.

The next proposition shows that the relations $\mathcal{R}_{T_j}(X)$ for \mathbb{Z}^d -systems with the unique closing parallelepiped property are trivial.

Proposition 20. *Let (X, T_1, \dots, T_d) be a \mathbb{Z}^d -system with the unique closing parallelepiped property. Then, for every $j \in [d]$ we have that $\mathcal{R}_{T_j}(X) = \Delta_X$.*

Proof. Take $(x, y) \in \mathcal{R}_{T_j}(X)$ for a given $j \in [d]$. Then, by definition, there exists $\mathbf{a}_* \in X_*^{[d-1]}$ such that $\mathbf{z}(x, y, \mathbf{a}_*, j) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. By Proposition 6 (4), we have that $(x, \mathbf{a}_*) \in \mathbf{Q}_{T_1, \dots, T_{j-1}, T_{j+1}, \dots, T_d}(X)$.

Define $\mathbf{w} \in X^{[d]}$ as

$$w_\varepsilon = \begin{cases} x & \text{if } \varepsilon = \emptyset, \\ x & \text{if } \varepsilon = \{j\}, \\ (\mathbf{a}_*)_\eta & \text{if } \varepsilon = \Psi_j^0(\eta) \vee \varepsilon = \Psi_j^1(\eta). \end{cases}$$

By Proposition 6 (5), we have that $\mathbf{w} \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ and therefore the unique closing parallelepiped property implies that $x = y$. \square

The proof of Proposition 20 in the case $d = 3$ and $j = 1$ is illustrated in the next figure.




FIGURE 2. Illustration of the proof of Proposition 20 in the case $d = 3$ for $j = 1$. We have the existence of a cube like in (a) because $(x, y) \in \mathcal{R}_{T_1}(X)$.

In the literature [19, 23, 7, 14, 15, 16, 13], when defining these types of relations one also looks for the following properties: (i) if $\pi: Y \rightarrow X$ is a factor map between the minimal \mathbb{Z}^d -systems (Y, T_1, \dots, T_d) and (X, T_1, \dots, T_d) we would like to have $\pi^{[d]}(\mathcal{R}_{T_j}(Y)) = \mathcal{R}_{T_j}(X)$ for every $j \in [d]$ and

$$\pi^{[d]}(\mathcal{R}_{T_1, \dots, T_d}(Y)) = \mathcal{R}_{T_1, \dots, T_d}(X);$$

and (ii) we also would like to have that $\mathcal{R}_{T_1, \dots, T_d}(X)$ is an equivalence relation to study the factor system $(X/\mathcal{R}_{T_1, \dots, T_d}(X), T_1, \dots, T_d)$. Nevertheless, for the moment we cannot prove these properties. In Section 5 we will prove that they hold for minimal distal systems.

5. CHARACTERIZING THE (T_1, \dots, T_d) -REGIONALLY PROXIMAL RELATION FOR DISTAL \mathbb{Z}^d -SYSTEMS

We study the properties of the (T_1, \dots, T_d) -regionally proximal relation for minimal distal systems that will be used in the proof of Theorem 1.

We start with the following theorem, which was proved in the case $d = 2$ by Donoso and Sun in [7].

Theorem 21. *Let (X, T_1, \dots, T_d) be a \mathbb{Z}^d -system where the gluing property holds. Consider $x, y \in X$. The following statements are equivalent:*

- (1) $(x, y) \in \mathcal{R}_{T_1, \dots, T_d}(X)$.
- (2) $(x, y_*^{[d]}) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$.
- (3) There exists $\mathbf{a}_* \in X_*^{[d]}$ such that $(x, \mathbf{a}_*), (y, \mathbf{a}_*) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$.
- (4) For every $\mathbf{a}_* \in X_*^{[d]}$, $(x, \mathbf{a}_*) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ if and only if $(y, \mathbf{a}_*) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$.
- (5) There exists $j \in [d]$ such that $(x, y) \in \mathcal{R}_{T_j}(X)$.

Proof. We start with the easy implications and then show the two main ones.

(1) \implies (5) follows from definition.

(2) \implies (1). Let $j \in [d]$ and $\mathbf{a}_* = y_*^{[d-1]} \in X_*^{[d-1]}$, then $\mathbf{z}(x, y, \mathbf{a}_*, j) = (x, y_*^{[d]}) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. Hence $(x, y) \in \mathcal{R}_{T_j}(X)$ and since j is arbitrary we get that $(x, y) \in \mathcal{R}_{T_1, \dots, T_d}(X)$.

(4) \implies (2). Since $y^{[d]} = (y, y_*^{[d]}) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$, we can use $\mathbf{a}_* = y_*^{[d]}$ in property (4) to get $(x, y_*^{[d]}) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$.

(2) \implies (3) It follows directly since $y^{[d]} = (y, y_*^{[d]})$ and $(x, y_*^{[d]}) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$.

(3) \implies (5) Let $\mathbf{a}_* \in X_*^{[d]}$ be such that $(x, \mathbf{a}_*), (y, \mathbf{a}_*) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. Then the d -th upper faces of (x, \mathbf{a}_*) and (y, \mathbf{a}_*) coincide. By Lemma 17 the point obtained by inserting the lower d -th face of (y, \mathbf{a}_*) into the d -th upper of (x, \mathbf{a}_*) also belongs to $\mathbf{Q}_{T_1, \dots, T_d}(X)$. But this point can be written as (x, b_*, y, b_*) for some $b_* \in X^{[d-1]}$. This implies that $(x, y) \in \mathcal{R}_{T_d}(X)$.

We now prove the main implications, namely (5) \implies (2) and (2) \implies (4).

(5) \implies (2). The idea of the proof is to use the gluing property to construct a sequence of points $\mathbf{z}^1, \dots, \mathbf{z}^d \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ so that we increase the number of times that y appears as a coordinate of \mathbf{z}^i until getting $\mathbf{z}^d = (x, y_*^{[d]}) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. We suggest the reader to look at the illustration of the proof in Figure 3 while reading it.

By the discussion right after Definition 19 we can assume for the moment that $(x, y) \in \mathcal{R}_{T_1}(X)$. So, there exists $\mathbf{a}_* \in X^{[d-1]}$ for which $\mathbf{z}(x, y, \mathbf{a}_*, 1) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. We call $\mathbf{z}^1 = \mathbf{z}(x, y, \mathbf{a}_*, 1)$.

Let \mathbf{v}^1 be the duplication of the 1-th upper face of \mathbf{z}^1 . Then \mathbf{z}^1 and \mathbf{v}^1 coincide in their 2-th upper face. By Lemma 17 we can insert the 2-th lower face of \mathbf{v}^1 into the 2-th upper face of \mathbf{z}^1 , obtaining a point \mathbf{z}^2 that belongs to $\mathbf{Q}_{T_1, \dots, T_d}(X)$. The point \mathbf{z}^2 has the property that

$$z_\varepsilon = y \text{ if } \varepsilon \neq \emptyset \text{ and } \varepsilon_j = 0 \text{ for all } j \geq 3.$$

Otherwise saying, the lower face of codimension $d - 2$ of \mathbf{z}^2 (*i.e.*, the one associated to the face $\{0, 1\}^2 \times \{0\} \times \dots \times \{0\}$) is (x, y, y, y) .

Now suppose that for $1 \leq k \leq d - 1$ we have constructed $\mathbf{z}^k \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ such that its lower face of codimension $d - k$ (*i.e.*, the face associated to $\{0, 1\}^k \times \{0\}^{d-k}$) coincides with $(x, y_*^{[k]})$. Let \mathbf{v}^k be the duplication of the k -th upper face of \mathbf{z}^k . Then \mathbf{v}^k and \mathbf{z}^k coincide in their $(k + 1)$ -th upper face. By Lemma 17 we can insert the $(k + 1)$ -th lower face of \mathbf{v}^k into the $(k + 1)$ -th upper face of \mathbf{z}^k , obtaining a point \mathbf{z}^{k+1} that belongs to $\mathbf{Q}_{T_1, \dots, T_d}(X)$. This point satisfies that its lower face of codimension $d - (k + 1)$ (*i.e.*, the one associated to the face $\{0, 1\}^{k+1} \times \{0\}^{d-(k+1)}$) is $(x, y_*^{[k+1]})$. With this procedure we build a sequence of points which ends up in $\mathbf{z}^d = (x, y_*^{[d]})$ as desired.

As was stated before, the assumption $(x, y) \in \mathcal{R}_{T_1}(X)$ is irrelevant. Indeed, modulo changing the order of the transformations we get that $(x, y_*^{[d]}) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. But, for every permutation $\sigma: [d] \rightarrow [d]$ we have that $\sigma_*(x, y_*^{[d]}) = (x, y_*^{[d]})$. Hence, this point also belongs to $\mathbf{Q}_{T_{\sigma(1)}, \dots, T_{\sigma(d)}}(X)$ by Lemma 14. This shows that the order considered for the transformations is not important.

We remark that a consequence of this implication is that $(x, y_*^{[d]}) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ is equivalent to $(y, x_*^{[d]}) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$.

(2) \implies (4). Here we also advice the reader to look at Figure 4 while reading the proof. Assume that $(x, y_*^{[d]}) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ and let $\mathbf{a}_* \in X_*^{[d]}$ be such that $\mathbf{z} = (y, \mathbf{a}_*) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. We have to show that $(x, \mathbf{a}_*) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. The other implication will follow by the symmetry implied by previous implication remarked above.

Consider the d points $\mathbf{w}^1, \dots, \mathbf{w}^d \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ such that for every $1 \leq k \leq d$ the point \mathbf{w}^k is the replication of the lower face of dimension k of \mathbf{z} , (the one associated to $\{0, 1\}^k \times \{0\}^{d-k}$ or to $[k]$ in the subset notation). That means that for every $\varepsilon = \varepsilon_1 \dots \varepsilon_k \varepsilon_{k+1} \dots \varepsilon_d \in \{0, 1\}^d$

$$w_\varepsilon^k = z_{\varepsilon_1 \dots \varepsilon_k 0 \dots 0}.$$

Such points can be obtained using Proposition 6(5). We remark that $\mathbf{w}^d = \mathbf{z}$.

Next we construct $d + 1$ points $\mathbf{v}^1, \dots, \mathbf{v}^{d+1} \in \mathbf{Q}_{T_1, \dots, T_d}(X)$, starting with $\mathbf{v}^1 = (x, y_*^{[d]})$ and transforming it step by step into $\mathbf{v}^{d+1} = (x, \mathbf{a}_*)$.

We have that the 1-th upper face of \mathbf{v}^1 and the 1-th lower face of \mathbf{w}^1 coincide (they are equal to $y^{[d-1]}$). Hence by Lemma 12, the point \mathbf{v}^2 obtained by gluing the 1-th lower face of \mathbf{v}^1 and the 1-th upper face of \mathbf{w}^1 belongs to $\mathbf{Q}_{T_1, \dots, T_d}(X)$. We claim that the point \mathbf{v}^2 is equal to \mathbf{w}^1 everywhere but in the $\vec{0}$ coordinate, where it is equal to x . To see this, note that in the lower 1-th face, *i.e.*, if $\varepsilon_1 = 0$ and $\varepsilon \neq \vec{0}$ we have

$$v_\varepsilon^2 = v_\varepsilon^1 = y = w_\varepsilon^1,$$

while if $\varepsilon_1 = 1$ we have

$$v_\varepsilon^2 = w_\varepsilon^1,$$

proving the claim.




FIGURE 3. Illustration of the proof of (5) \implies (2) of Theorem 21 for the case $d = 3$ and $j = 1$. Always the grey faces are duplicated and the color faces serve to pass to the next step using Lemma 17.

Now assume that we have constructed \mathbf{v}^k such that it coincides with \mathbf{w}^{k-1} up to the $\vec{0}$ coordinate, where it is equal to x , *i.e.*,

$$v_{\vec{0}}^k = x \quad \text{and} \quad v_{\varepsilon}^k = w_{\varepsilon}^{k-1} \text{ for all } \varepsilon \in \{0, 1\}^d \setminus \{\vec{0}\}.$$

We now construct \mathbf{v}^{k+1} satisfying the corresponding properties. We have that the k -th lower face of \mathbf{w}^k coincides with the k -th upper face of \mathbf{v}^k . To see this, note that

$$w_{\varepsilon_1 \dots \varepsilon_{k-1} 0 \varepsilon_{k+1} \dots \varepsilon_d}^k = z_{\varepsilon_1 \dots \varepsilon_{k-1} 0 \dots 0}$$

and

$$v_{\varepsilon_1 \dots \varepsilon_{k-1} 1 \varepsilon_{k+1} \dots \varepsilon_d}^k = w_{\varepsilon_1 \dots \varepsilon_{k-1} 1 \varepsilon_{k+1} \dots \varepsilon_d}^{k-1} = z_{\varepsilon_1 \dots \varepsilon_{k-1} 0 \dots 0}.$$

Hence the gluing property ensures the existence of a point $\mathbf{v}^{k+1} \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ whose k -th lower face is equal to the one of \mathbf{v}^k and its k -th upper face is equal to the one of \mathbf{w}^k . We now check that this point is equal to \mathbf{w}^k everywhere but in the $\vec{0}$ coordinate.


In the k -th lower face, *i.e.*, if $\varepsilon_k = 0$ and $\varepsilon \neq \vec{0}$ we have

$$v_{\varepsilon}^{k+1} = v_{\varepsilon}^k = w_{\varepsilon_1 \dots \varepsilon_k 0 \dots 0}^{k-1} = z_{\varepsilon_1 \dots \varepsilon_{k-1} 0 \dots 0} = z_{\varepsilon_1 \dots \varepsilon_{k-1} \varepsilon_k 0 \dots 0} = w_{\varepsilon}^k,$$

while in the k -th upper face, *i.e.*, if $\varepsilon_k = 1$, we have by definition that

$$v_{\varepsilon}^{k+1} = w_{\varepsilon}^k.$$

We end up with \mathbf{v}^{d+1} which is equal to $\mathbf{w}^d = \mathbf{z}$ everywhere but in the $\vec{0}$ coordinate, where it is equal to x . That is, $\mathbf{v}^{d+1} = (x, \mathbf{a}_*) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$, finishing the proof. \square


 FIGURE 4. Illustration of the proof of (2) \implies (4) in the case $d = 3$.

We can prove that $\mathcal{R}_{T_1, \dots, T_d}(X)$ is an equivalence relation in the minimal distal case.

Theorem 22. *Let (X, T_1, \dots, T_d) be a minimal distal \mathbb{Z}^d -system. Then, $\mathcal{R}_{T_1, \dots, T_d}(X)$ is a closed and invariant equivalence relation on X .*

Proof. It suffices to prove the transitivity of $\mathcal{R}_{T_1, \dots, T_d}(X)$. Let $(x, y), (y, z) \in \mathcal{R}_{T_1, \dots, T_d}(X)$. By Theorem 21(2) we have that $(y, z_*^{[d]}) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. Now, by Theorem 21(4) $(x, z_*^{[d]}) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ and thus $(x, z) \in \mathcal{R}_{T_1, \dots, T_d}(X)$. \square

We also have the following property which allows us to lift the relation $\mathcal{R}_{T_1, \dots, T_d}(X)$ by a factor map and to prove our main theorem.

Theorem 23. *Let $\pi: Y \rightarrow X$ be a factor map between the \mathbb{Z}^d -systems (Y, T_1, \dots, T_d) and (X, T_1, \dots, T_d) . If (X, T_1, \dots, T_d) is minimal and distal, then $\pi \times \pi(\mathcal{R}_{T_1, \dots, T_d}(Y)) = \mathcal{R}_{T_1, \dots, T_d}(X)$.*

Proof. The proof is similar to that of Theorem 6.4 in [23]. It is a direct consequence of the definition that $\pi \times \pi(\mathcal{R}_{T_1, \dots, T_d}(Y)) \subseteq \mathcal{R}_{T_1, \dots, T_d}(X)$ and so we are left to prove the reverse inclusion. Let $(x, z) \in \mathcal{R}_{T_1, \dots, T_d}(X)$. By Theorem 21 we have that $(x, \dots, x, z) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. Replacing Y by a minimal subsystem if needed, we may assume that (Y, T_1, \dots, T_d) is minimal.

By Proposition 8 there exists $\mathbf{y}^0 = (y_1, \mathbf{a}_*) \in \mathbf{Q}_{T_1, \dots, T_d}(Y)$ such that $\pi^{[d]}(\mathbf{y}^0) = (x, \dots, x, z)$. Let $\mathbf{y}_{\mathbf{I}}^0 = (y_\varepsilon^0 : \varepsilon_d = 0)$ and $\mathbf{y}_{\mathbf{II}}^0 = (y_\varepsilon^0 : \varepsilon_d = 1)$ the d -th lower and upper faces of \mathbf{y}^0 respectively. We have that $\mathbf{y}_{\mathbf{I}}^0 \in \mathbf{Q}_{T_1, \dots, T_{d-1}}(Y)$. Let $\mathcal{G}'_{T_1, \dots, T_{d-1}}$ denote the projection of the action of $\mathcal{G}_{T_1, \dots, T_d}$ onto $\mathbf{Q}_{T_1, \dots, T_{d-1}}(Y)$. That is, $\mathcal{G}'_{T_1, \dots, T_{d-1}}$ is generated by $\mathcal{G}_{T_1, \dots, T_{d-1}}$ and the diagonal action $G_{[d-1]}^\Delta$. By minimality of $(\mathbf{Q}_{T_1, \dots, T_d}(Y), \mathcal{G}_{T_1, \dots, T_d})$ we have that $(\mathbf{Q}_{T_1, \dots, T_{d-1}}(Y), \mathcal{G}'_{T_1, \dots, T_{d-1}})$ is minimal so there exists a sequence $(\mathcal{G}_i^0)_{i \geq 1}$ in $\mathcal{G}'_{T_1, \dots, T_{d-1}}$ such that $\mathcal{G}_i^0 \mathbf{y}_{\mathbf{I}}^0 \rightarrow y_1^{[d-1]}$. By compactness we can assume that $\mathcal{G}_i^0 \mathbf{y}_{\mathbf{II}}^0 \rightarrow \mathbf{z}_{\mathbf{II}}^0$ for some $\mathbf{z}_{\mathbf{II}}^0 \in \mathbf{Q}_{T_1, \dots, T_{d-1}}(Y)$. Let $\tilde{\mathcal{G}}_i^0 = (\mathcal{G}_i^0, \mathcal{G}_i^0)$, i.e., the transformation that consists in applying \mathcal{G}_i^0 to both the upper and lower d -th faces of a point in $\mathbf{Q}_{T_1, \dots, T_d}(Y)$. Thus, we have that $\tilde{\mathcal{G}}_i^0 \mathbf{y}^0 = (\mathcal{G}_i^0 \mathbf{y}_{\mathbf{I}}^0, \mathcal{G}_i^0 \mathbf{y}_{\mathbf{II}}^0)$ converges to a point \mathbf{y}^1 in $\mathbf{Q}_{T_1, \dots, T_d}(Y)$ whose d -th lower face is equal to $y_1^{[d-1]}$ and whose d -th upper face is equal to $\mathbf{z}_{\mathbf{II}}^0$. Notice that

$$\pi^{[d]}(\mathbf{y}^1) = \mathbf{x}^1 = (x, \dots, x, z_1),$$

for some $z_1 \in X$.

Claim: The point (x, z_1) belongs to $\overline{\mathcal{O}((x, z), G_{[2]}^\Delta)}$.

To prove the claim, remark that the projection of $\tilde{\mathcal{G}}_i^0 \mathbf{y}^0$ onto the coordinates $011\dots 1$ and $1\dots 1$ is of the form (gx, gz) for some $g \in G$. By construction these coordinates converge to (x, z_1) , proving the claim.

Now assume we have constructed points $\mathbf{y}^j \in \mathbf{Q}_{T_1, \dots, T_d}(Y)$ for $1 \leq j \leq d-1$ with $\pi^{[d]}(\mathbf{y}^j) = \mathbf{x}^j$ such that $y_\varepsilon^j = y_1$ if there exists some k with $d-j+1 \leq k \leq d$ and $\varepsilon_k = 0$ (i.e., \mathbf{y}^j coincides with $y_1^{[d]}$ in all k -th lower faces for $d-j+1 \leq k \leq d$), $z_j = x_{[d]}^j$, $x_\varepsilon^j = x$ for all $\varepsilon \neq [d]$ and $(x, z_j) \in \overline{\mathcal{O}((x, z_{j-1}), G_{[2]}^\Delta)}$.

Let $\mathbf{y}_\mathbf{I}^j = (y_\varepsilon^j: \varepsilon \in \{0, 1\}^d, \varepsilon_{d-j} = 0)$ and $\mathbf{y}_\mathbf{II}^j = (y_\varepsilon^j: \varepsilon \in \{0, 1\}^d, \varepsilon_{d-j} = 1)$ be the $(d-j)$ -th lower and upper faces of \mathbf{y}^j respectively. Let $\mathcal{G}'_{T_1, \dots, T_{d-j-1}, T_{d-j+1}, \dots, T_d}$ (excluding T_{d-j}) be the group generated by $\mathcal{G}_{T_1, \dots, T_{d-j-1}, T_{d-j+1}, \dots, T_d}$ and the diagonal $G_{[d-1]}^\Delta$. By a minimality argument (identical to the one written for $j=0$), there exists a sequence $(\mathcal{G}_i^j)_{i \geq 1}$ in $\mathcal{G}'_{T_1, \dots, T_{d-j-1}, T_{d-j+1}, \dots, T_{d-1}}$, such that $\mathcal{G}_i^j \mathbf{y}_\mathbf{I}^j \rightarrow y_1^{[d-1]}$ as i goes to infinity. By compactness we can assume that $\mathcal{G}_i^j \mathbf{y}_\mathbf{II}^j \rightarrow \mathbf{z}_\mathbf{II}^j$ for some $\mathbf{z}_\mathbf{II}^j \in \mathbf{Q}_{T_1, \dots, T_{d-j-1}, T_{d-j+1}, \dots, T_d}(Y)$. Let $\tilde{\mathcal{G}}_i^j$ be the transformation which consists in applying \mathcal{G}_i^j to both the upper and lower $(d-j)$ -th faces of a point in $\mathbf{Q}_{T_1, \dots, T_d}(Y)$. We have that $\tilde{\mathcal{G}}_i^j \mathbf{y}^j$ converges to a point $\mathbf{y}^{j+1} \in \mathbf{Q}_{T_1, \dots, T_d}(Y)$ whose $(d-j)$ -th lower face is equal to $y_1^{[d-1]}$ and whose $(d-j)$ -th upper face is equal to $\mathbf{z}_\mathbf{II}^j$.

Let us analyze the properties of \mathbf{y}^{j+1} . We show that \mathbf{y}^j coincides with $y_1^{[d]}$ in all k -lower faces for $d-(j+1)+1 = d-j \leq k \leq d$. By construction of \mathbf{y}^j , this is true for the $(d-j)$ -th lower face. For $d-j < k \leq d$, let $\varepsilon \in \{0, 1\}^d$ with $\varepsilon_k = 0$ and notice that y_ε^{j+1} equals to $y_{\varepsilon'}^{j+1}$, for all ε' differing from ε at most in the $d-j$ coordinate, where it is equal to 0 (i.e., $\varepsilon'_{d-j} = 0$). Since y_ε^{j+1} equals to a value in the $(d-j)$ -th lower face, we have that it is equal to y_1 . Set $\mathbf{x}^{j+1} = \pi^{[d]}(\mathbf{y}^{j+1})$ and notice that since \mathbf{x}^j is equal to (x, \dots, x, z_j) we have that \mathbf{x}^{j+1} is equal to (x, \dots, x, z_{j+1}) for some $z_{j+1} \in X$.

Claim: The point (x, z_{j+1}) belongs to $\overline{\mathcal{O}((x, z_j), G_{[2]}^\Delta)}$.

To see this, remark that the projection of $\tilde{\mathcal{G}}_i^j \mathbf{y}^j$ onto the coordinates $1\dots 0\dots 1$ (where the 0 is in the $d-j$ position) and $1\dots 1$ has the form (gx, gz_j) and these coordinates converge to (x, z_{j+1}) . This allows us to conclude.

Repeating this process we construct the points $\mathbf{y}^1, \dots, \mathbf{y}^d \in \mathbf{Q}_{T_1, \dots, T_d}(Y)$ and $\mathbf{x}^1, \dots, \mathbf{x}^d \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ such that $\pi^{[d]}(\mathbf{y}^j) = \mathbf{x}^j$ for all $j \in [d]$. The point \mathbf{y}^d satisfies that it is equal to y_1 in all k -lower faces, $k \in [d]$. That is, $y_\varepsilon^d = y_1$ if $\varepsilon_k = 0$ for some $k \in [d]$. That means that \mathbf{y}^d is equal to (y_1, \dots, y_1, y_2) for some $y_2 \in Y$. By Theorem 21, $(y_1, y_2) \in \mathcal{R}_{T_1, \dots, T_d}(Y)$.

Also, by construction we have that $\pi(y_2) = z_d$ and from the condition $(x, z_j) \in \overline{\mathcal{O}((x, z_{j-1}), G_{[2]}^\Delta)}$ for $j \in [d]$, we get that $(x, z_d) \in \overline{\mathcal{O}((x, z), G_{[2]}^\Delta)}$. Now, by distality, we have that $(x, z) \in \overline{\mathcal{O}((x, z_d), G_{[2]}^\Delta)}$, and thus there exists a sequence $(g_i)_{i \geq 1}$ in G such that $(g_i x, g_i z_d) \rightarrow (x, z)$. By compactness we may assume that $(g_i y_1, g_i y_2) \rightarrow (y'_1, y'_2)$ for some $y'_1, y'_2 \in Y$. Then, as $\mathcal{R}_{T_1, \dots, T_d}(Y)$ is closed and invariant, the point (y'_1, y'_2) belongs to $\mathcal{R}_{T_1, \dots, T_d}(Y)$. It is immediate that $\pi \times \pi(y'_1, y'_2) = (x, z)$, finishing the proof. \square

6. STRUCTURE OF SYSTEMS WITH THE UNIQUE CLOSING PARALLELEPIPED PROPERTY

In this section we describe the structure of a minimal \mathbb{Z}^d -system with the unique closing parallelepiped property (for a given choice of generators). To do so, we need first to introduce some terminology.

6.1. The $\mathbf{Q}_H(X)$ relations and classes Z_0^H . Let (X, T_1, \dots, T_d) be a \mathbb{Z}^d -system and let H be a subgroup of $\langle T_1, \dots, T_d \rangle$. We define

$$\mathbf{Q}_H(X) = \overline{\{(x, hx) : x \in X, h \in H\}}.$$

With this notation, slightly abusing of the notation, we have that $\mathbf{Q}_{T_j}(X) = \overline{\{(x, T_j^n x) : x \in X, n \in \mathbb{Z}\}}$ defined in Section 2.2 is nothing but $\mathbf{Q}_{\langle T_j \rangle}(X)$.

It is easy to see that $\mathbf{Q}_H(X)$ is a closed, H -invariant, reflexive and symmetric relation. In [11] Glasner proved that for a G_δ -dense subset X_0 of X , $(X_0 \times X_0) \cap \mathbf{Q}_H(X)$ is transitive. But in general this relation is not an equivalence relation (for an example see [25]). Nevertheless, transitivity of $\mathbf{Q}_H(X)$ is satisfied when the system is distal, because it is a particular case of the gluing property: if $(x, y), (y, z) \in \mathbf{Q}_H(X)$

then $(x, z) \in \mathbf{Q}_H(X)$. The proof of the gluing property in this context is almost a verbatim copy of the one of Lemma 12 so we omit it.

Proposition 24. *Let (X, T_1, \dots, T_d) be a minimal distal \mathbb{Z}^d -system. Then, for every subgroup H of $\langle T_1, \dots, T_d \rangle$ we have that $\mathbf{Q}_H(X)$ is a closed and invariant equivalence relation of X .*

We now define the following classes of \mathbb{Z}^d -systems that will be useful to characterize systems with the unique closing parallelepiped property. For H as before, let

$$\mathbf{Z}_0^H = \{(X, T_1, \dots, T_d) : H \text{ acts as the identity in } X\}.$$

This definition and notation is analogous to the one used by Austin in [3] when studying the L^2 convergence of multiple ergodic averages in the measure theoretical setting. Using a Zorn's Lemma argument it can be proved that every \mathbb{Z}^d -system possesses a maximal \mathbf{Z}_0^H -factor [2, Chapter 9]. That is, any factor of a given \mathbb{Z}^d -system in the class \mathbf{Z}_0^H factorizes through its maximal \mathbf{Z}_0^H -factor. In addition, this factor can be characterized precisely. To this purpose, for a minimal \mathbb{Z}^d -system (X, T_1, \dots, T_d) let $\sigma_H(X)$ denote the smallest closed and invariant equivalence relation containing $\mathbf{Q}_H(X)$.

Lemma 25. *Let (X, T_1, \dots, T_d) be a minimal \mathbb{Z}^d -system. Then, $(X/\sigma_H(X), T_1, \dots, T_d)$ is the maximal \mathbf{Z}_0^H -factor of (X, T_1, \dots, T_d) .*

Proof. Let $\pi : X \rightarrow X/\sigma_H(X)$ be the natural factor map. First we prove that $(X/\sigma_H(X), T_1, \dots, T_d) \in \mathbf{Z}_0^H$. Let $h \in H$, if $[\cdot]$ denotes the equivalence class of a point for the relation $\sigma_H(X)$, then $h[x] = [hx]$. Since $(x, hx) \in \sigma_H(X)$ we get that $h[x] = [x]$. So H acts trivially on $X/\sigma_H(X)$. Now, let (Z, T_1, \dots, T_d) be a factor of (X, T_1, \dots, T_d) that belongs to \mathbf{Z}_0^H and let $\pi' : X \rightarrow Z$ be the corresponding factor map. Let $R_{\pi'} = \{(x, y) \in X : \pi'(x) = \pi'(y)\}$. We have to prove that $\sigma_H(X) \subseteq R_{\pi'}$. In fact, since $R_{\pi'}$ is a closed and invariant equivalence relation, it suffices to prove that $\mathbf{Q}_H(X) \subseteq R_{\pi'}$. Since $(Z, T_1, \dots, T_d) \in \mathbf{Z}_0^H$ we have that for every $h \in H$, $\pi' \circ h = h \circ \pi' = \pi'$. Hence, for any $x \in X$, $\{(x, hx) : h \in H\} \subseteq R_{\pi'}$, and thus $\mathbf{Q}_H(X) \subseteq R_{\pi'}$ as desired. \square

In the distal case, by Proposition 24, $\mathbf{Q}_H(X)$ is an equivalence relation and thus $\mathbf{Q}_H(X) = \sigma_H(X)$. In this case Lemma 25 can be stated as,

Corollary 26. *Let (X, T_1, \dots, T_d) be a minimal distal \mathbb{Z}^d -system. Then, $(X/\mathbf{Q}_H(X), T_1, \dots, T_d)$ is the maximal \mathbf{Z}_0^H -factor of (X, T_1, \dots, T_d) .*

This result generalizes to iterated quotients.

Proposition 27. *Let (X, T_1, \dots, T_d) be a minimal distal \mathbb{Z}^d -system and let H, H_1 and H_2 be subgroups of $\langle T_1, \dots, T_d \rangle$ such that $H = H_1 H_2$. Let $Y = X/\mathbf{Q}_{H_1}(X)$. Then, $(X/\mathbf{Q}_H(X), T_1, \dots, T_d)$ is isomorphic to $(Y/\mathbf{Q}_{H_2}(Y), T_1, \dots, T_d)$.*

Proof. Let $\pi_1 : X \rightarrow Y (= X/\mathbf{Q}_{H_1}(X))$ and $\pi : X \rightarrow Y/\mathbf{Q}_{H_2}(Y)$ be the natural factor maps and $R_\pi = \{(x, x') : \pi(x) = \pi(x')\}$. We have that $Y/\mathbf{Q}_{H_2}(Y)$ is a factor of X where the action of both H_1 and H_2 are trivial. Hence, the action of H is trivial as well. It follows that $\mathbf{Q}_H(X) \subseteq R_\pi$. For the converse inclusion, assume that $(x, x') \in R_\pi$. Then $(\pi_1(x), \pi_1(x')) \in \mathbf{Q}_{H_2}(Y)$.

Claim: there exists $x'' \in X$ such that $(x, x'') \in \mathbf{Q}_{H_2}(X)$ and $\pi_1(x'') = \pi_1(x')$.

To prove the claim, consider $\epsilon > 0$ and let $\epsilon > \delta > 0$ so that $\pi(B(x, \epsilon))$ contains $B(\pi_1(x), \delta)$. This can be done by the openness of π_1 (recall the systems are distal). Since $(\pi_1(x), \pi_1(x')) \in \mathbf{Q}_{H_2}(Y)$, there exists $\tilde{x} \in X$ and $h_2 = h_2(\delta) \in H_2$ such that $\pi_1(\tilde{x})$ is δ close to $\pi_1(x)$ and $h_2 \pi_1(\tilde{x}) = \pi_1(h_2 \tilde{x})$ is δ close to $\pi_1(x')$. We can find a point \tilde{x}' that is ϵ close to x such that $\pi_1(\tilde{x}') = \pi_1(\tilde{x})$ and so $\pi_1(h_2 \tilde{x}')$ is δ close to $\pi_1(x')$. A compactness argument allows us to find x'' as a limit point of $h_2 \tilde{x}'$, satisfying then the claim statement.

Using the claim, notice that $(x, x'') \in \mathbf{Q}_{H_2}(X) \subseteq \mathbf{Q}_H(X)$ and $(x'', x') \in \mathbf{Q}_{H_1}(X) \subseteq \mathbf{Q}_H(X)$. Since $\mathbf{Q}_H(X)$ is an equivalence relation we conclude that $(x, x') \in \mathbf{Q}_H(X)$, finishing the proof. \square

To ease notations, we let $X/\mathbf{Q}_{H_1}/\mathbf{Q}_{H_2}$ denote the quotient $Y/\mathbf{Q}_{H_2}(Y)$, where $Y = X/\mathbf{Q}_{H_1}(X)$. We naturally extend this notation to iterated quotients $X/\mathbf{Q}_{H_1}/\mathbf{Q}_{H_2}/\dots/\mathbf{Q}_{H_n}$ that are defined in the obvious way.

Notice that if $\{j_1, \dots, j_k\} \subseteq [d]$ we have that

$$\mathbf{Q}_{\langle T_{j_1}, \dots, T_{j_k} \rangle}(X) = \overline{\{(x, hx) : x \in X, h \in \langle T_{j_1}, \dots, T_{j_k} \rangle\}}.$$

We warn the reader that the sets $\mathbf{Q}_{\langle T_{j_1}, \dots, T_{j_k} \rangle}(X)$ and $\mathbf{Q}_{T_{j_1}, \dots, T_{j_k}}(X)$ are different. The first one is a subset of X^2 while the second one is a subset of $X^{[k]}$.

Following the notation in [3] we denote by $Z_0^{e_j}$ the class $Z_0^{\langle T_j \rangle}$ and by $Z_0^{e_{j_1} \wedge e_{j_2}}$ the intersection of the classes $Z_0^{e_{j_1}}$ and $Z_0^{e_{j_2}}$, which corresponds to the class $Z_0^{\langle T_{j_1}, T_{j_2} \rangle}$.

6.2. Dynamical structure of systems with the unique closing parallelepiped property. We describe first a soft structure theorem for minimal systems with the unique closing parallelepiped property.

Proposition 28. *Let (X, T_1, \dots, T_d) be a minimal (not necessarily distal) \mathbb{Z}^d -system with the unique closing parallelepiped property. Then,*

- (1) *for each $j \in [d]$ there exists a \mathbb{Z}^d -system (Y_j, T_1, \dots, T_d) such that T_j is the identity transformation on Y_j ;*
- (2) *there exists a joining (Y, T_1, \dots, T_d) of the systems $(Y_1, T_1, \dots, T_d), \dots, (Y_d, T_1, \dots, T_d)$ which is an extension of (X, T_1, \dots, T_d) .*

Otherwise saying, (X, T_1, \dots, T_d) has an extension which is a joining of systems, where on each one a different transformation acts as the identity.




FIGURE 5. Structure of a minimal system (X, T_1, \dots, T_d) with the unique closing parallelepiped property

Proof. Let (X, T_1, \dots, T_d) be a minimal \mathbb{Z}^d -system and let $x_0 \in X$. Consider the \mathbb{Z}^d -system $(\mathbf{K}_{T_1, \dots, T_d}^{x_0}, T_1^{[d]}, \dots, T_d^{[d]})$ (recall the face transformations defined in Section 2.2). Such system is similar to the one considered in [7] when $d = 2$. Let $Y \subseteq \mathbf{K}_{T_1, \dots, T_d}^{x_0}$ be a minimal subsystem (note that in the distal case we have that $Y = \mathbf{K}_{T_1, \dots, T_d}^{x_0}$). Here we consider the natural factor map from $(Y, T_1^{[d]}, \dots, T_d^{[d]})$ to (X, T_1, \dots, T_d) given by the projection onto the last coordinate of points in Y . We analyze how Y looks like when (X, T_1, \dots, T_d) has the unique closing parallelepiped property and show that Y can be “naturally decomposed” as a joining of some factors where on each one the action of one transformation is the identity. That is, Y is a joining of lower dimensional actions.

We proceed as follows. Points in Y have $2^d - 1$ coordinates and since (X, T_1, \dots, T_d) has the unique closing parallelepiped property, the last one is determined by the others. Hence, we can think of points in Y having only $2^d - 2$ coordinates.

$$\begin{array}{cccccc|c}
 \text{A point in } Y : & y_{10\dots 0} & \cdots & y_{0\dots 01} & \cdots & y_{1\dots 10} & y_{1\dots 1} \\
 T_1^{[d]} : & T_1 & \cdots & \text{id} & \cdots & T_1 & T_1 \\
 & \vdots & \cdots & \vdots & \cdots & \vdots & \vdots \\
 T_{d-1}^{[d]} & \text{id} & \cdots & \text{id} & \cdots & T_{d-1} & T_{d-1} \\
 T_d^{[d]} & \text{id} & \cdots & T_d & \cdots & \text{id} & T_d
 \end{array}$$

FIGURE 6. How a point in Y and the face transformations look like.

In Figure 6.2, the dashed vertical line separates the last coordinate of a point in Y that is determined by the others. From now on we omit this coordinate. Now we describe how to build the factors Y_1, \dots, Y_d whose joining is Y .

For $j \in [d]$, let Y_j be the projection of Y onto the coordinates where $T_j^{[d]}$ is trivial, *i.e.*, onto the coordinates $\varepsilon \in \{0, 1\}^d \setminus \{\vec{0}\}$ such that $\varepsilon_j = 0$. We leave the reader to verify that Y_j is a subset of $\mathbf{K}_{T_1, \dots, T_{j-1}, T_{j+1}, \dots, T_d}^{x_0}$ (the face orbit where we forget the transformation T_j).

Because we are forgetting the last coordinate, for all coordinates of a point in Y at least one of the transformations acts as the identity. It follows then that the projections of a point $\mathbf{y} \in Y$ into the factors Y_j , $j \in [d]$, determine the point \mathbf{y} itself, *i.e.*, Y is a joining of the systems Y_j , $j \in [d]$. This finishes the proof. \square

Remark that by definition, for $j \in [d]$ the system (Y_j, T_1, \dots, T_d) belongs to the class $Z_0^{e_j}$. What next result shows is that in the distal case (Y_j, T_1, \dots, T_d) is actually the maximal factor of (Y, T_1, \dots, T_d) with respect to this class.

Proposition 29. *Let (X, T_1, \dots, T_d) be a minimal distal \mathbb{Z}^d -system and $x_0 \in X$. Then, for any $j \in [d]$, $(\mathbf{K}_{T_1, \dots, T_{j-1}, T_{j+1}, \dots, T_d}^{x_0}, T_1^{[d]}, \dots, T_d^{[d]})$ is isomorphic to $(\mathbf{K}_{T_1, \dots, T_d}^{x_0} / \mathbf{Q}_{T_j^{[d]}}(\mathbf{K}_{T_1, \dots, T_d}^{x_0}), T_1^{[d]}, \dots, T_d^{[d]})$. In particular, $(\mathbf{K}_{T_1, \dots, T_{j-1}, T_{j+1}, \dots, T_d}^{x_0}, T_1^{[d]}, \dots, T_d^{[d]})$ is the maximal $Z_0^{e_j}$ -factor of $(\mathbf{K}_{T_1, \dots, T_d}^{x_0}, T_1^{[d]}, \dots, T_d^{[d]})$.*

Proof. We follow the notation introduced in Proposition 28. We start recalling that in the distal case, by minimality we have that $Y = \mathbf{K}_{T_1, \dots, T_d}^{x_0}$ and $Y_j = \mathbf{K}_{T_1, \dots, T_{j-1}, T_{j+1}, \dots, T_d}^{x_0}$ for every $j \in [d]$. Let $j \in [d]$. We know by Corollary 26 that $(\mathbf{K}_{T_1, \dots, T_d}^{x_0} / \mathbf{Q}_{T_j^{[d]}}(\mathbf{K}_{T_1, \dots, T_d}^{x_0}), T_1^{[d]}, \dots, T_d^{[d]})$ is the maximal $Z_0^{e_j}$ -factor of $(\mathbf{K}_{T_1, \dots, T_d}^{x_0}, T_1^{[d]}, \dots, T_d^{[d]})$ and since $Y_j \in Z_0^{e_j}$ we only need to show that if $\mathbf{x}, \mathbf{y} \in \mathbf{K}_{T_1, \dots, T_d}^{x_0}$ are such that

$$(x_\varepsilon : \varepsilon \in \{0, 1\}^d \setminus \{\vec{0}\}, \varepsilon_j = 0) = (y_\varepsilon : \varepsilon \in \{0, 1\}^d \setminus \{\vec{0}\}, \varepsilon_j = 0)$$

(i.e., they have the same projection on Y_j), then $(\mathbf{x}, \mathbf{y}) \in \mathbf{Q}_{T_j^{[d]}}(\mathbf{K}_{T_1, \dots, T_d}^{x_0})$.

Let $\delta > 0$ and let $\pi : \mathbf{K}_{T_1, \dots, T_d}^{x_0} \rightarrow \mathbf{K}_{T_1, \dots, T_{j-1}, T_{j+1}, \dots, T_d}^{x_0}$ be the natural projection. By the openness of π (recall the systems are distal), we can find $0 < \delta' < \delta$ such that

$$(1) \quad B_{\rho_{X_*^{[d-1]}}}((x_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_j = 0), \delta') \subseteq \pi \left(B_{\rho_{X_*^{[d]}}}(\mathbf{x}, \delta) \right) \cap \pi \left(B_{\rho_{X_*^{[d]}}}(\mathbf{y}, \delta) \right).$$

Here we recall that if ρ is a metric on X , ρ_{X_*} is any metric that generates the product topology on $X_*^{[d]} = X^{2^d-1}$ (for instance the sum of the metrics in each coordinate).

By definition of $\mathbf{K}_{T_1, \dots, T_d}^{x_0}$, there exists $\mathbf{n} \in \mathbb{Z}^d$ such that

$$\rho_{X_*^{[d]}}((T_1^{n_1 \varepsilon_1} \dots T_d^{n_d \varepsilon_d} x_0)_{\varepsilon \in \{0, 1\}^d \setminus \{\vec{0}\}}, \mathbf{x}) < \delta'.$$

Call $\mathbf{z} = (T_1^{n_1 \varepsilon_1} \dots T_d^{n_d \varepsilon_d} x_0)_{\varepsilon \in \{0, 1\}^d \setminus \{\vec{0}\}}$. Then, we have that

$$\rho_{X_*^{[d-1]}}((z_\varepsilon : \varepsilon \in \{0, 1\}^d \setminus \{\vec{0}\}, \varepsilon_j = 0), (x_\varepsilon : \varepsilon \in \{0, 1\}^d \setminus \{\vec{0}\}, \varepsilon_j = 0)) < \delta'.$$

Thus, by (1), there exists $\mathbf{z}^1 \in \mathbf{K}_{T_1, \dots, T_d}^{x_0}$ such that

$$(z_\varepsilon^1 : \varepsilon \in \{0, 1\}^d \setminus \{\vec{0}\}, \varepsilon_j = 0) = (z_\varepsilon : \varepsilon \in \{0, 1\}^d \setminus \{\vec{0}\}, \varepsilon_j = 0)$$

and $\rho_{X_*^{[d]}}(\mathbf{z}^1, \mathbf{y}) < \delta$.

Let $0 < \delta'' < \delta'$ be such that $\delta' + \delta'' < \delta$ and for any $\mathbf{u}, \mathbf{v} \in \mathbf{K}_{T_1, \dots, T_d}^{x_0}$ we have

$$\rho_{X_*^{[d]}}(\mathbf{u}, \mathbf{v}) < \delta'' \implies \rho_{X_*^{[d]}}((T_j^{[d]})^{n_j} \mathbf{u}, (T_j^{[d]})^{n_j} \mathbf{v}) < \delta'.$$

By definition of $\mathbf{K}_{T_1, \dots, T_d}^{x_0}$, there exists $\mathbf{n}' \in \mathbb{Z}^d$ such that

$$\rho_{X_*^{[d]}}((T_1^{n'_1 \varepsilon_1} \dots T_d^{n'_d \varepsilon_d} x_0)_{\varepsilon \in \{0, 1\}^d \setminus \{\vec{0}\}}, \mathbf{z}^1) < \delta''.$$

Call $\mathbf{z}^2 = (T_1^{n'_1 \varepsilon_1} \dots T_d^{n'_d \varepsilon_d} x_0)_{\varepsilon \in \{0, 1\}^d \setminus \{\vec{0}\}}$ (i.e., $\rho_{X_*^{[d]}}(\mathbf{z}^2, \mathbf{z}^1) < \delta''$) and define $\mathbf{z}^3 = (T_j^{[d]})^{n_j - n'_j} \mathbf{z}^2$.

Notice that the j -th lower face of \mathbf{z}^3 coincides with the one of \mathbf{z}^2 which is $\delta'' + \delta$ close to the one of \mathbf{x} (here we are slightly abusing terminology, the j -th lower face should consider the coordinate $\vec{0}$, that we do not have in points of $\mathbf{K}_{T_1, \dots, T_d}^{x_0}$, but let us use this word to avoid introducing more terminology). On the other hand, the j -th upper face of $(T_j^{[d]})^{-n'_j} \mathbf{z}^2$ coincides with the j -lower face of \mathbf{z}^2 (with x_0 in the $\vec{0}$ coordinate) and then it is δ'' close to the j -th lower face of \mathbf{z}^1 and thus to the j -th lower face of \mathbf{z} . By the definition of δ'' , and using that the j -th upper face of \mathbf{z} is obtained by applying $(T_d^{[d]})^{n_d}$ to its j -th lower face we get that

$$\mathbf{z}^3 = (T_j^{[d]})^{n_j} (T_j^{[d]})^{-n'_j} \mathbf{z}^2 \text{ is } 3\delta \text{ close to } \mathbf{x}.$$

Now, remark that the point $(T_j^{[d]})^{n'_j - n_j} \mathbf{z}^3 = \mathbf{z}^2$ is $\delta'' + \delta < 2\delta$ close to \mathbf{y} . As $\delta > 0$ is arbitrary, we have shown that $(\mathbf{x}, \mathbf{y}) \in \mathbf{Q}_{T_j^{[d]}}(\mathbf{K}_{T_1, \dots, T_d}^{x_0})$ and therefore the factor map $\pi : \mathbf{K}_{T_1, \dots, T_d}^{x_0} / \mathbf{Q}_{T_j^{[d]}}(\mathbf{K}_{T_1, \dots, T_d}^{x_0}) \rightarrow \mathbf{K}_{T_1, \dots, T_{j-1}, T_{j+1}, \dots, T_d}^{x_0}$ is an isomorphism. \square

We use Theorem 21 to prove the following theorem, which gives deeper information about the structure of the systems with the unique closing parallelepiped property, introducing independence with respect to further factors.

To state the theorem, let us introduce another definition. Let (X, T_1, \dots, T_d) be a \mathbb{Z}^d -system. We say that $x_0 \in X$ is a continuity point if $\mathbf{K}_{T_1, \dots, T_d}^{x_0}$ coincides with the set of $\mathbf{x} \in X_*^{[d]}$ such that $(x_0, \mathbf{x}) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. The set of continuity points $x_0 \in X$ is a dense G_δ set of points of X [11, Lemma 4.5]. Note that if x_0 is a continuity point, then any point in its orbit, *i.e.*, $T_1^{n_1} \cdots T_d^{n_d} x_0$ with $n_1, \dots, n_d \in \mathbb{Z}$ is also a continuity point. For a continuity point x_0 , we have that $\mathbf{Q}_{T_j}(x_0) = \overline{\mathcal{O}(x_0, T_j)}$ for $j \in [d]$.

Theorem 30. *Let (X, T_1, \dots, T_d) be a minimal distal \mathbb{Z}^d -system and $x_0 \in X$ a continuity point. Suppose that X has the unique closing parallelepiped property. Then, $\mathbf{K}_{T_1, \dots, T_d}^{x_0}$ is a joining of its factors $\mathbf{K}_{T_1, \dots, T_{j-1}, T_{j+1}, \dots, T_d}^{x_0}$, $j \in [d]$, and it is relatively independent with respect to their maximal $\mathbf{Z}_0^{e_{j_1}} \wedge \mathbf{Z}_0^{e_{j_2}}$ -factors, for all $j_1, j_2 \in [d]$ with $j_1 < j_2$.*




FIGURE 7. Decomposition of the system $(\mathbf{K}_{T_1, \dots, T_d}^{x_0}, T_1^{[d]}, \dots, T_d^{[d]})$ as the joining of the systems $\mathbf{K}_{T_1, \dots, T_{j-1}, T_{j+1}, \dots, T_d}^{x_0}$. Theorem 30 asserts that the joining is relatively independent with respect to the factors in the “second level”, *i.e.*, $\mathbf{K}_{T_{i_1}, \dots, T_{i_{d-2}}}^{x_0}$, $1 \leq i_1 < \dots < i_{d-2} \leq d$. In the diagram further factors are also shown. A system in the k th-level is an action of \mathbb{Z}^{d-k} (k transformations are the identity, and $\mathbf{K}_{T_1, \dots, T_d}^{x_0}$ corresponds to the 0 level).

Proof. The fact that $\mathbf{K}_{T_1, \dots, T_d}^{x_0}$ is a joining of its factors $\mathbf{K}_{T_1, \dots, T_{j-1}, T_{j+1}, \dots, T_d}^{x_0}$, $j \in [d]$, was already shown in Proposition 28 and Proposition 29, so we are left to show the relative independence part.

Let $\mathbf{x}, \mathbf{y} \in X_*^{[d]}$ be such that:

- (1) $\mathbf{x} \in \mathbf{K}_{T_1, \dots, T_d}^{x_0}$.
- (2) For every $j \in [d]$, $(y_\varepsilon : \varepsilon \in \{0, 1\}^d \setminus \{\vec{0}\}, \varepsilon_j = 0) \in \mathbf{K}_{T_1, \dots, T_{j-1}, T_{j+1}, \dots, T_d}^{x_0}$.
- (3) For every $j_1, j_2 \in [d]$, $j_1 \neq j_2$,

$$(x_\varepsilon : \varepsilon \in \{0, 1\}^d \setminus \{\vec{0}\}, \varepsilon_{j_1} = 0, \varepsilon_{j_2} = 0) = (y_\varepsilon : \varepsilon \in \{0, 1\}^d \setminus \{\vec{0}\}, \varepsilon_{j_1} = 0, \varepsilon_{j_2} = 0),$$

i.e., \mathbf{x} and \mathbf{y} coincide in the coordinates associated to the maximal $\mathbf{Z}_0^{e_{j_1}} \wedge \mathbf{Z}_0^{e_{j_2}}$ -factor.

Roughly speaking, this condition says that given $\mathbf{x} \in \mathbf{K}_{T_1, \dots, T_d}^{x_0}$, one considers its projection into the systems of the “second level” and the relative independence means that one can freely change all the coordinates of \mathbf{x} of the form $[d] \setminus \{j\}$ with $j \in [d]$, as long as the resulting point (the point \mathbf{y} in the statement) has valid faces. We leave the reader to verify that these are the conditions we need to look at to check the relative independence.

So, given (i), (ii) and (iii), we want to show that $\mathbf{y} \in \mathbf{K}_{T_1, \dots, T_d}^{x_0}$. By (iii), the only coordinates $\varepsilon \subseteq [d]$, $\varepsilon \neq \emptyset$, such that $x_\varepsilon \neq y_\varepsilon$ are the coordinates $\varepsilon = [d]$ and $\varepsilon = [d] \setminus \{j\}$ for $j \in [d]$. The program of the

proof is to construct points $\mathbf{x}^0, \mathbf{x}^1, \dots, \mathbf{x}^d \in \mathbf{K}_{T_1, \dots, T_d}^{x_0}$ with $\mathbf{x}^0 = \mathbf{x}$ and such that for every $k \in [d]$

$$x_\varepsilon^k = \begin{cases} x_\varepsilon & \varepsilon \subseteq [d], \varepsilon \neq [d] \setminus \{j\} \text{ for some } j \in [d], \\ y_\varepsilon & \varepsilon = [d] \setminus \{j\} \text{ for some } j \in \{1, \dots, k\}, \\ a^k & \varepsilon = [d], \end{cases}$$

for some $a^k \in X$. Notice that since X has the unique closing parallelepiped property, the point a^k is unique. The point \mathbf{x}^d coincides with \mathbf{y} in all coordinates ε , $\varepsilon \neq [d]$, so the unique closing parallelepiped property implies that they are equal. This would conclude that $\mathbf{y} \in \mathbf{K}_{T_1, \dots, T_d}^{x_0}$. To construct the points mentioned above, assume we have constructed the point $\mathbf{x}^k \in \mathbf{K}_{T_1, \dots, T_d}^{x_0}$ for $0 \leq k < d$. Notice that $(x_\varepsilon^k: \varepsilon \in \{0, 1\}^d \setminus \{\vec{0}\}, \varepsilon_{k+1} = 0)$ and $(y_\varepsilon: \varepsilon \in \{0, 1\}^d \setminus \{\vec{0}\}, \varepsilon_{k+1} = 0)$ differ at most in the coordinate $[d] \setminus \{k+1\}$.

Claim 1. There exists $a \in X$ such that if we replace the coordinate $\varepsilon = [d] \setminus \{k+1\}$ of \mathbf{x}^k by the corresponding coordinate of \mathbf{y} and the coordinate $[d]$ of \mathbf{x}^k by a , we obtain a point \mathbf{x}^{k+1} that belongs to $\mathbf{K}_{T_1, \dots, T_d}^{x_0}$.

We suggest the reader to look at Figure 8, while reading the proof of the claim. There, we have illustrated the proof in the case $d = 3$. We show how to change the coordinate $[d] \setminus \{1\}$ of \mathbf{x} by the corresponding one of \mathbf{y} (paying the minor price of modifying the $[d]$ coordinate too).

Proof of the Claim 1. We first remark that it is enough to show the statement of the claim for $k = 0$ (i.e., for the first step). Indeed, by Lemma 14, after proving the case $k = 0$ we can permute the transformations and assume that now the second transformation is the first one, apply again the case $k = 0$ and repeat this argument as many times as needed. So, we have to show that from $\mathbf{x}^0 = \mathbf{x}$ we can change its $[d] \setminus \{1\}$ coordinate by the corresponding one of \mathbf{y} and obtain a point in $\mathbf{K}_{T_1, \dots, T_d}^{x_0}$. In the process we also allow us to change the $[d]$ coordinate of \mathbf{x} .

In this claim it will be convenient to work in $\mathbf{Q}_{T_1, \dots, T_d}(X)$ instead of in $\mathbf{K}_{T_1, \dots, T_d}^{x_0}$ so we identify $\mathbf{x}, \mathbf{y} \in \mathbf{K}_{T_1, \dots, T_d}^{x_0}$ with $(x_0, \mathbf{x}), (x_0, \mathbf{y}) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ respectively. This identification is harmless because of the unique closing parallelepiped property and avoids introducing more notation.

Consider the system (X, T_2, \dots, T_d) . Since the points $(x_\varepsilon: \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0)$ and $(y_\varepsilon: \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0)$ differ at most in one coordinate (the coordinate $[d] \setminus \{1\}$), by Theorem 21 (2) applied to the system (X, T_2, \dots, T_d) (see Remark 13) we have that

$$(2) \quad (x_{[d] \setminus \{1\}}, \dots, x_{[d] \setminus \{1\}}, y_{[d] \setminus \{1\}}) \in \mathbf{Q}_{T_2, \dots, T_d}(X).$$

Now, duplicating this point along the 1-th face, we obtain the point $\mathbf{v}^1 \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. This point is characterized by

$$v_\varepsilon^1 = \begin{cases} x_{[d] \setminus \{1\}} & \text{if } \varepsilon_1 = 0 \wedge \exists j \neq 1, \varepsilon_j = 0, \\ y_{[d] \setminus \{1\}} & \text{if } \varepsilon = [d] \setminus \{1\}, \\ x_{[d] \setminus \{1\}} & \text{if } \varepsilon_1 = 1 \wedge \exists j \neq 1, \varepsilon_j = 0, \\ y_{[d] \setminus \{1\}} & \text{if } \varepsilon = [d]. \end{cases}$$

Claim 2. There exists $a \in X$ such that the point \mathbf{u}^1 defined as

$$u_\varepsilon^1 = \begin{cases} x_{[d] \setminus \{1\}} & \text{if } \varepsilon_1 = 0 \wedge \exists j \neq 1, \varepsilon_j = 0, \\ y_{[d] \setminus \{1\}} & \text{if } \varepsilon = [d] \setminus \{1\}, \\ x_{[d]} & \text{if } \varepsilon_1 = 1 \wedge \exists j \neq 1, \varepsilon_j = 0, \\ a & \text{if } \varepsilon = [d], \end{cases}$$

belongs to $\mathbf{Q}_{T_1, \dots, T_d}(X)$.

Proof the Claim 2. To prove the claim, notice that $(x_{[d] \setminus \{1\}}, x_{[d]}) \in \mathbf{Q}_{T_1}(X)$, because the point \mathbf{x} belongs to $\mathbf{Q}_{T_1, \dots, T_d}(X)$. Now we analyze separately two cases, namely the case $\mathbf{Q}_{T_1}(x_{[d] \setminus \{1\}}) = \overline{\mathcal{O}(x_{[d] \setminus \{1\}}, T_1)}$ and the case $\mathbf{Q}_{T_1}(x_{[d] \setminus \{1\}}) \supsetneq \overline{\mathcal{O}(x_{[d] \setminus \{1\}}, T_1)}$. By Lemma 4.5 of [11], there is a G_δ -dense set of $x \in X$ such that $\mathbf{Q}_{T_1}(x) = \overline{\mathcal{O}(x, T_1)}$.

Case 1: $\mathbf{Q}_{T_1}(x_{[d] \setminus \{1\}}) = \overline{\mathcal{O}(x_{[d] \setminus \{1\}}, T_1)}$. Since $(x_{[d] \setminus \{1\}}, x_{[d]}) \in \mathbf{Q}_{T_1}(X)$, by the assumption of this case we can find a sequence $(n_i)_{i \in \mathbb{N}} \subseteq \mathbb{Z}$ such that

$$T_1^{n_i} x_{[d] \setminus \{1\}} \rightarrow x_{[d]}.$$

By compactness we can assume that $T_1^{n_i} y_{[d] \setminus \{1\}} \rightarrow a$ for some $a \in X$. Thus, applying the transformation $(T_1^{[d]})^{n_i}$ to the point \mathbf{v}^1 and taking the limit, we obtain a point \mathbf{u}^1 of the desired form that belongs to $\mathbf{Q}_{T_1, \dots, T_d}(X)$.

Case 2: $\mathbf{Q}_{T_1}(x_{[d]\setminus\{1\}}) \supseteq \overline{\mathcal{O}(x_{[d]\setminus\{1\}}, T_1)}$. We define the following projection maps.

- Let ϕ_1 be the projection from $\mathbf{K}_{T_1, \dots, T_d}^{x_0}$ onto the coordinates $\varepsilon \in \{0, 1\}^d$ where $\varepsilon_1 = 0$ and the coordinate $\varepsilon = [d]$. That is

$$\phi_1((z_\varepsilon : \varepsilon \in \{0, 1\}^d)) = (z_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0 \vee \varepsilon = [d])$$

- Let ϕ_2 be the projection from $\mathbf{K}_{T_1, \dots, T_d}^{x_0}$ onto the coordinates $\varepsilon \in \{0, 1\}^d$ where $\varepsilon_1 = 0$, and set $\phi_{1,2}$ the map from $\phi_1(\mathbf{K}_{T_1, \dots, T_d}^{x_0})$ onto $\phi_2(\mathbf{K}_{T_1, \dots, T_d}^{x_0})$ such that $\phi_{1,2} \circ \phi_1 = \phi_2$. That is,

$$\begin{aligned} \phi_2((z_\varepsilon : \varepsilon \in \{0, 1\}^d)) &= (z_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0), \\ \phi_{1,2}((z_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0 \vee \varepsilon = [d])) &= (z_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0). \end{aligned}$$

- Let ϕ_3 be the projection from $\mathbf{K}_{T_1, \dots, T_d}^{x_0}$ onto the coordinates $\varepsilon \in \{0, 1\}^d$ such that $\varepsilon_1 = 0$ and such that there exists $j \in [d]$, $j \neq 1$, such that $\varepsilon_j = 0$. Set $\phi_{2,3}$ the map from $\phi_2(\mathbf{K}_{T_1, \dots, T_d}^{x_0})$ onto $\phi_3(\mathbf{K}_{T_1, \dots, T_d}^{x_0})$ such that $\phi_{2,3} \circ \phi_2 = \phi_3$. That is,

$$\begin{aligned} \phi_3((z_\varepsilon : \varepsilon \in \{0, 1\}^d)) &= (z_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0 \wedge \exists j \neq 1, \varepsilon_j = 0), \\ \phi_{2,3}((z_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0)) &= (z_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0 \wedge \exists j \neq 1, \varepsilon_j = 0). \end{aligned}$$

Note that when we write an expression like $(z_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0)$ we implicitly assume that this point is the restriction of a point $\mathbf{z} = (z_\varepsilon : \varepsilon \in \{0, 1\}^d)$. By distality, all functions described above are open. Let $\delta > 0$. Because of the openness of $\phi_{1,2}$ and $\phi_{2,3}$, we can take $0 < \delta' < \delta$ such that

$$(3) \quad B((x_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0), \delta') \subseteq \phi_{1,2}(B((x_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0 \vee \varepsilon = [d]), \delta)),$$

and $0 < \delta'' < \delta'$ such that

$$(4) \quad \begin{aligned} &B((x_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0 \wedge \exists j \neq 1, \varepsilon_j = 0), \delta'') \\ &\subseteq \phi_{2,3}(B((x_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0), \delta')). \end{aligned}$$

Let $\mathbf{a} = (a_\varepsilon : \varepsilon \in \{0, 1\}^d)$ be such that $\mathbf{Q}_{T_1}(a_{[d]\setminus\{1\}}) = \overline{\mathcal{O}(a_{[d]\setminus\{1\}}, T_1)}$ and such that is δ'' close to $(x_\varepsilon : \varepsilon \in \{0, 1\}^d)$. This point exists because x_0 is a continuity point (and we may take $\mathbf{a} \in \mathbf{K}_{T_1, \dots, T_d}^{x_0}$ in the face orbit of (x_0, \dots, x_0) . See the remarks before the statement of Theorem 30)). The point $(a_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0 \wedge \exists j \neq 1, \varepsilon_j = 0)$ is δ'' close to $(x_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0 \wedge \exists j \neq 1, \varepsilon_j = 0)$ and then, by (4), there exists a point $(b_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0)$ such that $(b_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0)$ is δ' close to $(y_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0)$ and $(a_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0 \wedge \exists j \neq 1, \varepsilon_j = 0) = (b_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0 \wedge \exists j \neq 1, \varepsilon_j = 0)$. Therefore, the points $(a_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0)$ and $(b_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0)$ differ at most in one coordinate (the coordinate $[d] \setminus \{1\}$) and we can repeat the argument started in (2). The point $(a_\varepsilon : \varepsilon \in \{0, 1\}^d, \varepsilon_1 = 0)$ satisfies the assumption of Case 1 and thus we can construct $\mathbf{w} \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ such that

$$w_\varepsilon = \begin{cases} a_{[d]\setminus\{1\}} & \text{if } \varepsilon_1 = 0 \wedge \exists j \neq 1, \varepsilon_j = 0, \\ b_{[d]\setminus\{1\}} & \text{if } \varepsilon = [d] \setminus \{1\}, \\ a_{[d]} & \text{if } \varepsilon_1 = 1 \wedge \exists j \neq 1, \varepsilon_j = 0, \\ u_a & \text{if } \varepsilon = [d], \end{cases}$$

for some $u_a \in X$.

Letting $\delta \rightarrow 0$, by compactness we can assume that u_a converges to some $a \in X$. Thus we have that there exists $\mathbf{u}^1 \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ such that

$$u_\varepsilon^1 = \begin{cases} x_{[d]\setminus\{1\}} & \text{if } \varepsilon_1 = 0 \wedge \exists j \neq 1, \varepsilon_j = 0, \\ y_{[d]\setminus\{1\}} & \text{if } \varepsilon = [d] \setminus \{1\}, \\ x_{[d]} & \text{if } \varepsilon_1 = 1 \wedge \exists j \neq 1, \varepsilon_j = 0, \\ a & \text{if } \varepsilon = [d]. \end{cases}$$

This finishes the proof of Claim 2 in Case 2. \square

From \mathbf{u}^1 we aim to construct points $\mathbf{u}^2, \dots, \mathbf{u}^d$ in $\mathbf{Q}_{T_1, \dots, T_d}(X)$ where at each step they increasingly coincide with \mathbf{x} , but having their $[d] \setminus \{1\}$ coordinate equal to $y_{[d]\setminus\{1\}}$ and their $[d]$ coordinate equal to a . We suggest the reader to keep in mind that the $[d] \setminus \{1\}$ and $[d]$ coordinates are special.

Consider the face $(u_\varepsilon^1: \varepsilon_2 = 0)$. Since $\mathbf{x} \in \mathbf{Q}_{T_1, \dots, T_d}(X)$, duplicating the restriction of \mathbf{x} to the coordinates $[d] \setminus \{1, 2\}$, $[d] \setminus \{1\}$, $[d] \setminus \{2\}$ and $[d]$ we get the point $\mathbf{v}^2 \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ characterized by

$$v_\varepsilon^2 = \begin{cases} x_{[d] \setminus \{1, 2\}} & \text{if } \varepsilon_1 = 0 \wedge \varepsilon_2 = 0, \\ x_{[d] \setminus \{1\}} & \text{if } \varepsilon_1 = 0 \wedge \varepsilon_2 = 1, \\ x_{[d] \setminus \{2\}} & \text{if } \varepsilon_1 = 1 \wedge \varepsilon_2 = 0, \\ x_{[d]} & \text{if } \varepsilon_1 = 1 \wedge \varepsilon_2 = 1. \end{cases}$$

Hence, $(v_\varepsilon^2: \varepsilon \in \{0, 1\}^d, \varepsilon_2 = 1) = (u_\varepsilon^1: \varepsilon \in \{0, 1\}^d, \varepsilon_2 = 0)$. By Lemma 17, we can glue the 2-th lower face of \mathbf{v} with the 2-th upper face of \mathbf{u}^1 and obtain the point \mathbf{u}^2 that belongs to $\mathbf{Q}_{T_1, \dots, T_d}(X)$. The point \mathbf{u}^2 is characterized by

$$u_\varepsilon^2 = \begin{cases} x_{[d] \setminus \{1, 2\}} & \text{if } \varepsilon_1 = 0 \wedge \varepsilon_2 = 0, \\ x_{[d] \setminus \{1\}} & \text{if } \varepsilon_1 = 0 \wedge \varepsilon_2 = 1, \varepsilon \neq [d] \setminus \{1\}, \\ x_{[d] \setminus \{2\}} & \text{if } \varepsilon_1 = 1 \wedge \varepsilon_2 = 0, \\ x_{[d]} & \text{if } \varepsilon_1 = 1 \wedge \varepsilon_2 = 1, \\ y_{[d] \setminus \{1\}} & \varepsilon = [d] \setminus \{1\}, \\ a & \varepsilon = [d]. \end{cases}$$

The value of u_ε^2 for $\varepsilon \neq [d]$ and $\varepsilon \neq [d] \setminus \{1\}$ can only take the values $x_{[d] \setminus \{1, 2\}}$, $x_{[d] \setminus \{1\}}$, $x_{[d] \setminus \{2\}}$ or $x_{[d]}$. Furthermore, \mathbf{u}^2 coincides with \mathbf{x} in the coordinates $[d] \setminus \{1, 2\}$ and $[d] \setminus \{2\}$ while $u_{[d] \setminus \{1\}}^2 = y_{[d] \setminus \{1\}}$ and $u_{[d]}^2 = a$. Note that \mathbf{u}^2 does not depend on $\{3, \dots, d\}$, meaning that $u_\varepsilon^2 = x_{\varepsilon \cup \{3, \dots, d\}}$, for any ε , except for $\varepsilon = [d] \setminus \{1\}$ and $\varepsilon = [d]$.

Now assume that we have constructed $\mathbf{u}^k \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ for $2 \leq k < d$ such that \mathbf{u} coincides with \mathbf{x} in all coordinates of the form $[d] \setminus A$ where $A \subseteq \{1, \dots, k\}$, except for $A = \{1\}$ and $A = \emptyset$, where $u_{[d] \setminus \{1\}}^k = y_{[d] \setminus \{1\}}$ and $u_{[d]}^k = a$. Assume also that \mathbf{u}^k does not depend on $\{k+1, \dots, d\}$, meaning that for any ε (except for $\varepsilon = [d] \setminus \{1\}$ and $\varepsilon = [d]$)

$$(5) \quad u_\varepsilon^k = x_{\varepsilon \cup \{k+1, \dots, d\}}.$$

Consider the $(k+1)$ -th lower face of \mathbf{u}^k (i.e., $(u_\varepsilon^k: \varepsilon \in \{0, 1\}^d, \varepsilon_{k+1} = 0)$). Since $\mathbf{x} \in \mathbf{Q}_{T_1, \dots, T_d}(X)$, duplicating the restriction of \mathbf{x} to the coordinates $[d] \setminus A$, $A \subseteq \{1, \dots, k+1\}$ we get the point \mathbf{v}^{k+1} that belongs to $\mathbf{Q}_{T_1, \dots, T_d}(X)$. This point is characterized by

$$v_\varepsilon^{k+1} = x_{\varepsilon \cup \{k+2, \dots, d\}} \text{ for every } \varepsilon \in \{0, 1\}^d.$$

Note that if $\varepsilon_{k+1} = 1$ (or equivalently $k+1 \in \varepsilon$) then,

$$(6) \quad v_\varepsilon^{k+1} = x_{\varepsilon \cup \{k+2, \dots, d\}} = x_{(\varepsilon \setminus \{k+1\}) \cup \{k+1, k+2, \dots, d\}} = u_{\varepsilon \setminus \{k+1\}}^k,$$

where in the last equality we used (5) with $\varepsilon \setminus \{k+1\}$, which is different from $[d] \setminus \{1\}$ and $[d]$.

From (6) we deduce that $(v_\varepsilon^{k+1}: \varepsilon \in \{0, 1\}^d, \varepsilon_{k+1} = 1) = (u_\varepsilon^k: \varepsilon \in \{0, 1\}^d, \varepsilon_{k+1} = 0)$. By Lemma 12, we can glue the $(k+1)$ -th lower face of \mathbf{v}^{k+1} with the $(k+1)$ -th upper face of \mathbf{u}^k and obtain the point \mathbf{u}^{k+1} that belongs to $\mathbf{Q}_{T_1, \dots, T_d}(X)$. The point \mathbf{u}^{k+1} is characterized by

$$u_\varepsilon^{k+1} = \begin{cases} v_\varepsilon^{k+1} & \text{if } \varepsilon_{k+1} = 0, \\ u_\varepsilon^k & \text{if } \varepsilon_{k+1} = 1. \end{cases}$$

We check the point \mathbf{u}^{k+1} satisfies the conditions needed to continue the process, assuming obviously that $k+1 < d$. Let $A \subseteq \{1, \dots, k+1\}$ with $A \neq \{1\}$ and $A \neq \emptyset$. If $k+1 \notin A$ then $k+1 \in [d] \setminus A$ and $u_{[d] \setminus A}^{k+1} = u_{[d] \setminus A}^k = x_{([d] \setminus A) \cup \{k+1, \dots, d\}} = x_{[d] \setminus A}$. On the other hand, if $k+1 \in A$ then $k+1 \notin [d] \setminus A$ and $u_{[d] \setminus A}^{k+1} = v_{[d] \setminus A}^{k+1} = x_{([d] \setminus A) \cup \{k+2, \dots, d\}} = x_{[d] \setminus A}$. We deduce that \mathbf{u} coincides with \mathbf{x} in all coordinates $[d] \setminus A$, with $A \neq \{1\}$ and $A \neq \emptyset$. It is also immediate to check that $u_{[d] \setminus \{1\}}^{k+1} = u_{[d] \setminus \{1\}}^k = y_{[d] \setminus \{1\}}$ and $u_{[d]}^{k+1} = u_{[d]}^k = a$. We now check that \mathbf{u}^{k+1} does not depend on the coordinates $\{k+2, \dots, d\}$. For $\varepsilon \neq [d] \setminus \{1\}$ and $\varepsilon \neq [d]$ we have

$$u_\varepsilon^{k+1} = \begin{cases} v_\varepsilon^{k+1} = x_{\varepsilon \cup \{k+2, \dots, d\}} & \text{if } \varepsilon_{k+1} = 0, \\ u_\varepsilon^k = x_{\varepsilon \cup \{k+1, \dots, d\}} = x_{\varepsilon \cup \{k+2, \dots, d\}} & \text{if } \varepsilon_{k+1} = 1, \end{cases}$$

and we conclude that \mathbf{u}^{k+1} does not depend on the coordinates $\{k+2, \dots, d\}$.

We proceed with the process until we construct $\mathbf{u}^d \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. This point coincides with \mathbf{x} in all coordinates $[d] \setminus A$ for $A \subseteq \{1, \dots, d\} = [d]$ except for $A = \{1\}$ and $A = \emptyset$, where $u_{[d] \setminus \{1\}}^d = y_{[d] \setminus \{1\}}$ and $u_{[d]}^d = a$. We therefore conclude that \mathbf{u}^d coincides with \mathbf{x} everywhere but in the coordinates $[d] \setminus \{1\}$

and $[d]$. In particular $\mathbf{u}_{\vec{0}} = x_0$. Using that x_0 is a continuity point we get that $(u_\varepsilon : \varepsilon \neq \vec{0})$ belongs to $\mathbf{K}_{T_1, \dots, T_d}^{x_0}$, completing the proof of Claim 1. \square

To finish the proof of Theorem 30, use Claim 1 for $0 \leq k < d$ to construct the points $\mathbf{x}^1, \dots, \mathbf{x}^d \in \mathbf{K}_{T_1, \dots, T_d}^{x_0}$. As mentioned before, the point \mathbf{x}^d coincides everywhere with \mathbf{y} , except for the $[d]$ coordinate. The unique closing parallelepiped property implies that $\mathbf{y} = \mathbf{x}^d \in \mathbf{K}_{T_1, \dots, T_d}^{x_0}$. \square




FIGURE 8. Illustration of the proof of Claim 1 in Theorem 30 for the case $d = 3$. We change the $[d] \setminus \{1\} = \{2, 3\}$ coordinate of \mathbf{x} by the corresponding one of \mathbf{y} .

7. PROOF OF THEOREM 1

We have now all the ingredients to prove the structure Theorem 1. We restate it here for the reader's convenience. Interestingly, after the work of previous sections, all the \mathbb{Z}^d -systems that appear in the theorem can be described explicitly from the directional cube structures of (X, T_1, \dots, T_d) .

Theorem. *Let (X, T_1, \dots, T_d) be a minimal distal \mathbb{Z}^d -system. The following statements are equivalent:*

- (1) *(X, T_1, \dots, T_d) has the unique closing parallelepiped property, i.e., if $\mathbf{x}, \mathbf{y} \in \mathbf{Q}_{T_1, \dots, T_d}(X)$ have $2^d - 1$ coordinates in common, then $\mathbf{x} = \mathbf{y}$.*

(2) $\mathcal{R}_{T_1, \dots, T_d}(X) = \Delta_X$.

(3) The structure of (X, T_1, \dots, T_d) can be described as follows: (i) it is a factor of a minimal distal \mathbb{Z}^d -system (Y, T_1, \dots, T_d) which is a joining of \mathbb{Z}^d -systems $(Y_1, T_1, \dots, T_d), \dots, (Y_d, T_1, \dots, T_d)$, where for each $i \in \{1, \dots, d\}$ the action of T_i on Y_i is the identity; (ii) for each $i, j \in \{1, \dots, d\}$, $i < j$, there exists a \mathbb{Z}^d -system $(Y_{i,j}, T_1, \dots, T_d)$ which is a common factor of (Y_i, T_1, \dots, T_d) and (Y_j, T_1, \dots, T_d) and where T_i and T_j act as the identity; and (iii) Y is jointly relatively independent with respect to the systems $((Y_{i,j}, T_1, \dots, T_d) : i, j \in \{1, \dots, d\}, i < j)$.

More precisely, $Y = \mathbf{K}_{T_1, \dots, T_d}^{x_0}$ and $Y_j = \mathbf{K}_{T_1, \dots, T_{j-1}, T_{j+1}, \dots, T_d}^{x_0}$ with x_0 being a continuity point.

Relative independence of Y is with respect to their maximal $Z_0^{j_1} \wedge Z_0^{j_2}$ -factors, for all $j_1, j_2 \in [d]$ with $j_1 < j_2$.

Proof of Theorem 1. (1) \implies (2). This follows from Proposition 20.

(2) \implies (1). Suppose that (X, T_1, \dots, T_d) does not verify the unique closing parallelepiped property, then there exist $x, y \in X$ with $x \neq y$ and $\mathbf{a}_* \in X_*^{[d]}$ such that $(x, \mathbf{a}_*), (y, \mathbf{a}_*) \in \mathbf{Q}_{T_1, \dots, T_d}(X)$. By Proposition 21, we have that $(x, y) \in \mathcal{R}_{T_1, \dots, T_d}(X)$. Then, $x = y$, which is a contradiction.

(1) \implies (3). This is a consequence of Theorem 30.

(3) \implies (1). We show that the system (Y, T_1, \dots, T_d) given by (3) verifies the unique closing parallelepiped property.

Let $\mathbf{y} \in \mathbf{Q}_{T_1, \dots, T_j}(Y)$. By definition, for every $j \in [d]$ we have that $(y_{[d]}, y_{[d] \setminus \{j\}}) \in \mathbf{Q}_{T_j}(Y)$. But, T_j acts as the identity on the j -th coordinate of points in Y , then $((y_{[d]})_j, (y_{[d] \setminus \{j\}})_j) \in \mathbf{Q}_{T_j}(Y_j) = \Delta_{Y_j}$. This implies that, $(y_{[d]})_j = (y_{[d] \setminus \{j\}})_j$. Hence, $y_{[d]}$ can be determined from previous coordinates of \mathbf{y} , which proves the unique closing parallelepiped property for (Y, T_1, \dots, T_d) .

Since (1) is equivalent with (2) we have that $\mathcal{R}_{T_1, \dots, T_d}(Y) = \Delta_Y$. By Theorem 23, $\mathcal{R}_{T_1, \dots, T_d}(X) = \Delta_X$. This proves the unique closing parallelepiped property for (X, T_1, \dots, T_d) . \square

The following two corollaries can be deduced from Theorem 1 and Theorem 23. The first one was implicitly proved inside the last proof.

Corollary 31. *Let $\pi: Y \rightarrow X$ be a factor map between minimal distal \mathbb{Z}^d -systems (Y, T_1, \dots, T_d) and (X, T_1, \dots, T_d) . If (Y, T_1, \dots, T_d) has the unique closing parallelepiped property then (X, T_1, \dots, T_d) has it too.*

Corollary 32. *Let (X, T_1, \dots, T_d) be a minimal distal \mathbb{Z}^d -system. Then, $(X/\mathcal{R}_{T_1, \dots, T_d}(X), T_1, \dots, T_d)$ has the unique closing parallelepiped property. Moreover, this system is the maximal factor with this property, i.e., any other factor of (X, T_1, \dots, T_d) with the unique closing parallelepiped property factorizes through it.*

Proof. Observe that if (Z, T_1, \dots, T_d) is a factor of (X, T_1, \dots, T_d) with the unique closing parallelepiped property, then by Theorem 1 $\mathcal{R}_{T_1, \dots, T_d}(Z) = \Delta_Z$. Now, by Theorem 23, $\pi \times \pi(\mathcal{R}_{T_1, \dots, T_d}(X)) = \mathcal{R}_{T_1, \dots, T_d}(Z) = \Delta_Z$. That is, there exists a factor map from $(X/\mathcal{R}_{T_1, \dots, T_d}(X), T_1, \dots, T_d)$ to (Z, T_1, \dots, T_d) . It remains to prove that $\mathcal{R}_{T_1, \dots, T_d}(X/\mathcal{R}_{T_1, \dots, T_d}(X)) = \Delta_{X/\mathcal{R}_{T_1, \dots, T_d}(X)}$ (i.e., the quotient system has the unique closing parallelepiped property). Let $\pi: X \rightarrow X/\mathcal{R}_{T_1, \dots, T_d}(X)$ be the quotient map and take $(y_1, y_2) \in \mathcal{R}_{T_1, \dots, T_d}(X/\mathcal{R}_{T_1, \dots, T_d}(X))$. By Theorem 23, there exists $(x_1, x_2) \in \mathcal{R}_{T_1, \dots, T_d}(X)$ with $\pi(x_1) = y_1$ and $\pi(x_2) = y_2$. But $y_1 = \pi(x_1) = \pi(x_2) = y_2$, so $\mathcal{R}_{T_1, \dots, T_d}(X/\mathcal{R}_{T_1, \dots, T_d}(X))$ coincides with the diagonal relation of $X/\mathcal{R}_{T_1, \dots, T_d}(X)$, which proves the corollary. \square

8. RECURRENCE IN MINIMAL DISTAL \mathbb{Z}^d -SYSTEMS WITH THE UNIQUE CLOSING PARALLELEPIPED PROPERTY

As an application of previous work, in this section we study sets of return times for minimal distal \mathbb{Z}^d -systems with the unique closing parallelepiped property. In particular, we get a characterization of minimal distal systems with this property using return time ideas.

Definition 33. Let (X, T_1, \dots, T_d) be a \mathbb{Z}^d -system. Let $x \in X$ and U be an open neighborhood of x . The set of return times of x to U is defined as

$$N_{T_1, \dots, T_d}(x, U) = \{(n_1, \dots, n_d) \in \mathbb{Z}^d : T_1^{n_1} \dots T_d^{n_d} x \in U\}.$$

A subset B of \mathbb{Z}^d is a set of return times for a \mathbb{Z}^d -system if there exists a \mathbb{Z}^d -system (X, T_1, \dots, T_d) , $x \in X$ and an open neighborhood U of x such that $N_{T_1, \dots, T_d}(x, U) \subseteq B$.

We are able to characterize sets of return times for minimal distal \mathbb{Z}^d -systems via the unique closing parallelepiped property. For this we consider the following definition.

Definition 34. Let $d \geq 2$ be an integer and $B_1, \dots, B_d \subseteq \mathbb{Z}^{d-1}$. We define the d -joining of B_1, \dots, B_d as the set

$$B = \{(n_1, \dots, n_d) \in \mathbb{Z}^d : \forall i \in [d], (n_1, \dots, n_{i-1}, n_{i+1}, \dots, n_d) \in B_i\} \subseteq \mathbb{Z}^d.$$

We remark that the 2-joining of $B_1, B_2 \subseteq \mathbb{Z}$ is the Cartesian product $B_1 \times B_2$. For $d \geq 3$ the set B might be empty for general sets. For instance, let $B_1 = \{(n_1, n_2) \in \mathbb{Z}^2 : n_1 - n_2 \text{ is even}\}$, $B_2 = \{(n_1, n_2) \in \mathbb{Z}^2 : n_1 - n_2 \text{ is odd}\}$, $B_3 = B_1$. Then, $(n_1, n_2, n_3) \in B$ if and only if $n_2 - n_3$ is even, $n_1 - n_3$ is odd and $n_1 - n_2$ is even, which leads to a contradiction. However, if the building sets B_1, \dots, B_d are return times, then the set B is always non-empty (it contains the vector of 0's). With the help of this notion of joining of sets we can obtain a nice relation with the unique closing parallelepiped property in the minimal distal case.

Theorem 35. Let $d \geq 2$ be an integer. A subset $B \subseteq \mathbb{Z}^d$ contains a set of return times for a minimal distal \mathbb{Z}^d -system with the unique closing parallelepiped property if and only if B contains a d -joining of sets that are return times of minimal distal \mathbb{Z}^{d-1} -systems.

Proof. Let B be a subset of \mathbb{Z}^d that contains a set of return times of a minimal distal \mathbb{Z}^d -system (X, T_1, \dots, T_d) with the unique closing parallelepiped property. Let $x \in X$ and U an open neighborhood of x such that $N_{T_1, \dots, T_d}(x, U) = \{(n_1, \dots, n_d) \in \mathbb{Z}^d : T_1^{n_1} \dots T_d^{n_d} x \in U\} \subseteq B$. We want to show that B contains a d -joining of sets that are return times of minimal distal \mathbb{Z}^{d-1} -systems.

Let (Y, T_1, \dots, T_d) be an extension of (X, T_1, \dots, T_d) as in Theorem 1. Recall that (Y, T_1, \dots, T_d) is a joining of systems (Y_j, T_1, \dots, T_d) , where for each $j \in [d]$ the action of T_j on Y_j is the identity.

Let $(y_1, \dots, y_d) \in Y$ be such that $\pi(y_1, \dots, y_d) = x$ and let \tilde{U} be a neighborhood of (y_1, \dots, y_d) in Y such that $\pi(\tilde{U}) \subseteq U$. We may assume that $\tilde{U} = (\tilde{U}_1 \times \dots \times \tilde{U}_d) \cap Y$, where for each $j \in [d]$ the set \tilde{U}_j is an open neighborhood of y_j . We have,

$$N_{T_1, \dots, T_d}((y_1, \dots, y_d), \tilde{U}) = N_{T_1, \dots, T_d}((y_1, \dots, y_d), (\tilde{U}_1 \times \dots \times \tilde{U}_d) \cap Y) = \bigcap_{j \in [d]} N_{T_1, \dots, T_d}(y_j, \tilde{U}_j).$$

Recall that on each Y_j the action of T_j is the identity, so the action of T_1, \dots, T_d on Y_j can be seen as a \mathbb{Z}^{d-1} action and

$$\begin{aligned} (n_1, \dots, n_d) \in N_{T_1, \dots, T_d}((y_1, \dots, y_d), \tilde{U}) &\iff \\ (n_1, \dots, n_{j-1}, n_{j+1}, \dots, n_d) \in N_{T_1, \dots, T_{j-1}, T_{j+1}, \dots, T_d}(y_j, \tilde{U}_j) &\text{ for every } j \in [d]. \end{aligned}$$

Thus the sets $B_j = \{(n_1, \dots, n_{j-1}, n_{j+1}, \dots, n_d) \in N_{T_1, \dots, T_{j-1}, T_{j+1}, \dots, T_d}(y_j, \tilde{U}_j)\}$ are return times of minimal \mathbb{Z}^{d-1} -systems whose d -joining coincides with $N_{T_1, \dots, T_d}((y_1, \dots, y_d), \tilde{U})$. Since $\pi(\tilde{U}) \subseteq U$, we have that $N_{T_1, \dots, T_d}((y_1, \dots, y_d), \tilde{U}) \subseteq N_{T_1, \dots, T_d}(x, U) \subseteq B$, which serves to conclude.

Conversely, assume that a set $B \subseteq \mathbb{Z}^d$ contains a d -joining of sets B_j , $j \in [d]$, where each B_j is a set of return times of a minimal distal \mathbb{Z}^{d-1} -system $(Y_j, S_{j,1}, \dots, S_{j,d-1})$. We want to show that B contains the set of return times of a minimal distal \mathbb{Z}^d -system with the unique closing parallelepiped property.

For convenience, for each $j \in [d]$ we write $(Y_j, T_1, \dots, T_{j-1}, T_{j+1}, T_d)$ instead of $(Y_j, S_{j,1}, \dots, S_{j,d-1})$. By doing so, we stress the fact that we are viewing the \mathbb{Z}^{d-1} action as a \mathbb{Z}^d action where one of the transformations is the identity. We will see below that using the same set of transformations T_1, \dots, T_d for all systems will not be a notational problem since we will consider a product system.

Let $y_j \in Y_j$ and let U_j be an open neighborhood of y_j such that

$$(7) \quad N_{T_1, \dots, T_{j-1}, T_{j+1}, \dots, T_d}(y_j, U_j) \subseteq B_j.$$

We now construct a minimal distal \mathbb{Z}^d -system with the unique closing parallelepiped property and a set of return times for this system that is contained in B . Consider the product system $\prod_{j=1}^d Y_j$ and the

diagonal action of T_1, \dots, T_d on it. For the point $\mathbf{y} = (y_1, \dots, y_d) \in \prod_{j=1}^d Y_j$ let $Y = \overline{\mathcal{O}(\mathbf{y}, T_1, \dots, T_d)}$. The

\mathbb{Z}^d -system (Y, T_1, \dots, T_d) is minimal and distal (see Theorem 3), and contains the point $\mathbf{y} = (y_1, \dots, y_d)$. For any $j \in [d]$, the transformation T_j acts trivially in the j -th coordinate of Y and thus by Theorem 1 (Y, T_1, \dots, T_d) has the unique closing parallelepiped property. Now, consider the open neighborhood of \mathbf{y}

given by $U = \left(\prod_{j=1}^d U_j \right) \cap Y$ and note that $(n_1, \dots, n_d) \in N_{T_1, \dots, T_d}(\mathbf{y}, U)$ if and only if for every $j \in [d]$, $(n_1, \dots, n_{j-1}, n_{j+1}, \dots, n_d) \in N_{T_1, \dots, T_{j-1}, T_{j+1}, \dots, T_d}(y_j, U_j)$. That is, $N_{T_1, \dots, T_d}(\mathbf{y}, U)$ is the d -joining of the sets $N_{T_1, \dots, T_{j-1}, T_{j+1}, \dots, T_d}(y_j, U_j)$. Using (7) we conclude that $N_{T_1, \dots, T_d}(\mathbf{y}, U)$ is contained in B . \square

We denote by \mathcal{B}_d the family generated by sets of return times arising from minimal distal \mathbb{Z}^d -systems with the unique closing parallelepiped property and by \mathcal{B}_d^* the (dual) family of subsets of \mathbb{Z}^d which have nonempty intersection with every set in \mathcal{B}_d .

Lemma 36. *Let (X, T_1, \dots, T_d) be a minimal distal \mathbb{Z}^d -system. Suppose that $(x, y) \in \mathcal{R}_{T_1, \dots, T_d}(X)$. Let (Z, T_1, \dots, T_d) be a minimal distal \mathbb{Z}^d -system with the unique closing parallelepiped property and let (J, T_1, \dots, T_d) be a joining between (X, T_1, \dots, T_d) and (Z, T_1, \dots, T_d) . Then, for $z_0 \in Z$ we have that $(x, z_0) \in J$ if and only if $(y, z_0) \in J$.*

Proof. The proof is similar to the proof of Lemma 6.19 in [7], which is an adaptation of the corresponding statement in [20]. We provide it for completeness. Let $W = Z^{\mathbb{Z}}$ and $T_1^{\mathbb{Z}}, \dots, T_d^{\mathbb{Z}}: W \rightarrow W$ be the corresponding commuting transformations. Let $\omega^* \in W$ be the point satisfying $\omega^*(z) = z$ for all $z \in Z$ and $Z_\infty = \mathcal{O}(\omega^*, G^{\mathbb{Z}})$, where $G^{\mathbb{Z}}$ is the group generated by $T_1^{\mathbb{Z}}, \dots, T_d^{\mathbb{Z}}$. Then, Z_∞ is minimal and distal. So for any $\omega \in Z_\infty$ there exists $p \in E(Z, G)$ such that $\omega(z) = p\omega^*(z) = p(z)$ for any $z \in Z$. Since (Z, T_1, \dots, T_d) is minimal and distal, $E(Z, G)$ is a group and thus $p: Z \rightarrow Z$ is surjective. This implies that there exists $z_\omega \in Z$ such that $\omega(z_\omega) = z_0$.

Take a minimal subsystem $(A, T_1 \times T_1^{\mathbb{Z}}, \dots, T_d \times T_d^{\mathbb{Z}})$ of the product system $(X \times Z_\infty, T_1 \times T_1^{\mathbb{Z}}, \dots, T_d \times T_d^{\mathbb{Z}})$. Let $\pi_X: A \rightarrow X$ be the natural coordinate projection. Then, π_X is a factor map between two distal minimal systems. By Theorem 23, there exist $\omega^1, \omega^2 \in W$ such that $((x, \omega^1), (y, \omega^2)) \in \mathcal{R}_{\hat{T}_1, \dots, \hat{T}_d}(A)$, where $\hat{T}_j = T_j \times T_j^{\mathbb{Z}}$, for $j \in [d]$.

Let $z_1 \in Z$ be such that $\omega^1(z_1) = z_0$. Denote by $\pi: A \rightarrow X \times Z$, $\pi(u, \omega) = (u, \omega(z_1))$ for $(u, \omega) \in A$, $u \in X$ and $\omega \in W$. Consider the projection $B = \pi(A)$. Then, $(B, T_1 \times T_1, \dots, T_d \times T_d)$ is a minimal distal subsystem of $(X \times Z, T_1 \times T_1, \dots, T_d \times T_d)$ and since $\pi(x_0, \omega^1) = (x, z_0) \in B$ we have that J contains B . Suppose that $\pi(x, \omega^2) = (x, z_2)$. Then, $((x, z_0), (y, z_2)) \in \mathcal{R}_{T_1 \times T_1, \dots, T_d \times T_d}(B)$ and we conclude that $(z_0, z_2) \in \mathcal{R}_{T_1, \dots, T_d}(Z)$. Since $\mathcal{R}_{T_1, \dots, T_d}(Z) = \Delta_Z$ we have that $z_0 = z_2$ and thus $(y, z_0) \in B \subseteq J$. \square

Lemma 37. *Let (X, T_1, \dots, T_d) be a minimal distal \mathbb{Z}^d -system. Then, for $x, y \in X$, $(x, y) \in \mathcal{R}_{T_1, \dots, T_d}(X)$ if and only if $N_{T_1, \dots, T_d}(x, U) \in \mathcal{B}_d^*$ for any open neighborhood U of y .*

Proof. The proof is similar to the one of Theorem 6.20 in [7]. Suppose $N_{T_1, \dots, T_d}(x, U) \in \mathcal{B}_{T_1, \dots, T_d}^*$ for any open neighborhood U of y . Since (X, T_1, \dots, T_d) is distal, $\mathcal{R}_{T_1, \dots, T_d}(X)$ is an equivalence relation. Let π be the projection map $\pi: X \rightarrow Y = X/\mathcal{R}_{T_1, \dots, T_d}(X)$. By Corollary 32 we have that $\mathcal{R}_{T_1, \dots, T_d}(Y) = \Delta_Y$. We also have that the factor map π is open and $\pi(U)$ is an open neighborhood of $\pi(y)$. In particular, $N_{T_1, \dots, T_d}(x, U) \subseteq N_{T_1, \dots, T_d}(\pi(x), \pi(U))$. Let V be an open neighborhood of $\pi(x)$. By hypothesis, we have that $N_{T_1, \dots, T_d}(x, U) \cap N_{T_1, \dots, T_d}(\pi(x), \pi(U)) \neq \emptyset$, which implies that $N_{T_1, \dots, T_d}(\pi(x), \pi(U)) \cap N_{T_1, \dots, T_d}(\pi(x), V) \neq \emptyset$. This implies that $\pi(U) \cap V \neq \emptyset$. But this holds for every V , so we have that $\pi(x) \in \pi(U) = \pi(\bar{U})$. Finally, since this fact holds for every U we conclude that $\pi(x) = \pi(y)$. This shows that $(x, y) \in \mathcal{R}_{T_1, \dots, T_d}(X)$ as desired.

Conversely, suppose that $(x, y) \in \mathcal{R}_{T_1, \dots, T_d}(X)$. Let U be an open neighborhood of y and A be a \mathcal{B}_d^* set. Then, there exists a minimal distal system (Z, T_1, \dots, T_d) with $\mathcal{R}_{T_1, \dots, T_d}(Z) = \Delta_Z$, an open set $V \subseteq Z$ and $z_0 \in V$ such that $N_{T_1, \dots, T_d}(z_0, V) \subseteq A$. Let J be the orbit closure of (x, z_0) under $T_j \times T_j$ for $j \in [d]$. By distality we have that $(J, T_1 \times T_1, \dots, T_d \times T_d)$ is a minimal system and $(x, z_0) \in J$. By Theorem 23 we have that $(y, z_0) \in J$ and particularly there exists a sequence $(\mathbf{n}^i)_{i \in \mathbb{N}} \subseteq \mathbb{Z}^d$ such that $(T_1^{\mathbf{n}_1^i} \dots T_d^{\mathbf{n}_d^i} x, T_1^{\mathbf{n}_1^i} \dots T_d^{\mathbf{n}_d^i} z_0) \rightarrow (y, z_0)$. This implies that $N_{T_1, \dots, T_d}(x, U) \cap N_{T_1, \dots, T_d}(z_0, V) \neq \emptyset$ and the proof is achieved. \square

We get the following characterization of the unique closing parallelepiped property for minimal distal \mathbb{Z}^d -systems.

Corollary 38. *Let (X, T_1, \dots, T_d) be a minimal distal \mathbb{Z}^d -system. Then, (X, T_1, \dots, T_d) has the unique closing parallelepiped property if and only if for every $x \in X$ and every open neighborhood U of x , $N_{T_1, \dots, T_d}(x, U)$ contains a d -joining built from d sets of return times of minimal distal \mathbb{Z}^{d-1} -systems.*

Proof. The implication that $N_{T_1, \dots, T_d}(x, U)$ contains a d -joining of d sets of return times for minimal distal \mathbb{Z}^{d-1} -systems whenever (X, T_1, \dots, T_d) has the unique closing parallelepiped property follows immediately from Theorem 35. Then we only need to prove the other implication. Let us suppose that there exists

$(x, y) \in \mathcal{R}_{T_1, \dots, T_d}(X) \setminus \Delta_X$ and let U, V be open neighborhoods of x and y respectively such that $U \cap V = \emptyset$. By assumption $N_{T_1, \dots, T_d}(x, U)$ is a \mathcal{B}_d set and by Lemma 37 $N_{T_1, \dots, T_d}(x, V)$ has nonempty intersection with $N_{T_1, \dots, T_d}(x, U)$. This implies that $U \cap V \neq \emptyset$, a contradiction. We conclude that $\mathcal{R}_{T_1, \dots, T_d}(X) = \Delta_X$ and therefore (X, T_1, \dots, T_d) has the unique closing parallelepiped property. \square

As an application we get the following criterion for a minimal distal system (X, T) (so defined by a single transformation) for being topologically conjugate to an inverse limit of d -step nilsystems, *i.e.*, to an inverse limit of systems that can be written as $X = G/\Gamma$, where G is a d -step nilpotent Lie group and Γ is a cocompact subgroup of G , and T is a left translation by a fixed element in G . For an integer $d \geq 1$ let $\phi_d: \mathbb{Z}^d \rightarrow \mathbb{Z}$ be the morphism $(n_1, \dots, n_d) \mapsto n_1 + \dots + n_d$.

Theorem 39. *Let (X, T) be a minimal distal system. Then, (X, T) is topologically conjugate to an inverse limit of d -step nilsystems if and only if for all $x \in X$ and U an open set containing x the set of return times $N_T(x, U) = \{n \in \mathbb{Z} : T^n x \in U\}$ contains the image under ϕ_{d+1} of a $(d+1)$ -joining of $(d+1)$ sets of return times for distal \mathbb{Z}^d -systems.*

Proof. We regard (X, T) as the \mathbb{Z}^{d+1} -system (X, T, \dots, T) . Then, we have that (X, T, \dots, T) has the unique closing parallelepiped property if and only if for all $x \in X$ and all neighborhood U of x the set $\{(n_1, \dots, n_{d+1}) \in \mathbb{Z}^{d+1} : T^{n_1} T^{n_2} \dots T^{n_{d+1}} x \in U\} \subseteq \mathbb{Z}^{d+1}$ contains a $(d+1)$ -joining of $(d+1)$ sets of return times for \mathbb{Z}^d -distal systems. But, since all transformations are the same, $\{n \in \mathbb{Z} : T^n x \in U\}$ contains the image under ϕ_{d+1} of $\{(n_1, \dots, n_{d+1}) \in \mathbb{Z}^{d+1} : T^{n_1} T^{n_2} \dots T^{n_{d+1}} x \in U\}$. On the other hand, (X, T, \dots, T) has the unique closing parallelepiped property if and only if it is isomorphic to an inverse limit of d -step nilsystems [19, Theorem 1.2]. This finishes the proof. \square

9. EXAMPLES OF SYSTEMS WITH THE UNIQUE CLOSING PARALLELEPIPED PROPERTY

In this section we provide a family of examples of systems with the unique closing parallelepiped property.

9.1. Affine transformations in the torus. Let $r \geq 1$ be an integer. Consider different affine transformations $T_i: \mathbb{T}^r \rightarrow \mathbb{T}^r$, $x \mapsto A_i x + \alpha_i$, where A_i is an unipotent integer matrix (*i.e.*, $(A_i - I)^p = 0$ for some $p \in \mathbb{N}$) and $\alpha_i \in \mathbb{T}^r$ for every $i \in \{1, \dots, d\}$. The \mathbb{Z}^d -system $(\mathbb{T}^r, T_1, \dots, T_d)$ can be seen as a nilsystem as long as the matrices commute (we will not give all details on this fact but ideas can be found in [22]).

Let G be the group of transformations of \mathbb{T}^r generated by the matrices A_1, \dots, A_d and the translations of \mathbb{T}^r . Then, every element $g \in G$ is a map $x \mapsto A(g)x + \beta(g)$, where $A(g) = A_1^{m_1} \dots A_d^{m_d}$ with $m_1, \dots, m_d \in \mathbb{Z}$ and $\beta(g) \in \mathbb{T}^r$. A simple computation shows that if $g_1, g_2 \in G$ then the commutator $[g_1, g_2]$ is the map $x \mapsto x + (A(g_1) - I)\beta(g_2) - (A(g_2) - I)\beta(g_1)$ and thus it is a translation of \mathbb{T}^r . On the other hand, if $g \in G$ and $\beta \in \mathbb{T}^r$, then $[g, \beta]$ is the translation $x \mapsto x + (A(g) - I)\beta$. It follows that if $g_1, \dots, g_k \in G$ then the iterated commutator $[\dots [g_1, g_2], g_3] \dots g_k]$ belongs to \mathbb{T}^r and is contained in the image of $(A(g_3) - I) \dots (A(g_k) - I)$. If k is large enough, this product is trivial. So G is a nilpotent Lie group. The torus \mathbb{T}^r can be identified with G/Γ , where Γ is the stabilizer of 0, which is the group generated by the matrices A_1, \dots, A_d . We refer to $(\mathbb{T}^r, T_1, \dots, T_d)$ as an *affine nilsystem* with d transformations. It is worth noting that the transformations T_i and T_j commute if and only if $(A_i - I)\alpha_j = (A_j - I)\alpha_i$ in \mathbb{T}^r .

From results in [21] one can deduce that,

Proposition 40. *Let $(\mathbb{T}^r, T_1, \dots, T_d)$ be an affine nilsystem with d transformations. Then, the properties of transitivity, minimality, ergodicity and unique ergodicity under the action of $\langle T_1, \dots, T_d \rangle$ are equivalent.*

We consider some conditions on the commuting transformations T_1, \dots, T_d under which the system $(\mathbb{T}^r, T_1, \dots, T_d)$ has the unique closing parallelepiped property. We start by presenting the examples whose elementary proofs are deliberately omitted. Interested readers can find them in [4]. For the sake of clarity we first consider the case $d = 2$.

Lemma 41. *Let (\mathbb{T}^r, T_1, T_2) be an affine nilsystem with 2 commuting transformations, where $T_i x = A_i x + \alpha_i$ for $i \in \{1, 2\}$. Then, we have that for every $n, m \in \mathbb{Z}$,*

$$T_1^n T_2^m x = T_1^n x + T_2^m x - x$$

if and only if the following conditions hold

$$(8) \quad (A_1 - I)(A_2 - I) = 0,$$

$$(9) \quad (A_1 - I)\alpha_2 = (A_2 - I)\alpha_1 = 0.$$

In particular, if conditions (8) and (9) are satisfied we have that

$$\mathbf{Q}_{T_1, T_2}(X) = \overline{\{(x, T_1^n x, T_2^m x, T_1^n x + T_2^m x - x) : x \in \mathbb{T}^r, n, m \in \mathbb{Z}\}},$$

and thus (\mathbb{T}^r, T_1, T_2) has the unique closing parallelepiped property.

Example. Consider the following matrices,

$$A_1 = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 2 \\ 0 & 1 & 0 & 3 & 1 & 4 \\ 0 & 0 & 1 & 6 & 3 & 6 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad A_2 = \begin{pmatrix} 1 & 0 & 0 & 1 & 1 & 2 \\ 0 & 1 & 0 & 2 & 2 & 4 \\ 0 & 0 & 1 & 1 & 2 & 3 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

For the first matrix, the eigenspace (associated to the unique eigenvalue 1) is given by,

$$W_1(A_1) = \left\langle \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ -2 \\ 2 \\ 1 \end{pmatrix} \right\} \right\rangle.$$

For the second one, the eigenspace is given by,

$$W_1(A_2) = \left\langle \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ -1 \\ -1 \\ 1 \end{pmatrix} \right\} \right\rangle.$$

It is easy to see that $(A_1 - I)(A_2 - I) = 0$ and we can choose $\alpha_1 \in W_1(A_2)$ and $\alpha_2 \in W_1(A_1)$ such that (\mathbb{T}^6, T_1, T_2) has the unique closing parallelepiped property.

Conditions (8) and (9) can be generalized as follows. Again, the simple but tedious proof of this result can be found in [4].

Lemma 42. Let $(\mathbb{T}^r, T_1, \dots, T_d)$ be an affine nilsystem with d commuting transformations such that $T_i x = A_i x + \alpha_i$ for $i \in \{1, \dots, d\}$. Then, for every $(n_1, \dots, n_d) \in \mathbb{Z}^d$,

$$T_1^{n_1} \dots T_d^{n_d} x = (-1)^d \sum_{i=0}^{d-1} (-1)^{i+1} \sum_{\substack{I \subseteq [d] \\ |I|=i}} \bigcirc_{k \in I} T_k^{n_k} x$$

if and only if the following conditions hold

$$(10) \quad \prod_{i=1}^d (A_i - I) = 0,$$

$$(11) \quad \forall j \in \{1, \dots, d\}, \prod_{\substack{i=1 \\ i \neq j}}^d (A_i - I) \alpha_j = 0.$$

In particular, if conditions (10) and (11) hold the system $(\mathbb{T}^r, T_1, \dots, T_d)$ has the unique closing parallelepiped property.

REFERENCES

- [1] O. Antolin Camarena and B. Szegedy. Nilspaces, nilmanifolds and their morphisms. *ArXiv e-prints*, September 2010.
- [2] J. Auslander. *Minimal flows and their extensions*, volume 153. Elsevier, 1988.
- [3] T. Austin. Multiple recurrence and the structure of probability-preserving systems. page 101, 2010. Thesis (Ph.D.)–University of California, Los Angeles.
- [4] C. Cabezas. Cubos dinámicos direccionales para \mathbb{Z}^d -sistemas minimales. *Master Thesis, Universidad de Chile*, 2018.
- [5] P. Candela. Notes on nilspaces: algebraic aspects. *Discrete Anal.*, pages Paper No. 15, 59pp, 2017.
- [6] P. Candela. Notes on compact nilspaces. *Discrete Anal.*, pages Paper No. 16, 57pp, 2017.
- [7] S. Donoso and W. Sun. Dynamical cubes and a criteria for systems having product extensions. *J. Mod. Dyn.*, 9:365–405, 2015.
- [8] S. Donoso and W. Sun. A pointwise cubic average for two commuting transformations. *Israel J. Math.*, 216(2):657–678, 2016.

- [9] S. Donoso and W. Sun. Pointwise multiple averages for systems with two commuting transformations. *Ergodic Theory Dynam. Systems*, 38(6):2132–2157, 2018.
- [10] S. Donoso and W. Sun. Pointwise convergence of some multiple ergodic averages. *Adv. Math.*, 330:946 – 996, 2018.
- [11] E. Glasner. Topological ergodic decompositions and applications to products of powers of a minimal transformation. *J. Anal. Math.*, 64:241–262, 1994.
- [12] E. Glasner. The structure of tame minimal dynamical systems. *Ergodic Theory Dynam. Systems*, 27(6):1819–1837, 2007.
- [13] E. Glasner, Y. Gutman, and X. Ye. Higher order regionally proximal equivalence relations for general minimal group actions. *Adv. Math.*, 333:1004–1041, 2018.
- [14] Y. Gutman, F. Manners, and P. Varjú. The structure theory of nilspaces i. To appear in *J. Analyse Math.*, 2018.
- [15] Y. Gutman, F. Manners, and P. Varjú. The structure theory of nilspaces ii: Representation as nilmanifolds. To appear in *Trans. Amer. Math. Soc.*, 2018.
- [16] Y. Gutman, F. Manners, and P. Varjú. The structure theory of nilspaces iii: Inverse limit representations and topological dynamics. submitted, 2016.
- [17] B. Host. Ergodic seminorms for commuting transformations and applications. *Studia Math.*, 195(1):31–49, 2009.
- [18] B. Host and B. Kra. Nonconventional ergodic averages and nilmanifolds. *Ann. of Math. (2)*, 161(1):397–488, 2005.
- [19] B. Host, B. Kra, and A. Maass. Nilsequences and a structure theorem for topological dynamical systems. *Adv. Math.*, 224(1):103–129, 2010.
- [20] W. Huang, S. Shao, and X. Ye. Nil Bohr-sets and almost automorphy of higher order. *Mem. Amer. Math. Soc.*, 241(1143):v+83, 2016.
- [21] A. Leibman. Pointwise convergence of ergodic averages for polynomial sequences of translations on a nilmanifold. *Ergodic Theory Dynam. Systems*, 25(1):201–213, 2005.
- [22] W. Parry. Ergodic properties of affine transformations and flows on nilmanifolds. *Amer. J. Math.*, 91:757–771, 1969.
- [23] S. Shao and X. Ye. Regionally proximal relation of order d is an equivalence one for minimal systems and a combinatorial consequence. *Adv. Math.*, 231(3-4):1786–1817, 2012.
- [24] T. Tao. Norm convergence of multiple ergodic averages for commuting transformations. *Ergodic Theory Dynam. Systems*, 28(2):657–688, 2008.
- [25] S. Tu and X. Ye. Dynamical parallelepipeds in minimal systems. *J. Dynam. Differential Equations*, 25(3):765–776, 2013.

DEPARTAMENTO DE INGENIERÍA MATEMÁTICA, UNIVERSIDAD DE CHILE, AV. BEAUCHEF 851, SANTIAGO, CHILE
E-mail address: ccabez@dim.uchile.cl

INSTITUTO DE CIENCIAS DE LA INGENIERÍA, UNIVERSIDAD DE O’HIGGINGS, AV. LIB. BERNARDO O’HIGGINS 611, RANCAGUA, CHILE
E-mail address: sebastian.donos@uoh.cl

DEPARTAMENTO DE INGENIERÍA MATEMÁTICA AND CENTRO DE MODELAMIENTO MATEMÁTICO, UNIVERSIDAD DE CHILE & UMI-CNRS 2807, AV. BEAUCHEF 851, SANTIAGO, CHILE
E-mail address: amaass@dim.uchile.cl