Changes in Age and Geographic Distribution of the Risk of Chagas Disease in Chile from 1989 to 2017

Por: <u>Canals, M</u> (Canals, Mauricio)^[1]; <u>Canals, A</u> (Canals, Andrea)^[2]; <u>Ayala, S</u> (Ayala, Salvador)^[3]; <u>Valdebenito, J</u> (Valdebenito, Jorge)^[4]; <u>Alvarado, S</u> (Alvarado, Sergio)^[2,5]; <u>Caceres, D</u> (Caceres, Dante)^[1,5]

VECTOR-BORNE AND ZOONOTIC DISEASES

Volumen: 21 Número: 2 Páginas: 98-104

DOI: 10.1089/vbz.2020.2647
Fecha de publicación: FEB 1 2021
Acceso anticipado: NOV 2020
Tipo de documento:Article
Ver impacto de la revista

Abstract

The interruption of vector-borne transmission of Chagas disease was certified in Chile in 1999. Our goal was to determine the effects of the interruption of vector transmission on the age and spatial distributions of the risk of Chagas disease. We analyzed cases of Chagas disease by age and sex between 1989 and 2017, from notified disease reports of the Ministry of Health. Bayesian risk maps were constructed using the Besag-York-Mollie model. The reported cases of Chagas disease had a mean age of 45.9 +/- 17.6 years. Small changes in the age distribution were found among different periods (chi(2)(15) = 602.4, p < 0.001). These were explained mainly by numbers lower than those expected in age groups 0-39 years in the 2011-2017 period. Part of the observed reduction in the proportion of individuals in the lower strata could be explained by the aging of the Chilean population. An increase of reported cases was detected after the interruption of vector-borne transmission (F-1,F-327 = 4.24, p < 0.04), with regional differences (F-14,F-1308 = 4.35, p < 0.001). The regions of the north-central area that have the highest burden of Chagas tended to decrease the relative risk, while the regions of the south tended to increase and small risk areas appear in zones where there are no insect vectors. There is still no clear evidence of a reduction in the reported cases in Chile. This could be explained mainly by an improvement in the detection of cases, but it cannot be ruled out that vector transmission still exists. The changes in distribution suggest potential impact from human internal migration and blood transfusion. This study provides strong evidence supporting the idea that entomological surveillance and long-term follow-up of Chagas disease need to be maintained after certification of interruption in endemic countries.

Palabras clave

Palabras clave de autor: Chagas disease; Chile; distribution; relative risk

KeyWords Plus:TRYPANOSOMA-CRUZI

INFECTION; TRANSMISSION; VECTOR; INTERRUPTION; REDUVIDAE; PREVENTION; WIL

D

Información del autor

Dirección para petición de copias:

Universidad de Chile Univ Chile, Fac Med, Dept Med, Programa Salud Ambiental, Escuela Salud Publ, Independencia 939, Santiago 8380453, Chile.

Dirección correspondiente: Canals, M (autor correspondiente)

Univ Chile, Fac Med, Dept Med, Programa Salud Ambiental, Escuela Salud Publ, Independencia 939, Santiago 83804

Direcciones:

- [1] Univ Chile, Fac Med, Escuela Salud Publ, Programa Salud Ambiental, Santiago, Chile
- [2] Univ Chile, Fac Med, Escuela Salud Publ, Programa Bioestadist, Santiago, Chile
 - [3] Inst Salud Publ Chile, Dept Vigilancia Lab, Santiago, Chile
 - [4] Minist Salud, Dept Enfermedades Transmisibles, Santiago, Chile
- [5] Univ Tarapaca, Fac Ciencias Salud, Arica, Chile

Direcciones de correo electrónico: mcanals@uchile.cl

Financiación

Entidad financiadora	Número de concesión
Regular Fondecyt Project	1150514

Ver texto de financiación

Editorial

MARY ANN LIEBERT, INC, 140 HUGUENOT STREET, 3RD FL, NEW ROCHELLE, NY 10801 USA

Información de la revista

• Impact Factor: <u>Journal Citation Reports</u>

Categorías / Clasificación

Áreas de investigación: Public, Environmental & Occupational Health; Infectious Diseases Categorías de Web of Science: Public, Environmental & Occupational Health; Infectious Diseases

Información del documento

Idioma: English

Número de acceso: WOS:000592479600001

ID de PubMed: 33226892

ISSN: 1530-3667 eISSN: 1557-7759